Cu(0)/Selectfluor 体系催化的邻芳基磺酰亚胺的串联环化/芳构化反应： 一种简便合成 6H-菲啉的方法

郑立孟 施冬冬 鲍汉扬 刘运奎*

(浙江工业大学化学工程学院 绿色化学与技术国家重点实验室培育基地 杭州 310014)

摘要 发展了 Cu(0)/Selectfluor 体系催化的邻芳基磺酰亚胺的串联环化/芳构化反应, 于温和的反应条件下以中等到良好的产率简便、高效地构建了一系列 6H-菲啉类化合物。机理研究表明, 反应的关键步骤经历了由 Cu(0)/Selectfluor 体系现场原位产生 XCuOH (X=F, BF₄) 物种, 进而诱导对 C=N 键的羟铜化反应和分子内 C—H 键胺化反应, 从而合成了 6H-菲啉类化合物。

关键词 铜催化; Selectfluor; 6H-菲啉; 环化反应; C—H 键胺化

Copper(0)/Selectfluor System-Catalyzed Tandem Annulation/ Aromatization of *o*-Aryl Benzenesulfonylimides: A Facile Synthesis of 6H-Phenanthridines

Zheng, Limeng Shi, Dongdong Bao, Hanyang Liu, Yunkui*

(State Key Laboratory Breeding Base of Green Chemistry-Synthesis Technology, College of Chemical Engineering, Zhejiang University of Technology, Hangzhou 310014)

Abstract A facile and efficient method for the synthesis of 6H-phenanthridines has been successfully developed involving a copper(0)/Selectfluor system-catalyzed tandem annulation/aromatization of *o*-aryl benzenesulfonylimides. A variety of substituted 6H-phenanthridines were synthesized in moderate to good yields under mild reaction conditions. Mechanistic experiments revealed that the reaction might involve an oxycupration of C=N bond followed by an intramolecular C—H bond amination as the key steps triggered by an *in situ* generated copper species XCuOH (X=F or BF₄) from the Cu(0)/Selectfluor system.

Keywords copper catalysis; Selectfluor; 6H-phenanthridine; annulation reaction; C—H bond amination

1 Introduction

Phenanthridines and their derivatives are a class of important heterocyclic compounds that occur frequently in natural products as well as synthetic molecules, exhibiting a broad spectrum of biological activities, such as antimalarial, antitumor, cytotoxicity, antimycobacterial, *etc.*^[1,2] Besides, they have been found wide application in materials science due to their unique optoelectronic properties.^[3] Among the phenanthridine family, 6H-phenanthridines have received special attention due to their unique biological properties and pharmaceutical applications.^[2] Several representative and biologically active compounds contain-

ing 6H-phenanthridine scaffold are presented in Figure 1. To date, numerous methods have been established for the construction of 6-substituted phenanthridines^[4–8] involving the classic Pictet-Hubert condensation,^[5] metal-catalyzed cross-coupling/cyclization,^[6] radical-mediated cyclization of *ortho*-substituted biphenyls,^[4,7] and others.^[8] In contrast, efficient and convenient methods for the preparation of 6H-phenanthridines have been relatively underexplored.^[9] In addition, most of the existing methods for the synthesis of 6H-phenanthridines have one or more limitations with respect to harsh reaction conditions,^[9a–9e] expensive catalysts,^[9b–9d,9f–9h] multisteps,^[9i–9k] narrow scope of substrates,^[9i,9k–9n] and the starting materials being not

* Corresponding author. E-mail: ykuiliu@zjut.edu.cn

Received April 24, 2019; revised May 26, 2019; published online June 12, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 21772176, 21372201), the Opening Foundation of Zhejiang Key Course of Chemical Engineering and Technology, Zhejiang University of Technology.

国家自然科学基金(Nos. 21772176, 21372201)、浙江工业大学“省重中之重一级学科”开放基金资助项目。

Figure 1 Several biologically active compounds containing 6-*H*-phenanthridine scaffold

readily available.^[9a] Therefore, it is still highly desirable to develop novel and efficient methods for the construction of 6-*H*-phenanthridines from readily available starting materials by using inexpensive catalysts under mild conditions.

On the other hand, in the past decades, the nucleometallation-triggered difunctionalization of carbon-carbon multiple bonds has received great attention and became one of the most powerful tools for the construction of useful molecules including cyclic compounds.^[10] In this field, we independently disclosed that the combination of the copper powder and Selectfluor [1-(chloromethyl)-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate), F-TEDA-BF₄] in the presence of water enabled *in situ* generation of XCuOH species (X=F or BF₄) which is very active to undergo oxycupration and successive difunctionalization of carbon-carbon multiple bonds (Scheme 1a).^[11a–11c] On the basis of the above-mentioned findings, it was further envisioned that the *in situ* generated XCuOH might also readily undergo oxycupration towards C=N bond and thus induce successive difunctionalization of C=N bond.^[11f] To the best of our knowledge, the nucleometallation-triggered difunctionalization of C=N bond have never been reported. Herein the facile synthesis of 6-*H*-phenanthridines involving the copper(0)/Selectfluor system-catalyzed tandem annulation/aromatization of *o*-aryl benzenesulfonylimides is described by using the oxycupration of C=N bond as a key step (Scheme 1b).

2 Results and discussion

Initially, *o*-phenyl benzenesulfonamide **1a** as the model substrate for the optimization of reaction conditions was

selected (Table 1). When **1a** was allowed to react with Cu(0) powder (10 mol%) and Selectfluor (2.0 equiv.) in newly purchased acetonitrile at 50 °C for 24 h, phenanthridine **2a** was indeed formed in 63% yield (Entry 1, Table 1). When anhydrous MeCN was used, the reaction only gave 15% yield of **2a**, suggesting that the residual water in solvent may play some role in the reaction (Entry 2, Table 1).^[12] The yield of **2a** could be further improved to 84% by using MeCN/H₂O (*V*:*V*=500:1) solvent system (Entry 3, Table 1). It was found that either increasing or reducing the amount of water in the solvent system had no positive effect on the yield of **2a** (Entries 4, 5, Table 1). A series of other solvent systems including toluene, acetone, 1,4-dioxane, DMF, DMSO, 1,2-dichloroethane, and methanol were investigated for the reaction, and it was found that MeCN/H₂O (*V*:*V*=500:1) solvent system was the best choice (Entry 6 vs 3, Table 1). The catalytic activity of several copper salts (Cu(I) and Cu(II)) on the reaction was also investigated, and their catalytic performances were proven to be much inferior to that of Cu(0) powder (Entries 7~12 vs 3, Table 1). Control experiments indicated that both Cu(0) and Selectfluor were indispensable for the reaction (Entries 13, 14, Table 1). The screening experiments of oxidants showed that Selectfluor exhibited unique effectiveness on the formation of **2a** (Entries 15, 16 vs 3, Table 1). Increasing or lowering the reaction temperature only led to a reduced yield of **2a** (Entries 18, 19 vs 3, Table 1). Note that the use of 2 equiv. of Selectfluor is enough for getting satisfactory yield of **2a** while a deficient use of Selectfluor (1.0 equiv.) resulted in a lower yield of **2a** (Entry 20 vs 3, Table 1).

(a) Cu(0)/Selectfluor system-catalyzed difunctionalization of C-C multiple bonds (previous work)

(b) Concept for the synthesis of 6H-phenanthridine via oxycupration of C=N bond (This work)

Scheme 1 Cu(0)/Selectfluor system-catalyzed difunctionalization of carbon-carbon multiple bonds and carbon-nitrogen double bond

With the optimized reaction conditions, we then set out to investigate the reaction scope (Table 2). Firstly, the effect of functional groups in the aryl ring A (R^1) on the reaction was studied. As shown in Table 2, both electron-donating and electron-withdrawing substituents in the aryl ring A of **1** were compatible with the reaction conditions, among which alkyl, halo (F, Cl, Br), and trifluoromethoxy groups were included. It seemed that the electronic nature of R^1 had no obvious effect on the reaction outcome in terms of the yield. In most cases, the reaction could give moderate to good yields of desired products (60%~80%, **2a**~**2j**, Table 2). Substrate **1j** bearing a *me*-

ta-chloro group in ring A provided a single product (**2j**, 66%) with exclusive regio-selectivity at the more steric hindrance position which was in accordance with previous report.^[8h] Next, the effect of substituents (R^2) in aryl ring B of **1** on the reaction was further investigated. A range of substrates bearing electron-rich or electron-deficient substituents in ring B were workable for the reaction (**1k**~**1u**, Table 2). Similarly, the electronic properties of R^2 had no obvious effect on the reaction performance and the yield of products (**2k**~**2u**, Table 2). Unfortunately, 2-phenylnicotinaldehyde derived sulfonylimide **1w** failed to give the desired product while the starting material was almost recovered (**2w**, Table 2).

To gain insight into the mechanism of the Cu(0)/Selectfluor system-catalyzed annulation/aromatization of *o*-aryl benzenesulfonylimides, several mechanistic experiments were made (Scheme 2). Firstly, the measurement of the intermolecular KIE between **1a** and **1a-d₅** on the basis of the competitive dehydrogenative cross-coupling was carried out. The intermolecular k_H/k_D of **1a** to **1a-d₅** was determined to be 1.85, indicating that the cleavage of the C—H bond might be involved in the rate-determining step (Scheme 2). When **1a** was allowed to react under the standard reaction conditions except using the preparative HOCuX ($X=F$ or BF_4)^[11a,11f] as a catalyst, the reaction could also proceed smoothly and afford **2a** in 80% yield (Scheme 2). This result suggests that HOCuX ($X=F$ or BF_4) might play a key role in the catalytic cycle.

Scheme 2 Preliminary mechanistic studies

Based on the above experiments and previous literature,^[11,13,14] a possible mechanism for the Cu(0)/Selectfluor system-catalyzed annulation/aromatization of **1a** is proposed in Scheme 3. First, the redox reaction between Cu(0) and Selectfluor may produce a copper species $FCuBF_4$ (**5**) and release base **6**.^[11a] With the aid of **6**, **5** may be transferred into copper species $XCuOH$ ($X=F$ or BF_4 , **7**) in the presence of water.^[11] Then, an oxycupration of **7** to the C—N double bond in **1a** followed by dehydrogenative cross-coupling^[11e,13] would deliver intermediate **10**. Finally, **10** might undergo Julia-type olefination^[14] to give the aromatization product **2a**.

Table 1 Optimization of the reaction conditions^a

Entry	Catalyst	Oxidant	Solvent	Temp./°C	Yield ^b /%
1	Cu(0)	Selectfluor	MeCN ^c	50	63
2	Cu(0)	Selectfluor	Anhydrous MeCN	50	15
3	Cu(0)	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	84 (80) ^d
4	Cu(0)	Selectfluor	MeCN/H ₂ O (V : V=800 : 1)	50	82
5	Cu(0)	Selectfluor	MeCN/H ₂ O (V : V=250 : 1)	50	72
6	Cu(0)	Selectfluor	Solvent ^e /H ₂ O (V : V=500 : 1)	50	0
7	CuI	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	55
8	CuCl	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	63
9	CuBr	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	77
10	Cu ₂ O	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	72
11	CuCl ₂	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	47
12	Cu(OAc) ₂	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	48
13	—	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	0
14	Cu(0)	—	MeCN/H ₂ O (V : V=500 : 1)	50	0
15	Cu(0)	NFSI ^f	MeCN/H ₂ O (V : V=500 : 1)	50	0
16	Cu(0)	Other oxidant ^g	MeCN/H ₂ O (V : V=500 : 1)	50	0
17	Cu(0)	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	25	69
18	Cu(0)	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	80	80
19	Cu(0) ^h	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	76
20	Cu(0)	Selectfluor	MeCN/H ₂ O (V : V=500 : 1)	50	46 ⁱ , 82 ^j

^a Reaction conditions: **1a** (0.3 mmol), Cu(0) powder (10 mol% based on **1a**), oxidant (2.0 equiv.), solvent (3.5 mL), 50 °C under N₂ atmosphere for 24 h unless otherwise noted. ^b GC yield by using dodecane as an internal standard. ^c Newly purchased CH₃CN was used without dehydration. ^d Isolated yield. ^e Tested solvents include toluene, acetone, 1,4-dioxane, DMF, DMSO, 1,2-dichloroethane and methanol. ^f NFSI: *N*-fluorobenzenesulfonimide. ^g Other tested oxidants include PhI(OAc)₂, DDQ and K₂S₂O₈. ^h The amount of Cu(0) is 5 mol% based on **1a**. ⁱ The amount of Selectfluor is 1.0 equiv. ^j The amount of Selectfluor is 3.0 equiv.

Table 2 Synthesis of **2** via Cu(0)/Selectfluor system-catalyzed annulation/aromatization of *o*-aryl benzenesulfonylimides **1**^{a,b}

^a Reaction conditions: **1** (0.3 mmol), Cu(0) powder (10 mol% based on **1**), Selectfluor (2.0 equiv.), CH₃CN/H₂O (V/V=500 : 1) (3.5 mL), 50 °C under N₂ atmosphere for 24 h unless otherwise noted. ^b Isolated yield.

Scheme 3 Proposed mechanism for the annulation/aromatization of **1a**

3 Conclusions

In summary, a facile and efficient method for the preparation of 6*H*-phenanthridines involving Cu(0)/Selectfluor system-catalyzed tandem oxycupration/C—H amination/aromatization of *o*-aryl benzenesulfonylimides was described. The present protocol for the synthesis of 6*H*-phenanthridines has several advantages including mild reaction conditions (50 °C), the use of inexpensive copper catalyst, and easy availability of the starting materials.

4 Experimental section

4.1 General experimental information

Unless otherwise stated, all reagents were purchased from commercial suppliers and used without purifications. Melting points are uncorrected. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker ADVANCE III at 25 °C in CDCl₃ at 500 (or 600) MHz and 125 (or 150) MHz, respectively. GC-MS experiments were performed with an Agilent 6890N GC system equipped with a 5973N mass-selective detector. High resolution mass spectra (HRMS) were obtained on a water GCT premier TOF MS with EI or ESI source. CH₃CN is dehydrated by CaH₂ before preparation of the combined solvent CH₃CN/H₂O (*V* : *V* = 500 : 1).

4.2 General procedure for the synthesis of starting material 1

All *o*-aryl benzenesulfonylimides (**1**) were synthesized from the corresponding *o*-aryl benzaldehydes and benzenesulfonamide according to the literature procedure.^[15] The *o*-aryl benzaldehydes were synthesized by the standard Suzuki-Miyaura cross-coupling reaction between an *ortho*-bromobenzaldehyde derivative and an arylboronic acid.^[16]

4.3 General procedure for the synthesis of 2

1 (0.3 mmol), Cu(0) powder (1.92 mg, 10 mol%), Selectfluor (212.6 mg, 0.6 mmol, 2 equiv.), and CH₃CN/H₂O (*V* : *V* = 500 : 1) (3.5 mL) were added to a 15 mL flask. Then the reaction mixture was stirred at 50 °C for 24 h under N₂ atmosphere. Upon completion, the resulting mixture was diluted with CH₂Cl₂ (10 mL) and filtered through Celite. After evaporation of the solvent under vacuum, the residue was purified by column chromatography on silica gel (100~200 mesh) using petroleum ether-EtOAc (*V* : *V* = 10 : 1) as eluent to give pure **2**.

Phenanthridine (2a): White solid (43.0 mg, 80%), m.p. 109~111 °C (lit.^[17] 107~108 °C); ¹H NMR (CDCl₃, 500 MHz) δ: 9.27 (s, 1H), 8.53 (t, *J* = 8.5 Hz, 2H), 8.21 (d, *J* = 8.5 Hz, 1H), 8.00 (d, *J* = 7.5 Hz, 1H), 7.82~7.79 (m, 1H), 7.75~7.72 (m, 1H), 7.67~7.64 (m, 2H); ¹³C NMR (CDCl₃, 125 MHz) δ: 153.4, 144.4, 132.4, 130.9, 130.0, 128.61, 128.57, 127.4, 127.0, 126.3, 124.0, 122.1, 121.7.

3-Methylphenanthridine (2b): Yellow solid (34.8 mg, 60%), m.p. 80~81 °C (lit.^[18] 78~79 °C); ¹H NMR (CDCl₃, 500 MHz) δ: 9.27 (s, 1H), 8.56 (d, *J* = 8.0 Hz, 1H), 8.46 (d, *J* = 8.5 Hz, 1H), 8.03 (d, *J* = 8.0 Hz, 1H), 8.00 (s, 1H), 7.86~7.82 (m, 1H), 7.69~7.66 (m, 1H), 7.52 (dd, *J* = 8.5, 1.5 Hz, 1H), 2.61 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz) δ: 153.5, 144.5, 138.9, 132.6, 131.0, 130.7, 129.6, 128.8, 128.7, 127.0, 126.1, 122.0, 121.7, 21.5.

3-Isopropylphenanthridine (2c): Yellow liquid (42.5 mg, 64%); ¹H NMR (CDCl₃, 500 MHz) δ: 9.27 (s, 1H), 8.55 (d, *J* = 8.5 Hz, 1H), 8.48 (d, *J* = 8.5 Hz, 1H), 8.07 (d, *J* = 1.5 Hz, 1H), 8.01 (d, *J* = 8.0 Hz, 1H), 7.83~7.80 (m, 1H), 7.66 (t, *J* = 7.5 Hz, 1H), 7.58 (dd, *J* = 8.0, 1.5 Hz, 1H), 3.22~3.14 (m, 1H), 1.41 (d, *J* = 6.5 Hz, 6H); ¹³C NMR (CDCl₃, 125 MHz) δ: 153.4, 149.7, 144.7, 132.6, 130.9, 128.7, 127.0, 126.9, 126.5, 126.1, 122.1, 121.7, 34.1, 23.9; HRMS (EI) calcd for C₁₆H₁₅N 221.1204, found 221.1199.

3-(*tert*-Butyl)phenanthridine (2d): Yellow liquid (45.2 mg, 64%); ¹H NMR (CDCl₃, 500 MHz) δ: 9.27 (s, 1H), 8.55 (d, *J* = 8.5 Hz, 1H), 8.48 (d, *J* = 8.5 Hz, 1H), 8.22 (d, *J* = 2.0 Hz, 1H), 8.00 (d, *J* = 8.0 Hz, 1H), 7.83~7.80 (m, 1H), 7.76 (d, *J* = 8.5, 2.0 Hz, 1H), 7.67~7.64 (m, 1H), 1.49 (s, 9H); ¹³C NMR (CDCl₃, 125 MHz) δ: 153.3, 152.0, 144.4, 132.4, 130.8, 128.6, 127.7, 126.9, 126.1, 126.0, 125.3, 121.8, 121.6, 34.9, 31.3; HRMS (EI) calcd for C₁₇H₁₇N 235.1361, found 235.1367.

3-Phenylphenanthridine (2e): White solid (57.4 mg, 75%); m.p. 151~153 °C; ¹H NMR (CDCl₃, 500 MHz) δ: 9.33 (s, 1H), 8.63 (dd, *J* = 8.5, 4.5 Hz, 2H), 8.46 (d, *J* = 1.5 Hz, 1H), 8.07 (d, *J* = 7.5 Hz, 1H), 7.96 (dd, *J* = 8.5, 1.5 Hz, 1H), 7.90~7.87 (m, 1H), 7.82 (d, *J* = 7.5 Hz, 2H), 7.73 (t, *J* = 7.5 Hz, 1H), 7.54 (t, *J* = 7.5 Hz, 2H), 7.44 (t, *J* = 7.0 Hz, 1H); ¹³C NMR (CDCl₃, 125 MHz) δ: 154.0, 144.9, 141.4, 140.2, 132.4, 131.1, 129.0, 128.8, 128.0, 127.8, 127.5, 127.4, 126.4, 126.2, 123.1, 122.8, 121.9; HRMS (EI) calcd or C₁₉H₁₃N 255.1048, found 255.1043.

3-Fluorophenanthridine (2f): White solid (37.3 mg, 63%); m.p. 109~111 °C; ¹H NMR (CDCl₃, 500 MHz) δ:

9.28 (s, 1H), 8.54~8.50 (m, 2H), 8.04 (d, $J=8.0$ Hz, 1H), 7.87~7.83 (m, 2H), 7.72~7.68 (m, 1H), 7.45~7.41 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 162.6 (d, $J=247.5$ Hz), 154.7, 145.7 (d, $J=11.3$ Hz), 132.3, 131.4, 128.9, 127.3, 125.9, 124.1 (d, $J=8.8$ Hz), 121.6, 120.8, 116.1, (d, $J=23.8$ Hz), 114.6 (d, $J=21.3$ Hz); HRMS (EI) calcd for $\text{C}_{13}\text{H}_8\text{FN}$ 197.0641, found 197.0645.

3-Chlorophenanthridine (**2g**): White solid (39.7 mg, 62%); m.p. 106.8~108.2 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.21 (s, 1H), 8.43 (d, $J=8.5$ Hz, 1H), 8.37 (d, $J=9.0$ Hz, 1H), 8.12 (d, $J=1.5$ Hz, 1H), 7.98 (d, $J=7.5$ Hz, 1H), 7.82 (t, $J=7.0$ Hz, 1H), 7.68 (t, $J=7.5$ Hz, 1H), 7.56 (dd, $J=9.0$, 2.0 Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 154.5, 144.9, 134.2, 131.9, 131.3, 129.2, 128.8, 127.7, 127.5, 126.1, 123.4, 122.4, 121.6; HRMS (EI) calcd for $\text{C}_{13}\text{H}_8\text{ClN}$ 213.0345, found 213.0340.

3-Bromophenanthridine (**2h**): White solid (49.6 mg, 64%); m.p. 129~131 °C (lit.^[19] 110~112 °C); ^1H NMR (CDCl_3 , 500 MHz) δ : 9.27 (s, 1H), 8.54 (d, $J=8.0$ Hz, 1H), 8.40 (d, $J=8.5$ Hz, 1H), 8.34 (d, $J=2.0$ Hz, 1H), 8.05 (d, $J=8.0$ Hz, 1H), 7.89~7.86 (m, 1H), 7.77~7.72 (m, 2H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 154.6, 145.4, 132.6, 132.1, 131.4, 130.3, 128.9, 127.9, 126.3, 123.7, 122.9, 122.4, 121.7.

3-(Trifluoromethoxy)phenanthridine (**2i**): White solid (50.5 mg, 64%); m.p. 87~89 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.25 (s, 1H), 8.60 (dd, $J=12.0$, 9.0 Hz, 2H), 8.10 (d, $J=8.0$ Hz, 1H), 8.06 (s, 1H), 7.94~7.90 (m, 1H), 7.79~7.75 (m, 1H), 7.57 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 155.0, 149.2, 145.3, 132.1, 131.6, 129.0, 128.0, 126.4, 124.1, 122.7, 121.9, 120.9, 120.7 (q, $J=256.7$ Hz), 120.4; HRMS (EI) calcd for $\text{C}_{14}\text{H}_8\text{F}_3\text{NO}$ 263.0558, found 263.0554.

4-Chlorophenanthridine (**2j**): White solid (42.3 mg, 66%); m.p. 100~102 °C (lit.^[20] 98~102 °C); ^1H NMR (CDCl_3 , 500 MHz) δ : 9.25 (s, 1H), 8.50 (dd, $J=5.5$, 3.0 Hz, 2H), 8.11 (d, $J=8.5$ Hz, 1H), 8.05 (d, $J=8.0$ Hz, 1H), 7.89~7.86 (m, 1H), 7.75 (t, $J=7.5$ Hz, 1H), 7.68 (dd, $J=9.0$, 2.5 Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 153.8, 142.9, 133.1, 131.62, 131.56, 131.3, 129.2, 128.9, 128.2, 126.5, 125.2, 121.93, 121.90.

9-Methylphenanthridine (**2k**): Yellow solid (38.3 mg, 66%); m.p. 85~86 °C (lit.^[21] 82~84 °C); ^1H NMR (CDCl_3 , 500 MHz) δ : 9.21 (s, 1H), 8.53 (d, $J=8.5$ Hz, 1H), 8.34 (s, 1H), 8.18 (dd, $J=8.0$, 0.5 Hz, 1H), 7.90 (d, $J=8.0$ Hz, 1H), 7.74~7.71 (m, 1H), 7.67~7.63 (m, 1H), 7.49 (dd, $J=8.0$, 0.5 Hz, 1H), 2.63 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 153.2, 144.5, 141.4, 132.6, 133.0, 129.2, 128.52, 128.46, 126.8, 124.5, 124.0, 122.1, 121.4, 22.4.

9-Chlorophenanthridine (**2l**): White solid (45.5 mg, 71%); m.p. 113~115 °C (lit.^[22] 118.5 °C); ^1H NMR (CDCl_3 , 500 MHz) δ : 9.16 (s, 1H), 8.43 (s, 1H), 8.36 (d, $J=8.0$ Hz, 1H), 8.15 (d, $J=8.0$ Hz, 1H), 7.88 (d, $J=8.5$ Hz, 1H), 7.74 (t, $J=7.5$ Hz, 1H), 7.64 (t, $J=7.5$ Hz, 1H), 7.57 (dd, $J=8.5$, 1.5 Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 152.5, 144.5, 137.3, 133.5, 130.1, 130.0, 129.3,

128.0, 127.3, 124.4, 122.8, 122.1, 121.5.

9-Fluoro-3-isopropylphenanthridine (**2m**): White solid (46.7 mg, 65%); m.p. 57~59 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.20 (s, 1H), 8.33 (d, $J=8.0$ Hz, 1H), 8.11 (dd, $J=10.0$, 2.0 Hz, 1H), 8.04 (d, $J=2.0$ Hz, 1H), 8.00 (dd, $J=9.0$, 5.5 Hz, 1H), 7.57 (dd, $J=8.5$, 2.0 Hz, 1H), 7.39~7.35 (m, 1H), 3.21~3.13 (m, 1H), 1.40 (d, $J=7.0$ Hz, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 164.1 (d, $J=250.0$ Hz), 152.5, 150.6, 144.7, 134.8 (d, $J=10.0$ Hz), 131.4 (d, $J=10.0$ Hz), 127.0, 126.6, 123.1, 122.3, 121.6, 116.3 (d, $J=25.0$ Hz), 106.9 (d, $J=21.3$ Hz), 34.1, 23.9; HRMS (EI) calcd for $\text{C}_{16}\text{H}_{14}\text{FN}$ 239.1110, found 239.1115.

9-Chloro-3-isopropylphenanthridine (**2n**): White solid (46.8 mg, 61%); m.p. 169~171 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.16 (s, 1H), 8.43 (s, 1H), 8.36 (d, $J=8.0$ Hz, 1H), 8.15 (d, $J=8.0$ Hz, 1H), 7.88 (d, $J=8.5$ Hz, 1H), 7.74 (t, $J=7.5$ Hz, 1H), 7.64 (t, $J=7.5$ Hz, 1H), 7.57 (dd, $J=8.5$, 1.5 Hz, 1H), 3.23~3.14 (m, 1H), 1.40 (d, $J=7.0$ Hz, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 152.5, 144.5, 137.3, 133.5, 130.1, 130.0, 129.3, 128.0, 127.3, 124.4, 122.8, 122.1, 121.5, 34.1, 23.9; HRMS (EI) calcd for $\text{C}_{16}\text{H}_{14}\text{ClN}$ 255.0815, found 255.0811.

8-Chlorophenanthridine (**2o**): White solid (40.4 mg, 63%); m.p. 99~101 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.11 (s, 1H), 8.38 (dd, $J=7.5$, 5.0 Hz, 2H), 8.15 (d, $J=8.0$ Hz, 1H), 7.90 (d, $J=2.0$ Hz, 1H), 7.74~7.71 (m, 1H), 7.68 (dd, $J=9.0$, 2.5 Hz, 1H), 7.65~7.61 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 152.1, 144.2, 133.2, 131.3, 130.6, 130.2, 128.9, 127.5, 127.4, 127.0, 123.5, 123.4, 121.9; HRMS (EI) calcd for $\text{C}_{13}\text{H}_8\text{ClN}$ 213.0345, found 213.0342.

8-Chloro-3-isopropylphenanthridine (**2p**): White solid (46.0 mg, 60%); m.p. 49~51 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.21 (s, 1H), 8.50 (d, $J=1.5$ Hz, 1H), 8.38 (d, $J=8.5$ Hz, 1H), 8.04 (d, $J=1.5$ Hz, 1H), 7.94 (d, $J=8.5$ Hz, 1H), 7.61~7.58 (m, 2H), 3.23~3.14 (m, 1H), 1.40 (d, $J=7.0$ Hz, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 152.6, 150.6, 144.9, 132.8, 131.5, 131.0, 127.7, 127.1, 127.03, 126.94, 123.6, 122.0, 121.6, 34.1, 23.9; HRMS (EI) calcd for $\text{C}_{16}\text{H}_{14}\text{ClN}$ 255.0815, found 255.0812.

N-8-Chloro-3-fluorophenanthridine (**2q**): White solid (47.3 mg, 68%); m.p. 144~146 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.14 (s, 1H), 8.41 (dd, $J=9.0$, 6.0 Hz, 1H), 8.36 (d, $J=8.5$ Hz, 1H), 7.94 (d, $J=2.0$ Hz, 1H), 7.80 (dd, $J=9.5$, 2.5 Hz, 1H), 7.74 (dd, $J=8.5$, 2.0 Hz, 2H), 7.43~7.40 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 162.7 (d, $J=247.5$ Hz), 153.3, 145.6 (d, $J=12.5$ Hz), 133.1, 131.9, 130.5, 127.7, 126.6, 123.9 (d, $J=10.0$ Hz), 123.4, 120.2, 116.6 (d, $J=23.8$ Hz), 114.8 (d, $J=21.3$ Hz); HRMS (EI) calcd for $\text{C}_{13}\text{H}_7\text{ClFN}$ 231.0251, found 231.0255.

3-Chloro-8-fluorophenanthridine (**2r**): White solid (44.5 mg, 64%); m.p. 165~167 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 9.15 (s, 1H), 8.44 (dd, $J=9.0$, 5.0 Hz, 1H), 8.31 (d, $J=9.0$ Hz, 1H), 8.10 (d, $J=2.5$ Hz, 1H), 7.61 (dd, $J=8.0$, 2.5 Hz, 1H), 7.58~7.55 (m, 2H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 161.5 (d, $J=248.8$ Hz), 153.4 (d, $J=1.3$ Hz), 144.6, 134.1, 129.4, 128.6 (d, $J=1.3$ Hz), 127.9, 127.3 (d,

$J=8.8$ Hz), 124.4 (d, $J=8.8$ Hz), 123.2, 122.1, 120.6 112.8 (d, $J=21.3$ Hz); HRMS (EI) calcd for $C_{13}H_7ClFN$ 231.0251, found 231.0254.

8-Bromophenanthridine (**2s**): White solid (41.0 mg, 53%); m.p. 96~98 °C; 1H NMR ($CDCl_3$, 500 MHz) δ : 9.19 (s, 1H), 8.51 (dd, $J=8.0, 1.0$ Hz, 1H), 8.44 (d, $J=8.5$ Hz, 1H), 8.19 (dd, $J=8.0, 1.0$ Hz, 1H), 8.17 (d, $J=2.0$ Hz, 1H), 7.91 (dd, $J=8.5, 2.0$ Hz, 1H), 7.80~7.76 (m, 1H), 7.71~7.68 (m, 1H); ^{13}C NMR ($CDCl_3$, 125 MHz) δ : 152.1, 144.3, 134.1, 131.1, 130.9, 130.3, 129.1, 127.52, 127.48, 123.8, 123.5, 122.0, 121.3; HRMS (EI) calcd for $C_{13}H_8BrN$ 256.9840, found 256.9847.

8-Methylphenanthridine (**2t**): Yellow solid (35.9 mg, 62%); m.p. 80~82 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 9.23 (s, 1H), 8.55 (d, $J=8.4$ Hz, 1H), 8.51 (d, $J=8.4$ Hz, 1H), 8.19~8.17 (m, 1H), 7.83 (s, 1H), 7.73~7.65 (m, 3H), 2.61 (s, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 153.3, 144.3, 137.5, 132.8, 130.5, 130.1, 128.19, 128.15, 127.0, 126.6, 124.2, 122.0, 121.8, 21.5; HRMS (EI) calcd for $C_{14}H_{11}N$ 193.0891, found 193.0896.

7-Fluorophenanthridine (**2u**): White solid (33.1 mg, 56%); m.p. 114~116 °C; 1H NMR ($CDCl_3$, 500 MHz) δ : 9.49 (s, 1H), 8.37 (d, $J=8.5$ Hz, 1H), 8.19~8.14 (m, 2H), 7.72~7.59 (m, 3H), 7.23 (t, $J=8.0$ Hz, 1H); ^{13}C NMR ($CDCl_3$, 125 MHz) δ : 159.9 (d, $J=247.5$ Hz), 146.0 (d, $J=6.3$ Hz), 144.3, 134.0 (d, $J=2.5$ Hz), 131.2 (d, $J=8.8$ Hz), 130.2, 128.1, 127.3, 122.9, 122.3, 117.5 (d, $J=3.8$ Hz), 115.8 (d, $J=13.8$ Hz), 111.9 (d, $J=18.8$ Hz); HRMS (EI) calcd for $C_{13}H_8FN$ 197.0641, found 197.0644.

Benzo[*c*]phenanthridine (**2v**)^[23]: White solid (33.0 mg, 48%); m.p. 141~143 °C (lit.^[23] 136 °C); 1H NMR ($CDCl_3$, 500 MHz) δ : 9.50 (s, 1H), 9.43 (d, $J=8.5$ Hz, 1H), 8.68 (d, $J=8.5$ Hz, 1H), 8.56 (d, $J=9.0$ Hz, 1H), 8.16 (d, $J=8.0$ Hz, 1H), 8.05 (d, $J=9.0$ Hz, 1H), 8.01 (d, $J=8.0$ Hz, 1H), 7.92~7.88 (m, 1H), 7.82~7.78 (m, 1H), 7.76~7.70 (m, 2H); ^{13}C NMR ($CDCl_3$, 125 MHz) δ : 152.0, 141.5, 133.3, 132.9, 132.1, 130.8, 128.7, 127.9, 127.7, 127.4, 127.1, 127.0, 126.9, 124.8, 122.2, 121.1, 119.9; HRMS (EI) calcd for $C_{17}H_{11}N$ 229.0891, found 229.0896.

4.4 Intermolecular kinetic isotope effects (KIE) based on substrate **1a** and **1a-d₅**

1a (64.3 mg, 0.2 mmol), **1a-d₅** (65.3 mg, 0.2 mmol), Cu(0) powder (2.56 mg, 10 mol%), Selectfluor (283.5 mg, 0.8 mmol, 2 equiv.), and CH_3CN/H_2O ($V:V=500:1$, 4.5 mL) were added to a 15 mL flask. Then the reaction mixture was stirred at 50 °C for 12 h. Upon completion, the resulting mixture was diluted with CH_2Cl_2 (10 mL) and filtered through Celite. After evaporation of the solvent under vacuum, the residue was purified by column chromatography on silica gel (100~200 mesh) using petroleum ether-EtOAc ($V:V=10:1$) as eluent to give pure products. On the basis of 1H NMR spectra analysis, the intermolecular competitive KIE was calculated as $k_H/k_D \approx 1.85$.

4.5 Reaction of **1a** in the standard reaction conditions except using the preparative HOCuX as the catalyst

1a (0.3 mmol), HOCuX ($X=F$ or BF_4 , 10 mol%, prepared according to our previous work^[11a]), Selectfluor (212.6 mg, 0.6 mmol, 2 equiv.), and CH_3CN/H_2O ($V:V=500:1$) solvent (3.5 mL) were added to a 15 mL flask. Then the reaction mixture was stirred at 50 °C for 24 h. Upon completion, the resulting mixture was diluted with CH_2Cl_2 (10 mL) and filtered through Celite. After evaporation of the solvent under vacuum, the residue was purified by column chromatography on silica gel (100~200 mesh) using petroleum ether-EtOAc ($V:V=10:1$) as eluent to give pure **2a** (43.0 mg, 80%).

Supporting Information 1H NMR, ^{13}C NMR and HRMS of the products **2a**~**2w**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Cushman, M.; Mohan, P.; Smith, E. C. R. *J. Med. Chem.* **1984**, 27, 544.
(b) Fang, S. D.; Wang, L. K.; Hecht, S. M. *J. Org. Chem.* **1993**, 58, 5025.
(c) Lynch, M. A.; Duval, O.; Sukhanova, A.; Devy, J.; Mackay, S. P.; Waigh, R. D.; Nabiev, I. *Bioorg. Med. Chem. Lett.* **2001**, 11, 2643.
(d) Bernardo, P. H.; Wan, K. F.; Sivaraman, T.; Xu, J.; Moore, F. K.; Hung, A. W.; Mok, H. Y. K.; Yu, V. C.; Chai, C. L. L. *J. Med. Chem.* **2008**, 51, 6699.
(e) Zhu, S.; Ruchelman, A. L.; Zhou, N.; Liu, A.; Liu, L. F.; LaVoie, E. J. *Bioorg. Med. Chem.* **2005**, 13, 6782.
(f) Li, D.; Zhao, B.; Sim, S.-P.; Li, T.-K.; Liu, A.; Liu, L. F.; Edmond, J.; LaVoie, E. J. *Bioorg. Med. Chem.* **2003**, 11, 521.
(g) Tsukamoto, H.; Kondo, S.; Mukudai, Y.; Nagumo, T.; Yasuda, A.; Kurihara, Y.; Kamatani, T.; Shintani, S. *Anticancer Res.* **2011**, 31, 2841.
- [2] (a) Cushman, M.; Mohan, P.; Smith, E. C. R. *J. Med. Chem.* **1984**, 27, 544.
(b) Fang, S. D.; Wang, L. K.; Hecht, S. M. *J. Org. Chem.* **1993**, 58, 5025.
(c) Lynch, M. A.; Duval, O.; Sukhanova, A.; Devy, J.; Mackay, S. P.; Waigh, R. D.; Nabiev, I. *Bioorg. Med. Chem. Lett.* **2001**, 11, 2643.
(d) Bernardo, P. H.; Wan, K. F.; Sivaraman, T.; Xu, J.; Moore, F. K.; Hung, A. W.; Mok, H. Y. K.; Yu, V. C.; Chai, C. L. L. *J. Med. Chem.* **2008**, 51, 6699.
(e) Zhu, S.; Ruchelman, A. L.; Zhou, N.; Liu, A.; Liu, L. F.; LaVoie, E. J. *Bioorg. Med. Chem.* **2005**, 13, 6782.
(f) Li, D.; Zhao, B.; Sim, S.-P.; Li, T.-K.; Liu, A.; Liu, L. F.; Edmond, J.; LaVoie, E. J. *Bioorg. Med. Chem.* **2003**, 11, 521.
(g) Tsukamoto, H.; Kondo, S.; Mukudai, Y.; Nagumo, T.; Yasuda, A.; Kurihara, Y.; Kamatani, T.; Shintani, S. *Anticancer Res.* **2011**, 31, 2841.
- [3] (a) Stevens, N.; O'Connor, N.; Vishwasrao, H.; Samaroo, D.; Kandell, E. R.; Akins, D. L.; Drain, C. M.; Rurro, N. J. *J. Am. Chem. Soc.* **2008**, 130, 7182.
(b) Bondarev, S. L.; Knyukshto, V. N.; Tikhomirov, S. A.; Pyrko, A. N. *Opt. Spectrosc.* **2006**, 100, 386.
(c) Zhang, J.; Lakowicz, J. R. *J. Phys. Chem. B* **2005**, 109, 8701.
- [4] For selected recent reviews, see: (a) Zhang, B.; Studer, A. *Chem. Soc. Rev.* **2015**, 44, 3505.

- (b) Hayashi, H.; Kaga, A.; Chiba, S. *J. Org. Chem.* **2017**, *82*, 11981.
 (c) Hu, B.; DiMaggio, S. G. *Org. Biomol. Chem.* **2015**, *13*, 3844.
- [5] (a) Pictet, A.; Hubert, A. *Ber. Dtsch. Chem. Ges.* **1896**, *29*, 1182.
 (b) Morgan, T.; Walls, L. P. *J. Chem. Soc.*, 1931, 2447.
 (c) Chinnagolla, R. K.; Jeganmohan, M. *Chem. Commun.* **2014**, *50*, 2442.
- [6] (a) Ge, J.; Wang, X.; Liu, T.; Shi, Z.; Xiao, Q.; Yin, D. *RSC Adv.* **2016**, *6*, 19571.
 (b) Sahoo, M. K.; Midya, S. P.; Landge, V. G.; Balaraman, E. *Green Chem.* **2017**, *19*, 2111.
 (c) Candito, D. A.; Lautens, M. *Angew. Chem. Int. Ed.* **2009**, *48*, 6713.
 (d) Xu, Z.; Hang, Z.; Liu, Z.-Q. *Org. Lett.* **2016**, *18*, 4470.
 (e) Li, Z.; Fan, F.; Yang, J.; Liu, Z.-Q. *Org. Lett.* **2014**, *16*, 3396.
 (f) Xu, Z.; Yan, C.; Liu, Z.-Q. *Org. Lett.* **2014**, *16*, 5670.
- [7] (a) Dai, Q.; Yun, J.-T.; Feng, X.; Jiang, Y.; Yang, H.; Cheng, J. *Adv. Synth. Catal.* **2014**, *356*, 3341.
 (b) Sha, W.; Yu, J.-T.; Jiang, Y.; Yang, H.; Cheng, J. *Chem. Commun.* **2014**, *50*, 9179.
 (c) Tu, H.-Y.; Liu, Y.-R.; Chu, J.-J.; Hu, B.-L.; Zhang, X.-G. *J. Org. Chem.* **2014**, *79*, 9907.
 (d) Jiang, H.; An, X.; Tong, K.; Zhang, Y.; Yu, S. *Angew. Chem. Int. Ed.* **2015**, *54*, 4055.
 (e) Cao, J.-J.; Zhu, T.-H.; Wang, S.-Y.; Gu, Z.-Y.; Wang, X.; Ji, S.-J. *Chem. Commun.* **2014**, *50*, 6439.
 (f) Zhang, B.; Mück-Lichtenfeld, C.; Daniliuc, C. G.; Studer, A. *Angew. Chem. Int. Ed.* **2013**, *52*, 10792.
 (g) Sun, X.; Yu, S. *Chem. Commun.* **2016**, *52*, 10898.
 (h) Wang, Y.-F.; Lonca, G. H.; Runigo, M. L.; Chiba, S. *Org. Lett.*, **2014**, *16*, 4272.
 (i) Lu, L.; Zhou, B.; Jin, H.; Liu, Y. *Chin. J. Org. Chem.* **2019**, *39*, 515 (in Chinese).
 (陆露露, 周丙伟, 金红卫, 刘运奎, 有机化学, **2019**, *39*, 515.)
 (j) Shi, D.; Bao, H.; Xu, Z.; Liu, Y. *Chin. J. Org. Chem.* **2017**, *37*, 1290 (in Chinese).
 (施冬冬, 鲍汉扬, 徐峥, 刘运奎, 有机化学, **2017**, *37*, 1290.)
- [8] (a) Evoniuk, C. J.; dos Passos Gomes, Hill, G.; S. P.; Fujita, S.; Hanson, K.; Alabugin, I. V. *J. Am. Chem. Soc.* **2017**, *139*, 16210.
 (b) Evoniuk, C. J.; Hill, S. P.; Hanson, K.; Alabugin, I. V. *Chem. Commun.* **2016**, *52*, 7138.
 (c) Zhang, L.; Ang, G. Y.; Chiba, S. *Org. Lett.* **2010**, *12*, 3682.
- [9] (a) Liu, Y.-Y.; Song, R.-J.; Wu, C.-Y.; Gong, L.-B.; Hu, M.; Wang, Z.-Q.; Xie, Y.-X.; Li, J.-H. *Adv. Synth. Catal.* **2012**, *354*, 347.
 (b) Borah, A.; Gogoi, P. *Eur. J. Org. Chem.* **2016**, 2200.
 (c) Han, W.; Zhou, X.; Yang, S.; Xiang, G.; Cui, B.; Chen, Y. *J. Org. Chem.* **2015**, *80*, 11580.
 (d) Maestri, G.; Larraufie, M.-H.; Derat, É.; Ollivier, C.; Fensterbank, L.; Lacôte, E.; Malacria, M. *Org. Lett.* **2010**, *12*, 5692.
 (e) Portela-Cubillo, F.; Scott, J. S.; Walton, J. C. *J. Org. Chem.* **2008**, *73*, 5558.
 (f) Jiang, H.; An, X.; Tong, K.; Zheng, T.; Zhang, Y.; Yu, S. *Angew. Chem. Int. Ed.* **2015**, *54*, 4055.
 (g) An, X.-D.; Yu, S. *Org. Lett.* **2015**, *17*, 2692.
 (h) Ghosh, M.; Ahmed, A.; Singha, R.; Ray, J. K. *Tetrahedron Lett.* **2015**, *56*, 353.
 (i) Portela, C. F.; Scanlan, E. M.; Scott, J. S.; Walton, J. C. *Chem. Commun.* **2008**, 4189.
 (j) Tummatorn, J.; Krajangsri, S.; Norseeda, K.; Thongsornkleeb, C.; Ruchirawat, S. *Org. Biomol. Chem.* **2014**, *12*, 5077.
- (k) Budén, M.; Dorn, V. B.; Gamba, M.; Pierini, A. B.; Rossi, R. A. *J. Org. Chem.* **2010**, *75*, 2206.
 (l) Linsenmeier, A. M.; Williams, C. M.; Bräse, S. *J. Org. Chem.* **2011**, *76*, 9127.
 (m) McBurney, R. T.; Slawin, A. M. Z.; Smart, L. A.; Yu, Y.; Walton, J. C. *Chem. Commun.* **2011**, 47, 7974.
 (n) Hofstra, J. L.; Grassbaugh, B. R.; Tran, Q. M.; Armada, N. R.; de Lijser, H. J. P. *J. Org. Chem.* **2015**, *80*, 256.
 (o) Chen, W.-L.; Chen, C.-Y.; Chen, Y.-F.; Hsieh, J.-C. *Org. Lett.* **2015**, *17*, 1613.
- [10] Selected reviews:
 (a) Yin, G.; Mu, X.; Liu, G. *Acc. Chem. Res.* **2016**, *49*, 2413.
 (b) Egami, H.; Sodeoka, M. *Angew. Chem. Int. Ed.* **2014**, *53*, 8294.
 (c) Shimizu, Y.; Kanai, M. *Tetrahedron Lett.* **2014**, *55*, 3727.
 (d) McDonald, R.; Liu, G.; Stahl, S. S. *Chem. Rev.* **2011**, *111*, 2981.
 (e) Zeni, G.; Larock, R. C. *Chem. Rev.* **2006**, *106*, 4644.
 (f) Vlaar, T.; Ruijter, E.; Orru, R. *Adv. Synth. Catal.* **2011**, *353*, 809.
 (g) Giri, R.; Shekhar, K. C. *J. Org. Chem.* **2018**, *83*, 3013.
 (h) Chemler, S. R.; Bovino, M. T. *ACS Catal.* **2013**, *3*, 1076.
- [11] Addition of XCuOH to carbon-carbon multiple bonds, see:
 (a) Zhang, W.; Zhang, J.; Liu, Y.; Xu, Z. *Synlett* **2013**, *24*, 2709.
 (b) Zhang, J.; Wu, D.; Chen, X.; Liu, Y.; Xu, Z. *J. Org. Chem.* **2014**, *79*, 4799.
 (c) Zhang, J.; Wang, H.; Ren, S.; Zhang, W.; Liu, Y. *Org. Lett.* **2015**, *17*, 2920.
 (d) Zhang, J.; Zhang, H.; Shi, D.; Jin, H.; Liu, Y. *Eur. J. Org. Chem.* **2016**, 5545.
 (e) Bao, H.; Xu, Z.; Wu, D.; Zhang, H.; Jin, H.; Liu, Y. *J. Org. Chem.* **2017**, *82*, 109.
- [12] Addition of XCuOH to C=O bonds, see: (f) Zhang, J.; Shi, D.; Zhang, H.; Xu, Z.; Bao, H.; Jin, H.; Liu, Y. *Tetrahedron* **2017**, *73*, 154.
- [13] (a) Takamatsu, K.; Hirano, K.; Satoh, T.; Miura, M. *J. Org. Chem.* **2015**, *80*, 3243.
 (b) Takamatsu, K.; Hirano, K.; Satoh, T.; Miura, M. *Org. Lett.* **2014**, *16*, 2892.
 (c) Zhou, W.; Liu, Y.; Yang, Y.; Deng, G.-J. *Chem. Commun.* **2012**, *48*, 10678.
- [14] (a) Srimani, D.; Leitun, G.; Ben-David, Y.; Milstein, D. *Angew. Chem. Int. Ed.* **2014**, *53*, 11092, and references cited therein.
 (b) Julia, M.; Paris, J.-M. *Tetrahedron Lett.* **1973**, *14*, 4833.
 (c) Blakemore, P. R. *J. Chem. Soc. Perkin Trans* **2002**, 2563.
 (d) Chatterjee, B.; Bera, S.; Mondal, D. *Tetrahedron: Asymmetry* **2014**, *25*, 1.
- [15] Yamanaka, M.; Nishida, A.; Nakagawa, M. *J. Org. Chem.* **2003**, *68*, 3112.
- [16] Qiu, J.; Wang, L.; Liu, M.; Shen, Q.; Tang, J. *Tetrahedron Lett.* **2011**, *52*, 6489.
- [17] Siddiqui, M. A.; Snieckus, V. *Tetrahedron Lett.* **1988**, *29*, 5463.
- [18] Keene, B. R. T. *J. Chem. Soc.* **1965**, 3032.
- [19] Badger, G. M.; Sasse, W. F. H. *J. Chem. Soc.* **1957**, 4.
- [20] Maestri, G.; Larraufie, M.-H.; Derat, E.; Ollivier, C.; Fensterbank, L.; Lacôte, E.; Malacria, M. *Org. Lett.* **2010**, *12*, 5692.
- [21] Coombs, M. M. *J. Chem. Soc.* **1958**, 3454.
- [22] Arcus, C. L.; Coombs, M. M.; Evans, J. V. *J. Chem. Soc.* **1956**, 1498.
- [23] Kessar, S. V.; Gupta, Y. P.; Balakrishnan, P.; Sawal, K. K.; Mohammad, T.; Dutt, M. *J. Org. Chem.* **1988**, *53*, 1708.

(Li, L.; Fan, Y.)