钯催化噻唑定位基导向的单选择性 C(sp²)—H 键碘化反应

邢立浩 邵玲艳* 付小盼 邓克钻 杨金月 冀亚飞*

(华东理工大学药学院 上海 200237)

摘要 发展了一种钯催化, 4-芳基噻唑化合物邻位 C(sp²)—H 键碘化的反应体系. 通过定位基筛选以及反应条件优化得到最高效的单选择性邻位碘化反应条件, 并应用于一系列含不同取代基的 4-(2-碘代芳基)噻唑的合成中. 该催化体系呈现出良好的底物适应性. 通过简单的后续衍生化即可将碘化产物转化为多种基团取代产物, 为该方法学研究提供了更高的应用价值. 通过氘代动力学研究以及自由基抑制实验, 提出了合理的催化循环机理.

关键词 钯催化; 4-芳基噻唑; 单选择性; C(sp²)—H 键; 碘化

Palladium-Catalyzed Thiazole-Directed *mono*-Selective C(sp²)—H Bond Iodination Reaction

Xing, Lihao Shao, Lingyan* Fu, Xiaopan Deng, Kezuan Yang, Jinyue Ji, Yafei*

(School of Pharmacy, East China University of Science & Technology, Shanghai 200237)

Abstract A palladium-catalyzed *ortho*-C(sp²)—H bond iodination of 4-arylthiazoles has been developed. Through screening of directing groups and optimization of reaction parameters, the most efficient reaction conditions for *mono-ortho*-position iodination were obtained, which were applied to synthesize a series of 4-(2-iodoaryl)thiazoles with broad scope of 4-arylthiazole substrates. Furthermore, the iodine group can be easily transformed into other organic functional groups, which improved the application value of this methodology. At last, plausible mechanism was proposed based on an intermolecular deuterium labeling kinetic experiment and radical inhibition experiments.

Keywords palladium-catalyzed; 4-arylthiazoles; *mono*-selectivity; C(sp²)—H bond; iodination

1 Introduction

Aryl iodides represent widespread structural motifs distributed in a range of pharmaceuticals, agrochemicals and natural products.^[1] Over the past few decades, aryl iodides have been extensively used as aromatic synthons for constructing complex organic structures *via* nucleophilic substitution^[2] or a variety of cross-coupling reactions (such as Suzuki,^[3] Heck^[4] and Ullmann reaction,^[5] *etc.*). The traditional strategies to access aryl iodides are direct electrophilic substitution, Sandmeyer reaction and *ortho*-lithiation/halogenation, which were limited in application by poor regioselectivities, harsh reaction conditions and tedious long procedures.^[6]

To overcome these above mentioned disadvantages, diverse C—I bond formation reactions promoted by rear-metal catalysts or more recently investigated base-metal catalysts^[7] have been developed. Ever since the first Pd-catalyzed C(sp²)—H bond iodination reaction disclosed

by Sanford,^[8] a series of protocols using Suárez reagent,^[9] *N*-iodosuccinimide (NIS),^[10] sole I₂^[11] or other iodine sources^[12] have emerged to introduce iodine atom to different aryl substrates. A variety of directing groups have also been employed for *ortho*-C—H bond activation/iodination.^[13] However, the thiazole based auxiliary-directed iodination has rarely been reported. Recently, Petel *et al.*^[14] disclosed Pd-catalyzed benzothiazole-directed C(sp²)—H bond halogenation, in which only four disubstituted examples of iodination were included and the *mono*-selectivity was not obtained (Scheme 1A). Considering that highly functionalized thiazoles serve as important pharmacophores in drug discovery^[15], it is highly desirable to exploit the non-fused thiazole-directed C(sp²)—H bond *mono*-selective iodination reaction (Scheme 1B).

2 Results and discussion

Given our recent success in thiazole-directed C(sp²)—H

* Corresponding authors. E-mail: shaoly520@163.com; jyf@ecust.edu.cn

Received April 25, 2019; revised June 2, 2019; published online June 19, 2019.

Project supported by the National Science Foundation of China (Nos. 21676088, 21476074).

国家自然科学基金(Nos. 21676088, 21476074)资助项目.

Scheme 1 Previous related works and present work

bond activation/acyloxylation,^[16] our research was initiated by briefly evaluating the substituted pattern of various thiazole directing groups. As indicated in Table 1, direct electrophilic iodination preferred to occurring on the thiazole bearing active site^[17] (**DG₁**), which restrained the aimed *ortho*-iodination. Introducing methyl group to C-5 site of thiazole drastically promoted the reaction both on reaction rate and yield (**DG₂**). Replacing methyl group with ethyl group (**DG₃**) performed similar efficiency. However the larger steric hinderance may potentially limit substrate scope, which means **DG₂** is still the suitable directing group. To our surprise, changing carboxylate group on the C-2 site of thiazole to methyl (**DG₄**) notably inhibited the reaction, indicating that carboxylate is vital in this catalytic system. Furthermore, **DG₅** resulted in inferior efficiencies and **DG₆** was inefficient, both demonstrating

Table 1 Screening of directing groups^{a,b}

^a Reaction condition: **1** (0.3 mmol), NIS (135.0 mg, 0.6 mmol), Pd(OAc)₂ (6.7 mg, 10 mol%), DCE/HFIP (V:V=4:1, 3.0 mL), 100 °C, 9 h; ^b Isolated yield; ^c Direct oxidation iodination on thiazole observed, failed to isolate the pure iodination product; ^d Reaction completed within 4 h; ^e Decarboxylation observed; NIS = *N*-Iodosuccinimide; DCE = 1,2-Dichloroethane; HFIP = Hexafluoroisopropanol.

that the appropriate electron density of **DG₂** could promote the C—H bond activation. Thus, the fully substituted **DG₂** was selected as the optimal directing group for further iodination investigations.

Upon the preferable directing group confirmed, the optimization of reaction conditions utilizing **1b** as the model substrate was followed (Table 2). Firstly, a control experiment revealed that the C(sp²)—H bond activation/iodination process does not occur in the absence of the catalyst (Entry 1). Compared to Pd(OAc)₂, other Pd salts led to lower efficiency (Entries 2 vs 3~6). Lowering the loading of Pd(OAc)₂ to 5 mol% distinctly depressed the reaction (Entry 7), howbeit further increasing the dosage of Pd(OAc)₂ did not result in significantly better performance

Table 2 Optimization of the reaction conditions^a

Entry	Catalyst	NIS (equiv.)	Solvent	Yield ^b /%
1	None	2.0	DCE/HFIP (V:V=4:1)	0
2	Pd(OAc) ₂	2.0	DCE/HFIP (V:V=4:1)	76
3	Pd(CH ₃ CN) ₂ Cl ₂	2.0	DCE/HFIP (V:V=4:1)	64
4	Pd(TFA) ₂	2.0	DCE/HFIP (V:V=4:1)	57
5	PdCl ₂	2.0	DCE/HFIP (V:V=4:1)	38
6	Pd(PPh ₃) ₂ Cl ₂	2.0	DCE/HFIP (V:V=4:1)	27
7 ^c	Pd(OAc) ₂	2.0	DCE/HFIP (V:V=4:1)	58
8 ^d	Pd(OAc) ₂	2.0	DCE/HFIP (V:V=4:1)	75
9	Pd(OAc) ₂	2.0	DCE	86
10	Pd(OAc) ₂	2.0	HFIP	69
11	Pd(OAc) ₂	2.0	DMF	85
12	Pd(OAc) ₂	2.0	DCM	62
13	Pd(OAc) ₂	2.0	TCM	21
14	Pd(OAc) ₂	1.5	DCE	76
15	Pd(OAc)₂	2.5	DCE	93
16	Pd(OAc) ₂	3.0	DCE	92
17 ^e	Pd(OAc) ₂	2.5	DCE	78
18 ^f	Pd(OAc) ₂	2.5	DCE	90

^a Reaction conditions: **1b** (74.1 mg, 0.3 mmol), NIS, catalyst (10 mol%), solvent (3 mL, 0.1 mol·L⁻¹), 100 °C, 4 h; ^b NMR yield using CH₂Br₂ as the internal standard; ^c Pd(OAc)₂ (5 mol%); ^d Pd(OAc)₂ (15 mol%); ^e DCE (0.05 mol·L⁻¹); ^f DCE (0.15 mol·L⁻¹); NIS = *N*-Iodosuccinimide; DCE = 1,2-Dichloroethane; HFIP = Hexafluoroisopropanol; DMF = *N,N*-Dimethylformamide; DCM = Dichloromethane; TCM = Trichloromethane; DMSO = Dimethyl sulfoxide.

(Entry 8). Subsequently, screening of solvents, including DCE/HFIP ($V:V=4:1$), DCE, HFIP, DMF, DCM and TCM uncovered that DCE exhibited the best result (Entries 9 vs 2, 10~13). Afterwards, adjusting the amount of NIS showed 2.5 equiv. is more favorable and economic loading (Entries 15 vs 14, 16). Finally, different scales of solvent were applied to this reaction, and $0.1 \text{ mol}\cdot\text{L}^{-1}$ was the appropriate concentration (Entries 15 vs 17~18). Hence, Entry 15 was selected as the standard reaction conditions for the subsequent functionalization.

With the optimized conditions for *mono*-selective $\text{C}(\text{sp}^2)\text{—H}$ bond activation/iodination established, we set about testing the generality of this methodology (Table 3). The iodination reaction with NIS catalyzed by $\text{Pd}(\text{OAc})_2$ was found to be perfectly compatible with a wide range of functional groups, including OMe (**4e**, **4l**, **4u**), Me (**4f**, **4n**, **4t**), Et (**4m**), F (**4c**, **4g**, **4o**, **4v**, **4w**), Cl (**4h**, **4p**), Br (**4i**, **4q**, **4s**) and CF_3 (**4j**, **4r**).

However, most *ortho*-substituted 4-arylthiazoles failed to generate corresponding products, only with the

exception of sterically less hindered F-substituted substrate (**4b**~**4d**). It is speculated that steric hindrance between the *ortho*-substituents and the C-5-methyl group on thiazole ring impedes the free rotation of benzene ring, thus preventing the formation of five-member cyclopalladium intermediate. *meta*-Substituents were well tolerated, and the desired products were obtained with up to 96% yield (**4f**). All iodination took place at the sterically less hindered *ortho*-positions (**4e**~**4j**). Notably, substrate containing NO_2 group performed no react activity (**4k**). This can be attributed to the strong coordination of NO_2 to Pd center, which blocked the subsequent C—H bond activation^[18]. Not surprisingly, di-iodination was not observed when subjecting *para*-substituted substrates to this catalytic system, due to aforementioned steric-hindrance theory.

Mono-iodinated thiophene (**4s**), furan (**4t**) and naphthalene (**4u**) derivatives were successfully obtained in moderate to excellent yields. Furthermore, this protocol was also applicable in the cases of disubstituted masked

Table 3 Scope of 4-phenylthiazoles^{a,b}

^a Reaction conditions: **1** (0.3 mmol), NIS (167.8 mg, 0.75 mmol), $\text{Pd}(\text{OAc})_2$ (6.7 mg, 10 mol%), DCE (3 mL), 100 °C, 4 h; ^b Isolated yields.

4-benzothiazoles (**4v**, **4w**).

To further expand the versatility of this newly developed C—H bond functionalization reaction, iodination product **4a** was successfully converted to different functionalized 4-arylthiazoles through a series of reactions. As demonstrated in Scheme 2, compounds with various functional groups, including aryl, cyano and thiol were conveniently prepared in moderate to good yields.

Reaction conditions: (a) **4a** (111.9 mg, 0.30 mmol), 3-methoxybenzoboric acid (59.3 mg, 0.39 mmol), I_2 (15.2 mg, 0.06 mmol), K_2CO_3 (82.9 mg, 0.60 mmol), PEG, 140 °C, 24 h; (b) **4a** (111.9 mg, 0.30 mmol), DDQ (68.1 mg, 0.30 mmol), $Cu(OAc)_2$ (60.0 mg, 0.30 mmol), Ag_2O (36.9 mg, 0.30 mmol), NMP, 120 °C, 22 h; (c) **4a** (111.9 mg, 0.30 mmol), 3-chlorothiophenol (56.2 mg, 0.39 mmol), CuI (1.4 mg, 2.5 mol%), K_2CO_3 (45.6 mg, 0.33 mmol), 100 °C, 16 h; ^b Isolated yields

Scheme 2 Diversification of the iodination products

This mono-selective iodination reaction was also performed on gram scale in 88% yield (Scheme 3), illustrating that this protocol is still efficient on larger scale in the aspect of selectivity and productivity.

Scheme 3 Gram scale reaction

To gain insight into the mechanism of this catalytic system, a deuterium labeling isotope experiment was carried out to probe the kinetics of Pd-catalyzed $C(sp^2)$ —H iodination (Scheme 4). The intermolecular competition reactions between **3a** and **3a'** with NIS in one vessel performed the kinetic isotopic effect (KIE) value of 3.8, suggesting that the $C(sp^2)$ —H bond cleavage might be involved in the rate-limiting step.

The radical inhibition experiment was then carried out to

Scheme 4 Kinetic isotope effect study

further understand the reaction path way. The use of 2,2,6,6-tetramethylpiperidinoxy (TEMPO) or 2,6-di-*tert*-butyl-4-methylphenol (BHT) as radical scavengers completely restrained the formation of the desired product **4a** (Scheme 5), implying radical may generated in this catalytic system. However, some control experiments showed that NIS reacted with TEMPO and BHT when no substrates were subjected into the reaction system. And for the difficulty of isolating, the products of radical inhibition experiments were failed to be obtained. Consequently, the oxidative addition/reductive elimination path still can not be excluded.

Scheme 5 Radical inhibition experiment

Based on these experimental investigations and the well-documented reports^[10c,10g,14,19], a plausible mechanism was proposed to account for this palladium-catalyzed *mono*-selective iodination reaction (Scheme 6). Firstly, $Pd(OAc)_2$ coordinates with nitrogen atom on thiazole ring to generate cyclopalladium intermediate **A**. Acetate then participates in aromatic proton abstraction to generate Pd^{II} intermediate **B**. There are two possible pathways for the reaction of intermediate **B**. The first pathway proceeds via oxidative addition of NIS to intermediate **B** to form Pd^{IV} intermediate **C**. The subsequent reductive elimination of **C** furnishes the *ortho*-iodinated product **4a** and regenerates the Pd^{II} catalyst. The second pathway b involves the generating of iodine radical, radical attack to intermediate **B** and the forming of Pd^{IV} intermediate **D**, which undergoes reductive elimination to produce **4a** and the Pd^{II} catalyst.

Scheme 6 Plausible reaction mechanism

3 Conclusions

In summary, the first Pd-catalyzed non-fused thiazole-directed C(sp²)—H bond activation/iodination reaction was realized. Various 4-(2-iodoaryl)thiazoles were obtained with excellent ortho- and mono-selectivities with up to 94% yield. The broad substrate scope and the subsequent diverse convenient transformations of iodination products both increase the value of this methodology. A series of control experiments revealed a plausible Pd^{II}/Pd^{IV} mechanism.

4 Experimental section

4.1 General methods

All reactions were carried out in oven-dried glassware and monitored by thin layer chromatography (TLC, pre-coated silica gel plates containing HF254). NMR spectra were determined on a Bruker AV400 in CDCl₃ with TMS as internal standard for ¹H NMR (400 MHz) and ¹³C NMR (100 MHz), respectively. HRMS were measured on a QSTAR Pulsar I LC/TOF MS mass spectrometer or Micromass GCTM gas chromatograph-mass spectrometer.

4.2 General procedure for screening of directing groups

A mixture of substrate **1** (0.3 mmol), Pd(OAc)₂ (6.7 mg, 10 mol%), NIS (135.0 mg, 0.6 mmol) and DCE/HFIP (*V*:*V*=4:1, 3 mL) was stirred at 100 °C for 12 h. Upon completion of the reaction (monitored by TLC), the mixture was transferred into saturated brine (20 mL), then extracted with ethyl acetate (20 mL×3). The combined organic layers were dried over anhydrous MgSO₄ and concentrated *in vacuo* to provide a crude product, which was purified *via* a column chromatography on silica gel (eluent: petroleum ether/ethyl acetate) to supply the desired product **2**, which was identified by ¹H NMR, ¹³C NMR and HRMS.

Ethyl 4-(2-iodophenyl)-5-methylthiazole-2-carboxylate (**2b**): 85 mg, 76% yield. Yellow solid, m.p. 71~72 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.89 (dd, *J*=8.0, 0.9 Hz, 1H), 7.38 (td, *J*=7.5, 1.1 Hz, 1H), 7.30 (dd, *J*=7.6, 1.7 Hz, 1H), 7.07 (td, *J*=7.8, 1.8 Hz, 1H), 4.45 (q, *J*=7.1 Hz, 2H), 2.35 (s, 3H), 1.40 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.3, 156.3, 154.1, 139.2, 139.1, 137.6, 131.1, 130.23, 128.1, 99.6, 62.5, 14.3, 12.8; HRMS (EI) calcd for C₁₃H₁₂¹²⁷INO₂S: 372.9633, found 372.9635.

Ethyl 5-ethyl-4-(2-iodophenyl)thiazole-2-carboxylate (**2c**): 86 mg, 74% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.92 (d, *J*=7.8 Hz, 1H), 7.40 (t, *J*=7.4 Hz, 1H), 7.31 (d, *J*=7.4 Hz, 1H), 7.09 (t, *J*=7.2 Hz, 1H), 4.48 (q, *J*=6.9 Hz, 2H), 2.72 (q, *J*=7.3 Hz, 2H), 1.43 (t, *J*=7.0 Hz, 3H), 1.27 (t, *J*=7.3 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.2, 155.2, 154.0, 145.4, 139.5, 139.0, 131.0, 130.2, 128.0, 99.9, 62.4, 21.0, 16.1, 14.4; HRMS (EI) calcd for C₁₄H₁₄¹²⁷INO₂S: 386.9790, found 386.9791.

4-(2-Iodophenyl)-2,5-dimethylthiazole (**2d**): 9 mg, 9% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.92 (dd, *J*=8.0, 1.0 Hz, 1H), 7.39 (td, *J*=7.4, 1.0 Hz, 1H), 7.31 (dd, *J*=7.6, 1.7 Hz, 1H), 7.06 (td, *J*=7.7, 1.7 Hz, 1H), 2.69 (s, 3H), 2.25 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.9, 153.0, 140.3, 139.1, 131.0, 129.7, 129.3, 128.0, 99.8, 19.2, 12.2; HRMS (EI) calcd for C₁₁H₁₀¹²⁷INS: 314.9579, found 314.9581.

4-(2-iodophenyl)-5-methylthiazole (**2e'**): 33 mg, 36% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 8.60 (s, 1H), 7.88 (dd, *J*=8.0, 1.1 Hz, 1H), 7.34 (td, *J*=7.5, 1.2 Hz, 1H), 7.24 (dd, *J*=7.6, 1.7 Hz, 1H), 7.02 (ddd, *J*=7.9, 7.5, 1.8 Hz, 1H), 2.26 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 154.4, 149.2, 139.9, 139.3, 131.0, 130.0, 129.9, 128.1, 99.6, 12.1; HRMS (EI) calcd for C₁₀H₈¹²⁷INS: 300.9422, found 300.9424.

4.3 General procedure for iodination reaction

A mixture of substrate **3** (0.3 mmol), Pd(OAc)₂ (6.7 mg, 10 mol%), NIS (167.8 mg, 0.75 mmol) and DCE (3 mL)

was stirred at 100 °C for 4 h. Upon completion of the reaction (monitored by TLC), the mixture was transferred into saturated brine (20 mL), then extracted with ethyl acetate (20 mL × 3). The combined organic layers were dried over anhydrous MgSO₄ and concentrated *in vacuo* to provide a crude product, which was purified *via* a column chromatography on silica gel (eluent: petroleum ether/ethyl acetate, *V*: *V*=75:1) to supply the desired product **4**.

Ethyl 4-(2-iodophenyl)-5-methylthiazole-2-carboxylate (**4a**): 97 mg, 87% yield. Yellow solid, m.p. 71~72 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.89 (dd, *J*=8.0, 0.9 Hz, 1H), 7.38 (td, *J*=7.5, 1.1 Hz, 1H), 7.30 (dd, *J*=7.6, 1.7 Hz, 1H), 7.07 (td, *J*=7.8, 1.8 Hz, 1H), 4.45 (q, *J*=7.1 Hz, 2H), 2.35 (s, 3H), 1.40 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.3, 156.3, 154.1, 139.2, 139.1, 137.6, 131.1, 130.23, 128.1, 99.6, 62.5, 14.3, 12.8; HRMS (EI) calcd for C₁₃H₁₂¹²⁷INO₂S: 372.9633, found 372.9635.

Ethyl 4-(2-fluoro-6-iodophenyl)-5-methylthiazole-2-carboxylate (**4c**): 43 mg, 37% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.73 (dd, *J*=6.6, 2.4 Hz, 1H), 7.18~7.09 (m, 2H), 4.48 (q, *J*=7.1 Hz, 2H), 2.37 (s, 3H), 1.43 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.2 (t, ¹*J*_{CF}=126.0 Hz), 154.8, 150.2, 139.3, 134.7 (d, ⁴*J*_{CF}=3.6 Hz), 131.9 (d, ³*J*_{CF}=8.7 Hz), 130.5 (d, ³*J*_{CF}=8.2 Hz), 127.3 (d, ²*J*_{CF}=18.4 Hz), 124.2 (d, ⁴*J*_{CF}=3.6 Hz), 115.7 (d, ²*J*_{CF}=22.5 Hz), 62.5, 14.3, 12.3; HRMS (EI) calcd for C₁₃H₁₁F¹²⁷INO₂S: 390.9539, found 390.9543.

Ethyl 4-(2-iodo-5-methoxyphenyl)-5-methylthiazole-2-carboxylate (**4e**): 88 mg, 73% yield. Yellow solid, m.p. 101~103 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.76 (d, *J*=8.8 Hz, 1H), 6.90 (d, *J*=3.0 Hz, 1H), 6.71 (dd, *J*=8.8, 3.0 Hz, 1H), 4.48 (q, *J*=7.1 Hz, 2H), 3.79 (s, 3H), 2.39 (s, 3H), 1.43 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.2, 159.7, 156.2, 154.1, 140.0, 139.6 (2C), 137.6, 117.0, 116.7, 88.0, 62.5, 55.5, 12.8; HRMS (EI) calcd for C₁₄H₁₄¹²⁷INO₃S: 402.9739, found 402.9740.

Ethyl 4-(2-iodo-5-methylphenyl)-5-methylthiazole-2-carboxylate (**4f**): 111 mg, 96% yield. White solid, m.p. 81~82 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.73 (d, *J*=8.1 Hz, 1H), 7.13 (d, *J*=1.8 Hz, 1H), 6.88 (dd, *J*=8.1, 1.8 Hz, 1H), 4.43 (q, *J*=7.1 Hz, 2H), 2.35 (s, 3H), 2.27 (s, 3H), 1.39 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 159.2, 155.4, 152.9, 137.8, 137.7, 137.1, 136.4, 131.0, 130.2, 94.3, 61.4, 19.8, 13.3, 11.8; HRMS (EI) calcd for C₁₄H₁₄¹²⁷INO₂S: 386.9790, found 386.9789.

Ethyl 4-(5-fluoro-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4g**): 89 mg, 76% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.86 (dd, *J*=8.7, 5.5 Hz, 1H), 7.10 (dd, *J*=8.8, 3.0 Hz, 1H), 6.88 (td, *J*=8.5, 3.0 Hz, 1H), 4.48 (q, *J*=7.1 Hz, 2H), 2.40 (s, 3H), 1.43 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.62 (d, ¹*J*_{CF}=249.3 Hz), 160.1, 155.1, 154.5, 141.0 (d, ³*J*_{CF}=7.9 Hz), 140.4 (d, ³*J*_{CF}=7.9 Hz), 138.0, 118.6 (d, ²*J*_{CF}=22.5 Hz), 117.8 (d, ²*J*_{CF}=21.7 Hz), 92.6 (d, ⁴*J*_{CF}=3.4 Hz), 62.6, 14.3, 12.8; HRMS (EI) calcd for C₁₃H₁₁F¹²⁷INO₂S: 390.9539, found 390.9543.

Ethyl 4-(5-chloro-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4h**): 100 mg, 82% yield. Yellow solid, m.p. 69~71 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.83 (d, *J*=8.5 Hz, 1H), 7.35 (d, *J*=2.5 Hz, 1H), 7.11 (dd, *J*=8.5, 2.5 Hz, 1H), 4.48 (q, *J*=7.1 Hz, 2H), 2.40 (s, 3H), 1.43 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.0, 154.9, 154.5, 140.8, 140.1, 138.0, 134.4, 131.2, 130.5, 96.7, 62.6, 14.3, 12.8; HRMS (EI) calcd for C₁₃H₁₁Cl¹²⁷INO₂S: 406.9244, found 406.9241.

Ethyl 4-(5-bromo-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4i**): 128 mg, 94% yield. Yellow solid, m.p. 87~89 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.77 (d, *J*=8.5 Hz, 1H), 7.49 (d, *J*=2.4 Hz, 1H), 7.24 (dd, *J*=8.5, 2.4 Hz, 1H), 4.48 (q, *J*=7.1 Hz, 2H), 2.40 (s, 3H), 1.43 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.0, 154.8, 154.5, 141.1, 140.3, 138.1, 134.0, 133.4, 122.3, 97.6, 62.6, 14.3, 12.9; HRMS (EI) calcd for C₁₃H₁₁Br¹²⁷INO₂S: 452.8718, found 452.8721.

Ethyl 4-(2-iodo-5-(trifluoromethyl)phenyl)-5-methylthiazole-2-carboxylate (**4j**): 85 mg, 64% yield. Yellow solid, m.p. 70~72 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.1 (d, *J*=8.3 Hz, 1H), 7.6 (s, 1H), 7.4 (d, *J*=7.2 Hz, 1H), 4.5 (q, *J*=7.1 Hz, 2H), 2.4 (s, 3H), 1.4 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.0, 154.8, 154.7, 140.2, 139.8, 138.2, 130.8 (q, ²*J*_{CF}=33.2 Hz), 127.8 (q, ³*J*_{CF}=3.6 Hz), 126.7 (dd, ³*J*_{CF}=7.4, 3.6 Hz), 126.4 (q, ¹*J*_{CF}=270.6 Hz), 104.1 (d, ⁴*J*_{CF}=1.3 Hz), 62.7, 14.3, 12.8; HRMS (EI) calcd for C₁₄H₁₁F₃¹²⁷INO₂S: 440.9507, found 440.9509.

Ethyl 4-(2-iodo-4-methoxyphenyl)-5-methylthiazole-2-carboxylate (**4l**): 52 mg, 43% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.37 (d, *J*=2.6 Hz, 1H), 7.15 (d, *J*=8.5 Hz, 1H), 6.87 (dd, *J*=8.5, 2.6 Hz, 1H), 4.40 (q, *J*=7.1 Hz, 2H), 3.75 (s, 3H), 2.30 (s, 3H), 1.35 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 159.3, 158.9, 155.2, 152.7, 136.7, 130.4, 130.4, 123.1, 113.1, 98.7, 61.4, 54.6, 13.3, 11.8; HRMS (EI) calcd for C₁₄H₁₄¹²⁷INO₃S: 402.9739, found 402.9740.

Ethyl 4-(4-ethyl-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4m**): 88 mg, 73% yield. Yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.76 (s, 1H), 7.23 (d, *J*=0.9 Hz, 2H), 4.47 (q, *J*=7.1 Hz, 2H), 2.64 (q, *J*=7.6 Hz, 2H), 2.38 (s, 3H), 1.42 (t, *J*=7.1 Hz, 3H), 1.24 (t, *J*=7.6 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.2, 156.4, 153.8, 146.7, 138.4, 137.5, 136.4, 130.8, 127.7, 99.6, 62.4, 28.1, 15.4, 14.3, 12.8; HRMS (EI) calcd for C₁₅H₁₆¹²⁷INO₂S: 400.9946, found 400.9950.

Ethyl 4-(2-iodo-4-methylphenyl)-5-methylthiazole-2-carboxylate (**4n**): 93 mg, 79% yield. Yellow solid, m.p. 70~72 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.69 (s, 1H), 7.13 (s, 2H), 4.40 (q, *J*=7.1 Hz, 2H), 2.30 (s, 3H), 2.28 (s, 3H), 1.35 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.3, 156.4, 153.9, 140.4, 139.5, 137.5, 136.2, 130.7, 128.9, 99.5, 62.4, 20.7, 14.3, 12.8; HRMS (EI) calcd for C₁₃H₁₁F¹²⁷INO₂S: 390.9539, found 390.9540.

Ethyl 4-(4-fluoro-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4o**): 99 mg, 84% yield. Yellow solid, m.p. 65~67 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.65 (dd, *J*=

8.1, 2.6 Hz, 1H), 7.31 (dd, $J=9.2, 5.2$ Hz, 1H), 7.14 (td, $J=8.3, 2.6$ Hz, 1H), 4.48 (q, $J=7.1$ Hz, 2H), 2.38 (s, 3H), 1.43 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.0 (d, $^1J_{\text{CF}}=253.4$ Hz), 160.1, 155.3, 154.1, 137.9, 135.4 (d, $^4J_{\text{CF}}=3.6$ Hz), 131.9 (d, $^3J_{\text{CF}}=8.4$ Hz), 126.0 (d, $^2J_{\text{CF}}=23.8$ Hz), 115.3 (d, $^2J_{\text{CF}}=21.2$ Hz), 99.2 (d, $^3J_{\text{CF}}=8.2$ Hz), 62.5, 14.3, 12.7; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{11}\text{F}^{127}\text{INO}_2\text{S}$: 390.9539, found 390.9540.

Ethyl 4-(4-chloro-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4p**): 79 mg, 65% yield. Yellow solid, m.p. 93~95 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.90 (d, $J=2.1$ Hz, 1H), 7.37 (dd, $J=8.2, 2.1$ Hz, 1H), 7.23 (d, $J=8.2$ Hz, 1H), 4.45 (q, $J=7.1$ Hz, 2H), 2.35 (s, 3H), 1.40 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.1, 155.1, 154.3, 138.4, 137.4, 137.9, 135.1, 131.7, 128.4, 99.6, 62.5, 14.3, 12.8; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{11}\text{Cl}^{127}\text{INO}_2\text{S}$: 406.9244, found 406.9243.

Ethyl 4-(4-bromo-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4q**): 103 mg, 76% yield. Yellow solid, m.p. 103~104 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.08 (d, $J=1.9$ Hz, 1H), 7.55 (dd, $J=8.2, 1.9$ Hz, 1H), 7.20 (d, $J=8.2$ Hz, 1H), 4.47 (d, $J=7.1$ Hz, 2H), 2.38 (s, 3H), 1.43 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.0, 155.1, 154.3, 141.1, 138.2, 137.9, 132.0, 131.3, 123.2, 100.1, 62.5, 14.3, 12.8; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{11}\text{Br}^{127}\text{INO}_2\text{S}$: 450.8739, found 452.8720.

Ethyl 4-(2-iodo-4-(trifluoromethyl)phenyl)-5-methylthiazole-2-carboxylate (**4r**): 99 mg, 75% yield. Yellow solid, m.p. 80~82 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.68 (dd, $J=8.0, 1.0$ Hz, 1H), 7.46 (d, $J=7.9$ Hz, 1H), 4.48 (q, $J=7.1$ Hz, 2H), 2.40 (s, 3H), 1.44 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.0, 154.8 (d, $^3J_{\text{CF}}=14.9$ Hz), 143.0, 138.0, 135.9 (q, $^4J_{\text{CF}}=3.8$ Hz), 132.2 (q, $^2J_{\text{CF}}=33.0$ Hz), 131.4, 129.1, 125.0 (q, $^4J_{\text{CF}}=3.6$ Hz), 123.9 (q, $^1J_{\text{CF}}=271.3$ Hz), 99.4, 62.6, 14.3, 12.7; HRMS (EI) calcd for $\text{C}_{14}\text{H}_{11}\text{F}_3^{127}\text{INO}_2\text{S}$: 440.9507, found 440.9509.

Ethyl 4-(5-bromo-3-iodothiophen-2-yl)-5-methylthiazole-2-carboxylate (**4s**): 77 mg, 56% yield. Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.23 (s, 1H), 4.47 (q, $J=7.1$ Hz, 2H), 2.51 (s, 3H), 1.43 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.8, 154.7, 146.4, 140.6, 137.2, 136.8, 115.3, 82.0, 62.7, 14.3, 13.5; HRMS (EI) calcd for $\text{C}_{11}\text{H}_9\text{Br}^{127}\text{INO}_2\text{S}_2$: 458.8282, found 458.8285.

Ethyl 4-(3-iodo-5-methylfuran-2-yl)-5-methylthiazole-2-carboxylate (**4t**): 70 mg, 62% yield. White solid, m.p. 92~94 °C; ^1H NMR (400 MHz, CDCl_3) δ : 6.21 (s, 1H), 4.46 (q, $J=7.1$ Hz, 2H), 2.63 (s, 3H), 2.35 (s, 3H), 1.43 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.0, 154.6, 154.1, 146.7, 143.9, 139.2, 115.2, 66.8, 62.5, 14.3, 13.6, 13.6; HRMS (EI) calcd for $\text{C}_{12}\text{H}_{12}\text{I}^{127}\text{INO}_3\text{S}$: 376.9583, found 372.9584.

Ethyl 4-(3,5-diiodo-6-methoxynaphthalen-2-yl)-5-methylthiazole-2-carboxylate (**4u**): 154 mg, 89% yield. Yellow solid, m.p. 123~125 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.72 (s, 1H), 7.73 (d, $J=8.9$ Hz, 1H), 7.70 (s, 1H), 7.19 (d, $J=8.8$ Hz, 1H), 4.46 (q, $J=7.0$ Hz, 2H), 4.00 (s, 3H),

2.37 (s, 3H), 1.41 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.2, 157.6, 155.8, 154.0, 141.6, 138.2, 136.9, 134.2, 130.6, 130.3, 128.7, 113.8, 99.8, 85.4, 62.5, 57.2, 14.4, 13.0; HRMS (EI) calcd for $\text{C}_{18}\text{H}_{15}\text{I}_2^{127}\text{NO}_3\text{S}$: 578.8862, found 578.8861.

Ethyl 4-(2,4-difluoro-6-iodophenyl)-5-methylthiazole-2-carboxylate (**4v**): 48 mg, 39% yield. White solid, m.p. 62~64 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.23 (q, 1H), 7.17~7.09 (m, 1H), 4.48 (q, $J=7.1$ Hz, 2H), 2.37 (s, 3H), 1.43 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.1, 154.5, 154.3 (d, $^3J_{\text{CF}}=1.9$ Hz), 151.4 (dd, $^1,^3J_{\text{CF}}=245.0, 13.7$ Hz), 149.7 (dd, $^1,^3J_{\text{CF}}=253.4, 15.1$ Hz), 138.1, 136.3 (d, $^3J_{\text{CF}}=4.0$ Hz), 126.6 (dd, $^2,^2J_{\text{CF}}=6.5, 3.7$ Hz), 117.1 (d, $^2J_{\text{CF}}=18.0$ Hz), 88.9 (d, $^2J_{\text{CF}}=21.7$ Hz), 62.6, 14.3, 12.7; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{10}\text{F}_2^{127}\text{INO}_2\text{S}$: 408.9445, found 408.9446.

Ethyl 4-(3,5-difluoro-2-iodophenyl)-5-methylthiazole-2-carboxylate (**4w**): 90 mg, 73% yield. White solid, m.p. 71~73 °C; ^1H NMR (400 MHz, CDCl_3) δ : 6.96 (d, $J=8.4$ Hz, 1H), 6.90 (td, $J=8.3, 2.6$ Hz, 1H), 4.47 (q, $J=7.1$ Hz, 2H), 2.38 (s, 3H), 1.42 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.9 (dd, $^1,^3J_{\text{CF}}=249.6, 12.2$ Hz), 162.1 (dd, $^1,^3J_{\text{CF}}=246.2, 12.6$ Hz), 160.0, 154.7, 153.8, 142.1 (dd, $^3,^3J_{\text{CF}}=9.4, 3.1$ Hz), 138.1, 114.4 (dd, $^2,^4J_{\text{CF}}=22.3, 3.4$ Hz), 104.3 (dd, $^2,^2J_{\text{CF}}=28.5, 25.8$ Hz), 81.2 (dd, $^2,^4J_{\text{CF}}=25.9, 4.3$ Hz), 62.6, 14.3, 12.7; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{10}\text{F}_2^{127}\text{INO}_2\text{S}$: 408.9445, found 408.9448.

4.4 Procedures for derivatizations of iodinated products

A mixture of **4a** (111.9 mg, 0.3 mmol), 3-boronoanisole (59.3 mg, 0.39 mmol), I_2 (82.9 mg, 0.06 mmol), K_2CO_3 (82.9 mg, 0.06 mmol) and PEG (3 mL) was stirred in a sealed tube at 140 °C for 24 h. Upon completion of the reaction (as monitored by TLC), the mixture was transferred into saturated brine (20 mL) then extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were dried over anhydrous MgSO_4 and concentrated *in vacuo* to provide a crude product, which was purified *via* a column chromatography on silica gel (eluent: petroleum ether/ethyl acetate, $V:V=100:1$) to supply the desired product 4-(3'-methoxy-[1,1'-biphenyl]-2-yl)-5-methylthiazole (**5a**), 62 mg, 73% yield. Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.61 (s, 1H), 7.54~7.39 (m, 4H), 7.16 (t, $J=7.9$ Hz, 1H), 6.83~6.74 (m, 2H), 6.63 (dd, $J=2.4, 1.7$ Hz, 1H), 3.64 (s, 3H), 1.88 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.1, 152.0, 149.1, 142.7, 141.7, 133.2, 131.1, 130.0, 129.3, 129.1, 128.7, 127.4, 121.5, 114.1, 113.0, 55.1, 11.6; HRMS (EI) calcd for $\text{C}_{17}\text{H}_{15}\text{NOS}$ 281.0874, found 281.0872.

A mixture of **4a** (111.9 mg, 0.3 mmol), DDQ (68.1 mg, 0.3 mmol), $\text{Cu}(\text{OAc})_2$ (60 mg, 0.3 mmol), AgO (36.9 mg, 0.3 mmol) and NMP (3 mL) was stirred in a sealed tube at 120 °C for 22 h. Upon completion of the reaction (as monitored by TLC), the mixture was transferred into saturated brine (20 mL), then extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were dried over

anhydrous MgSO_4 and concentrated *in vacuo* to provide a crude product, which was purified *via* a column chromatography on silica gel (eluents: petroleum ether/ethyl acetate, $V:V=30:1$) to supply the desired product ethyl 4-(2-cyanophenyl)-5-methylthiazole-2-carboxylate (**5b**), 63 mg, 77% yield. White solid, m.p. 100~101 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.73 (dd, $J=7.8, 0.9$ Hz, 1H), 7.64 (td, $J=7.7, 1.3$ Hz, 1H), 7.57 (dd, $J=7.8, 0.9$ Hz, 1H), 7.48 (td, $J=7.6, 1.3$ Hz, 1H), 4.44 (q, $J=7.1$ Hz, 2H), 2.50 (s, 3H), 1.39 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.9, 155.1, 150.7, 138.9, 137.6, 133.1, 132.8, 131.3, 129.0, 117.8, 113.0, 62.6, 14.3, 12.9; HRMS (EI) calcd for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$: 272.0619, found 272.0620.

A mixture of **4a** (111.9 mg, 0.3 mmol), 3-chlorothiophenol (56.2 mg, 0.39 mmol), CuI (1.4 mg, 0.0075 mmol), K_2CO_3 (45.6 mg, 0.33 mmol) and NMP (3 mL) was stirred in a sealed tube at 100 °C for 16 h. Upon completion of the reaction (as monitored by TLC), the mixture was transferred into saturated brine (20 mL), then extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were dried over anhydrous MgSO_4 and concentrated *in vacuo* to provide a crude product, which was purified *via* a column chromatography on silica gel (eluents: petroleum ether/ethyl acetate, $V:V=100:1$) to supply the desired product 5-methyl-4-(2-(phenylthio)phenyl)thiazole (**5c**), 52 mg, 55% yield. Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.65 (s, 1H), 7.34 (dd, $J=5.9, 3.3$ Hz, 1H), 7.29 (dd, $J=6.4, 2.9$ Hz, 3H), 7.25~7.17 (m, 4H), 2.34 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 151.1, 149.3, 136.9, 135.4, 133.7, 133.5, 133.4 (2C), 131.3, 131.0, 130.4, 129.3 (2C), 129.1, 126.9, 12.1; HRMS (EI) calcd for $\text{C}_{16}\text{H}_{12}\text{S}_2$: 319.0070, found 319.0072.

References

- [1] Gribble, G. W. *Acc. Chem. Res.* **1998**, *31*, 141.
- [2] Lindley, J. *Tetrahedron* **1984**, *40*, 1433.
- [3] Suzuki, A. *J. Organomet. Chem.* **1999**, *576*, 147.
- [4] Crisp, T. G. *Chem. Soc. Rev.* **1998**, *27*, 427.
- [5] Sambriago, C.; Marsden, S. P.; Blacker, A. J.; McGowan, P. C. *Chem. Soc. Rev.* **2014**, *43*, 3525.
- [6] Merkushev, E. B. *Synthesis* **1988**, 923.
- [7] (a) Li, B.; Liu, B.; Shi, B. F. *Chem. Commun.* **2015**, *51*, 5093.
(b) Pal, P.; Singh, H.; Panda, A. B.; Ghosh, S. C. *Asian J. Org. Chem.* **2015**, *4*, 879.
(c) Zhan, B. B.; Liu, Y. H.; Hu, F.; Shi, B. F. *Chem. Commun.* **2016**, *52*, 4934.
(d) Aihara, Y.; Chatani, N. *ACS Catal.* **2016**, *6*, 4323.
(e) Khan, B.; Kant, R.; Koley, D. *Adv. Synth. Catal.* **2016**, *358*, 2352.
(f) Kommagalla, Y.; Yamazaki, K.; Yamaguchi, T.; Chatani, N. *Chem. Commun.* **2018**, *54*, 1359.
(g) Singh, H.; Sen, C.; Sahoo, T.; Ghosh, S. C. *Eur. J. Org. Chem.* **2018**, *34*, 4748.
(h) Du, Y.; Liu, Y. Y.; Wan, J. P. *J. Org. Chem.* **2018**, *83*, 3403.
- [8] Kalyani, D.; Dick, A. R.; Anani, W. Q.; Sanford, M. S. *Org. Lett.* **2006**, *8*, 2523.
- [9] (a) Giri, R.; Chen, X.; Yu, J. Q. *Angew. Chem., Int. Ed.* **2005**, *44*, 2112.
(b) Li, J. J.; Mei, T. S.; Yu, J. Q. *Angew. Chem., Int. Ed.* **2008**, *47*, 6452.
(c) Mei, T. S.; Giri, R.; Mangel, N.; Yu, J. Q. *Angew. Chem., Int. Ed.* **2008**, *47*, 5215.
(d) Mei, T. S.; Wang, D. H.; Yu, J. Q. *Org. Lett.* **2010**, *12*, 3140.
(e) Nack, W. A.; Wang, X.; Wang, B.; He, G.; Chen, G. *Beilstein J. Org. Chem.* **2016**, *12*, 1243.
(f) Zhu, R. Y.; Liu, L. Y.; Yu, J. Q. *J. Am. Chem. Soc.* **2017**, *139*, 12394.
(g) Zhu, R. Y.; Saint-Denis, T. G.; Shao, Y.; He, J.; Sieber, J. D.; Senanayake, C. H.; Yu, J. Q. *J. Am. Chem. Soc.* **2017**, *139*, 5724.
- [10] (a) Dudnik, A. S.; Chernyak, N.; Huang, C.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2010**, *49*, 8729.
(b) Du, B.; Jiang, X.; Sun, P. J. *Org. Chem.* **2013**, *78*, 2786.
(c) Sadhu, P.; Alla, S. K.; Punniyamurthy, T. *J. Org. Chem.* **2013**, *78*, 6104.
(d) Pascanu, V.; Carson, F.; Solano, M. V.; Su, J.; Zou, X.; Johansson, M. J.; Martin-Matute, B. *Chem.-Eur. J.* **2016**, *22*, 3729.
(e) Testa, C.; Gigot, E.; Genc, S.; Decreau, R.; Roger, J.; Hierso, J. C. *Angew. Chem., Int. Ed.* **2016**, *55*, 5555.
(f) Yang, X.; Sun, Y.; Sun, T. Y.; Rao, Y. *Chem. Commun.* **2016**, *52*, 6423.
(g) Das, R.; Kapur, M. *J. Org. Chem.* **2017**, *82*, 1114.
(h) Dubost, E.; Babin, V.; Benoist, F.; Hebert, A.; Barbey, P.; Chollet, C.; Bouillon, J. P.; Manrique, A.; Pieters, G.; Fabis, F.; Cailly, T. *Org. Lett.* **2018**, *20*, 6302.
(i) Tang, R. J.; Milcent, T.; Crousse, B. *J. Org. Chem.* **2018**, *83*, 930.
- [11] (a) Chu, L.; Wang, X. C.; Moore, C. E.; Rheingold, A. L.; Yu, J. Q. *J. Am. Chem. Soc.* **2013**, *135*, 16344.
(b) Wang, X. C.; Hu, Y.; Bonacorsi, S.; Hong, Y.; Burrell, R.; Yu, J. Q. *J. Am. Chem. Soc.* **2013**, *135*, 10326.
(c) Chu, L.; Xiao, K. J.; Yu, J. Q. *Science* **2014**, *346*, 451.
- [12] (a) Lu, C.; Zhang, S. Y.; He, G.; Nack, W. A.; Chen, G. *Tetrahedron* **2014**, *70*, 4197.
(b) Chu, L.; Shang, M.; Tanaka, K.; Chen, Q.; Pissarnitski, N.; Streckfuss, E.; Yu, J. Q. *ACS Cent. Sci.* **2015**, *1*, 394.
(c) Sun, X.; Yao, X.; Zhang, C.; Rao, Y. *Chem. Commun.* **2015**, *51*, 10014.
(d) Fan, X. M.; Guo, Y.; Li, Y. D.; Yu, K. K.; Liu, H. W.; Liao, D. H.; Ji, Y. F. *Asian J. Org. Chem.* **2016**, *5*, 499.
- [13] (a) Das, R.; Kapur, M. *Asian J. Org. Chem.* **2018**, *7*, 1524.
(b) Liao, G.; Shi, B. F. *Acta Chim. Sinica* **2015**, *73*, 1283 (in Chinese).
(廖港, 史炳烽, 化学学报, **2015**, *73*, 1283.)
- [14] Santra, S. K.; Banerjee, A.; Khatun, N.; Samanta, A.; Patel, B. K. *RSC Adv.* **2015**, *5*, 11960.
- [15] (a) Al-Ghorbani, M.; Alghamdi, H. A.; Khanum, S. A. *Eur. J. Bio-med. Pharm. Sci.* **2018**, *5*, 1.
(b) Attri, C.; Bhatia, P.; Kumar, P. *J. Mod. Chem. Chem. Technol.* **2018**, *9*, 19.
(c) Liaras, K.; Fesatidou, M.; Geronikaki, A. *Molecules* **2018**, *23*, 6851.
(d) de Siqueira, L. R. P.; de Moraes Gomes, P. A. T.; de Lima Ferreira, L. P.; de Melo Rego, M. J. B.; Leite, A. C. L. *Eur. J. Med. Chem.* **2019**, *170*, 237.
(e) Scarim, C. B.; Jornada, D. H.; Machado, M. G. M.; Ferreira, C. M. R.; dos Santos, J. L.; Chung, M. C. *Eur. J. Med. Chem.* **2019**, *162*, 378.
- [16] Yu, K. K.; Guo, Y.; Hu, Y. H.; Xu, Z.; Liu, H. W.; Liao, D. H.; Ji, Y. F. *Asian J. Org. Chem.* **2016**, *5*, 1219.
- [17] Bergström, M.; Suresh, G.; Naidu, V. R.; Unelius, C. R. *Eur. J. Org. Chem.* **2017**, *2017*, 3234.
- [18] Qiao, H. J.; Yang, F.; Wang, S. W.; Leng, Y. T.; Wu, Y. J. *Tetrahedron* **2015**, *71*, 9258.
- [19] Qiu, F. C.; Yang, W. C.; Chang, Y. Z.; Guan, B. T. *Asian J. Org. Chem.* **2017**, *6*, 1361.

(Li, L.; Fan, Y.)