微波辅助下通过 Fe/O₂ 催化 C—H 键官能团化合成
喹啉-2-甲醛类化合物谢庭辉^{a,b} 蒋筱莹^a 米治胜^a 李 雪^a 徐小河^a 白仁仁^a
帅 棋^a 谢媛媛^{*,a}^(a) 浙江工业大学药学院 长三角绿色制药协同创新中心 杭州 310014)^(b) 台州科技职业学院 浙江台州 318020)

摘要 在微波辅助下成功开发了一种由 Fe³⁺/O₂ 催化一锅法合成喹啉-2-甲醛类化合物的方法。以 2-甲基喹啉为原料, O₂ 为醛基中氧的合成子, 得到一系列底物适应性较广的喹啉-2-甲醛类化合物, 产率 48%~80%。初步机理研究表明, 该反应可能经历了一个自由基历程。此方法具有操作简便、反应时间短、选择性好等优点。

关键词 喹啉-2-甲醛; 铁催化; C—H 键氧化; 微波辐射

Iron/O₂-Promoted C—H Bond Functionalization for the Exclusive
Synthesis of 2-Quinoline Carboxaldehydes under
Microwave IrradiationXie, Tinghui^{a,b} Jiang, Xiaoying^a Mi, Zhisheng^a Li, Xue^a Xu, Xiaohe^a
Bai, Renren^a Shuai, Qi^a Xie, Yuanyuan^{*,a}^(a) Collaborative Innovation Center of Yangtze River Delta Region Green Pharmaceutical, College of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014)^(b) College of Agriculture and Bioengineering, Taizhou Vocational College of Science and Technology, Taizhou, Zhejiang 318020)

Abstract An one-pot iron-catalyzed oxidative formylation of 2-methylquinolines to produce 2-quinoline carboxaldehydes under microwave irradiation has been achieved by employing O₂ as the oxygen donor. The reaction was general for the substrates with a wide range of functional groups, providing a yield of 48%~80%. The preliminary mechanistic studies revealed that the reaction underwent a radical pathway. Advantages of this method include the easy operation, short reaction time and good selectivity.

Keywords 2-quinoline carboxaldehydes; iron catalysis; C—H oxidation; microwave irradiation

1 Introduction

2-Methylquinoline is an important scaffold, which is widely used in organic synthesis. With its active methyl, it could be not only prefunctionalized in short reaction steps with high atom economy,^[1] but also represents a potential synthon for the access of functionalized compounds, such as aldehydes,^[2] nitriles^[3] as well as halomethylation products.^[4] 2-Quinoline carboxaldehydes commonly used as pharmaceutical intermediates, have a profound influence

on the biological activities, such as antineoplastic and antimalarial activities of the following drugs^[5] (Figure 1).

The preparation of 2-quinoline carboxaldehydes has been extensively studied and reviewed.^[6] Holzapfel and co-workers^[7] reported the direct formylation of aryl halides into 2-quinoline carboxaldehydes in the presence of carbon monoxide and hydrogen using palladium as a catalyst. Ding *et al.*^[8] described a method to synthesize 2-quinoline carboxaldehydes by treating 2-methylquinolines with SeO₂ at 170 °C. Moreover, trioxane-promoted formylation of

* Corresponding author. E-mail: xyycz@zjut.edu.cn

Received March 4, 2019; revised April 25, 2019; published online July 9, 2019.

Project supported by the National Natural Science Foundation of China (No. 21576239).

国家自然科学基金(No. 20150281)资助项目.

Figure 1 Compounds derived from 2-quinoline carboxaldehydes

quinolines was established for the synthesis of 2-quinoline carboxaldehydes by Minisci *et al.*^[9] Wang *et al.*^[10] found that 2-methylquinolines could be oxidized in hours by O₂ in the presence of copper catalyst, generating the corresponding 2-quinoline carboxaldehydes. However, the generality of these methods was severely hampered by their drawbacks such as high pressure, high toxicity, limited environmental acceptability in view of the demand of industry and long reaction time.

Over the past decade, ferric salts have attracted considerable attention as an inexpensive and commercially available substance to promote a diverse number of reactions.^[11] In most of these reports, Fe(NO₃)₃•9H₂O was demonstrated as a good catalyst or oxidant exhibiting high activity in oxidative process.^[12] In light of the recently developed applications of 2-quinoline carboxaldehydes, herein, we report the iron(III)/O₂-promoted oxidative formylation of 2-methyl quinolines, yielding various 2-quinoline carboxaldehydes under microwave irradiation through a radical mechanism. This protocol has a lot of advantages, such as short reaction time, easy operation, non-toxicity and good selectivity.

2 Results and discussion

As an initial test, 2-methylquinoline was chosen as the model substrate in order to ascertain the feasibility and optimize the reaction conditions (Table 1).

Iron(III) was found to be crucial for this reaction, while iron(II) failed to give any desired product (Entry 2 vs Entry 1). Notably, other kind of iron(III) also succeeded in producing the expected product, and Fe(NO₃)₃•9H₂O turned out to be the optimal to afford **2a** in 65% yield (Entries 3 and 4). Changing the ratio of iron(III) to **1a** led to a decrease in reaction yield (Entries 4~6). The effect of solvent on the reaction was also examined. The reaction was completed in 10 mins with a good yield of 65% in dimethyl sulfoxide (DMSO). Other solvents such as *N,N*-dimethylformamide (DMF), toluene, dioxane, and *N,N*-dimethylacetamide (DMAC) gave a lower yield of the product with longer reaction time (Entries 4, 7~10). The temperature of this conversion was also screened, 150 °C was found to be the optimal temperature, producing **2a** in 78% yield in 5 min (Entries 4, 11~13). The materials decomposed at a higher temperature. The results obtained by changing reaction time were similar to those obtained by changing reaction temperature, 5 min being the best choice for this transformation (Entries 12, 14~15). Other trials, such as performing the reaction under N₂ atmosphere, decrease the yield compared with that under air (Entries 16 and 17), indicating that O₂ was essential for the present conversion.

Based on the above results, the optimized conditions for the conversion of 2-methylquinolines (1.0 mmol) into 2-quinoline carboxaldehydes were as follows: iron nitrate

Table 1 Optimization of the reaction conditions^a

Entry	Promoter	Fe ³⁺ / 1a (molar ratio)	Solvent	Temp./°C	Time ^b /min	Yield ^c %
1	FeCl ₂	1/1	DMSO	130	30	Trace
2	FeCl ₃	1/1	DMSO	130	10	60
3	FeCl ₃ •6H ₂ O	1/1	DMSO	130	10	62
4	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	130	10	65
5	Fe(NO ₃) ₃ •9H ₂ O	0.5/1	DMSO	130	10	46
6	Fe(NO ₃) ₃ •9H ₂ O	1.5/1	DMSO	130	10	58
7	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMF	130	15	61
8	Fe(NO ₃) ₃ •9H ₂ O	1/1	Toluene	130	30	53
9	Fe(NO ₃) ₃ •9H ₂ O	1/1	Dioxane	130	30	48
10	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMAC	130	15	60
11	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	140	8	70
12	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	150	5	78
13	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	160	5	71
14	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	150	1	37
15	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	150	10	76
16 ^d	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	150	30	45
17 ^e	Fe(NO ₃) ₃ •9H ₂ O	1/1	DMSO	150	30	18

^a Reaction conditions: **1a** (1 mmol), solvent (10 mL), under microwave irradiation in O₂; ^b Monitored by TLC; ^c Isolated yield; ^d The reaction was heated in an oil bath in Air; ^e The reaction was heated in N₂

nonahydrate (1.0 equiv.) and DMSO (10 mL) in a sealed tube full of O₂ under microwave irradiation.

With the optimized reaction conditions in hand, the scope of a variety of 2-methylquinolines was then tested to investigate the impact of electronic as well as steric factors on the reaction profile. The results were summarized in Table 2. Delightedly, compared with other oxidative methods, the major drawback of overoxidation of aldehydes was not observed under current reaction conditions. What's more, we found that the reaction was selective and only occurred on the methyl group at 2-position in the presence of other methyl groups on the quinoline ring (Entries 4, 5, and 6). It was observed that 2-methylquinolines with both electron-donating groups (Entries 2~6) and electron-withdrawing groups (Entries 7~10) gave the corresponding products in moderate to good yield. Interestingly, more iron(III) and time were needed with electron-donating substituents viz. 6-OMe (**1b**), 7-OMe (**1c**), 3-Me (**1d**), 6-Me (**1e**), 8-Me (**1f**). It was noteworthy that disubstituted ones viz. 3-Me (**1d**), 8-Me (**1f**), 8-Cl (**1g**) underwent difficulty in providing the corresponding 2-quinoline carboxaldehydes in modest yields (48%~54%), which might be due to the steric effect at C(3) or C(8) position of 2-methylquinolines. Furthermore, 2-methylquinoxaline (**1k**) was also compatible to access aldehyde derivative **2k** in a good yield of 80% (Entries 11).

According to the optimization of process conditions, the amplification experiments to investigate the impurities were studied with the reaction condition of **1a** (10 mmol), Fe(NO₃)₃·9H₂O (10 mmol), DMSO (10 mL), MW 150 °C, power 150 W in O₂. The impurities **3** and **4** were determined in the yields of 10.8% and 3.5% (Figure 2).

To explore the reaction mechanism, the radical trapping experiment was carried out by using 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) as a radical scavenger (Scheme 1). As expected, the reaction failed to give 2-quinoline carboxaldehydes. This result indicated that the

Figure 2 Impurities of **3** and **4**

reaction might involve a radical pathway.

Scheme 1 Radical trapping experiment with TEMPO

On the basis of above experiment and literature examples,^[13] a plausible mechanism was deduced (Scheme 2). In this hypothesis, it is assumed that **1a** can be isomerized to a nonaromatic enamine intermediate **A**, which then coordinates with iron(III) nitrate nonahydrate to generate iron-enamide species **B**. Fe³⁺ replaces the nitro to react firstly with molecular oxygen to form a Fe³⁺ peroxo species **C**, subsequent isomerization gives peroxide **D**. Elimination of [Fe²⁺]OH affords 2-quinoline carboxaldehydes **2a**.

3 Conclusions

In summary, the oxidative formylation of various quinoline derivatives was presented in the presence of iron(III) nitrate nonahydrate and DMSO under O₂ atmosphere promoted by microwave irradiation. Availability, stability and non-toxicity of the reagents, mild reaction conditions, relatively short reaction time, promising yields, and easy work-up are advantages which make this methodology be an important alternative for the preparation of 2-quinoline carboxaldehydes.

Table 2 Synthesis of 2-quinoline carboxaldehydes^a

Entry	X	R	Time ^b /min	Fe ³⁺ /1 (molar ratio)	Product	Yield ^c /%
1	CH	H	5	1.0	2a	78
2	CH	6-OMe	10	1.5	2b	60
3	CH	7-OMe	10	1.5	2c	64
4	CH	3-Me	10	1.5	2d	54
5	CH	6-Me	10	1.5	2e	63
6	CH	8-Me	10	1.5	2f	48
7	CH	8-Cl	5	1.0	2g	52
8	CH	6-NO ₂	5	1.0	2h	81
9	CH	6-F	5	1.0	2i	73
10	CH	6-Cl	5	1.0	2j	67
11	N	H	5	1.0	2k	80

^a Reaction conditions: **1** (1 mmol), DMSO (10 mL), MW 150 °C, power 150 W in O₂. ^b Monitored by TLC. ^c Isolated yield.

Scheme 2 Plausible mechanism

4 Experimental section

4.1 General information

All reagents were obtained from commercial suppliers and used without further purification, unless otherwise indicated. TLC analysis was performed using pre-coated glass plates. Silica gel for column chromatography was purchased from Qingdao Haiyang Chemical Co., Ltd. Microwave reactor was CEM Discover 908010 microwave reactor with IR-monitored temperature control. Melting points were determined using a Büchi B-540 capillary melting point apparatus. ^1H NMR and ^{13}C NMR spectra were measured on a Varian 400 (400 MHz) spectrometer (chemical shifts in δ) using tetramethylsilane as internal standard.

4.2 Synthesis of 2-quinoline carboxaldehydes

2-methylquinolines (**1**, 1 mmol), $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ (1 mmol) and DMSO (10 mL) were mixed in an 80 mL microwave tube full of O_2 . The mixture was stirred at $150\text{ }^\circ\text{C}$ for 5 min under microwave irradiation using a CEM Discover microwave reactor. The progress of the reaction was monitored by TLC. After completion, the reaction mixture was filtered firstly, then washed by sodium bicarbonate solution and filtered. The aqueous layer extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo* and the crude residue was purified by flash chromatography on silica gel using hexane/EtOAc as eluent to give the product **2a**~**2k**.

Quinoline-2-carboxaldehyde (**2a**): Yield 78%. White solid, m.p. $68\sim69\text{ }^\circ\text{C}$ (lit.^[14] $70\sim72\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.21 (s, 1H), 8.29 (d, $J=8.4\text{ Hz}$, 1H), 8.23 (d, $J=8.4\text{ Hz}$, 1H), 8.01 (d, $J=8.4\text{ Hz}$, 1H), 7.88 (d, $J=8.0\text{ Hz}$, 1H), 7.83~7.79 (m, 1H), 7.69~7.66 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 192.8, 152.0, 147.4, 137.0, 130.1, 130.0, 129.7, 128.8, 127.5, 117.1.

6-Methoxyquinoline-2-carboxaldehyde (**2b**): Yield 60%. Yellow solid, m.p. $101\sim102\text{ }^\circ\text{C}$ (lit.^[15] $105\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.16 (s, 1H), 8.15 (d, $J=8.4\text{ Hz}$, 1H), 8.11 (d, $J=9.2\text{ Hz}$, 1H), 7.98 (d, $J=8.4\text{ Hz}$, 1H), 7.45 (dd, $J=9.2\text{ Hz}$, 2.8 Hz, 1H), 7.11 (d, $J=2.8\text{ Hz}$, 1H), 3.97 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 192.4, 159.2, 149.9, 143.3, 135.2, 131.4, 131.2, 123.2, 117.5, 104.8, 55.9.

7-Methoxyquinoline-2-carboxaldehyde (**2c**): Yield 64%.

Yellow solid, m.p. $103\sim105\text{ }^\circ\text{C}$ (lit.^[16]); ^1H NMR (400 MHz, CDCl_3) δ : 10.15 (s, 1H), 8.15~8.08 (m, 2H), 7.98~7.94 (m, 1H), 7.45~7.42 (m, 1H), 7.10 (s, 1H), 3.96 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 193.2, 159.7, 150.4, 143.8, 135.4, 131.7, 131.4, 123.4, 117.7, 104.9, 55.7.

3-Methylquinoline-2-carbaldehyde (**2d**): Yield 54%. Yellow solid, m.p. $111\sim112\text{ }^\circ\text{C}$ (lit.^[17] $115\sim116\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.30 (s, 1H), 8.17 (d, $J=8.4\text{ Hz}$, 1H), 7.99 (s, 1H), 7.78 (d, $J=8.4\text{ Hz}$, 1H), 7.75~7.71 (m, 1H), 7.65~7.61 (m, 1H), 2.78 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 195.0, 150.6, 146.2, 138.0, 130.8, 129.7, 129.2, 129.0, 126.6, 19.2.

6-Methylquinoline-2-carbaldehyde (**2e**): Yield 63%. Yellow solid, m.p. $105\sim106\text{ }^\circ\text{C}$ (lit.^[18] $105\sim106\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.18 (s, 1H), 8.18 (d, $J=8.4\text{ Hz}$, 1H), 8.11 (d, $J=8.8\text{ Hz}$, 1H), 7.97 (d, $J=8.4\text{ Hz}$, 1H), 7.64~7.63 (m, 2H), 2.58 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 192.6, 151.2, 145.9, 139.1, 136.0, 132.3, 129.7, 129.5, 126.2, 117.1, 22.3.

8-Methylquinoline-2-carbaldehyde (**2f**): Yield 48%. Yellow solid, m.p. $78\sim79\text{ }^\circ\text{C}$ (lit.^[15] $81\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.21 (s, 1H), 8.23 (d, $J=8.4\text{ Hz}$, 1H), 7.98 (d, $J=8.4\text{ Hz}$, 1H), 7.70 (d, $J=8.8\text{ Hz}$, 1H), 7.64~7.62 (m, 1H), 7.56~7.52 (m, 1H), 2.88 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 193.9, 151.4, 146.8, 138.7, 137.2, 130.2, 130.0, 128.8, 125.6, 116.8, 17.9.

8-Chloroquinoline-2-carbaldehyde (**2g**): Yield 52%. Yellow solid, m.p. $143\sim145\text{ }^\circ\text{C}$ (lit.^[19] $129\sim131\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.28 (s, 1H), 8.33 (d, $J=8.4\text{ Hz}$, 1H), 8.08 (d, $J=8.4\text{ Hz}$, 1H), 7.92 (d, $J=8.0\text{ Hz}$, 1H), 7.82 (d, $J=8.0\text{ Hz}$, 1H), 7.59 (t, $J=8.0\text{ Hz}$, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 192.3, 152.2, 143.7, 137.4, 134.4, 130.9, 130.1, 128.6, 126.4, 117.7.

6-Nitro-quinoline-2-carboxaldehyde (**2h**): Yield 81%. Yellow solid, m.p. $190\sim192\text{ }^\circ\text{C}$ (lit.^[20] $188\sim189\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, DMSO) δ : 10.12 (s, 1H), 9.17 (d, $J=2.4\text{ Hz}$, 1H), 8.92 (d, $J=8.4\text{ Hz}$, 1H), 8.55 (dd, $J=9.2\text{ Hz}$, 2.4 Hz, 1H), 8.40 (d, $J=9.6\text{ Hz}$, 1H), 8.13 (d, $J=8.4\text{ Hz}$, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 193.2, 154.4, 148.7, 146.2, 140.2, 131.4, 128.5, 124.9, 123.6, 118.4.

6-Fluoro-quinoline-2-carboxaldehyde (**2i**): Yield 73%. White solid, m.p. $116\sim117\text{ }^\circ\text{C}$ (lit.^[21] $122\sim124\text{ }^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 10.18 (s, 1H), 8.26~8.23 (m, 2H), 8.03 (d, $J=8.8\text{ Hz}$, 1H), 7.61~7.56 (m, 1H), 7.51 (dd, $J=8.4\text{ Hz}$, 2.8 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3)

δ : 192.3, 161.2 (d, $^1J_{\text{CF}}=249.4$ Hz), 151.5, 144.4, 136.1 (d, $^4J_{\text{CF}}=5.4$ Hz), 132.6 (d, $^3J_{\text{CF}}=9.3$ Hz), 130.5 (d, $^3J_{\text{CF}}=10.5$ Hz), 120.6 (d, $^2J_{\text{CF}}=25.7$ Hz), 117.7, 110.6 (d, $^2J_{\text{CF}}=21.8$ Hz).

6-Chloroquinoline-2-carbaldehyde (**2j**): Yield 67%; Yellow solid, m.p. 136~137 °C (lit.^[15] 170~171 °C); ^1H NMR (400 MHz, CDCl_3) δ : 10.18 (s, 1H), 8.22~8.16 (m, 2H), 8.03 (d, $J=8.4$ Hz, 1H), 7.78 (s, 1H), 7.74 (d, $J=8.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 192.2, 152.1, 145.7, 135.9, 134.7, 131.5, 131.0, 130.1, 126.1, 117.9.

Quinoxaline-2-carboxaldehyde (**2k**): Yield 80%. Yellow solid, m.p. 97~98 (lit.^[22] 107~108 °C); ^1H NMR (400 MHz, CDCl_3) δ : 10.20 (s, 1H), 9.35 (s, 1H), 8.20~8.11 (m, 2H), 7.92~7.82 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 191.6, 145.0, 141.9, 132.8, 132.4, 131.4, 130.7, 130.0, 129.2.

Quinoline (**3**): Yield 10.8%. Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.90 (d, $J=8.4$ Hz, 1H), 8.23 (d, $J=3.2$ Hz, 1H), 8.12 (t, $J=9.6$ Hz, 1H), 7.80 (d, $J=8.0$ Hz, 1H), 7.71 (t, $J=7.2$ Hz, 1H), 7.53 (t, $J=7.2$ Hz, 1H), 7.38 (dd, $J=8.4, 4.4$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 150.6, 148.4, 136.3, 129.7, 129.6, 128.5, 128.0, 126.7, 121.2; HRMS (ESI) calcd for $[\text{M}+\text{H}]^+$ $\text{C}_9\text{H}_8\text{N}$, m/z 130.0651, found 130.0652.

Di-2-quinolinylmethanone (**4**): Yield 3.5%, yellow solid. ^1H NMR (400 MHz, CDCl_3) δ : 8.31 (d, $J=8.4$ Hz, 2H), 8.18 (d, $J=8.4$ Hz, 2H), 7.90 (d, $J=8.0$ Hz, 2H), 7.85 (t, $J=7.6$ Hz, 2H), 7.73~7.69 (m, 4H); HRMS (ESI) calcd for $[\text{M}+\text{H}]^+$ $\text{C}_{19}\text{H}_{13}\text{N}_2\text{O}$, m/z 285.1022, found 285.1030.

Supporting Information ^1H NMR and ^{13}C NMR spectra copies of the products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Li, Z.; Wu, S.-S.; Luo, Z.-G.; Liu, W.-K.; Feng, C.-T.; Ma, S.-T. *J. Org. Chem.* **2016**, *81*, 4386.
(b) Xu, L.-B.; Shao, Z.-Z.; Wang, L.; Zhao, H.-L.; Xiao, J. *Tetrahedron Lett.* **2014**, *55*, 6856.
(c) Thirunavukkarasu, V. S.; Kozhushkov, S. I.; Ackermann, L. *Chem. Commun.* **2014**, *50*, 29.
- [2] (a) Derong, D.; Linda, P. D.; Peter, A. C. *Tetrahedron Lett.* **2013**, *54*, 5211.
(b) Jiang, L.; Huang, Y.-Y.; Yan, Y.-Y.; Xie, Y.-Y. *Tetrahedron Lett.* **2016**, *57*, 4149.
- [3] Guo, S.-J.; Wan, G.; Sun, S.; Jiang, Y.; Yu, J.-T.; Cheng, J. *Chem. Commun.* **2015**, *51*, 5085.
- [4] Xie, Y.-Y.; Li, L.-H. *Tetrahedron Lett.* **2014**, *55*, 3892.
- [5] (a) Adsule, S.; Barve, V.; Chen, D.; Ahmed, F.; Dou, Q. P.; Padhye, S.; Sarkar, F. H. *J. Med. Chem.* **2006**, *49*, 7242.
(b) Teguh, S. C.; Klonis, N.; Duffy, S.; Lucantoni, L.; Avery, V. M.; Hutton, C. A.; Baell, J. B.; Tilley, L. *J. Med. Chem.* **2013**, *56*, 6200.
- [6] (a) Pokhrel, L.; Kim, Y.; Nguyen, T. D. T.; Prior, A. M.; Lu, J. Y.; Chang, K. O.; Hua, D. H. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 3480.
(b) Dai, Q.; Yu, J.-T.; Feng, X.-M.; Jiang, Y.; Yang, H.-T.; Cheng, J. *Adv. Synth. Catal.* **2014**, *356*, 3341.
(c) Gopinath, V. S.; Pinjari, J.; Dere, R. T.; Verma, A.; Vishwakarma, P.; Shivahare, R.; Moger, M.; Goud, P. S. K.; Ramanathan, V.; Bose, P.; Rao, M. V. S.; Gupta, S.; Puri, S. K.; Launay, D.; Martin, D. *Eur. J. Med. Chem.* **2013**, *69*, 527.
- [7] Holzapfel, C. W.; Ferreira, A. C.; Marais, W. J. *Chem. Res., Synop.* **2002**, *5*, 218.
- [8] Ding, D.; Dwoskin, L. P.; Crooks, P. A. *Tetrahedron Lett.* **2013**, *54*, 5211.
- [9] Minisci, F.; Vismara, E.; Levi, S. *J. Org. Chem.* **1986**, *51*, 536.
- [10] Zheng, G.; Liu, H.; Wang, M. *Chin. J. Chem.* **2016**, *34*, 519.
- [11] (a) Sindhu, K. S.; Abi, T. G.; Mathai, G.; Anilkumar, G. *Polyhedron* **2019**, *158*, 270.
(b) Wusiman, A.; Hudabaierdi, R. *Tetrahedron Lett.* **2019**, *60*, 681.
(c) Wang, X.-Z.; Zeng, C.-C. *Tetrahedron* **2019**, *75*, 1425.
(d) Ding, X.-Y.; Xu, F. *Chin. J. Org. Chem.* **2018**, *38*, 3345 (in Chinese).
(丁晓友, 徐凡, 有机化学, **2018**, *38*, 3345.)
- [12] (a) Ma, S.-M.; Liu, J.-X.; Li, S.-H.; Chen, B.; Cheng, J.-J.; Kuang, J.-Q.; Liu, Y.; Wan, B.-Q.; Wang, Y.-L.; Ye, J.-T.; Yu, Q.; Yuan, W. M.; Yu, S.-C. *Adv. Synth. Catal.* **2011**, *353*, 1005.
(b) Silva, M. J.; Carari, D. M. *Catal. Lett.* **2014**, *144*, 615.
(c) Tanaka, S.; Kon, Y.; Uesaka, Y.; Morioka, R.; Tamura, M.; Sato, K. *Chem. Lett.* **2016**, *45*, 188.
(d) Xu, X.-H.; Sun, J.; Cheng, J.-Y.; Li, P.-P.; Jiang, X.-Y.; Bai, R.-R.; Xie, Y.-Y. *Eur. J. Org. Chem.* **2017**, *47*, 7160.
(e) Amaya, T.; Fujimoto, H. *Tetrahedron Lett.* **2018**, *59*, 2657.
(f) Wen, J.; Zhang, J.; Chen, S.-Y.; Li, J.; Yu, X.-Q. *Angew. Chem. Int. Ed.* **2008**, *47*, 8897.
(g) Nambodiri, V. V.; Polshettiwar, V.; Varma, R. S. *Tetrahedron Lett.* **2007**, *48*, 8839.
- [13] (a) Jiang, K.; Pi, D.-W.; Zhou, H.-F.; Liu, S.-S.; Zou, K. *Tetrahedron* **2014**, *70*, 3056.
(b) Rao, N. N.; Meshram, H. M.; *Tetrahedron Lett.* **2013**, *54*, 1315.
(c) Li, Q.; Huang, Y.; Chen, T.-Q.; Zhou, Y.-B.; Xu, Q.; Yin, S.-F.; Han, L.-B. *Org. Lett.* **2014**, *16*, 3672.
- [14] Yasunari, M.; Kanoko, Y.; Takashi, T.; Takayuki, A.; Tomohiro, M.; Kosaku, H.; Hironao, S. *Heterocycles* **2010**, *80*, 737.
- [15] Mathes, W.; Sauermilch, W. *Chem. Ber.* **1957**, *90*, 758.
- [16] Chen, X.-Y.; Shi, J.; Li, Y.-M.; Wang, F.-L.; Wu, X.; Guo, Q.-X.; Liu, L. *Org. Lett.* **2009**, *11*, 4421.
- [17] Brown, B. R.; Hammick, D. L. *J. Chem. Soc.* **1950**, 628.
- [18] Buehler, C. A.; Edwards, S. P. *J. Am. Chem. Soc.* **1952**, *74*(4), 977.
- [19] Ballesteros, G. R.; Leroux, F. R.; Ballesteros, R. *Tetrahedron* **2009**, *65*(22), 4410.
- [20] Janina, B. *PL 56421*, **1968** [*Chem. Abstr.* **1968**, *70*, 115023].
- [21] Wang, L.; Hou, X.-B.; Fu, H.-S.; Pan, X.-L.; Xu, W.-F.; Tang, W.-P.; Fang, H. *Bioorg. Med. Chem.* **2015**, *15*(23), 4364.
- [22] Leese, C. L.; Rydon, H. N. *J. Chem. Soc.* **1956**, 303.

(Li, L.; Fan, Y.)