1,4-二氮杂二环[2.2.2]辛烷(DABCO)催化联烯的[4+2]串联反应: 合成苯并噻吩并吡喃衍生物

贾继茹 于爱敏 刘旭光 孟祥太*

(天津市有机太阳能电池与光化学转换重点实验室 天津理工大学化学化工学院 天津 300384)

摘要 发展了 1,4-二氮杂二环[2.2.2]辛烷(DABCO)催化硫代橙酮类似物与联烯酸酯之间的串联环化反应,以甲醇为溶剂,以较好的选择性和收率获得两类苯并噻吩并吡喃衍生物.同时,该反应条件温和、底物适应性广,吸电子基或供电子基取代的硫代橙酮类似物都能以较好的选择性和收率生成目标产物.该反应为苯并噻吩并吡喃衍生物的合成提供了一种便利的方法.

关键词 串联反应; 联烯酸酯; 苯并噻吩衍生物

1,4-Diazabicyclo[2.2.2]octane (DABCO)-Catalyzed [4+2] Domino Reaction of Allenates: Synthesis of Benzo[4,5]thieno-[3,2-*b*]pyran Derivatives

Jia, Jiru Yu, Aimin Liu, Xuguang Meng, Xiangtai*

(* Tianjin Key Laboratory of Organic Solar Cells and Photochemical Conversion, School of Chemistry & Chemical Engineering, Tianjin University of Technology, Tianjin 300384 China)

Abstract A 1,4-diazabicyclo[2.2.2]octane (DABCO)-catalyzed [4+2] annulation reaction between 2-alkylidenbenzothiophene-3(2*H*)-ones and allenates has been developed. The substrate scope includes both electron-withdrawing and electron-donating groups on the benzothiophene moiety. This method can be carried out under mild conditions and gives a wide range of highly functionalized benzothiophene-fused γ -pyran derivatives in good yields with moderate selectivity.

Keywords domino reaction; allenates; benzothiophene derivatives

1 Introduction

Benzothiophene-fused γ -pyran derivatives are important structural skeletons in biologically active compounds.^[1] For example, some benzo[*b*]thieno[3,2-*b*]pyran derivatives have pronounced anti-asthmatic activity.^[2] This has motivated considerable efforts in exploring various strategies for their synthesis. Some reports include a C—H bond activation approach from benzothiophene and carboxylic acids as substrates.^[3] In another report, 2,3-substituted benzothiophene derivatives were converted into benzothiophene-fused pyran derivatives.^[4] Although many methods have been described to synthesize these skeletons, there are only a few reports of the efficient synthesis of these target

molecules using domino reaction.^[5] We recently reported many domino reactions based on thioaurones. For example, the reaction of thioaurones bearing a hydroxyl group with allenates gave benzothiophene-fused dioxabicyclo[3.3.1]nonane derivatives (Scheme 1a).^[6] Moreover, when the thioaurone analogs reacted with γ -benzyl allenates, a series of benzothieno[3,2-*b*]pyran derivatives were obtained (Scheme 1b).^[7] In line with our program aiming toward the development of domino reactions for the efficient construction of benzothiophene-fused heterocyclic compounds,^[8] we reported a [4+2] pericyclic reaction between thioaurone analogs and non-substituted allenates.^[9–12] Two new types of benzothiophene-fused pyran derivatives were obtained (Scheme 1c).

* Corresponding author. E-mail: xtmeng@tjut.edu.cn

Received April 30, 2019; revised June 25, 2019; published online July 9, 2019.

Dedicated to the 100th anniversary of the birth of Professor Ruyi Chen.

Project supported by the National Natural Science Foundation of China (No. 21403154), the Natural Science Foundation of Tianjin City (No. 17JCZDJC37700) and the Tianjin Municipal Education Commission (No. 20180KJ137).

国家自然科学基金(No. 21403154)、天津市自然科学基金(No. 17JCZDJC37700)、天津市教委(No. 20180KJ137)资助项目.

Scheme 1 Domino reactions between thioaurones with allenates

2 Results and discussion

To optimize the reaction conditions, Reactions of methyl (*Z*)-2-(5-methyl-3-oxobenzo-*[b]*thiophen-2(3*H*)-ylidene) acetate (**1a**) with allenate **2** was selected in the presence of DABCO (10 mol%) in CHCl_3 at 60 °C for 2 h. The reaction proceeded smoothly and obtained the product **3a** in 46% yield along with the product **4a** in 15% yield (Table 1, Entry 1). The structure of **3a** was determined using single-crystal X-ray diffraction (Figure 1),^[13] NMR, and HRMS. The structure of **4a** was determined using 2D-NMR. After confirming the structures of the two new products, the effect of the solvent on the yield and regioselectivity were then examined. From Entries 2~11, it was found that when MeOH was used as the solvent, the yield increased slightly and the ratio of **3a** and **4a** increased to 5 : 1 (Entry 2). However, when the reaction was carried out in EtOH, moderate yields and regioselectivity were observed (Entry 3). No improvement in yields was obtained when the reactions were carried out in the strong polar solvents of DMSO and DMF (Entries 4 and 5). When the reactions were performed in toluene or 1,4-dioxane, only moderate yields and regioselectivity were obtained in both solvents (Entries 6 and 7). With THF as the solvent, the reaction proceeded smoothly to produce **3a** and **4a** in 62% yield with 1 : 1 regioselectivity (Entry 8). Therefore,

MeOH was selected as the solvent and the effect of the catalysts was determined. When 4-dimethylaminopyridine (DMAP) was used as catalyst, the starting materials were consumed but a very complex mixture was obtained (Entry 9). Phosphines were also tested, but the reaction was still complex, such as PPh_3 and $^n\text{Bu}_3\text{P}$ (Entries 10 and 11). Therefore, DABCO was selected as the catalyst.

Table 1 Optimization of the domino reaction^a

Entry ^a	Solvent	Catalyst	Time/h	Yield ^b /%	
				3a	4a
1	CHCl_3	DABCO	2.0	46	15
2	MeOH	DABCO	2.2	64	13
3	EtOH	DABCO	2.0	32	32
4	DMSO	DABCO	1.2	21	3
5	DMF	DABCO	2.0	16	16
6	Toluene	DABCO	2.5	34	17
7	1,4-Dioxane	DABCO	2.2	34	17
8	THF	DABCO	2.2	31	31
9	MeOH	DMAP	3.0	—	—
10	MeOH	PPh_3	3.0	—	—
11	MeOH	$^n\text{Bu}_3\text{P}$	3.0	—	—

^a Unless otherwise noted, reactions of **1a** (0.2 mmol) with **2** (0.24 mmol) were performed in the presences of catalyst (0.1 equiv.) in 2 mL of solvent at 60 °C.

^b Isolated yields.

Figure 1 X-ray crystal structure of **3a**

Having established the optimal reaction conditions, we explored the substrate scope of the domino reactions with regard to the substitution at **1** (Table 2). The effect of the ester group (R^2) was first examined when R^1 was a methyl group at the C5 position. Substrate **1** bearing an ethyl ester gave **3b** and **4b** in 79% yield with a 2 : 1 regioselectivity. Similarly, benzyl-substituted substrate **1c** led to the corresponding products **3c** and **4c** in 72% yield with 4 : 1 regioselectivity. When a halogen group was installed in the C5 position (R^1), the corresponding products **3d**~**3f** and **4d**~**4f** were obtained in 49%~72% yields with 3 : 1~4 : 1

Table 2 Scope of the [4+2] domino reaction^{a,b}

^a Reaction conducted on a 0.2 mmol scale of **1** in 2 mL of MeOH. ^b Isolated yields.

regioselectivity. When a methoxyl group was substituted at the C6 position (R¹), the domino reaction produced the corresponding products **3g~3i** and **4g~4i** in 51%~74% yields with 2 : 1 to 9 : 1 regioselectivity. The non-substituted substrates **1j~1l** also reacted well regardless of the ester groups. In addition, substrates bearing benzyl groups were also suitable (**1m** and **1n**), producing the corresponding products **3m** and **3n** in 53% and 67% yields. However, in these reactions, the regioselectivity of **3** and **4** is poor.

A mechanism for this domino reaction is proposed on the basis of previous reports in the literature (Scheme 2). The DABCO catalyst acts as a nucleophilic trigger and attacks the β carbon of the allenoate to produce intermedi-

ates **A** and **A'**. The allylic carbanion **A** binds to substrate **1a** to produce intermediate **C**. Intermediate **D**, which is a resonance structure of **C**, undergoes an intramolecular umpolung addition from an oxygen anion to a γ carbon of the allenoate to produce intermediate **E**, which then becomes **F** via elimination of the DABCO catalyst. Finally, intermediate **F** isomerizes to **3a**. Similarly, another intermediate **A'**, produced by allenoate, attacks the double-bond position of substrate **1a** to produce intermediate **C'**. **D'** is the resonance structure of **C'** and generates intermediate **E'** through an intramolecular addition reaction. With the departure of the catalyst (DABCO), product **4a** is finally formed.

Scheme 2 Proposed mechanisms.

3 Conclusions

In conclusion, a novel DABCO-catalyzed [4+2] annulation reaction between allenolate and 2-alkylidenebenzothiophene-3(2*H*)-ones has been developed. A number of functionalized benzothiophene-fused γ -pyran derivatives were formed in moderate yields. Further applications of this domino reaction in complex molecular syntheses are currently under study.

4 Experimental section

4.1 Experimental details

All reactions were performed under Ar atmosphere in

oven dried glassware with magnetic stirring. Unless otherwise stated, all reagents were purchased from commercial suppliers and used without further purification. All solvents were purified and dried according to standard methods prior to use. All reactions were monitored by thin-layer chromatography (TLC) with silica gel-coated plates. Flash column chromatography was performed using 200~300 mesh silica gel. ¹H NMR and ¹³C NMR spectra were recorded at ambient temperature on a Bruker 400 instruments. All spectra were referenced to CDCl₃ (¹H NMR δ 7.26 and ¹³C NMR δ 77.00. HRMS were obtained on a Waters Xevo Q-TOF MS in ESI mode. Melting points were measured on an RY-I apparatus and reported uncorrected. Substrates **1** were synthesized according to literature procedures.^[8a]

4.2 Typical procedure for the preparation of **3** and **4**

Under the protection of argon, DABCO (0.02 mmol, 10 mmol%) and allenolate (0.24 mmol, 1.2 equiv.) were added to a suspension of thioaurone **1** (0.2 mmol, 1.0 equiv.) in MeOH (2 mL). The mixture was stirred for 3 h at 60 °C. After the TLC detection reaction was completed, the reaction mixture was evaporated to remove the solvent and purified by flash silica gel column chromatography eluted with ethyl acetate/hexane (*V* : *V* = 1 : 10) to provide pure samples **3** and **4**.

3-Benzyl-4-methyl-2,8-dimethyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3a**): Yellow solid, m.p. 123~124 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.53~7.61 (m, 2H), 7.33~7.40 (m, 5H), 7.18 (d, *J* = 8.2 Hz, 1H), 5.29 (d, *J* = 12.3 Hz, 1H), 5.12 (d, *J* = 12.3 Hz, 1H), 4.92 (s, 1H), 3.58 (s, 3H), 2.60 (s, 3H), 2.47 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 171.46, 166.38, 162.39, 140.42, 135.81, 134.57, 133.67, 129.68, 128.51, 128.51, 128.22, 128.20, 128.19, 127.24, 122.31, 119.95, 111.92, 99.51, 66.49, 52.40, 41.49, 21.38, 19.65; IR (KBr) ν : 3031, 2958, 1751, 1715, 1640, 1595, 1454, 1436, 1358, 1310, 1232, 1200, 1172, 1059, 972, 809, 756, 699 cm⁻¹. ESI-HRMS calcd for C₂₃H₂₀O₅SNa [M + Na]⁺ 431.0924, found 431.0947.

3-Benzyl-4-ethyl-2,8-dimethyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3b**): White solid, m.p. 85~86 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.54~7.62 (m, 2H), 7.31~7.41 (m, 5H), 7.18 (d, *J* = 9.6 Hz, 1H), 5.25 (d, *J* = 12.4 Hz, 1H), 5.15 (d, *J* = 12.4 Hz, 1H), 4.90 (s, 1H), 4.07~4.16 (m, 1H), 3.96~4.04 (m, 1H), 2.60 (s, 3H), 2.47 (s, 3H), 1.18 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 170.95, 166.43, 162.32, 140.37, 135.83, 134.51, 133.66, 129.70, 128.48, 128.15, 127.17, 122.28, 119.92, 112.06, 99.52, 66.47, 61.55, 41.61, 21.37, 19.62, 14.05; IR (KBr) ν : 3063, 3033, 2985, 2961, 2883, 1748, 1713, 1639, 1595, 1455, 1359, 1329, 1306, 1252, 1231, 1204, 1182, 1088, 807, 773, 726, 691 cm⁻¹; ESI-HRMS calcd for C₂₄H₂₂O₅SNa [M + Na]⁺ 445.1080, found 445.1105.

Dibenzyl-2,8-dimethyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3c**): Yellow solid, m.p. 86~87 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.54~7.61 (m, 2H),

7.27~7.37 (m, 10H), 7.18 (d, $J=8.4$ Hz, 1H), 5.05~5.13 (m, 3H), 4.93~4.98 (m, 2H), 2.60 (s, 3H), 2.47 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.87, 166.32, 162.47, 140.47, 135.80, 135.40, 134.54, 133.69, 129.65, 128.50, 128.44, 128.29, 128.22, 128.13, 127.23, 122.31, 119.95, 111.83, 99.43, 67.32, 66.41, 41.62, 21.37, 19.63; IR (KBr) ν : 3033, 2952, 1748, 1694, 1649, 1454, 1377, 1305, 1253, 1199, 1166, 1089, 1071, 960, 752, 699 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{29}\text{H}_{25}\text{O}_5\text{S}$ $[\text{M}+\text{H}]^+$ 485.1417, found 485.1421.

3-Benzyl-4-ethyl-8-fluoro-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3d**): White solid, m.p. 139~140 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.65 (dd, $J=8.8$, 4.4 Hz, 1H), 7.32~7.45 (m, 6H), 7.10 (m, 1H), 5.26 (d, $J=12.4$ Hz, 1H), 5.15 (d, $J=12.4$ Hz, 1H), 4.91 (s, 1H), 4.08~4.17 (m, 1H), 3.96~4.05 (m, 1H), 2.59 (s, 3H), 1.18 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.69, 166.27, 162.16, 160.94 (d, $J=243.5$ Hz), 140.42 (d, $J=4.2$ Hz), 135.75, 131.80, 130.50 (d, $J=9.7$ Hz), 128.52, 128.21, 124.02 (d, $J=9.2$ Hz), 114.49 (d, $J=7.0$ Hz), 114.20, 106.09 (d, $J=24.3$ Hz), 99.61, 66.59, 61.71, 41.66, 19.58, 14.06; IR (KBr) ν : 3028, 2987, 2905, 1742, 1716, 1646, 1604, 1446, 1242, 1228, 1193, 1083, 1046, 961, 853, 817, 737 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{F}-\text{O}_5\text{S}$ $[\text{M}+\text{H}]^+$ 427.1010, found 427.1014.

3-Benzyl-4-ethyl-8-chloro-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3e**): White solid, m.p. 104~105 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.72 (s, 1H), 7.62 (d, $J=8.4$ Hz, 1H), 7.27~7.42 (m, 6H), 5.26 (d, $J=12.0$ Hz, 1H), 5.15 (d, $J=12.0$ Hz, 1H), 4.90 (s, 1H), 3.96~4.16 (m, 2H), 2.59 (s, 3H), 1.18 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.60, 166.21, 162.09, 139.98, 135.72, 134.55, 131.08, 130.54, 128.51, 128.20, 125.94, 123.72, 119.82, 114.01, 99.62, 66.58, 61.71, 41.57, 19.54, 14.04; IR (KBr) ν : 3080, 3009, 2981, 2938, 1742, 1711, 1638, 1595, 1436, 1353, 1326, 1303, 1247, 1227, 1190, 1085, 1071, 809, 744, 696 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{ClO}_5\text{S}$ $[\text{M}+\text{H}]^+$ 443.0714, found 443.0733.

3-Benzyl-4-ethyl-8-bromo-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3f**): Yellow solid, m.p. 106~107 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.89 (s, 1H), 7.58 (d, $J=8.4$ Hz, 1H), 7.45 (dd, $J=8.6$, 1.8 Hz, 1H), 7.33~7.39 (m, 5H), 5.25 (d, $J=12.4$ Hz, 1H), 5.15 (d, $J=12.4$ Hz, 1H), 4.91 (s, 1H), 4.08~4.15 (m, 1H), 3.98~4.05 (m, 1H), 2.59 (s, 3H), 1.18 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.0, 166.4, 162.3, 140.7, 136.5, 135.8, 129.5, 128.5, 128.2, 125.5, 124.6, 122.6, 120.0, 112.0, 99.6, 66.5, 61.6, 41.6, 19.6, 14.1; IR (KBr) ν : 3064, 2980, 2901, 1740, 1715, 1642, 1599, 1430, 1349, 1249, 1226, 1189, 1162, 967, 808, 749, 695 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{BrO}_5\text{S}$ $[\text{M}+\text{H}]^+$ 487.0215, found 487.0235.

3-Benzyl-4-methyl-7-methoxy-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3g**): White solid, m.p. 141~142 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.62 (d, $J=8.8$ Hz, 1H), 7.29~7.44 (m, 5H), 7.19 (d, $J=2.4$ Hz, 1H), 7.00 (dd, $J=8.8$, 2.0 Hz, 1H), 5.28 (d, $J=12.4$ Hz, 1H), 5.12 (d, $J=12.0$ Hz, 1H), 4.88 (s, 1H), 3.86 (s,

3H), 3.58 (s, 3H), 2.59 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.52, 166.40, 162.38, 158.29, 140.43, 138.07, 135.82, 128.51, 128.19, 123.45, 120.75, 114.48, 108.89, 105.46, 99.56, 66.48, 55.60, 52.37, 41.40, 19.65; IR (KBr) ν : 3006, 1733, 1688, 1649, 1601, 1372, 1323, 1280, 1264, 1204, 1171, 1090, 1024, 856, 759, 699 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{O}_6\text{SNa}$ $[\text{M}+\text{Na}]^+$ 447.0873, found 447.0898.

3-Benzyl-4-ethyl-7-methoxy-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3h**): Red solid, m.p. 105~106 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.62 (d, $J=8.8$ Hz, 1H), 7.32~7.39 (m, 5H), 7.19 (d, $J=2.0$ Hz, 1H), 7.00 (dd, $J=8.8$, 2.4 Hz, 1H), 5.25 (d, $J=12.4$ Hz, 1H), 5.14 (d, $J=12.4$ Hz, 1H), 4.86 (s, 1H), 4.08~4.17 (m, 1H), 3.95~4.04 (m, 1H), 3.86 (s, 3H), 2.59 (s, 3H), 1.18 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.02, 166.43, 162.32, 158.19, 140.34, 138.02, 135.80, 128.47, 128.13, 123.43, 120.70, 114.42, 108.98, 105.37, 99.51, 66.45, 61.51, 55.56, 41.48, 19.63, 14.06; IR (KBr) ν : 3035, 2970, 2903, 1749, 1712, 1647, 1602, 1369, 1243, 1183, 1163, 1076, 1020, 825, 810, 697 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{24}\text{H}_{22}\text{O}_6\text{SNa}$ $[\text{M}+\text{Na}]^+$ 461.1029, found 461.1044.

Dibenzyl-7-methoxy-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3i**): Red solid, m.p. 54~55 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=8.8$ Hz, 1H), 7.32~7.41 (m, 10H), 7.20 (d, $J=2.4$ Hz, 1H), 7.00 (dd, $J=8.8$, 2.0 Hz, 1H), 5.25 (d, $J=12.4$ Hz, 1H), 5.14 (d, $J=12.4$ Hz, 1H), 4.86 (s, 1H), 4.08~4.12 (m, 1H), 3.96~4.04 (m, 1H), 3.86 (s, 3H), 2.59 (s, 3H), 1.18 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.93, 166.34, 162.46, 158.27, 140.48, 138.09, 135.81, 135.43, 128.50, 128.44, 128.28, 128.22, 128.13, 123.42, 120.74, 114.47, 108.79, 105.45, 99.48, 67.28, 66.40, 55.59, 41.53, 19.63; IR (KBr) ν : 2960, 2938, 2891, 1748, 1710, 1644, 1600, 1371, 1242, 1175, 1167, 1068, 1023, 743, 699 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{29}\text{H}_{24}\text{O}_6\text{SNa}$ $[\text{M}+\text{Na}]^+$ 523.1186, found 523.1192.

3-Benzyl-4-methyl-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3j**): Red solid, m.p. 50~51 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.70~7.78 (m, 2H), 7.31~7.42 (m, 7H), 5.29 (d, $J=12.4$ Hz, 1H), 5.12 (d, $J=12.4$ Hz, 1H), 4.94 (s, 1H), 3.59 (s, 3H), 2.61 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.39, 166.35, 162.39, 140.72, 136.49, 135.80, 129.46, 128.52, 128.24, 128.21, 125.53, 124.61, 122.68, 120.06, 111.84, 99.56, 66.52, 52.43, 41.45, 19.65; IR (KBr) ν : 3030, 3008, 2951, 1747, 1720, 1645, 1605, 1434, 1373, 1332, 1247, 1228, 1178, 1166, 1083, 961, 762, 735 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{22}\text{H}_{18}\text{O}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 417.0767, found 417.0778.

3-Benzyl-4-ethyl-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate (**3k**): White solid, m.p. 137~138 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.71~7.78 (m, 2H), 7.32~7.42 (m, 7H), 5.23~5.30 (m, 1H), 5.11~5.18 (m, 1H), 4.92 (d, $J=8.0$ Hz, 1H), 3.84~4.27 (m, 2H), 2.61 (s, 3H), 1.18 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.89, 166.39, 162.30, 140.67, 136.49, 135.82, 129.49, 128.49, 128.16, 125.46, 124.56, 122.64, 120.03,

111.98, 99.57, 66.49, 61.59, 41.57, 19.61, 14.05; IR (KBr) ν : 2980, 2936, 1740, 1718, 1645, 1604, 1456, 1373, 1332, 1300, 1229, 1188, 1081, 1021, 761, 738, 729 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{O}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 431.0924, found 431.0941.

Dibenzyl-2-methyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3,4-dicarboxylate; Black solid (**3l**): m.p. 60~61 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.71~7.75 (m, 2H), 7.28~7.43 (m, 12H), 5.08~5.19 (m, 3H), 4.97~5.00 (m, 2H), 2.63 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.71, 166.17, 162.34, 140.66, 136.41, 135.73, 135.31, 129.34, 128.44, 128.38, 128.23, 128.17, 128.07, 125.43, 124.51, 122.58, 119.96, 111.67, 99.41, 67.27, 66.35, 41.51, 19.54; IR (KBr) ν : 3032, 2963, 1749, 1719, 1643, 1597, 1454, 1373, 1301, 1253, 1227, 1181, 1165, 1079, 956, 941, 755, 700 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{28}\text{H}_{22}\text{O}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 493.1080, found 493.1051.

Benzyl-8-chloro-2-methyl-4-phenyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3-carboxylate (**3m**): White solid, m.p. 123~124 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.74 (d, $J=2.0$ Hz, 1H), 7.53 (d, $J=8.4$ Hz, 1H), 7.28 (m, 3H), 7.18~7.26 (m, 6H), 7.04~7.09 (m, 2H), 5.27 (s, 1H), 4.98~5.09 (m, 2H), 2.57 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.77, 160.13, 144.70, 138.47, 135.73, 134.71, 130.90, 128.62, 128.39, 128.11, 128.01, 127.52, 127.11, 125.39, 123.83, 121.77, 119.79, 104.22, 66.35, 40.74, 19.49; IR (KBr) ν : 3062, 3028, 2950, 1716, 1638, 1597, 1455, 1434, 1351, 1315, 1250, 1225, 1084, 926, 804, 757, 701 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{26}\text{H}_{19}\text{ClO}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 469.0636, found 469.0658.

Benzyl-8-bromo-2-methyl-4-phenyl-4*H*-benzo[4,5]thieno[3,2-*b*]pyran-3-carboxylate (**3n**): White solid, m.p. 133~134 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.90 (d, $J=2.0$ Hz, 1H), 7.48 (d, $J=8.4$ Hz, 1H), 7.38 (dd, $J=8.6$, 1.8 Hz, 1H), 7.28 (m, 3H), 7.18~7.26 (m, 5H), 7.06 (dd, $J=6.4$, 2.8 Hz, 2H), 5.27 (s, 1H), 4.98~5.09 (m, 2H), 2.57 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.77, 160.13, 144.69, 138.33, 135.72, 135.22, 131.25, 128.63, 128.40, 128.12, 128.02, 128.00, 127.53, 127.12, 124.12, 122.85, 121.61, 118.51, 104.22, 66.36, 40.70, 19.50; IR (KBr) ν : 3081, 3029, 2922, 1705, 1634, 1590, 1455, 1349, 1317, 1252, 1226, 1075, 809, 744, 697 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{BrO}_3\text{S}$ $[\text{M}+\text{H}]^+$ 491.0311, found 491.0336.

Methyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-8-methyl-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4a**): red solid, m.p. 111~112 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=8.4$ Hz, 1H), 7.54 (s, 1H), 7.31~7.42 (m, 5H), 7.19 (d, $J=8.4$ Hz, 1H), 5.91 (s, 1H), 5.17~5.25 (m, 2H), 3.91~4.03 (m, 2H), 3.76 (s, 3H), 3.62 (dd, $J=15.4$, 5.8 Hz, 1H), 2.47 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.82, 166.77, 164.85, 141.90, 136.26, 134.41, 133.78, 129.99, 128.51, 128.07, 127.23, 122.31, 119.84, 110.56, 100.68, 65.82, 52.62, 37.12, 25.53, 21.39; IR (KBr) ν : 3037, 2963, 2868, 1743, 1707, 1653, 1449, 1352, 1250, 1214, 1121, 1062, 1024, 841, 801, 697 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{O}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 431.0924, found 431.0940.

Ethyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-8-methyl-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4b**): Red solid, m.p. 101~102 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=8.4$ Hz, 1H), 7.54 (s, 1H), 7.33~7.41 (m, 5H), 7.19 (d, $J=8.8$ Hz, 1H), 5.90 (s, 1H), 5.19~5.23 (m, 2H), 4.17~4.25 (m, 2H), 3.91~4.00 (m, 2H), 3.58~3.65 (m, 1H), 2.47 (s, 3H), 1.27 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.31, 166.81, 165.05, 141.87, 136.29, 134.39, 133.80, 130.04, 128.52, 128.07, 127.19, 122.32, 119.83, 110.78, 100.62, 65.81, 61.79, 37.34, 25.50, 21.39, 14.08; IR (KBr) ν : 3034, 2962, 1732, 1709, 1656, 1450, 1376, 1351, 1247, 1213, 1122, 1021, 847, 803, 698 cm^{-1} ; SI-HRMS: calcd for $\text{C}_{24}\text{H}_{22}\text{O}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 445.1080, found 445.1075.

Benzyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-8-methyl-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4c**): yellow solid, m.p. 96~97 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.60 (d, $J=8.4$ Hz, 1H), 7.54 (s, 1H), 7.28~7.43 (m, 10H), 7.19 (d, $J=7.2$ Hz, 1H), 5.89 (s, 1H), 5.12~5.25 (m, 4H), 4.00~4.10 (m, 2H), 3.50~3.59 (m, 1H), 2.47 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.21, 166.75, 164.79, 141.97, 136.27, 135.20, 134.43, 133.79, 130.01, 128.55, 128.53, 128.38, 128.31, 128.08, 128.06, 127.25, 122.34, 119.88, 110.47, 100.75, 67.50, 65.81, 37.32, 25.46, 21.40; IR (KBr) ν : 3032, 2961, 2923, 1740, 1713, 1655, 1450, 1349, 1276, 1255, 1199, 1121, 1048, 1022, 843, 757, 696 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{29}\text{H}_{24}\text{O}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 507.1237, found 507.1255.

Ethyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-8-fluoro-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4d**): red solid, m.p. 128~129 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.66 (dd, $J=8.8$, 4.4 Hz, 1H), 7.31~7.42 (m, 6H), 7.11 (m, 1H), 5.91 (s, 1H), 5.21 (d, $J=3.2$ Hz, 2H), 4.20~4.26 (m, 2H), 3.99 (t, $J=6.4$ Hz, 1H), 3.92 (dd, $J=15.6$, 6.8 Hz, 1H), 3.66 (dd, $J=15.4$, 5.8 Hz, 1H), 1.29 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.06, 166.69, 164.54, 160.84 (d, $J=241.7$ Hz), 141.88 (d, $J=4.3$ Hz), 136.18, 131.92, 130.77 (d, $J=9.7$ Hz), 130.83, 130.73, 128.54, 128.13, 128.09, 124.01 (d, $J=9.2$ Hz), 114.31 (d, $J=25.2$ Hz), 113.26, 105.93 (d, $J=24.3$ Hz), 101.06, 65.90, 61.94, 37.41, 25.38, 14.08; IR (KBr) ν : 3088, 2961, 2923, 1740, 1709, 1650, 1459, 1448, 1379, 1347, 1196, 1122, 841, 822, 757, 698 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{19}\text{FO}_5\text{SNa}$ $[\text{M}+\text{Na}]^+$ 449.0829, found 449.0835.

Ethyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-8-chloro-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4e**): Red solid, m.p. 132~133 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.72 (d, $J=1.6$ Hz, 1H), 7.64 (d, $J=8.4$ Hz, 1H), 7.30~7.40 (m, 6H), 5.91 (s, 1H), 5.15~5.27 (m, 2H), 4.16~4.29 (m, 2H), 3.99 (t, $J=6.4$ Hz, 1H), 3.90 (dd, $J=15.4$, 6.6 Hz, 1H), 3.66 (dd, $J=15.4$, 5.8 Hz, 1H), 1.28 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.00, 166.65, 164.44, 141.52, 136.18, 134.72, 130.94, 128.55, 128.11, 125.93, 123.75, 119.77, 112.78, 101.15, 65.93, 61.98, 37.37, 25.36, 14.09; IR (KBr) ν : 2961, 2921, 1740, 1709, 1651, 1448, 1436, 1380, 1344, 1258, 1168,

1125, 1049, 843, 817, 698 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{19}\text{ClO}_5\text{SNa} [\text{M}+\text{Na}]^+$ 465.0534, found 465.0551.

Ethyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-8-bromo-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4f**): Yellow solid, m.p. 131~133 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.88 (d, $J=2.0$ Hz, 1H), 7.58 (d, $J=8.8$ Hz, 1H), 7.45 (dd, $J=8.4$, 2.0 Hz, 1H), 7.32~7.41 (m, 5H), 5.90 (s, 1H), 5.21 (d, $J=3.2$ Hz, 2H), 4.18~4.27 (m, 2H), 3.99 (t, $J=6.4$ Hz, 2H), 3.90 (dd, $J=15.2$, 6.8 Hz, 1H), 3.66 (dd, $J=15.6$, 6.0 Hz, 1H), 1.28 (t, $J=7.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 169.96, 166.61, 164.40, 141.37, 136.18, 135.21, 131.29, 128.53, 128.08, 124.00, 122.81, 118.53, 112.62, 101.15, 65.90, 61.96, 37.33, 25.34, 14.07; IR (KBr) ν : 2961, 1739, 1709, 1651, 1448, 1380, 1343, 1334, 1298, 1257, 1167, 1124, 1048, 844, 698 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{19}\text{BrO}_5\text{SNa} [\text{M}+\text{Na}]^+$ 509.0029, found 509.0047.

Methyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-7-methoxy-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4g**): Red solid, m.p. 131~132 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=8.8$ Hz, 1H), 7.30~7.43 (m, 5H), 7.20 (d, $J=2.0$ Hz, 1H), 7.00 (dd, $J=8.8$, 2.4 Hz, 1H), 5.90 (s, 1H), 5.16~5.25 (m, 2H), 3.91~4.00 (m, 2H), 3.86 (s, 3H), 3.76 (s, 3H), 3.55~3.64 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.95, 166.77, 164.85, 158.31, 141.94, 138.19, 136.26, 128.51, 128.06, 123.79, 120.69, 114.40, 107.42, 105.41, 100.75, 65.83, 55.61, 52.60, 37.04, 25.51; IR (KBr) ν : 2960, 2924, 1737, 1707, 1375, 1268, 1212, 1179, 1122, 1037, 854, 830, 698 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{O}_6\text{SNa} [\text{M}+\text{Na}]^+$ 447.0873, found 447.0882.

Ethyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-7-methoxy-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4h**): Red solid, m.p. 92~93 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=8.8$ Hz, 1H), 7.32~7.42 (m, 5H), 7.20 (d, $J=2.4$ Hz, 1H), 7.00 (dd, $J=8.8$, 2.0 Hz, 1H), 5.89 (s, 1H), 5.16~5.24 (m, 2H), 4.16~4.28 (m, 2H), 3.91~3.98 (m, 2H), 3.86 (s, 3H), 3.58 (dd, $J=17.4$, 7.8 Hz, 1H), 1.27 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.42, 166.79, 165.05, 158.28, 141.88, 138.18, 136.28, 128.51, 128.06, 123.83, 120.67, 114.38, 107.63, 105.41, 100.69, 65.81, 61.75, 55.61, 37.23, 25.46, 14.08; IR (KBr) ν : 2957, 1741, 1707, 1655, 1604, 1367, 1352, 1266, 1173, 1128, 1036, 822, 699 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{24}\text{H}_{22}\text{O}_6\text{SNa} [\text{M}+\text{Na}]^+$ 461.1029, found 461.1038.

Benzyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-7-methoxy-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4i**): Red solid, m.p. 108~109 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=8.8$ Hz, 1H), 7.32~7.41 (m, 10H), 7.20 (d, $J=2.4$ Hz, 1H), 6.97~7.02 (m, 1H), 5.88 (s, 1H), 5.11~5.23 (m, 4H), 3.98~4.10 (m, 2H), 3.86 (s, 3H), 3.53 (dd, $J=15.1$, 4.7 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.32, 166.72, 164.79, 158.32, 141.99, 138.19, 136.26, 135.23, 128.54, 128.52, 128.36, 128.29, 128.07, 128.05, 123.79, 120.71, 114.42, 107.32, 105.43, 100.81, 67.46, 65.81, 55.62, 37.22, 25.42; IR (KBr) ν : 3036, 2946, 1739, 1699, 1647, 1605, 1366, 1351, 1264, 1214, 1126,

1033, 854, 828, 725, 695 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{29}\text{H}_{24}\text{O}_6\text{SNa} [\text{M}+\text{Na}]^+$ 523.1186, found 523.1205.

Methyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4j**): Red solid, m.p. 61~63 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.72~7.76 (m, 2H), 7.31~7.42 (m, 7H), 5.92 (s, 1H), 5.16~5.25 (m, 2H), 3.92~4.05 (m, 2H), 3.77 (s, 3H), 3.63 (dd, $J=15.4$, 5.8 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.81, 166.77, 164.73, 142.25, 136.63, 136.27, 129.81, 128.54, 128.09, 125.51, 124.49, 122.69, 120.00, 110.48, 100.88, 65.87, 52.67, 37.14, 25.54; IR (KBr) ν : 2959, 2923, 1750, 1712, 1651, 1372, 1352, 1252, 1167, 1117, 1021, 971, 728, 696 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{22}\text{H}_{18}\text{O}_5\text{SNa} [\text{M}+\text{Na}]^+$ 417.0767, found 417.0786.

Ethyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4k**): White solid, m.p. 115~116 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.71~7.77 (m, 2H), 7.30~7.42 (m, 7H), 5.92 (s, 1H), 5.15~5.26 (m, 2H), 4.17~4.31 (m, 2H), 3.92~4.04 (m, 2H), 3.62 (dd, $J=14.8$, 5.2 Hz, 1H), 1.28 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.25, 166.76, 164.90, 142.16, 136.59, 136.25, 129.80, 128.51, 128.07, 125.43, 124.43, 122.66, 119.95, 110.66, 100.77, 65.82, 61.83, 37.28, 25.45, 14.08; IR (KBr) ν : 2961, 2920, 1738, 1709, 1645, 1372, 1256, 1176, 1122, 1046, 1022, 837, 696 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{O}_5\text{SNa} [\text{M}+\text{Na}]^+$ 431.0924, found 431.0927.

Benzyl-(*E*)-2-(2-(benzyloxy)-2-oxoethylidene)-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-4-carboxylate (**4l**): Red solid, m.p. 93~94 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.72~7.76 (m, 2H), 7.29~7.44 (m, 12H), 5.91 (s, 1H), 5.12~5.25 (m, 4H), 4.01~4.12 (m, 2H), 3.52~3.61 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.71, 166.17, 162.34, 140.66, 136.41, 135.73, 135.31, 129.34, 128.44, 128.38, 128.23, 128.17, 128.07, 125.43, 124.51, 122.58, 119.96, 111.67, 99.41, 67.27, 66.35, 41.51, 19.54; IR (KBr) ν : 2963, 2923, 1741, 1711, 1653, 1372, 1352, 1277, 1256, 1199, 1171, 1121, 756, 696 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{28}\text{H}_{22}\text{O}_5\text{SNa} [\text{M}+\text{Na}]^+$ 493.1080, found 493.1084.

Benzyl-(*E*)-2-(8-chloro-4-phenyl-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-2-ylidene)acetate (**4m**): White solid, m.p. 133~134 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.75 (d, $J=2.0$ Hz, 1H), 7.60 (d, $J=8.4$ Hz, 1H), 7.27~7.40 (m, 9H), 7.19~7.28 (m, 2H), 5.88 (s, 1H), 5.09~5.18 (m, 2H), 4.30~4.35 (m, 1H), 3.90 (dd, $J=15.2$, 6.0 Hz, 1H), 3.61 (dd, $J=15.2$, 7.6 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.64, 165.57, 141.27, 141.11, 136.19, 134.48, 131.24, 130.91, 128.81, 128.52, 128.08, 128.02, 127.64, 127.49, 125.54, 123.86, 120.05, 119.65, 100.71, 65.78, 37.67, 31.73; IR (KBr) ν : 3063, 3031, 2962, 1708, 1654, 1448, 1432, 1379, 1342, 1171, 1122, 1076, 1019, 872, 842, 796, 695 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{26}\text{H}_{19}\text{ClO}_3\text{SNa} [\text{M}+\text{Na}]^+$ 469.0636, found 469.0627.

Benzyl-(*E*)-2-(8-bromo-4-phenyl-3,4-dihydro-2*H*-benzo[4,5]thieno[3,2-*b*]pyran-2-ylidene)acetate (**4n**): White solid, m.p. 133~134 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.90 (d, $J=2.0$ Hz, 1H), 7.48 (d, $J=8.4$ Hz, 1H), 7.38 (dd, $J=$

8.6, 1.8 Hz, 1H), 7.28 (m, 3H), 7.18~7.26 (m, 5H), 7.06 (dd, $J=6.4, 2.8$ Hz, 2H), 5.27 (s, 1H), 4.98~5.09 (m, 2H), 2.57 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.77, 160.13, 144.69, 138.33, 135.72, 135.22, 131.25, 128.63, 128.40, 128.12, 128.02, 128.00, 127.53, 127.12, 124.12, 122.85, 121.61, 118.51, 104.22, 66.36, 40.70, 19.50; IR (KBr) ν : 3081, 3029, 2922, 1705, 1634, 1590, 1455, 1349, 1317, 1252, 1226, 1075, 809, 744, 697 cm^{-1} ; ESI-HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{BrO}_3\text{S}$ $[\text{M} + \text{H}]^+$ 491.0317, found 491.0336.

Supporting Information NMR spectra of all new compound (PDF) and crystallographic data for compound **3a** (CIF). The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Scala, A.; Micale, N.; Piperno, A.; Rescifina, A.; Schirmeister, T. Kesselring, J.; Grassi, G. *RSC Adv.* **2016**, *6*, 30628.
(b) Mishra, A.; Ma, C.-Q.; Bäuerle, P. *Chem. Rev.* **2009**, *109*, 1141.
(c) Rasmussen, S. C.; Evenson, S. J.; McCausland, C. B. *Chem. Commun.* **2015**, *51*, 4528.
(d) Gopalsamy, A.; Aplasca, A.; Ciszewski, G.; Park, K.; Ellingboe, J. W.; Orłowski, M.; Feld, B. *Med. Chem. Lett.* **2006**, *16*, 457.
- [2] (a) Wright, J. B.; Johnson, H. G. *J. Med. Chem.* **1973**, *16*, 861.
(b) Gabbutt, C. D.; Heoworth, J. D.; Heron, B. M.; Thomas, J. L. *Chem. Commun.* **1999**, 541.
- [3] (a) Wang, Z.-G.; Yang, M.-F.; Yang, Y.-D. *Org. Lett.* **2018**, *20*, 3001.
(b) Tan, G.-Y.; You, Q.-L.; Lan, J.-B.; You, J.-S. *Angew. Chem., Int. Ed.* **2018**, *57*, 6309.
(c) Qin, X.-R.; Li, X.-Y.; Huang, Q.; Liu, H.; Wu, D.; Guo, Q.; Lan, J.-B.; Wang, R.-L.; You, J.-S. *Angew. Chem., Int. Ed.* **2015**, *54*, 7167.
- [4] Pradhan, T. K.; De, A. *Tetrahedron Lett.* **2005**, *46*, 1493.
- [5] (a) Caruana, L.; Forch, M.; Bernardi, L. *Synlett.* **2017**, *28*, 1530.
(b) Evans, C. S.; Davis, L. O. *Molecules* **2018**, *23*, 33.
(c) Chauhan, P.; Mahajan, S.; Enders, D. *Acc. Chem. Res.* **2017**, *50*, 2809.
(d) Volla, C. M.; Atodiresei, L.; Rueping, M. *Chem. Rev.* **2014**, *114*, 2390.
(e) Zheng, J.; He, M.; Xie, B.-H.; Yang, L.; Hu, Z.; Zhou, H.-B.; Dong, C. *Org. Biomol. Chem.* **2018**, *16*, 472.
- [6] Ma, S.; Yu, A.; Zhang, L.; Meng, X. *J. Org. Chem.* **2018**, *83*, 5410.
- [7] Ma, S.; Yu, A.; Meng, X. *Org. Biomol. Chem.* **2018**, *16*, 2885.
- [8] (a) Jia, J.; Yu, A.; Ma, S.; Zhang, Y.; Li, K.; Meng, X. *Org. Lett.* **2017**, *19*, 6084.
(b) Zhang, Y.; Yu, A.; Jia, J.; Ma, S.; Li, K.; Wei, Y.; Meng, X. *Chem. Commun.* **2017**, *53*, 10672.
(c) Ding W.; Zhang, Y.; Yu, A.; Zhang, L.; Meng, X. *J. Org. Chem.* **2018**, *83*, 13821.
- [9] Allenoate as one carbon synthon:
(a) Meng, X.; Huang, Y.; Chen, R. *Org. Lett.* **2009**, *11*, 137.
(b) Xu, S.; Zhou, L.; Ma, R.; Song, H.; He, Z. *Org. Lett.* **2010**, *12*, 544.
- [10] Selected examples of allenoate as two carbon synthon:
(a) Gu, Y.; Li, F.; Hu, P.; Liao, D.; Tong, X. *Org. Lett.* **2015**, *17*, 1106.
(b) Yang, L.-J.; Li, S.; Wang, S.; Nie, J.; Ma, J.-A. *J. Org. Chem.* **2014**, *79*, 3547.
(c) Pei, C.-K.; Jiang, Y.; Wei, Y.; Shi, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 11328.
(d) Liu, Y.; Du, Y.; Yu, A.; Qin, D.; Meng, X. *Tetrahedron* **2015**, *71*, 7706.
- [11] Selected reviews for allenoates:
(a) Li, Y.; Chen, X.; Chen, X.; Shen, X. *Molecules* **2018**, *23*, 3022.
(b) Chuang, S.-C.; Nallapati, B. S. *Asian J. Org. Chem.* **2018**, *7*, 1743.
(c) Wang, Q.-N.; Liu, Z. Q.; Lou, J.; Yu, Z. *Org. Lett.* **2018**, *20*, 6007.
(d) Yu, Z.-Y.; Jin, Z.-C.; Duan, M.; Bai, R.; Lu, Y.; Lan, Y. *J. Org. Chem.* **2018**, *83*, 9729.
(e) Vaishnav, N. K.; Gupta, A. K.; Kant, R.; Mohanan, K. *J. Org. Chem.* **2018**, *83*, 8759.
- [12] Selected examples of allenoate as four carbon synthon:
(a) Zhu, X.-F.; Lan, J.; Kwon, O. *J. Am. Chem. Soc.* **2003**, *125*, 4716.
(b) Villa, R. A.; Xu, Q.; Kwon, O. *Org. Lett.* **2012**, *14*, 4634.
(c) Wang, T.; Ye, S. *Org. Lett.* **2010**, *12*, 4168.
(d) Yuan, C.; Zhou, L.; Xia, M.; Sun, Z.; Wang, D.; Guo, H. *Org. Lett.* **2016**, *18*, 5644.
- [13] CCDC-1913056 (**3a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

(Li, L.; Fan, Y.)