碱金属盐催化 1,2,4-三唑与 α,β -不饱和酮及二酰亚胺的氮杂 Michael 反应马 奔^{*,a} 王刚刚^a 周红艳^{a,b} 杨靖亚^{*,a}^(a) 西北师范大学化学化工学院 兰州 730070)^(b) 甘肃农业大学理学院 兰州 730070)

摘要 发展了一种高效的碱金属盐催化 1,2,4-三唑与 α,β -不饱和酮及 α,β -不饱和二酰亚胺的氮杂 Michael 加成反应的新方法,以中等到优异的产率得到目标产物. 该方法原料易得,底物普适性好,反应条件温和,易实现克级规模的制备. 产物容易转化为相应的 γ -氨基醇.

关键词 合成方法; Michael 加成; 烯酮; α,β -不饱和二酰亚胺; 含氮杂环; 1,2,4-三唑

Alkali Salt-Catalyzed Aza-Michael Addition of 1,2,4-Triazole to α,β -Unsaturated Ketones and ImidesMa, Ben^{*,a} Wang, Ganggang^a Zhou, Hongyan^{a,b} Yang, Jingya^{*,a}^(a) College of Chemistry and Chemical Engineering, Northwest Normal University, Lanzhou 730070)^(b) College of Science, Gansu Agricultural University, Lanzhou 730070)

Abstract An alkali salt-catalyzed highly efficient aza-Michael addition of 1,2,4-triazole to α,β -unsaturated ketones and imides has been developed, giving the desired products in moderate to excellent yields. The salient features of this reaction involve readily available starting materials, good substrate scope, mild condition, high efficiency and ease of scale-up. The product can be transformed into corresponding γ -aminoalcohol.

Keywords synthetic methods; Michael addition; enones; α,β -unsaturated imides; nitrogen heterocycles; 1,2,4-triazole

1 Introduction

Over the past decades, the addition of various nucleophiles to electron-deficient alkenes (conjugate addition reaction) represents one of the most useful synthetic tools for construction of C—C or C—heteroatom bonds. Aza-Michael addition has been witnessed as one of the most efficient and straightforward synthetic strategy for constructing C—N bond.^[1] In this transformation, valuable synthetic intermediates are usually obtained, such as β -amino carbonyl derivatives, of which β -amino acids are the most representative examples. A wide range of nitrogen nucleophiles (amines, carbamates, hydrazines, azides or nitrogen-containing heterocycles) and Michael acceptors (conjugated aldehydes, ketones, esters, amides, nitriles, nitro olefins, vinyl sulfones or vinyl phosphonates) are good partners in this reaction.^[2] Additionally, the process

can be catalyzed by bases, acids, transition-metals, and even sometimes without catalyst. All of these features together with its simplicity and atom economy make the aza-Michael reaction to form C—N bond.^[3] However, the intermolecular aza-Michael reactions of aromatic nitrogen heterocycles with low nucleophilicity are still challenging and highly attractive.^[4]

1,2,4-Triazole is an important structural motif presenting in a large number of functionalized molecules, and its derivatives are widely existed in nature. Besides applications in materials science, organocatalysis and coordination chemistry,^[5] 1,2,4-triazoles have been specially paid increasing attention because of their extensively medicinal applications as antimicrobial,^[6] anti-cancer, anti-inflammatory, cannabinoid antagonist, protein inhibitor receptor, antagonist, anti-chagasic agent, anti-diabetic *etc.* (Figure 1).^[7] The importance of this heterocyclic moiety has

* Corresponding authors. E-mail: yangjy@nwnu.edu.cn; maben@nwnu.edu.cn

Received July 29, 2019; revised August 26, 2019; published online September 13, 2019.

Project supported by the National Natural Science Foundation of China (No. 21362034).

国家自然科学基金(No. 21362034)资助项目.

prompted the development of many practical synthetic routes to 1,2,4-triazole derivatives, typically involving multistep reaction sequences.^[8]

Figure 1 1,2,4-Triazole subunit in pharmaceuticals

The catalytic aza-Michael reaction of 1,2,4-triazole with electron-deficient alkenes has emerged as a promising approach to access *N*¹-substituted 1,2,4-triazoles. However, only a few intermolecular aza-Michael reactions of 1,2,4-triazole have been successfully established due to the low nucleophilicity. In general, the reactions mainly focus on highly active Michael acceptors. The group of Lin^[9] reported the *N*-methylimidazole catalyzed aza-Michael reaction of 1,2,4-triazole with methyl acrylate and methyl vinyl ketone. Alternatively, the group of Kotsuki^[10] demonstrated the high pressure promoted aza-Michael reaction of 1,2,4-triazole with cyclic enones. The catalytic asymmetric aza-Michael additions of to α,β -unsaturated aldehydes were also reported.^[11] However, the asymmetric Michael addition of 1,2,4-triazole with chalcone reported by Ye *et al.*^[12] afforded the desired adduct in only 43% yield even after a prolonged reaction time. On the other hand, α,β -unsaturated imides can be used as the equivalents of the unsaturated carboxylic acids and represent an important class of Michael acceptors.^[13] However, to our best knowledge, the aza-Michael reactions of triazole with α,β -unsaturated imides are not found as yet. Therefore, it would be extremely desirable to develop effective protocols to access aza-Michael reaction of 1,2,4-triazole with aromatic enones and α,β -unsaturated imides (Scheme 1).

Recently, a novel and efficient cesium carbonate catalyzed aza-Michael addition of pyrazole to α,β -unsaturated ketones has been developed.^[14] However, the catalytic system limits the utilization in the aza-Michael reaction of 1,2,4-triazole with α,β -unsaturated ketones. Thus, developing an efficient protocol to realize the aza-Michael reaction of 1,2,4-triazole with α,β -unsaturated ketones and imides is highly desirable. Herein we report our recent progress on alkali salt catalyzed aza-Michael reaction of 1,2,4-triazole with α,β -unsaturated ketones and imides.

Scheme 1 Aza-Michael reactions of triazole

2 Results and discussion

Our initial study was to examine the reaction of 1,2,4-triazole with chalcone (**1a**) in the presence of various catalysts (Table 1). Firstly, several commercially organic bases were screened (Table 1, Entries 1~3). The desired conjugated addition product **2a** was obtained in 13% NMR yield with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU)^[15] (Table 1, Entry 3). After further investigation of several alkali salts (Table 1, Entries 4~9), to our delight, **2a** was obtained in 76% isolated yield when LiCl^[16] was chosen as the catalyst (Table 1, Entry 9). Next, the effect of solvent on the reaction was evaluated (Table 1, Entries 10~14). Except for dimethyl sulfoxide (DMSO), other employed solvents afforded the desired **2a** with varying yields, but

Table 1 Optimization of reaction conditions^a

Entry	Catalyst	Solvent	Temp./°C	Yield ^b /%
1	Et ₃ N	MeCN	Reflux	0
2	DMAP	MeCN	Reflux	0
3	DBU	MeCN	Reflux	13
4	NaOAc	MeCN	Reflux	23
5	K ₂ CO ₃	MeCN	Reflux	35
6	Li ₂ CO ₃	MeCN	Reflux	41
7	CsCl	MeCN	Reflux	15
8	Cs ₂ CO ₃	MeCN	Reflux	68
9	LiCl	MeCN	Reflux	80 (76)
10	LiCl	DCM	Reflux	10
11	LiCl	THF	Reflux	48
12	LiCl	EtOH	Reflux	11
13	LiCl	DMSO	120	0
14	LiCl	Dioxane	Reflux	94 (92)
15	LiCl	Dioxane	90	75 (72)
16 ^c	LiCl	Dioxane	Reflux	65
17	—	Dioxane	Reflux	<5

^a Reaction conditions: **1a** (0.1 mmol), 1,2,4-triazole (0.12 mmol, 1.2 equiv.), catalyst (5 mol%), solvent (1 mL). ^b NMR yield, the number in parenthesis is isolated yield. ^c 5 h.

the best result was obtained in dioxane. Further investigation showed that a lower reaction temperature (Table 1, Entry 15) and a shortening reaction time (Table 1, Entry 16) are unfavorable to the reaction. Not surprisingly, none of the desired product **2a** was obtained in the absence of any catalyst (Table 1, Entry 17). Thus, the best conditions for this reaction have been established as 5 mol% LiCl as catalyst in 1,4-dioxane at reflux temperature.

To explore the substrate scope, a variety of α,β -unsaturated ketones (**1a**~**1x**) were examined (Table 2). The reactions of chalcone derivatives (**1a**~**1u**) reacted well with 1,2,4-triazole under the established conditions, affording the corresponding adducts **2a**~**2u** in good to excellent yields. For most of the substrates, the reaction was completed within 6~10 h. It is noteworthy that the electron effect of the substituents was slightly affected this reaction, but there is no obvious regularity. To our delight, heteroaryl and aliphatic enones (**1v**~**1x**) were also suitable substrates for the present protocol and gave the products **2v**~**2x** in moderate to excellent yields. In addition, the structure of **2i** was confirmed by single-crystal X-ray diffraction analysis (CCDC 1916254).

Table 2 Scope with respect to α,β -unsaturated ketones^a

Entry	R ¹	R ²	Time/h	Product	Yield ^b /%
1	Ph	Ph	8	2a	92
2	4-MeC ₆ H ₄	Ph	10	2b	87
3	4-MeOC ₆ H ₄	Ph	10	2c	83
4	4-CF ₃ C ₆ H ₄	Ph	8	2d	89
5	4-O ₂ NC ₆ H ₄	Ph	6	2e	90
6	4-ClC ₆ H ₄	Ph	9	2f	93
7	4-BrC ₆ H ₄	Ph	7	2g	88
8	3-MeC ₆ H ₄	Ph	10	2h	83
9	3-O ₂ NC ₆ H ₄	Ph	6	2i	90
10	2-ClC ₆ H ₄	Ph	10	2j	89
11	Ph	4-MeOC ₆ H ₄	8	2k	85
12	Ph	4-ClC ₆ H ₄	9	2l	87
13	Ph	4-BrC ₆ H ₄	8	2m	86
14	4-MeOC ₆ H ₄	4-MeC ₆ H ₄	10	2n	77
15	4-MeC ₆ H ₄	4-MeC ₆ H ₄	12	2o	83
16	4-CF ₃ C ₆ H ₄	4-ClC ₆ H ₄	10	2p	90
17	4-FC ₆ H ₄	4-MeOC ₆ H ₄	10	2q	83
18	4-FC ₆ H ₄	4-ClC ₆ H ₄	9	2r	88
19	4-FC ₆ H ₄	4-BrC ₆ H ₄	9	2s	85
20	4-FC ₆ H ₄	3-MeOC ₆ H ₄	9	2t	82
21	4-ClC ₆ H ₄	4-MeC ₆ H ₄	8	2u	83
22	Furan-2-yl	Ph	10	2v	65
23	Thiophen-2-yl	Ph	10	2w	63
24	Me	Et	9	2x	85

^a Reaction conditions: **1** (0.5 mmol), 1,2,4-triazole (0.6 mmol, 1.2 equiv), LiCl (5 mol%), dioxane (3 mL). ^b Isolated yield.

Inspired by the above results, we then tried to extend

this aza-Michael addition to α,β -unsaturated imides. *N*-Cinnamoyl benzamide (**3a**) was initially selected as a model substrate to optimize the reaction condition. Pleasingly, the desired conjugated addition product **4a** was obtained in 23% isolated yield under the optimal conditions for the reactions of α,β -unsaturated ketones. After many attempts, the aza-Michael addition of α,β -unsaturated imides was finally achieved by the use of cesium carbonate as the catalyst in acetonitrile. The reaction scope was then investigated by variation of α,β -unsaturated imides (Table 3). In general, the α,β -unsaturated imides with both electron-donating and electron-withdrawing substituents on the benzene ring reacted smoothly with 1,2,4-triazole to give the desired products **4a**~**4m** in good to excellent yields (67%~94%). However, the aromatic α,β -unsaturated imides deliver the corresponding product in relatively longer reaction time probably due to the poor dissolution in acetonitrile.

Table 3 Scope with respect to α,β -unsaturated imides^a

Entry	R ³	Ar	Time/h	Product	Yield ^b /%
1	Ph	Ph	12	4a	78
2	4-MeOC ₆ H ₄	Ph	12	4b	77
3	4-FC ₆ H ₄	Ph	9	4c	88
4	3-O ₂ NC ₆ H ₄	Ph	8	4d	86
5	Furan-2-yl	Ph	9	4e	67
6	Et	Ph	6	4f	88
7	ⁿ Pr	Ph	7	4g	94
8	ⁱ Pr	Ph	7	4h	93
9	ⁱ Bu	Ph	6	4i	83
10	Decyl	Ph	6	4j	89
11	ⁿ Pr	3-BrC ₆ H ₄	9	4k	81
12	ⁿ Pr	2-ClC ₆ H ₄	8	4l	83
13	ⁿ Pr	3-EtOC ₆ H ₄	9	4m	80

^a Reaction conditions: **3** (0.5 mmol), 1,2,4-triazole (0.6 mmol, 1.2 equiv), Cs₂CO₃ (5 mol%), MeCN (3 mL). ^b Isolated yield.

Methyl acrylate and nitrostyrene are also good Michael acceptors. Gratifyingly, when both of them were subjected to the standard conditions, the reactions proceeded smoothly and led to the conjugated addition products **5** and **6** in excellent yields (Scheme 2).

Scheme 2 Aza-Michael reaction of 1,2,4-triazole with methyl acrylate and nitrostyrene

To clarify the practicality and utility of this alkali salt-catalyzed aza-Michael reaction of 1,2,4-triazole, an up-

scaling experiment of **1a** and further transformation of the product **2a** were performed (Scheme 3). The gram-scale reaction of **1a** and 1,2,4-triazole was carried out with a much lower catalyst loading, furnishing the corresponding product **2a** in 90% isolated yield. In view of the fact that carbonyl usually served as versatile functional group for synthetic transformations, the reduction of **2a** was conducted and led to the corresponding alcohol **7** in 92% yield.

Scheme 3 Gram-scale experiment of **1a** and the transformation of the product **2a**

3 Conclusions

In conclusion, we have developed a novel and efficient alkali salt-catalyzed aza-Michael reaction of 1,2,4-triazole with α,β -unsaturated ketones and imides. Using a cheap and readily available alkali salt as the catalyst, a variety of aromatic and aliphatic α,β -unsaturated ketones and imides were successfully converted into the corresponding products with moderate to excellent yields and high regioselectivity. The established protocol has advantages of readily available starting materials, high efficiency, broad substrate scope, and ease of scale-up. We anticipate that this study will promote the utilization of aza-Michael addition in more plentiful and efficient organic synthesis. The application of the current reaction to the synthesis of medicine containing heterocycles is now under investigation.

4 Experimental section

4.1 General procedure for aza-Michael addition of 1,2,4-triazole to α,β -unsaturated ketones

α,β -Unsaturated ketones **1** (0.5 mmol), 1,2,4-triazole (0.6 mmol, 41.4 mg), LiCl (0.025 mmol, 1.06 mg, 5 mol%) and dioxane (3.0 mL) were successively added into a dry Schlenk tube equipped with cold finger. The reaction mixture was stirred at reflux temperature until the reaction was completed (monitored by TLC). Then, H₂O (4 mL) was added and the resulting mixture was extracted with ethyl acetate (8 mL $\times$ 2). The combined organic phase was washed with saturated brine, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using petroleum ether/ethyl acetate as the eluent to afford the desired product **2**.

1,3-Diphenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one

(**2a**): colorless solid, yield 127.6 mg (92%). m.p. 84~86 °C; ¹H NMR (400 MHz, CDCl₃) δ : 8.19 (s, 1H), 7.97~7.88 (m, 3H), 7.54 (t, *J*=7.2 Hz, 1H), 7.44~7.28 (m, 7H), 6.19 (dd, *J*=8.8, 4.4 Hz, 1H), 4.42 (dd, *J*=17.8, 9.0 Hz, 1H), 3.62 (dd, *J*=17.8, 4.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 195.6, 151.6, 143.4, 138.7, 135.9, 133.5, 128.9, 128.5, 128.6, 127.9, 126.8, 58.8, 43.6; HRMS (ESI) calcd for C₁₇H₁₆N₃O [M+H]⁺ 278.1288, found 278.1293.

1-Phenyl-3-(*p*-tolyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2b**): colorless solid, yield 126.7 mg (87%). m.p. 131~133 °C; ¹H NMR (600 MHz, CDCl₃) δ : 8.15 (s, 1H), 7.94 (dd, *J*=8.4, 1.2 Hz, 2H), 7.89 (s, 1H), 7.56 (t, *J*=7.2 Hz, 1H), 7.44 (t, *J*=7.8 Hz, 2H), 7.30 (d, *J*=7.8 Hz, 2H), 7.17 (d, *J*=8.4 Hz, 2H), 6.15 (dd, *J*=8.4, 4.8 Hz, 1H), 4.40 (dd, *J*=17.8, 8.7 Hz, 1H), 3.62 (dd, *J*=17.8, 4.9 Hz, 1H), 2.33 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ : 195.8, 151.7, 143.4, 138.5, 136.2, 135.7, 133.6, 129.7, 128.7, 128.1, 126.9, 58.8, 43.7, 21.1; HRMS (ESI) calcd for C₁₈H₁₈N₃O [M+H]⁺ 292.1444, found 292.1445.

3-(4-Methoxyphenyl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2c**): colorless solid, yield 127.6 mg (83%). m.p. 149~150 °C; ¹H NMR (600 MHz, CDCl₃) δ : 8.14 (s, 1H), 7.94 (dd, *J*=8.4, 1.2 Hz, 2H), 7.89 (s, 1H), 7.56 (t, *J*=7.5 Hz, 1H), 7.45 (t, *J*=7.8 Hz, 2H), 7.35 (d, *J*=8.4 Hz, 2H), 6.88 (d, *J*=8.8 Hz, 2H), 6.13 (dd, *J*=8.4, 5.4 Hz, 1H), 4.37 (dd, *J*=18.0, 9.0 Hz, 1H), 3.78 (s, 3H), 3.62 (dd, *J*=18.0, 4.8 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ : 195.9, 159.7, 151.8, 143.4, 136.2, 133.6, 130.7, 128.7, 128.3, 128.1, 114.3, 58.5, 55.3, 43.8; HRMS (ESI) calcd for C₁₈H₁₈N₃O₂ [M+H]⁺ 308.1394, found 308.1399.

1-Phenyl-3-(1*H*-1,2,4-triazol-1-yl)-3-(4-(trifluoromethyl)phenyl)propan-1-one (**2d**): colorless solid, yield 153.7 mg (89%). m.p. 119~120 °C; ¹H NMR (600 MHz, CDCl₃) δ : 8.22 (s, 1H), 7.93~7.92 (m, 3H), 7.61 (d, *J*=7.8 Hz, 2H), 7.58~7.53 (m, 3H), 7.44 (t, *J*=7.8 Hz, 2H), 6.26 (dd, *J*=8.5, 4.9 Hz, 1H), 4.40 (dd, *J*=17.9, 8.6 Hz, 1H), 3.65 (dd, *J*=17.9, 4.8 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ : 195.3, 152.1, 143.7, 142.7, 135.9, 133.8, 130.8 (q, *J*=32.85 Hz), 128.7, 128.1, 127.4, 125.9 (q, *J*=3.75 Hz), 123.7 (q, *J*=225.625 Hz), 58.4, 43.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -63.16; HRMS (ESI) calcd for C₁₈H₁₅F₃N₃O [M+H]⁺ 346.1162, found 346.1161.

3-(4-Nitrophenyl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2e**): colorless solid, yield 145.0 mg (90%). m.p. 164~165 °C; ¹H NMR (600 MHz, CDCl₃) δ : 8.26 (s, 1H), 8.19 (d, *J*=8.4 Hz, 2H), 7.94~7.91 (m, 3H), 7.61~7.56 (m, 3H), 7.45 (t, *J*=7.8 Hz, 2H), 6.32 (dd, *J*=8.1, 5.1 Hz, 1H), 4.38 (dd, *J*=18.0, 8.4 Hz, 1H), 3.69 (dd, *J*=17.7, 5.1 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ : 195.0, 152.3, 147.8, 145.7, 143.8, 135.7, 133.9, 128.8, 128.1, 128.0, 124.1, 58.1, 43.7; HRMS (ESI) calcd for C₁₇H₁₅N₄O₃ [M+H]⁺ 323.1139, found 323.1144.

3-(4-Chlorophenyl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2f**): colorless solid, yield 144.9 mg (93%). m.p. 117~119 °C; ¹H NMR (400 MHz, CDCl₃) δ : 8.20 (s, 1H), 7.91 (dd, *J*=6.4, 4.8 Hz, 3H), 7.56 (t, *J*=8.0 Hz,

1H), 7.43 (t, $J=7.6$ Hz, 2H), 7.37~7.30 (m, 4H), 6.17 (dd, $J=8.5, 4.9$ Hz, 1H), 4.36 (dd, $J=17.9, 8.6$ Hz, 1H), 3.61 (dd, $J=17.9, 4.9$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 195.4, 151.9, 143.5, 137.3, 135.9, 134.5, 133.7, 129.1, 128.7, 128.3, 128.0, 58.2, 43.7; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{ClN}_3\text{O} [\text{M}+\text{H}]^+$ 312.0898, found 312.0904.

3-(4-Bromophenyl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2g**): colorless solid, yield 156.3 mg (88%). m.p. 133~135 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.93~7.90 (m, 3H), 7.57 (t, $J=7.5$ Hz, 1H), 7.48~7.43 (m, 4H), 7.29 (d, $J=8.4$ Hz, 2H), 6.15 (dd, $J=8.7, 5.1$ Hz, 1H), 4.36 (dd, $J=17.7, 8.2$ Hz, 1H), 3.61 (dd, $J=17.7, 5.1$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.5, 151.9, 143.6, 137.8, 136.0, 133.8, 132.1, 128.8, 128.7, 128.1, 122.7, 58.4, 43.7; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{BrN}_3\text{O} [\text{M}+\text{H}]^+$ 356.0393, found 356.0400.

1-Phenyl-3-(*m*-tolyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2h**): colorless solid, yield 120.9 mg (83%). m.p. 102~103 °C. ^1H NMR (600 MHz, CDCl_3) δ : 8.16 (s, 1H), 7.94 (d, $J=7.8$ Hz, 2H), 7.90 (s, 1H), 7.56 (t, $J=7.2$ Hz, 1H), 7.44 (t, $J=7.8$ Hz, 2H), 7.24 (d, $J=7.8$ Hz, 1H), 7.19 (d, $J=7.8$ Hz, 2H), 7.13 (d, $J=7.2$ Hz, 1H), 6.15 (dd, $J=8.7, 4.5$ Hz, 1H), 4.42 (dd, $J=17.7, 8.7$ Hz, 1H), 3.60 (dd, $J=17.4, 4.8$ Hz, 1H), 2.33 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.8, 151.7, 143.5, 138.8, 138.7, 136.2, 133.6, 129.3, 128.9, 128.7, 128.1, 127.6, 123.9, 58.9, 43.7, 21.4; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{18}\text{N}_3\text{O} [\text{M}+\text{H}]^+$ 292.1444, found 292.1445.

3-(3-Nitrophenyl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2i**): colorless solid, yield 145.0 mg (90%). m.p. 161~162 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.37 (s, 1H), 8.27 (s, 1H), 8.19 (dd, $J=8.4, 1.2$ Hz, 1H), 7.95~7.93 (m, 3H), 7.79 (d, $J=7.8$ Hz, 1H), 7.60~7.55 (m, 2H), 7.47 (t, $J=7.8$ Hz, 2H), 6.32 (dd, $J=8.4, 5.0$ Hz, 1H), 4.41 (dd, $J=17.9, 8.5$ Hz, 1H), 3.71 (dd, $J=17.9, 5.0$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.1, 152.4, 148.5, 143.7, 140.8, 135.8, 134.0, 133.2, 130.1, 128.8, 128.1, 123.6, 122.3, 58.1, 43.8; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{N}_4\text{O}_3 [\text{M}+\text{H}]^+$ 323.1139, found 323.1145.

3-(2-Chlorophenyl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2j**): yellow oil, yield 138.73 mg (89%). ^1H NMR (600 MHz, CDCl_3) δ : 8.27 (s, 1H), 7.93 (dd, $J=8.4, 1.2$ Hz, 2H), 7.89 (s, 1H), 7.56~7.53 (m, 1H), 7.44~7.38 (m, 4H), 7.27~7.24 (m, 2H), 6.70 (dd, $J=9.6, 3.6$ Hz, 1H), 4.43 (dd, $J=18.0, 9.6$ Hz, 1H), 3.58 (dd, $J=18.0, 3.6$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.2, 151.6, 144.1, 136.3, 135.9, 133.5, 132.4, 129.8, 129.6, 128.6, 128.1, 128.0, 127.5, 55.0, 42.1; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{ClN}_3\text{O} [\text{M}+\text{H}]^+$ 312.0898, found 312.0899.

1-(4-Methoxyphenyl)-3-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2k**): colorless solid, yield 130.6 mg (85%). m.p. 141~143 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.17 (s, 1H), 7.92~7.89 (m, 3H), 7.39~7.37 (m, 2H), 7.34 (t, $J=7.4$ Hz, 2H), 7.30 (t, $J=7.2$ Hz, 1H), 6.90 (dd, $J=6.6, 4.8$ Hz, 2H), 6.18 (dd, $J=9.0, 4.8$ Hz, 1H), 4.35 (dd, $J=17.7, 8.7$ Hz, 1H), 3.83 (s, 3H), 3.56 (dd, $J=17.4, 4.8$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.1, 163.8, 151.7, 143.5,

138.9, 130.4, 128.9, 128.5, 126.9, 113.8, 109.7, 59.0, 55.4, 43.3; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{18}\text{N}_3\text{O}_2 [\text{M}+\text{H}]^+$ 308.1394, found 308.1399.

1-(4-Chlorophenyl)-3-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2l**): yellow oil, yield 135.6 mg (87%). ^1H NMR (600 MHz, CDCl_3) δ : 8.15 (s, 1H), 7.87 (dd, $J=12.9, 6.3$ Hz, 3H), 7.41~7.30 (m, 7H), 6.16 (dd, $J=8.7, 4.5$ Hz, 1H), 4.39 (dd, $J=18.0, 9.0$ Hz, 1H), 3.56 (dd, $J=17.7, 4.5$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.6, 151.7, 143.5, 140.1, 138.7, 134.4, 129.5, 129.0, 128.9, 128.6, 126.8, 58.9, 43.7; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{ClN}_3\text{O} [\text{M}+\text{H}]^+$ 312.0898, found 312.0899.

1-(4-Bromophenyl)-3-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2m**): colorless solid, yield 152.7 mg (86%). m.p. 142~143 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.15 (s, 1H), 7.89 (s, 1H), 7.79 (d, $J=8.4$ Hz, 2H), 7.57 (d, $J=8.4$ Hz, 2H), 7.39~7.30 (m, 5H), 6.16 (dd, $J=8.7, 4.5$ Hz, 1H), 4.39 (dd, $J=17.7, 9.0$ Hz, 1H), 3.55 (dd, $J=18.0, 4.8$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.8, 151.7, 143.5, 138.6, 134.8, 132.0, 129.6, 129.0, 128.9, 128.6, 126.8, 58.9, 43.6; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{BrN}_3\text{O} [\text{M}+\text{H}]^+$ 356.0393, found 356.0399.

3-(4-Methoxyphenyl)-1-(*p*-tolyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2n**): colorless solid, yield 123.7 mg (77%). m.p. 148~150 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.13 (s, 1H), 7.88 (s, 1H), 7.84 (d, $J=8.4$ Hz, 2H), 7.36~7.34 (m, 2H), 7.24 (d, $J=7.8$ Hz, 2H), 6.88 (d, $J=9.0$ Hz, 2H), 6.13 (dd, $J=8.4, 4.8$ Hz, 1H), 4.33 (dd, $J=17.7, 8.7$ Hz, 1H), 3.78 (s, 3H), 3.59 (dd, $J=17.7, 5.1$ Hz, 1H), 2.39 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.5, 159.7, 151.8, 144.6, 143.3, 133.7, 130.8, 129.4, 128.4, 128.2, 114.3, 58.6, 55.3, 43.6, 21.7; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{N}_3\text{O}_2 [\text{M}+\text{H}]^+$ 322.1550, found 322.1552.

1,3-Dip-tolyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2o**): colorless solid, yield 126.7 mg (83%). m.p. 141~143 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.15 (s, 1H), 7.88 (s, 1H), 7.83 (d, $J=7.8$ Hz, 2H), 7.29 (d, $J=8.4$ Hz, 2H), 7.23 (d, $J=8.4$ Hz, 2H), 7.16 (d, $J=7.8$ Hz, 2H), 6.14 (dd, $J=9.0, 4.8$ Hz, 1H), 4.35 (dd, $J=17.7, 8.7$ Hz, 1H), 3.59 (dd, $J=17.7, 5.1$ Hz, 1H), 2.38 (s, 3H), 2.32 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.4, 151.7, 144.5, 143.4, 138.4, 135.8, 133.7, 129.6, 129.3, 128.2, 126.9, 58.8, 43.5, 21.6, 21.0; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{N}_3\text{O} [\text{M}+\text{H}]^+$ 306.1601, found 306.1606.

1-(4-Chlorophenyl)-3-(1*H*-1,2,4-triazol-1-yl)-3-(4-(trifluoromethyl)phenyl)propan-1-one (**2p**): colorless solid, yield 170.7 mg (90%). m.p. 136~137 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.21 (s, 1H), 7.92 (s, 1H), 7.86 (d, $J=8.4$ Hz, 2H), 7.61 (d, $J=8.4$ Hz, 2H), 7.52 (d, $J=8.4$ Hz, 2H), 7.41 (d, $J=8.4$ Hz, 2H), 6.24 (dd, $J=8.7, 4.5$ Hz, 1H), 4.38 (dd, $J=18.0, 9.0$ Hz, 1H), 3.59 (dd, $J=18.0, 4.8$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.2, 152.1, 143.7, 142.6, 140.4, 134.2, 130.8 (q, $J=32.55$ Hz), 129.5, 129.1, 127.3, 126.0 (q, $J=3.75$ Hz), 123.7 (q, $J=210.75$ Hz), 58.3, 43.7; ^{19}F NMR (376 MHz, CDCl_3) δ : -63.18; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{ClF}_3\text{N}_3\text{O} [\text{M}+\text{H}]^+$ 380.0772, found 380.0779.

3-(4-Fluorophenyl)-1-(4-methoxyphenyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2q**): colorless solid, yield 135.0 mg (83%). m.p. 94~95 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.16 (s, 1H), 7.90 (d, *J*=9.0 Hz, 3H), 7.39 (dd, *J*=8.1, 5.1 Hz, 2H), 7.02 (dd, *J*=11.4, 4.8 Hz, 2H), 6.90~6.89 (m, 2H), 6.16 (dd, *J*=8.4, 4.8 Hz, 1H), 4.29 (dd, *J*=17.7, 8.4 Hz, 1H), 3.83 (d, *J*=1.2 Hz, 3H), 3.56 (dd, *J*=17.4, 5.4 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 193.9, 163.9, 162.5 (d, *J*=246.45 Hz), 151.9, 143.4, 134.8 (d, *J*=3.3 Hz), 130.4, 129.1, 128.8 (d, *J*=8.4 Hz), 115.8 (d, *J*=21.45 Hz), 113.8, 58.4, 55.4, 43.4; ¹⁹F NMR (376 MHz, CDCl₃) δ: -113.50~-113.43 (m); HRMS (ESI) calcd for C₁₈H₁₇FN₃O₂ [M+H]⁺ 326.1299, found 326.1298.

1-(4-Chlorophenyl)-3-(4-fluorophenyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2r**): colorless solid, yield 145.1 mg (88%). m.p. 100~101 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.16 (s, 1H), 7.89 (s, 1H), 7.86~7.85 (m, 2H), 7.41~7.38 (m, 4H), 7.05~7.02 (m, 2H), 6.14 (dd, *J*=8.4, 4.8 Hz, 1H), 4.34 (dd, *J*=17.7, 8.7 Hz, 1H), 3.55 (dd, *J*=17.7, 5.1 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 194.4, 162.6 (d, *J*=247.05 Hz), 151.9, 143.4, 140.2, 134.5 (d, *J*=3.3 Hz), 134.3, 129.5, 129.0, 128.8 (d, *J*=8.25 Hz), 115.9 (d, *J*=21.6 Hz), 58.2, 43.8; ¹⁹F NMR (376 MHz, CDCl₃) δ: -113.15~-113.08 (m); HRMS (ESI) calcd for C₁₇H₁₄-ClFN₃O [M+H]⁺ 330.0804, found 330.0808.

1-(4-Bromophenyl)-3-(4-fluorophenyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2s**): colorless solid, yield 159.0 mg (85%). m.p. 93~94 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.18 (s, 1H), 7.93~7.89 (m, 3H), 7.56 (t, *J*=7.2 Hz, 1H), 7.45~7.40 (m, 3H), 7.03 (t, *J*=8.7 Hz, 2H), 6.17 (dd, *J*=8.4, 4.8 Hz, 1H), 4.37 (dd, *J*=17.7, 8.7 Hz, 1H), 3.62 (dd, *J*=17.7, 5.1 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 195.6, 162.6 (d, *J*=246.75 Hz), 151.9, 143.4, 136.0, 134.6 (d, *J*=3.3 Hz), 133.7, 128.8 (d, *J*=8.4 Hz), 128.7, 128.0, 115.9 (d, *J*=21.6 Hz), 58.2, 43.8; ¹⁹F NMR (376 MHz, CDCl₃) δ: -113.35~-113.28 (m); HRMS (ESI) calcd for C₁₇H₁₄BrFN₃O [M+H]⁺ 374.0299, found 374.0304.

3-(4-Fluorophenyl)-1-(3-methoxyphenyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2t**): colorless solid, yield 133.4 mg (82%). m.p. 97~98 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.17 (s, 1H), 7.90 (s, 1H), 7.52 (d, *J*=7.8 Hz, 1H), 7.42~7.39 (m, 3H), 7.34 (t, *J*=7.8 Hz, 1H), 7.10 (dd, *J*=8.4, 2.4 Hz, 1H), 7.03 (dd, *J*=12.0, 4.8 Hz, 2H), 6.16 (dd, *J*=8.4, 5.4 Hz, 1H), 4.34 (dd, *J*=17.7, 8.7 Hz, 1H), 3.81 (s, 3H), 3.61 (dd, *J*=18.0, 4.8 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 195.4, 162.6 (d, *J*=246.6 Hz), 159.9, 151.9, 143.4, 137.4, 134.6 (d, *J*=3.3 Hz), 129.7, 128.8 (d, *J*=8.25 Hz), 120.6, 120.2, 115.9 (d, *J*=21.45 Hz), 112.3, 58.3, 55.4, 44.0; ¹⁹F NMR (376 MHz, CDCl₃) δ: -113.33~-113.25 (m); HRMS (ESI) calcd for C₁₈H₁₇FN₃O₂ [M+H]⁺ 326.1299, found 326.1298.

3-(4-Chlorophenyl)-1-(*p*-tolyl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2u**): colorless solid, yield 135.2 mg (83%). m.p. 91~92 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.18 (s, 1H), 7.90 (s, 1H), 7.82 (d, *J*=8.4 Hz, 2H), 7.33 (dd, *J*=22.2, 8.4 Hz, 4H), 7.23 (d, *J*=8.4 Hz, 2H), 6.16

(dd, *J*=8.4, 4.8 Hz, 1H), 4.32 (dd, *J*=18.0, 8.4 Hz, 1H), 3.59 (dd, *J*=18.0, 4.8 Hz, 1H), 2.38 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 195.0, 151.9, 144.7, 137.4, 134.5, 133.5, 129.4, 129.1, 128.4, 128.2, 58.4, 43.6, 21.6; HRMS (ESI) calcd for C₁₈H₁₇ClN₃O [M+H]⁺ 326.1055, found 326.1059.

3-(Furan-2-yl)-1-phenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2v**): colorless solid, yield 86.9 mg (65%). m.p. 127~128 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.23 (s, 1H), 7.95~7.91 (m, 3H), 7.59~7.56 (m, 1H), 7.47~7.44 (m, 2H), 7.29 (dd, *J*=4.8, 1.2 Hz, 1H), 7.10 (dd, *J*=4.8, 1.2 Hz, 1H), 6.97 (dd, *J*=5.4, 4.8 Hz, 1H), 6.48 (dd, *J*=9.0, 5.4 Hz, 1H), 4.37 (dd, *J*=18.0, 9.0 Hz, 1H), 3.74 (dd, *J*=11.4, 4.8 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 195.3, 152.1, 141.1, 136.0, 133.8, 128.8, 128.1, 126.9, 126.2, 126.1, 54.6, 44.7; HRMS (ESI) calcd for C₁₅H₁₄-N₃O₂ [M+H]⁺ 268.1081, found 268.1084.

1-Phenyl-3-(thiophen-2-yl)-3-(1*H*-1,2,4-triazol-1-yl)propan-1-one (**2w**): colorless solid, yield 89.2 mg (63%). m.p. 130~131 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.22 (s, 1H), 7.93~7.90 (m, 3H), 7.57~7.54 (m, 1H), 7.45~7.43 (m, 2H), 7.27 (dd, *J*=5.4, 1.2 Hz, 1H), 7.10~7.09 (m, 1H), 6.95 (dd, *J*=4.8, 3.6 Hz, 1H), 6.47 (dd, *J*=9.0, 5.4 Hz, 1H), 4.36 (dd, *J*=17.4, 8.4 Hz, 1H), 3.73 (dd, *J*=18.0, 5.4 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 195.3, 152.0, 143.4, 141.1, 135.9, 133.8, 128.7, 128.1, 126.9, 126.2, 126.1, 54.6, 44.6; HRMS (ESI) calcd for C₁₅H₁₄N₃O₂S [M+H]⁺ 284.0852, found 284.0855.

5-(1*H*-1,2,4-Triazol-1-yl)hexan-3-one (**2x**): colorless oil, yield 71.1 mg (85%). ¹H NMR (600 MHz, CDCl₃) δ: 8.08 (s, 1H), 7.87 (s, 1H), 4.98~4.92 (m, 1H), 3.17 (dd, *J*=17.4, 7.8 Hz, 1H), 2.76 (dd, *J*=17.4, 5.4 Hz, 1H), 2.40~2.26 (m, 2H), 1.52 (d, *J*=7.2 Hz, 3H), 0.97 (t, *J*=7.2 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 207.8, 151.8, 142.6, 51.3, 47.8, 36.4, 20.7, 7.4; HRMS (ESI) calcd for C₈H₁₄N₃O [M+H]⁺ 168.1131, found 168.1136.

4.2 General procedure for aza-Michael addition of 1,2,4-triazole to α,β-unsaturated imides

α,β-Unsaturated imides **3** (0.5 mmol), 1,2,4-triazole (0.6 mmol, 41.4 mg, 1.2 equiv.), Cs₂CO₃ (0.025 mmol, 8.14 mg, 5 mol%) and CH₃CN (3.0 mL) were successively added into a dry Schlenk tube equipped with cold finger. The reaction mixture was stirred at reflux temperature until the reaction was completed (monitored by TLC). Then, H₂O (4 mL) was added and the resulting mixture was extracted with ethyl acetate (8 mL×2). The combined organic phase was washed with saturated brine, dried over anhydrous Na₂SO₄, and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using petroleum ether/ethyl acetate as the eluent to afford the desired product **4**.

N-(3-Phenyl-3-(1*H*-1,2,4-triazol-1-yl)propanoyl)benzamide (**4a**): colorless solid, yield 124.9 mg (78%). m.p. 203~205 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 11.11 (s, 1H), 8.74 (s, 1H), 7.98 (s, 1H), 7.86 (d, *J*=7.2 Hz, 2H), 7.61 (t, *J*=7.4 Hz, 1H), 7.49 (t, *J*=7.6 Hz, 2H), 7.38~

7.29 (m, 5H), 6.14 (dd, $J=9.6, 4.8$ Hz, 1H), 3.95 (dd, $J=17.6, 9.6$ Hz, 1H), 3.62 (dd, $J=17.6, 4.8$ Hz, 1H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 171.4, 166.9, 151.8, 144.5, 139.9, 133.4, 133.3, 129.1, 128.8, 128.5, 127.3, 58.7, 42.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 321.1346, found 321.1349.

N-(3-(4-Methoxyphenyl)-3-(1*H*-1,2,4-triazol-1-yl)propanoyl)benzamide (**4b**): colorless solid, yield 134.9 mg (77%). m.p. 105~107 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.83 (s, 1H), 8.10 (s, 1H), 7.91 (s, 1H), 7.82 (d, $J=7.8$ Hz, 2H), 7.61 (t, $J=7.2$ Hz, 1H), 7.49 (t, $J=7.8$ Hz, 2H), 7.34 (d, $J=9.0$ Hz, 2H), 6.89 (d, $J=9.0$ Hz, 2H), 6.01 (dd, $J=9.3, 4.5$ Hz, 1H), 4.35 (dd, $J=18.0, 9.6$ Hz, 1H), 3.79 (s, 3H), 3.72 (dd, $J=18.0, 4.8$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.1, 165.6, 159.8, 151.7, 143.2, 133.5, 132.3, 130.1, 129.0, 128.5, 127.7, 114.4, 58.7, 55.3, 43.1; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{19}\text{N}_4\text{O}_3$ $[\text{M}+\text{H}]^+$ 351.1452, found 351.1459.

N-(3-(4-Fluorophenyl)-3-(1*H*-1,2,4-triazol-1-yl)propanoyl)benzamide (**4c**): colorless solid, yield 148.9 mg (88%). m.p. 152~153 °C; ^1H NMR (600 MHz, CDCl_3) δ : 9.08 (s, 1H), 8.13 (s, 1H), 7.90 (s, 1H), 7.82 (d, $J=7.2$ Hz, 2H), 7.61 (t, $J=7.8$ Hz, 1H), 7.49 (dd, $J=8.4, 4.2$ Hz, 2H), 7.40~7.38 (m, 2H), 7.04 (t, $J=8.4$ Hz, 2H), 6.04 (dd, $J=8.7, 3.9$ Hz, 1H), 4.35 (dd, $J=18.3, 9.3$ Hz, 1H), 3.71 (dd, $J=18.0, 4.2$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.1, 165.8, 162.7 (d, $J=246.75$ Hz), 151.8, 143.3, 134.0 (d, $J=3.3$ Hz), 133.5, 132.2, 129.0, 128.9, 127.8, 116.0 (d, $J=21.6$ Hz), 58.4, 43.1; ^{19}F NMR (376 MHz, CDCl_3) δ : -113.10~-113.03 (m); HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{FN}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 339.1252, found 339.1253.

N-(3-(3-Nitrophenyl)-3-(1*H*-1,2,4-triazol-1-yl)propanoyl)benzamide (**4d**): colorless solid, yield 157.1 mg (86%). m.p. 138~139 °C; ^1H NMR (600 MHz, CDCl_3) δ : 9.17 (s, 1H), 8.31 (s, 1H), 8.24 (s, 1H), 8.18 (d, $J=7.8$ Hz, 1H), 7.94 (s, 1H), 7.83 (d, $J=7.8$ Hz, 2H), 7.78 (d, $J=7.8$ Hz, 1H), 7.61 (t, $J=7.2$ Hz, 1H), 7.55 (t, $J=8.1$ Hz, 1H), 7.49 (t, $J=7.5$ Hz, 2H), 6.19 (dd, $J=8.7, 3.9$ Hz, 1H), 4.38 (dd, $J=18.3, 9.3$ Hz, 1H), 3.79 (dd, $J=18.0, 4.2$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 171.7, 165.8, 152.3, 148.4, 140.3, 133.6, 133.2, 132.0, 130.1, 129.0, 127.8, 123.7, 122.4, 58.3, 43.1; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{N}_5\text{O}_4$ $[\text{M}+\text{H}]^+$ 366.1197, found 366.1200.

N-(3-(Furan-2-yl)-3-(1*H*-1,2,4-triazol-1-yl)propanoyl)benzamide (**4e**): colorless solid, yield 104.0 mg (67%). m.p. 171~172 °C; ^1H NMR (600 MHz, DMSO- d_6) δ : 11.16 (s, 1H), 8.71 (s, 1H), 7.95 (s, 1H), 7.89~7.88 (m, 2H), 7.62~7.59 (m, 2H), 7.51~7.48 (m, 2H), 6.46 (d, $J=3.0$ Hz, 1H), 6.43 (dd, $J=3.0, 1.8$ Hz, 1H), 6.21 (dd, $J=9.0, 6.0$ Hz, 1H), 3.84 (dd, $J=17.4, 8.4$ Hz, 1H), 3.74 (dd, $J=17.4, 6.0$ Hz, 1H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 171.1, 166.9, 151.9, 151.7, 144.6, 143.6, 133.4, 133.3, 128.9, 128.8, 127.9, 111.1, 108.3, 52.6; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_4\text{O}_3$ $[\text{M}+\text{H}]^+$ 311.1139, found 311.1142.

N-(3-(1*H*-1,2,4-Triazol-1-yl)pentanoyl)benzamide (**4f**): colorless solid, yield 119.8 mg (88%). m.p. 150~151 °C;

^1H NMR (400 MHz, CDCl_3) δ : 9.37 (s, 1H), 8.10 (s, 1H), 7.85 (s, 1H), 7.81 (d, $J=7.8$ Hz, 2H), 7.58 (t, $J=7.2$ Hz, 1H), 7.46 (t, $J=7.8$ Hz, 2H), 4.77~4.72 (m, 1H), 3.75 (dd, $J=18.3, 9.3$ Hz, 1H), 3.39 (dd, $J=18.0, 3.6$ Hz, 1H), 2.08~2.01 (m, 1H), 1.92~1.85 (m, 1H), 0.78 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 173.0, 165.9, 151.8, 143.9, 133.3, 132.4, 128.8, 127.8, 57.5, 42.5, 27.7, 10.4; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{17}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 273.1346, found 273.1350.

N-(3-(1*H*-1,2,4-Triazol-1-yl)hexanoyl)benzamide (**4g**): colorless solid, yield 134.5 mg (94%). m.p. 112~113 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.04 (s, 1H), 8.03 (s, 1H), 7.80~7.76 (m, 3H), 7.50 (t, $J=7.2$ Hz, 1H), 7.39 (t, $J=7.4$ Hz, 2H), 4.82~4.77 (m, 1H), 3.67 (dd, $J=18.2, 9.4$ Hz, 1H), 3.31 (dd, $J=18.2, 3.4$ Hz, 1H), 2.01~1.92 (m, 1H), 1.76~1.67 (m, 1H), 1.15~0.98 (m, 2H), 0.81 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 173.1, 166.1, 151.5, 143.7, 133.0, 132.4, 128.5, 127.9, 55.7, 42.7, 36.3, 18.9, 13.2; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{19}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 287.1503, found 287.1504.

N-(4-Methyl-3-(1*H*-1,2,4-triazol-1-yl)pentanoyl)benzamide (**4h**): colorless solid, yield 133.6 mg (93%). m.p. 135~137 °C; ^1H NMR (600 MHz, CDCl_3) δ : 9.73 (s, 1H), 8.04 (s, 1H), 7.79 (t, $J=7.2$ Hz, 3H), 7.54 (t, $J=7.5$ Hz, 1H), 7.45~7.41 (m, 2H), 4.56~4.52 (m, 1H), 3.81 (dd, $J=18.0, 10.8$ Hz, 1H), 3.43 (dd, $J=18.3, 2.7$ Hz, 1H), 2.22~2.16 (m, 1H), 0.99 (d, $J=7.2$ Hz, 3H), 0.73 (d, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 173.4, 166.1, 151.4, 144.2, 133.1, 132.4, 131.8, 128.7, 127.9, 127.3, 61.6, 39.8, 32.3, 19.4, 18.8; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{19}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 287.1503, found 287.1504.

N-(5-Methyl-3-(1*H*-1,2,4-triazol-1-yl)hexanoyl)benzamide (**4i**): colorless solid, yield 124.6 mg (83%). m.p. 97~98 °C; ^1H NMR (600 MHz, CDCl_3) δ : 9.54 (s, 1H), 8.10 (s, 1H), 7.84~7.77 (m, 3H), 7.57 (t, $J=7.2$ Hz, 1H), 7.47~7.38 (m, 2H), 4.98~4.94 (m, 1H), 3.72 (dd, $J=18.3, 9.3$ Hz, 1H), 3.33 (dd, $J=18.0, 3.6$ Hz, 1H), 2.08~2.02 (m, 1H), 1.58 (d, $J=4.8$ Hz, 1H), 1.22 (t, $J=9.6$ Hz, 1H), 0.90 (d, $J=6.6$ Hz, 3H), 0.83 (d, $J=6.6$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 173.1, 165.9, 151.7, 143.9, 133.3, 132.4, 131.9, 128.8, 128.5, 127.9, 127.3, 54.0, 43.1, 24.4, 22.8, 21.4; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{21}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 301.1659, found 301.1663.

N-(3-(1*H*-1,2,4-Triazol-1-yl)tetradecanoyl)benzamide (**4j**): colorless solid, yield 177.4 mg (89%). m.p. 123~125 °C; ^1H NMR (600 MHz, CDCl_3) δ : 9.73 (s, 1H), 8.06 (s, 1H), 7.82 (d, $J=7.8$ Hz, 3H), 7.54 (s, 1H), 7.43 (s, 2H), 4.81 (s, 1H), 3.72 (dd, $J=18.0, 9.0$ Hz, 1H), 3.35 (d, $J=18.0$ Hz, 1H), 2.00 (s, 1H), 1.79 (s, 1H), 1.19 (d, $J=11.4$ Hz, 18H), 0.83 (dd, $J=7.2, 6.0$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 173.2, 166.1, 151.6, 143.8, 133.2, 132.4, 131.8, 128.7, 128.4, 127.9, 127.3, 56.0, 42.7, 34.4, 31.7, 29.3, 28.9, 25.8, 22.5, 13.9; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{35}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 399.2755, found 399.2758.

N-(3-(1*H*-1,2,4-Triazol-1-yl)hexanoyl)-3-bromobenzamide (**4k**): colorless oil, yield 147.5 mg (81%). ^1H NMR (600 MHz, CDCl_3) δ : 9.43 (s, 1H), 8.16 (s, 1H), 8.02 (s,

1H), 7.90 (s, 1H), 7.76 (d, $J=7.8$ Hz, 1H), 7.72 (d, $J=7.8$ Hz, 1H), 7.37 (t, $J=7.8$ Hz, 1H), 4.93~4.88 (m, 1H), 3.76 (dd, $J=18.0, 9.6$ Hz, 1H), 3.36 (dd, $J=18.0, 3.3$ Hz, 1H), 2.10~2.04 (m, 1H), 1.84~1.79 (m, 1H), 1.23~1.19 (m, 1H), 1.15~1.08 (m, 1H), 0.90 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 173.0, 164.6, 151.8, 136.2, 134.4, 131.0, 130.4, 126.5, 122.9, 55.8, 42.8, 36.5, 19.1, 13.4; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{BrN}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 365.0608, found 365.0609.

N-(3-(1*H*-1,2,4-Triazol-1-yl)hexanoyl)-2-chlorobenzamide (**4l**): colorless solid, yield 132.9 mg (83%). m.p. 93~94 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.86 (s, 1H), 8.12 (s, 1H), 7.88 (s, 1H), 7.58 (d, $J=7.8$ Hz, 1H), 7.46~7.42 (m, 2H), 7.38~7.35 (m, 1H), 4.89~4.84 (m, 1H), 3.70 (dd, $J=18.0, 9.0$ Hz, 1H), 3.33 (dd, $J=18.0, 2.4$ Hz, 1H), 2.10~2.03 (m, 1H), 1.84~1.79 (m, 1H), 1.25~1.08 (m, 2H), 0.90 (t, $J=7.5$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 171.5, 165.2, 151.9, 143.9, 133.4, 132.7, 130.9, 130.6, 130.0, 127.3, 55.7, 42.8, 36.5, 19.1, 13.4; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{ClN}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 321.1113, found 321.1117.

N-(3-(1*H*-1,2,4-Triazol-1-yl)hexanoyl)-2-ethoxybenzamide (**4m**): colorless solid, yield 132.2 mg (80%). m.p. 112~113 °C; ^1H NMR (400 MHz, CDCl_3) δ : 10.50 (s, 1H), 8.16~8.11 (m, 2H), 7.90 (s, 1H), 7.51~7.47 (m, 1H), 7.07 (t, $J=7.6$ Hz, 1H), 6.95 (d, $J=8.4$ Hz, 1H), 4.93~4.87 (m, 1H), 4.20 (q, $J=6.8$ Hz, 2H), 3.74 (dd, $J=18.0, 9.2$ Hz, 1H), 3.36 (dd, $J=18.2, 3.8$ Hz, 1H), 2.11~2.01 (m, 1H), 1.85~1.76 (m, 1H), 1.52 (t, $J=7.0$ Hz, 3H), 1.24~1.04 (m, 2H), 0.89 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.4, 163.8, 157.1, 151.9, 143.89, 134.89, 132.8, 121.6, 119.8, 112.6, 65.3, 55.8, 43.3, 36.5, 19.1, 14.5, 13.4; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{23}\text{N}_4\text{O}_3$ $[\text{M}+\text{H}]^+$ 331.1765, found 331.1764.

4.3 Procedure for gram-scale synthesis of 2a

α,β -Unsaturated ketones **1a** (5.05 mmol, 1.05 g), 1,2,4-triazole (6.06 mmol, 418 mg), LiCl (0.05 mmol, 2.12 mg, 1 mol%) and dioxane (20 mL) were successively added into a dry Schlenk tube equipped with cold finger. The reaction mixture was stirred at reflux temperature until the reaction completed (monitored by TLC). Then, H_2O (10 mL) was added and the resulting mixture was extracted with ethyl acetate (20 mL $\times$ 2). The combined organic phase was washed with saturated brine, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using petroleum ether/ethyl acetate as the eluent to afford the desired product **2a**, 1.26 g, 90% yield.

4.4 Procedure for aza-Michael addition of 1,2,4-triazole to methyl acrylate

Methyl acrylate (2 mmol, 172.0 mg), 1,2,4-triazole (34.5 mg, 0.5 mmol), LiCl (1.06 mg, 0.025 mmol, 5 mol%) and dioxane (3.0 mL) was successively added into a dry Schlenk tube equipped with cold finger. The reaction mixture was stirred at reflux temperature until the reaction

completed (monitored by TLC). Then, H_2O (4 mL) was added and the resulting mixture was extracted with ethyl acetate (8 mL $\times$ 2). The combined organic phase was washed with saturated brine, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using petroleum ether/ethyl acetate as the eluent to afford the desired product **5**.

N-(3-(1*H*-1,2,4-Triazol-1-yl)hexanoyl)benzamide (**5**): colorless oil, yield 76.1 mg (98%). ^1H NMR (600 MHz, CDCl_3) δ : 8.08 (d, $J=8.7$ Hz, 1H), 7.81 (s, 1H), 4.36 (t, $J=5.7$ Hz, 2H), 3.55 (s, 3H), 2.80 (t, $J=6.3$ Hz, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 170.7, 151.7, 143.5, 51.8, 44.6, 33.6; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{20}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 156.0768, found 156.0768.

4.5 Procedure for aza-Michael addition of 1,2,4-triazole to nitrostyrene

Nitrostyrene (0.5 mmol, 74.6 mg), 1,2,4-triazole (41.4 mg, 0.6 mmol), LiCl (1.06 mg, 0.025 mmol, 5 mol%) and dioxane (3.0 mL) were successively added into a dry Schlenk tube equipped with cold finger. The reaction mixture was stirred at reflux temperature until the nitrostyrene completed (monitored by TLC). Then, H_2O (4 mL) was added and the resulting mixture was extracted with ethyl acetate (8 mL $\times$ 2). The combined organic phase was washed with saturated brine, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using petroleum ether/ethyl acetate as the eluent to afford the desired product **6**.

1-(2-Nitro-1-phenylethyl)-1*H*-1,2,4-triazole (**6**): yellow oil, yield 101.4 mg (93%). ^1H NMR (600 MHz, CDCl_3) δ : 8.11 (s, 1H), 7.94 (s, 1H), 7.36~7.32 (m, 5H), 6.18 (dd, $J=9.9, 4.5$ Hz, 1H), 5.51 (dd, $J=14.7, 9.9$ Hz, 1H), 4.87 (dd, $J=14.7, 4.5$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 152.1, 143.5, 133.9, 129.6, 129.3, 126.9, 76.5, 60.2; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{11}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 219.0877, found 219.0879.

4.6 Procedure for the transformation of the product 2a

2a (138.7 mg, 0.5 mmol) was dissolved in MeOH (5 mL) and cooled to 0 °C, then NaBH_4 (0.53 mmol, 20 mg) was slowly added to the mixture. After stirring 2 h at room temperature, **2a** was consumed complete by TLC analysis. The mixture was quenched by distilled water (1 mL) under 0 °C, extracted with EtOAc (15 mL) three times, concentrated, and purified by column chromatography to afford **7** in 92% yield.

1,3-Diphenyl-3-(1*H*-1,2,4-triazol-1-yl)propan-1-ol (**7**): Colorless oil, yield 134.1 mg (96%), 1 : 1.61 *dr*. ^1H NMR (600 MHz, CDCl_3) δ : 7.97 (d, $J=4.8$ Hz, 1H), 7.80~7.78 (m, 2H), 7.73~7.72 (m, 2H), 7.32~7.20 (m, 25 H), 5.74 (dd, $J=10.8, 4.8$ Hz, 1H), 5.49 (dd, $J=8.4, 4.4$ Hz, 1H), 4.40 (t, $J=4.8$ Hz, 1H), 4.24 (d, $J=10.2$ Hz, 1H), 2.83~2.73 (m, 3H), 2.57~2.52 (m, 2H), 2.38~2.34 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 151.5, 151.3, 144.2, 143.8,

143.3, 142.2, 138.9, 138.2, 128.9, 128.7, 128.5, 128.4, 128.3, 128.2, 127.7, 127.5, 127.2, 126.8, 125.7, 125.5, 70.5, 69.7, 61.5, 60.6, 44.2, 44.0; HRMS (ESI) calcd for $C_{17}H_{18}N_3O[M+H]^+$ 280.1444, found 280.1442.

Supporting Information Optimization of the reaction conditions of imides with 1,2,4-triazole, NMR spectra of all products as well as the crystal structure of the product **2i**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Zheng, K.; Liu, X.; Feng, X. *Chem. Rev.* **2018**, *118*, 7586.
(b) Gmach, J.; Błażewska, K. M. *Synthesis* **2016**, *48*, 2681.
(c) Sánchez-Roselló, M.; Aceña, J. L.; Simón-Fuentes, A.; del Pozo, C. *Chem. Soc. Rev.* **2014**, *43*, 7430.
(d) Amara, Z.; Caron, J.; Joseph, D. *Nat. Prod. Rep.* **2013**, *30*, 1211.
(e) Wang, J.; Li, P.; Choy, P. Y.; Chan, A. S. C.; Kwong, F. Y. *ChemCatChem* **2012**, *4*, 917.
(f) Enders, D.; Wang, C.; Liebich, J. X. *Chem. Eur. J.* **2009**, *15*, 11058.
(g) Xu, L. W.; Xia, C. G. *Eur. J. Org. Chem.* **2005**, 633.
- [2] (a) Roy, T. K.; Parhi, B.; Ghorai, P. *Angew. Chem., Int. Ed.* **2018**, *57*, 9397.
(b) Barát, V.; Csókás, D.; Bates, R. W. *J. Org. Chem.* **2018**, *83*, 9088.
(c) Wu, L.; Jin, R.; Li, L.; Hu, X.; Cheng, T.; Liu, G. *Org. Lett.* **2017**, *19*, 3047.
(d) Yang, X.; Chen, Z.; Cai, Y.; Huang, Y. Y.; Shibata, N. *Green Chem.* **2014**, *16*, 4530.
(e) Amara, Z.; Caron, J.; Joseph, D. *Nat. Prod. Rep.* **2013**, *30*, 1211.
(f) Mikami, K.; Fustero, S.; Sánchez-Roselló, M.; Aceña, J. L.; So-loshonok, V.; Sorochinsky, A. *Synthesis* **2011**, 3045.
(g) Weiner, B.; Szymański, W.; Janssen, D. B.; Minnaard, A. J.; Feringa, B. L. *Chem. Soc. Rev.* **2010**, *39*, 1656.
(h) Vinogradov, M. G.; Turovaa, O. V.; Zlotin, S. G. *Org. Biomol. Chem.* **2019**, *17*, 3670.
- [3] (a) Arend, M.; Westermann, B.; Risch, N. *Angew. Chem., Int. Ed.* **1998**, *37*, 1044.
(b) Bartoli, G.; Bosco, M.; Marcantoni, E.; Petrini, M.; Sambri, L.; Torregiani, E. *J. Org. Chem.* **2001**, *66*, 9052.
(c) Reboule, I.; Gil, R.; Collin, J. *Tetrahedron Lett.* **2005**, *46*, 7761.
(d) Loh, T. P.; Wei, L. L. *Synlett* **1998**, 975.
(e) Wabnitz, T. C.; Spencer, J. B. *Tetrahedron Lett.* **2002**, *43*, 3891.
(f) Azizi, N.; Saidi, M. R. *Tetrahedron* **2004**, *60*, 383.
(g) Stanley, L. M.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2009**, *48*, 7841.
(h) Sukhorukov, A. Y.; Sukhanova, A. A.; Zlotin, S. G. *Tetrahedron* **2016**, *72*, 6191.
- [4] (a) Gandelman, M.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2005**, *44*, 2393.
(b) Wang, J.; Li, H.; Zu, L.; Wang, W. *Org. Lett.* **2006**, *8*, 1391.
(c) Uria, U.; Vicario, J. L.; Badia, D.; Carrillo, L. *Chem. Commun.* **2007**, 2509.
(d) Wang, J.; Zu, L.; Li, H.; Xie, H.; Wang, W. *Synthesis* **2007**, 2576.
(e) Madhavan, N.; Takatani, T.; Sherrill, C. D.; Weck, M. *Chem. Eur. J.* **2009**, *15*, 1186.
(f) Luo, G. S.; Zhang, S. L.; Duan, W. H.; Wang, W. *Synthesis* **2009**, 1564.
(g) Lv, J.; Wu, H.; Wang, Y. *Eur. J. Org. Chem.* **2010**, *11*, 2073.
(h) Zhou, Y.; Li, X.; Li, W.; Wu, C.; Liang, X.; Ye, J. *Synlett* **2010**, 2357.
(i) Guo, H. M.; Yuan, T. F.; Niu, H. Y.; Liu, J. Y.; Mao, R. Z. D.; Li, Y.; Qu, G. R. *Chem. Eur. J.* **2011**, *17*, 4095.
(j) Lee, S. J.; Youn, S. H.; Cho, C. W. *Org. Biomol. Chem.* **2011**, *9*, 7734.
(k) Li, H.; Zhao, J.; Zeng, L.; Hu, W. *J. Org. Chem.* **2011**, *76*, 8064.
(l) Uria, U.; Reyes, E.; Vicario, J. L.; Badia, D.; Carrillo, L. *Org. Lett.* **2011**, *13*, 336.
(m) Wang, J.; Wang, W.; Liu, X.; Hou, Z.; Lin, L.; Feng, X. *Eur. J. Org. Chem.* **2011**, *11*, 2039.
(n) Wu, H.; Tian, Z.; Zhang, L.; Huang, Y.; Wang, Y. *Adv. Synth. Catal.* **2012**, *354*, 2977.
(o) Lee, S. J.; Ahn, J. G.; Cho, C. W. *Tetrahedron: Asymmetry* **2014**, *25*, 1383.
(p) Chen, S. W.; Zhang, G. C.; Lou, Q. X.; Cui, W.; Zhang, S. S.; Hu, W. H.; Zhao, J. L. *ChemCatChem* **2015**, *7*, 1935.
(q) Kim, S.; Kang, S.; Kim, G.; Lee, Y. *J. Org. Chem.* **2016**, *81*, 4048.
(r) Bhagat, U. K.; Peddinti, R. K. *Synlett* **2018**, *29*, 99.
(s) Chittimalla, S. K.; Nakka, S.; Koodalingam, M.; Bandi, C. *Synlett* **2018**, *29*, 57.
(t) Bhagat, U. K.; Peddinti, R. K. *J. Org. Chem.* **2018**, *83*, 793.
- [5] (a) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2001**, *40*, 2004.
(b) Herrmann, W. A. *Angew. Chem., Int. Ed.* **2002**, *41*, 1290.
(c) Bräse, S.; Gil, C.; Knepper, K.; Zimmermann, V. *Angew. Chem., Int. Ed.* **2005**, *44*, 5188.
(d) Towns, A. D. *Dyes Pigm.* **1999**, *42*, 3.
(e) Hughes, G.; Bryce, M. R. *J. Mater. Chem.* **2005**, *15*, 94.
(f) Holmes, G.; Rice, A.; Snyder, K. C. *R. J. Mater. Sci.* **2006**, *41*, 4105.
(g) Kusama, H.; Sugihara, H.; Sayama, K. *J. Phys. Chem. C* **2009**, *113*, 20764.
- [6] Ghannoum, M. A.; Rice, L. B. *Clin. Microbiol. Rev.* **1999**, *12*, 50.
- [7] (a) Cardillo, G.; Tomasini, C. *Chem. Soc. Rev.* **1996**, *25*, 117.
(b) Seebach, D.; Matthews, J. L. *Chem. Commun.* **1997**, 2015.
(c) Jagerovic, N.; Hernandez-Folgado, L.; Alkorta, I.; Goya, P.; Navarro, M.; Serrano, A.; de Fonseca, F. R.; Dannert, M. T.; Al-sasua, A.; Suardiaz, M.; Pascual, D.; Martin, M. I. *J. Med. Chem.* **2004**, *47*, 2939.
(d) Papadopoulou, M. V.; Trunz, B. B.; Bloomer, W. D.; McKenzie, C.; Wilkinson, S. R.; Prasittichai, C.; Brun, R.; Kaiser, M.; Torreele, E. *J. Med. Chem.* **2011**, *54*, 8214.
(e) Moulin, A.; Demange, L.; Ryan, J.; Mousseaux, D.; Sanchez, P.; Bergé, G.; Gagne, D.; Perrissoud, D.; Locatelli, V.; Torsello, A.; Galleyrand, J. C.; Fehrentz, J. A.; Martinez, J. *J. Med. Chem.* **2008**, *51*, 689.
(f) Fustero, S.; Pina, B.; Salavert, E.; Navarro, A.; de Arellano, M. C. R.; Fuentes, A. S. *J. Org. Chem.* **2002**, *67*, 4667.
- [8] (a) Huntsman, E.; Balsells, J. *Eur. J. Org. Chem.* **2005**, 3761.
(b) Larsen, S. D.; DiPaolo, B. A. *Org. Lett.* **2001**, *3*, 3341.
(c) Stocks, M. J.; Cheshire, D. R.; Reynold, R. *Org. Lett.* **2004**, *6*, 2969.
(d) Zhang, J. P.; Lin, Y. Y.; Huang, X. C.; Chen, X. M. *J. Am. Chem. Soc.* **2005**, *127*, 5495.
(e) Ueda, S.; Nagasawa, H. *J. Am. Chem. Soc.* **2009**, *131*, 15080.
(f) Guru, M. M.; Punniyamurthy, T. *J. Org. Chem.* **2012**, *77*, 5063.
(g) Tam, A.; Armstrong, I. S.; La Cruz, T. E. *Org. Lett.* **2013**, *15*, 3586.
(h) Xu, H.; Ma, S.; Xu, Y.; Bian, L.; Ding, T.; Fang, X.; Zhang, W.; Ren, Y. *J. Org. Chem.* **2012**, *80*, 1758.
(i) Huang, H.; Guo, W.; Wu, W.; Li, C.; Jiang, H. *Org. Lett.* **2015**, *17*, 2894.
(j) Song, L.; Tian, X.; Lv, Z.; Li, E.; Wu, J.; Liu, Y.; Yu, W.; Chang, J. *J. Org. Chem.* **2015**, *80*, 7219.
(k) Chen, Z.; Li, H.; Dong, W.; Miao, M.; Ren, H. *Org. Lett.* **2016**, *18*, 1334.
(l) Yu, W.; Huang, Y.; Li, J.; Tang, X.; Wu, W.; Jiang, H. *J. Org. Chem.* **2018**, *83*, 9334.
(m) Li, H.; Wu, X.; Hao, W.; Li, H.; Zhao, Y.; Wang, Y.; Lian, P.; Zheng, Y.; Bao, X.; Wan, X. *Org. Lett.* **2018**, *20*, 5224.

- [9] Liu, B. K.; Wu, Q.; Qian, X. Q.; Lv, D. S.; Lin, X. F. *Synthesis* **2007**, 2653.
- [10] Uddin, M. I.; Nakano, K.; Ichikawa, Y.; Kotsuki, H. *Synlett* **2008**, 1402.
- [11] Dinér, P.; Nielsen, M.; Marigo, M.; Jørgensen, K. A. *Angew. Chem., Int. Ed.* **2007**, *46*, 1983.
- [12] Zhou, Y. F.; Li, X.; Li, W. L.; Wu, C. L.; Liang, X. M.; Ye, J. X. *Synlett* **2008**, 1402.
- [13] (a) Sammis, G. M.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2003**, *125*, 4442.
(b) Sammis, G. M.; Danjo, H.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2004**, *126*, 9928.
(c) Madhavan, N.; Weck, M. *Adv. Synth. Catal.* **2008**, *350*, 419.
(d) Mazet, C.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2008**, *47*, 1762.
(e) Madhavan, N.; Sommer, W.; Weck, M. *J. Mol. Catal. A: Chem.* **2011**, *334*, 1.
- [14] Yang, J.; Ma, B.; Zhou, H.; Zhan, B.; Li, Z. *Chin. J. Org. Chem.* **2015**, *35*, 121 (in Chinese).
(杨靖亚, 马奔, 周红艳, 占宝华, 李政, 有机化学, **2015**, *35*, 121.)
- [15] Yang, J.; Bao, Y.; Zhou, H.; Li, T.; Li, N.; Li, Z. *Synthesis* **2016**, *48*, 1139.
- [16] Zhou, H.; Xiang, X.; Ma, B.; Wang, G.; Zhang, Z.; Yang, J. *Synthesis* **2019**, *51*, 3142.

(Cheng, F.)