新型丁烯内酯衍生物的设计合成及诱导胃癌细胞凋亡研究

徐海伟 李媛媛 董航歧 孟夏 赵玲洁
吕春涛 王振亚* 金成允*(郑州大学药学院 药物关键制备技术教育部重点实验室 新药创制与药物安全性评价河南省协同创新中心
药物质量控制与评价河南省重点实验室 郑州 450001)

摘要 开发了一条合成天然产物 Uncinine 的新方法, 基于此设计合成了一系列新型的丁烯内酯衍生物. 通过噻唑蓝 (MTT) 法评价了目标化合物对胃癌细胞的增殖抑制活性, 分析了其构效关系. 其中, 3-吗啉甲基-4-(4-叔丁基苯基) 亚基丁烯内酯(**9I**)对 MGC803 的 IC_{50} 为 $2.9 \mu\text{mol/L}$, 对胃癌细胞 MGC803、HGC27 以及 SGC7901 具有明显的选择性增殖抑制作用, 而对正常的胃粘膜上皮细胞 GES1 具有较小的毒性. 初步的作用机制研究表明, 化合物 **9I** 诱导胃癌细胞 MGC803 凋亡依赖 Caspase 9/3 激活.

关键词 丁烯内酯; 合成; 胃癌, 诱导凋亡; Caspase

Design and Synthesis of Novel Butenolide Derivatives and Inducing Apoptosis in Gastric Cancer Cells

Xu, Haiwei Li, Yuanyuan Dong, Hangqi Meng, Xia Zhao, Lingjie
Lü, Chuntao Wang, Zhenya* Jin, Chengyun*

(Key Laboratory of Drug Preparation Technologies, Ministry of Education, Key Laboratory of Henan Province for Drug Quality Control and Evaluation, Collaborative Innovation Center of New Drug Research and Safety Evaluation, School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001)

Abstract A new method of preparation for natural product Uncinine was reported. According to the method, a series of novel butenolides derivatives were designed and synthesized based on the natural product Uncinine. Most of the synthetic compounds showed significant anti proliferation activity against MGC-803. Among of them, (Z)-5-(4-(*tert*-butyl)-benzylidene)-3-(morpholinomethyl)furan-2(5*H*)-one (**9I**) showed potent anticancer effect with IC_{50} of $2.9 \mu\text{mol/L}$ in gastric cancer cell MGC803 and showed good selectivity to gastric cancer cells MGC803, HGC27, SGC7901 and less cytotoxicity in normal gastric epithelial cell line (GES1). The research on the molecular level suggests that the mechanism of compound **9I** inducing apoptosis in gastric cancer cells is partially dependent on activation of caspase-9/3.

Keywords butenolide; synthesis; gastric cancer; apoptosis, caspase

1 Introduction

The development of novel therapeutic drugs from medicinal plants has been one of the most important resource to drug development. Natural butenolides, such as aspulvione,^[1] uncinine^[2] and so on including a similar structures BH3I-1^[3] and obaloclast^[4] (Figure 1), have a wide range of biological activities, such as antibacterial, antiviral, anticancer, antimalarial, antibiotic and phospholipase A2 inhibition *etc.*^[5]

In connection with previous studies concerning the synthesis and bioactivities of butenolide,^[6,7] we are interested in discovery novel anticancer compounds based on the natural product uncinine. Gastric cancer (GC) is a highly aggressive malignant tumor,^[8] especially in East Asia, such as China, South Korea and Japan. Its high mortality rate prompts the urgent need for novel therapeutic agents. Focusing on GC, our group designed and synthesized novel butenolides and assayed their anti tumor activities. Most of

* Corresponding authors. E-mail: cyjin@zhu.edu.cn; barton118@163.com

Received July 12, 2019; revised August 14, 2019; published online September 18, 2019.

Project supported by the Natural Science Foundation of Henan Province (No. 182300410367), and the Scientific Innovation Talent Award from the Department of Education of Henan Province (No. 15HASTIT036).

河南省自然科学基金(No. 182300410367)和河南省高校科技创新人才(No. 15HASTIT036)资助项目.

Figure 1 Structures of BH31-1, obatoclax and selected natural butenolides

synthetic compounds showed significant anti proliferation activity against gastric cancer cells. Among of them, compound **9l** was discovered with good and selective inhibition of proliferation on cancer cells by triggering apoptotic cell death. At the molecular level, compound **9l** induced apoptosis in gastric cancer cells through dependent caspase activation.

2 Results and discussion

2.1 Chemistry

Uncinine (**6**) as a natural bioactive butenolide, could be synthesized according to literature procedures that is the only one work about its synthesis so far.^[2] However the method is deficient in low yield and many steps. In order to get more uncinine (**6**) and its analogues to research, a new route for preparation of Uncinine was designed and carried out via 4 steps with 30.4% yield shown in Scheme 1. Bromolactone **3** was obtained from the commercially available tulipalin (**1**) by bromination reaction followed by regioselective elimination. The bromination reaction was done in the present of $\text{Me}_3\text{PhN}^+\text{Br}^-$ (1.2 equiv.) at room temperature. The crude product **2** was treated with 5 equiv. of $\text{LiBr}/\text{Li}_2\text{CO}_3$ in the present of TBAB in CH_3CN . By the above two steps reaction, compound **3** was obtained with a yield of 76%. Compounds **4** and **5** were isolated as a mixture after completion of a nucleophilic substitution reaction of bromolactone **3** and ethyl 4-aminobutanoate hydrochloride. The mixture was stirred in acetone and methanol for 72 h at room temperature and purified by silicon chroma-

Reagents and conditions: (a) $\text{Me}_3\text{PhNBr}_3$, dioxane, r.t., 10 h; (b) $\text{LiBr}/\text{Li}_2\text{CO}_3$, TBAB, CH_3CN , r.t., 20 h; (c) ethyl 4-aminobutanoate hydrochloride, Et_3N , dry THF, 0 °C ~ r.t., 26 h; (d) acetone, CH_3OH , r.t., 72 h

Scheme 1 Synthesis of compound **6**

tography to give Uncinine (**6**) with yield of 40% as a white solid.

Based on the intermediate **3**, 3-(alkylamino)methyl-5-(propan-2-ylidene)furan-2(5H)-ones (**8a**~**8g**) were synthesized to investigate the structure activity relationship (SAR) of the pyrrolidin-2-ones moiety in the Uncinine (**6**) by keeping the lactone and isopropyl group (Scheme 2). Different alkenes were introduced to C-5 of **7a** to replace the isopropanylidene group to prepare 5-benzylidene-3-(morpholinomethyl)furan-2(5H)-ones (**9a**~**9o**) and compound **9p** in Scheme 3. The configurations of double bonds on C-5 of compounds **9a**~**9o** were *cis*-double bonds according to NMR and the previous works.^[9,10]

Reagents and conditions: (a) various substituted amine, $\text{K}_2\text{CO}_3/\text{Et}_3\text{N}$, THF, r.t., 2 h; (b) Acetone, CH_3OH , reflux, 1 h

Scheme 2 Synthesis of compound **8**

Reagents and conditions: (a) 5-methylfuran-2-carbaldehyde, ethylenediamine, CH_3OH , reflux; (b) cyclohexanone, ethylenediamine, CH_3OH , reflux; (c) substituted benzaldehyde, ethylenediamine, CH_3OH , reflux

Scheme 3 Synthesis of compound **9**

Among of synthetic compounds, compound **9l** showed potent anticancer effect in gastric cancer MGC803 and less cytotoxicity in normal gastric epithelial cell line (GES1). Compound **9l** has a 4-isopropyl phenyl group in its structure. So two another compounds with 4-isopropyl phenyl methyl indene indol-2-ones were synthesized (Scheme 4).

Reagents and conditions: (a) 4-butylbenzaldehyde, ethylenediamine, CH_3OH , reflux; (b) 4-bromomethyl-tetrahydro-2H-pyran, DMF, NaH, $0\text{ }^\circ\text{C} \sim \text{r.t.}$

Scheme 4 Synthesis of compounds **11** and **12**

2.2 Biology

2.2.1 Evaluation of antiproliferation on MGC803

Anti-proliferation of compounds **8a**~**8g** is evaluated *in vitro* on MGC803. The results are reported in Table 1. Just compound **8a** shows inhibition on the tested cancer cells with IC_{50} of $42.7\text{ }\mu\text{mol/L}$. Uncinine (**6**) and compounds **8b**~**8g** show no significant anti-proliferative activity on MGC803. Based on the above results, compounds **9a**~**9p** are synthesized and assayed to optimize the structure of compound **8a**. The anti-proliferation of compounds **9a**~**9p** are listed in Table 1, and most of them give better anti-proliferation than compound **8a**. Compounds **9a1**~**9a3** with furan have no anti-proliferation on MGC803 cell. Among them compound **9l** is the best one with IC_{50} of $2.9\text{ }\mu\text{mol/L}$. Compounds **11** and **12** with similar substituted phenyl group can inhibit proliferation of MGC803 with IC_{50} of 7.5 and $12.3\text{ }\mu\text{mol/L}$ which are less than compound **9l**. Therefore compound **9l** is chosen to evaluate further.

The SAR of synthetic compounds for anti-proliferation on MGC803 suggests that morpholine aminomethyl group on carbon-3 of butenolide is beneficial to the bioactivity (such as compound **8a**). As for the substitution on carbon-5 butenolide, phenyl groups on the structures of compounds **9b**~**9q**, play more important role than the furan heterocyclic groups on the structures of compounds **9a1**~**9a3**.

Among the substituted groups on the phenyl group for compounds **9b**~**9q**, hydrophilic groups on carbon-4 of phenyl are better than others, such as compounds **9l** and **9m**.

2.2.2 Caspase-dependent apoptosis was induced by compound **9l** in MGC803 cells

Compound **9l** was proved hypotoxicity in normal human gastric epithelial cell line (GES1) and showed a selectivity to gastric cancer cells (MGC803, SGC7901 and HGC27) comparing with GES1 cells (Figure 2A). Based on the above results, compound **9l** was prioritized to perform further experiment for evaluating its anti-cancer potential in gastric cancer.

Morphological changes of MGC803 cells were determined by use of 4,6-diamidino-2-phenylindole (DAPI) staining, as shown in Figure 2B. An increasing number of cells with chromatin condensation were observed in compound **9l**-treated MGC803 cells in a dose-dependent manner. Chromatin condensation was not observed in the untreated control cells, suggesting that compound **9l**-induced cell death is related to apoptosis. We evaluated apoptotic induction in MGC803 by compound **9l** with annexin V and propidium iodide (PI) staining. It was found that the number of annexin V-positive cells showed a dose-dependent increase from 8.6% to 49.5% in MGC803 cells, whereas treatment of control cells only resulted in a 5.2% increase in annexin V-positive stained cells (Figure 2C). These data suggested that apoptosis was induced by compound **9l** in MGC803 cells. The apoptosis-related proteins were detected by western blot analysis in cells treated with compound **9l**. Caspase-3, caspase-9, and poly(ADP-ribose) polymerase (PARP-1) proteins were down regulated, whereas cleaved caspase-9 protein was up regulated by compound **9l** compared with control MGC803 cells (Figure 2D). To further evaluate the significance of caspase activation in compound **9l**-induced apoptosis, z-VAD-fmk ($100\text{ }\mu\text{mol/L}$), an irreversible pan-caspase inhibitor, was used to pretreat MGC803 cells for 1 h, followed by treatment with $20\text{ }\mu\text{mol/L}$ compound **9l** for 24 h. Morphological changes of MGC803 cells were then determined by DAPI staining, as shown in Figure 2E. A significant number of cells with chromatin condensation were observed in cells treated with compound **9l**, and these features were almost invisible in cells pretreated with z-VAD-fmk (100

Table 1 Antiproliferative activity of synthetic compounds on MGC803

Compd.	$\text{IC}_{50}^a/(\mu\text{mol}\cdot\text{L}^{-1})$	Compd.	$\text{IC}_{50}^a/(\mu\text{mol}\cdot\text{L}^{-1})$	Compd.	$\text{IC}_{50}^a/(\mu\text{mol}\cdot\text{L}^{-1})$	Compd.	$\text{IC}_{50}^a/(\mu\text{mol}\cdot\text{L}^{-1})$
8a	42.7 ± 3.1	9a1	>50	9f	28.1 ± 3.1	9m	5.6 ± 0.8
8b	>128	9a2	44.4 ± 1.6	9g	14.7 ± 2.1	9n	17.1 ± 1.2
8c	>128	9a3	>128	9h	13.3 ± 2.1	9o	26.7 ± 1.4
8d	>128	9b	23.1 ± 2.9	9i	26.3 ± 2.7	9p	23.0 ± 1.4
8e	>128	9c	18.6 ± 2.5	9j	19.3 ± 2.2	11	7.5 ± 0.9
8f	>128	9d	10.9 ± 1.7	9k	20.4 ± 1.2	12	12.4 ± 1.1
8g	119.8 ± 3.5	9e	23.2 ± 2.9	9l	2.9 ± 0.1		
5-Fu ^b	7.1 ± 1.3	Uncinine	>128				

^a MTT assay: All data are mean values from triplicate experiments. ^b Positive control: fluorouracil.

Figure 2 Apoptosis induced by compound **9I** in MGC803 cells

(A) The cytotoxicity of compound **9I** by MTT assay. Cells were treated with indicated concentrations for 24 h. $**P < 0.01$ vs. untreated group. (B) MGC803 cells were incubated with compound **9I** (20 μmol/L) for 24 h, and then stained with DAPI. The stained nuclei were observed under a fluorescent microscope (magnification, $\times 200$). (C) A dose-dependent induction of apoptosis by compound **9I** was demonstrated in MGC803 cells by flow cytometric analysis of annexin V/PI stain assay. (D) The effects of compound **9I**-induced (20 μmol/L) protein changes were detected by Western blot analysis for 24 h in MGC803 cells. (E) The effects of caspase inhibitor Z-VAD-fmk (100 μmol/L) on compound **9I**-induced (20 μmol/L) apoptosis of MGC803 cells were shown through DAPI solution (magnification, $\times 200$). (F) Annexin V/PI stained cells showed the effects of caspase inhibitor Z-VAD-fmk (100 μmol/L) on compound **9I**-induced (20 μmol/L) apoptosis of MGC803 cells

μmol/L) followed by treatment with 20 μmol/L compound **9I**. In Figure 2F, compound **9I**-induced apoptosis was reduced from 47.8% to 17.2% by pretreatment with z-VAD-fmk (100 μmol/L), according to flow cytometry analysis.

The activity of caspases is the molecular hallmark of apoptosis and is a controlling factor in the apoptotic cascade.^[11,12] Thus, caspase activity is considered in compound **9I**-induced apoptosis. Caspase-9 is the initiating enzyme of the apoptotic pathway, and caspase-3 is the key

executioner caspase and is activated through cleavage of caspase-9.^[13] The activity of these factors can be indirectly measured at the protein expression by Western blot analysis. Fulda *et al.*^[14] found that resveratrol-induced apoptosis was triggered by the activation of caspases and reversed by pretreatment with the caspase inhibitor Z-VAD-fmk. In this study, the results showed that the expressions of caspase-3 and caspase-9 were decreased by increasing compound **9I** concentrations. Compound **9I** induced apoptosis in gastric cancer cells through dependent caspase ac-

tivation.

3 Conclusions

In summary, based on the active natural products butenolide, an effective anti cancer agent compound **9I** in gastric cancer was found. The results suggested that apoptosis was involved in compound **9I**-induced gastric cancer cell death. The potential mechanisms of compound **9I** have resulted from the induction of apoptosis as evidenced by partially dependent caspase activation. It deserves further investigation into its utility for prevention and treatment of gastric cancer.

4 Experimental

4.1 Chemistry

All solvents and reagents were purchased from commercial sources, generally with no need of drying and/or purification unless necessary. Melting points were determined on a Beijing Fukai X-5 apparatus and were uncorrected. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker DPX-400 spectrometer at 400 and 100 MHz with TMS as internal standard. HRMS was recorded on a Q-ToF Micro HRMS of waters.

4.1.1 Synthesis of 3-(bromomethyl)furan-2(5H)-one (**3**)

Thiomethylphenylammonium tribromide ($\text{Me}_3\text{PhN}^+\text{Br}_3^-$) (13.8 g, 36.7 mmol) was added to the solution of α -methylene- γ -butene lactone (3 g, 30.6 mmol) in dioxane (90 mL). The reaction mixture was stirred for 10 h at room temperature. The reaction was monitored by TLC until all α -methylene- γ -butene lactone was consumed. Upon completion, the reaction mixture was diluted with 250 mL of diethyl ether and filtered. The filtrate was concentrated under reduced pressure. Without purification, the above crude product, Li_2CO_3 (11.1 g, 150 mmol), LiBr (15.6 g, 150 mmol) and tetrabutylammonium bromide (0.966 g, 3 mmol) were suspended in CH_3CN (90 mL). The mixture was stirred at 85°C for 20 h. The reaction was monitored by TLC. After filtering and removing the spare solvent under vacuum, the residue was dissolved in 300 mL of diethyl ether and washed with saturated salt water (150 mL). The organic phase was dried over anhydrous Na_2SO_4 and then concentrated. The concentrate was purified by column chromatography (petroleum ether/acetic ether, $V:V=5:1$) to afford the product **3**.^[15] Yellow oil, yield 4.09 g (76%). ^1H NMR (400 MHz, CDCl_3) δ : 7.55 (s, 1H), 4.89 (d, $J=1.7$ Hz, 2H), 4.13 (d, $J=1.4$ Hz, 2H).

4.1.2 Synthesis of compounds **4** and **5**

Compound **3** (0.5 g, 2.82 mmol) in dry tetrahydrofuran (THF) (5 mL) was added dropwise in four portions to a solution of ethyl- γ -aminobutyrate hydrochloride (0.72 g, 7.23 mmol) and triethylamine (0.86 g, 8.74 mmol) in dry THF (10 mL) at 0°C over 2 h. The reaction mixture was then warmed to room temperature, stirred for another 24 h. After the reaction completion, the solid was filtered off,

and the solvent was removed on a rotary evaporator. The crude product was mixture of compounds **4** and **5**, which were used directly to the next step.

4.1.3 Synthesis of unciniine (**6**)

The mixture of **4**, **5** (0.576 g) and acetone (5 mL) was added to a solution of methyl alcohol (10 mL), and the reaction mixture was stirred at room temperature for 72 hours. The reaction system was filtered and the solvent was evaporated *in vacuo*. The reaction mixture was then diluted with diethyl ether (30 mL $\times$ 3), and washed with H_2O (50 mL). The organic phase was dried over anhydrous Na_2SO_4 , and the solvent was removed. The crude product was purified by column chromatography on silica gel [$V(\text{petroleum ether}):V(\text{acetone})=6:1$] to afford the product **6**^[2] as a white crystalline substance (0.249 g, white solid). Yield 40%. ^1H NMR (400 MHz, CDCl_3) δ : 7.49 (s, 1H), 4.19 (s, 2H), 3.48 (m, 2H), 2.40 (m, 2H), 2.08~2.01 (m, 2H), 2.00 (s, 3H), 1.92 (s, 3H).

4.1.4 Synthesis of compounds **7a**~**7e**

Nitrogen compounds (3 mmol) were diluted with 1 mL of THF and then slowly dropped into the THF solution (1 mL) of compound **3** (400 mg, 2.27 mmol) with a lot of white solid precipitated. The reaction mixture was stirred at room temperature. After completion of the reaction, the reaction system was filtered, concentrated under vacuum and crystallized from ethyl acetate to yield the pure products **7a**~**7e**, which directly used for the next reaction.

4.1.5 General procedure for the synthesis of unciniine analogues **8a**~**8e**

The corresponding compounds **7a**~**7e** were dissolved in methanol and then refluxed with appropriate acetone at 65°C for 1 h. The reaction was monitored by TLC. After the reaction completed, the reaction mixture was evaporated *in vacuo* and purified by column chromatography to afford the pure product **8a**~**8e**.

3-(Morpholinomethyl)-5-(propan-2-ylidene)furan-2(5H)-one (**8a**): Light yellow solid, yield 60%. ^1H NMR (400 MHz, CDCl_3) δ : 6.10 (s, 1H), 3.76~3.67 (m, 4H), 3.44 (s, 2H), 2.55~2.46 (m, 4H), 2.04 (s, 3H), 2.05 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ : 144.8, 134.6, 129.9, 127.6, 122.5, 113.4, 39.9, 29.7, 20.4, 18.7. HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{18}\text{NO}_3$ [$\text{M}+\text{H}$] $^+$ 224.1281, found 224.1292.

3-(Piperidin-1-ylmethyl)-5-(propan-2-ylidene)furan-2(5H)-one (**8b**): Light yellow solid, yield 50%. ^1H NMR (400 MHz, CDCl_3) δ : 7.54 (s, 1H), 3.32 (s, 2H), 2.48 (brs, 4H), 2.02 (s, 3H), 1.96 (s, 3H), 1.63 (brs, 4H), 1.50~1.42 (m, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.8, 144.9, 136.1, 128.6, 121.9, 54.5, 53.1, 29.7, 25.8, 24.1, 18.6. HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{20}\text{NO}_2$ [$\text{M}+\text{H}$] $^+$ 222.1489, found 222.1483.

3-((4-Methylpiperazin-1-yl)methyl)-5-(propan-2-ylidene)furan-2(5H)-one (**8c**): Light yellow solid, yield 63%. ^1H NMR (400 MHz, CDCl_3) δ : 7.48 (s, 1H), 3.35 (s, 2H), 2.57 (brs, 8H), 2.32 (s, 3H), 2.01 (s, 3H), 1.95 (s, 3H). HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{21}\text{N}_2\text{O}_2$ [$\text{M}+\text{H}$] $^+$ 237.1598, found

237.1595.

5-(Propan-2-ylidene)-3-((*p*-tolylamino)methyl)furan-2(5*H*)-one (**8d**): White solid, yield 50%. ^1H NMR (400 MHz, CDCl_3) δ : 7.42 (s, 1H), 6.81 (d, $J=8.9$ Hz, 2H), 6.62 (d, $J=8.9$ Hz, 2H), 4.11 (s, 2H), 2.43 (s, 3H), 2.01 (s, 3H), 1.91 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.1, 152.7, 144.7, 141.2, 134.6, 129.9, 122.5, 114.9, 114.7, 55.7, 40.5, 29.7, 18.7. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 244.1332, found 244.1314.

3-(((4-Methoxyphenyl)amino)methyl)-5-(propan-2-ylidene)furan-2(5*H*)-one (**8e**): Light yellow solid, yield 55%. ^1H NMR (400 MHz, CDCl_3) δ : 7.42 (s, 1H), 7.09 (d, $J=8.9$ Hz, 2H), 6.72 (d, $J=8.9$ Hz, 2H), 4.11 (s, 2H), 3.77 (s, 3H), 2.01 (s, 3H), 1.91 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.5, 144.8, 136.4, 129.6, 127.7, 122.8, 122.7, 114.9, 114.7, 66.7, 52.7, 29.4, 18.7. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 260.1281, found 260.1273.

3-(((4-Methylbenzyl)amino)methyl)-5-(propan-2-ylidene)furan-2(5*H*)-one (**8f**): White solid, yield 70%. ^1H NMR (400 MHz, CDCl_3) δ : 7.42 (s, 1H), 7.02 (d, $J=8.1$ Hz, 2H), 6.57 (d, $J=8.4$ Hz, 2H), 4.13 (s, 2H), 3.79 (s, 2H), 2.26 (s, 3H), 2.01 (s, 3H), 1.90 (s, 3H). HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{20}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 258.1489, found 258.1476.

3-(((4-Methoxybenzyl)amino)methyl)-5-(propan-2-ylidene)furan-2(5*H*)-one (**8g**): Yellow solid, yield 75%. ^1H NMR (400 MHz, CDCl_3) δ : 7.42 (s, 1H), 6.81 (d, $J=8.9$ Hz, 2H), 6.62 (d, $J=8.9$ Hz, 2H), 4.28 (s, 2H), 4.11 (s, 2H), 3.76 (s, 3H), 2.01 (s, 3H), 1.91 (s, 3H). HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{20}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 274.1438, found 274.1467.

4.1.6 General procedure for the synthesis of compounds **9a**~**9p**

In a round-bottomed flask with a magneton, compound **7a** (183 mg, 1 mmol), substituted aldehyde or ketone (1 mmol), ethanediamine (6 mg, 0.1 mmol) and methanol (0.4 mL) were added. The reaction system was stirred at room temperature for 36 h. The progress was detected by thin-layer chromatography (TLC). Upon completion, a lot of solid were filtered, dried and crystallized from methanol.

(*Z*)-3-(Morpholinomethyl)-5-(thiophen-2-ylmethylene)-furan-2(5*H*)-one (**9a1**): Yellow solid, yield 11%. ^1H NMR (400 MHz, CDCl_3) δ : 7.49 (d, $J=5.1$ Hz, 1H), 7.37 (d, $J=3.7$ Hz, 1H), 7.33 (s, 1H), 7.07 (dd, $J=5.0$, 3.8 Hz, 1H), 6.28 (s, 1H), 3.77~3.69 (m, 4H), 3.36 (s, 2H), 2.56~2.48 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.6, 145.6, 140.0, 136.2, 131.0, 130.4, 128.7, 127.8, 106.8, 66.9, 53.6, 52.9. HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{16}\text{NO}_3\text{S}$ $[\text{M}+\text{H}]^+$ 278.0845, found 278.0856.

(*Z*)-3-(Morpholinomethyl)-5-(thiophen-3-ylmethylene)-furan-2(5*H*)-one (**9a2**): Yellow solid. ^1H NMR (400 MHz, CDCl_3) δ : 7.72 (t, $J=3.9$ Hz, 1H), 7.52 (dd, $J=11.2$, 7.0 Hz, 1H), 7.34 (dd, $J=4.3$, 2.1 Hz, 1H), 7.33 (s, 1H), 6.09 (s, 1H), 3.77~3.70 (overlap, 4H), 3.36 (s, 2H), 2.52 (brs, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.0, 146.4, 140.8, 134.4, 128.8, 128.5, 128.2, 126.2, 107.4, 66.9, 53.6, 52.8.

HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{16}\text{NO}_3\text{S}$ $[\text{M}+\text{H}]^+$ 278.0845, found 278.0857.

(*Z*)-5-((5-Methylfuran-2-yl)methylene)-3-(morpholinomethyl)furan-2(5*H*)-one (**9a3**): Brown solid, yield 75%. m.p. 114~115 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.05 (s, 1H), 6.46 (d, $J=3.2$ Hz, 1H), 6.34 (s, 1H), 6.11 (d, $J=3.1$ Hz, 1H), 3.79~3.76 (m, 4H), 3.43 (s, 2H), 2.61~2.56 (m, 4H), 2.42 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.6, 155.3, 147.9, 146.1, 138.4, 116.6, 108.9, 102.6, 66.9, 53.6, 52.9, 14.8. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 276.1230, found 276.1238.

(*Z*)-5-Benzylidene-3-(morpholinomethyl)furan-2(5*H*)-one (**9b**): White solid, yield 30%. m.p. 145~146 °C; ^1H NMR (400 MHz, CDCl_3) 7.79 (s, 1H), 7.35~7.38 (overlap, 5H), 6.02 (s, 1H), 3.80~3.73 (m, 4H), 3.40 (s, 2H), 2.61~2.52 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.2, 147.3, 141.4, 133.0, 130.4, 129.1, 128.9, 128.9, 128.7, 113.5, 66.9, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{18}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 272.1281, found 272.1273.

(*Z*)-5-(2-Chlorobenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9c**): White solid, yield 35%. m.p. 147~148 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.34~7.36 (overlap, 5H), 6.51 (s, 1H), 3.87~3.68 (m, 4H), 3.40 (s, 2H), 2.66~2.45 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.9, 148.4, 141.5, 134.3, 132.0, 130.9, 130.0, 129.6, 127.3, 108.5, 66.9, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{17}\text{ClNO}_3$ $[\text{M}+\text{H}]^+$ 306.0891/308.0862, found 306.0886/308.0851.

(*Z*)-5-(3-Chlorobenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9d**): White solid, yield 78%. m.p. 114~115 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (s, 1H), 7.67 (dd, $J=7.0$, 1.6 Hz, 1H), 7.40~7.31 (m, 3H), 5.95 (s, 1H), 3.81~3.73 (m, 4H), 3.40 (s, 2H), 2.61~2.53 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.8, 148.1, 141.1, 134.7, 130.0, 129.6, 129.0, 128.5, 111.7, 66.9, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{17}\text{ClNO}_3$ $[\text{M}+\text{H}]^+$ 306.0891/308.0862, found 306.0886/308.0851.

(*Z*)-5-(4-Chlorobenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9e**): White solid, yield 60%. m.p. 113~114 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.73 (d, $J=8.6$ Hz, 2H), 7.38 (d, $J=8.6$ Hz, 2H), 7.29 (s, 1H), 5.98 (s, 1H), 3.82~3.65 (m, 4H), 3.40 (s, 2H), 2.56 (brs, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.9, 147.6, 141.2, 135.1, 131.6, 129.1, 112.0, 66.9, 61.0, 56.3, 54.0, 52.8. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{17}\text{ClNO}_3$ $[\text{M}+\text{H}]^+$ 306.0891/308.0862, found 306.0886/308.0851.

(*Z*)-5-(2-Methoxybenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9f**): Light yellow solid, yield 40%. m.p. 98~99 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.73 (d, $J=8.6$ Hz, 2H), 7.38 (overlap, 3H), 5.97 (s, 1H), 3.83 (s, 3H), 3.81~3.73 (m, 4H), 3.40 (s, 2H), 2.64~2.51 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.1, 159.8, 147.5, 141.3, 134.2, 129.8, 128.8, 123.3, 115.3, 113.3, 66.9, 55.4, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 302.1387, found 302.1373.

(*Z*)-5-(3-Methoxybenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9g**): Yellow solid, yield 67%. m.p. 96~

97 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.32~7.36 (overlap, 4H), 6.91 (d, $J=8.1$ Hz, 1H), 5.99 (s, 1H), 3.87 (s, 3H), 3.81~3.71 (m, 4H), 3.39 (s, 2H), 2.61~2.52 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.1, 159.8, 147.5, 141.3, 134.2, 129.8, 128.8, 123.3, 115.3, 113.3, 66.9, 55.4, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 302.1387, found 302.1373.

(*Z*)-5-(4-methoxybenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9h**): Light yellow solid, yield 60%. m.p. 95~96 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.76(d, $J=8.6$ Hz, 2H), 7.36 (d, $J=8.6$ Hz, 2H), 6.93 (s, 1H), 5.97 (s, 1H), 3.83 (s, 3H), 3.82~3.65 (m, 4H), 3.40 (s, 2H), 2.56 (brs, 4H). ^{13}C NMR (101 MHz, CDCl_3) δ : 170.6, 160.4, 145.4, 141.6, 132.3, 127.1, 125.4, 113.6, 66.9, 61.0, 56.3, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 302.1387, found 302.1373.

(*Z*)-3-(morpholinomethyl)-5-(3,4,5-trimethoxybenzylidene)furan-2(5*H*)-one (**9i**): Yellow solid, yield 50%. m.p. 120~121 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.32 (s, 1H), 7.02 (s, 2H), 5.91 (s, 1H), 3.90 (d, $J=6.7$ Hz, 9H), 3.71~3.74 (overlap, 4H), 3.37 (s, 2H), 2.57~2.48 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.1, 153.3, 146.8, 141.3, 139.4, 128.5, 128.1, 113.5, 107.9, 66.9, 61.0, 56.3, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{NO}_6$ $[\text{M}+\text{H}]^+$ 362.1598, found 362.1587.

(*Z*)-5-(2,3-dimethoxybenzylidene)-3-(morpholinomethyl)furan-2(5*H*)-one (**9j**): White solid, yield 46%. m.p. 108~109 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.84 (d, $J=8.1$ Hz, 1H), 7.42 (s, 1H), 7.12 (t, $J=8.1$ Hz, 1H), 6.93 (d, $J=7.0$ Hz, 1H), 6.52 (s, 1H), 3.91 (s, 3H), 3.89 (s, 3H), 3.79~3.74 (m, 4H), 3.40 (s, 2H), 2.60~2.53 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.3, 152.6, 147.9, 141.8, 128.5, 127.2, 123.1, 113.3, 107.2, 61.4, 55.8, 53.6, 52.8, 29.7. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{22}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 332.1492, found 332.1490.

(*Z*)-5-(4-fluorobenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9k**): Light yellow solid, yield 77%. m.p. 116~117 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.79 (m, 2H), 7.37 (s, 1H), 7.11 (m, 2H), 5.98 (s, 1H), 3.81~3.70 (m, 4H), 3.39 (s, 2H), 2.61~2.50 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.1, 164.2, 161.7, 147.0, 141.2, 132.5, 129.3, 128.6, 116.1, 115.9, 112.1, 66.9, 53.6, 52.8. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{17}\text{FNO}_3$ $[\text{M}+\text{H}]^+$ 290.1187, found 290.1171.

(*Z*)-5-(4-(*tert*-Butyl)benzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9l**): White solid, yield 55%. m.p. 145~146 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.81~7.67 (d, $J=8.5$ Hz, 2H), 7.44 (d, $J=8.5$ Hz, 2H), 7.29 (s, 1H), 6.09 (s, 1H), 3.92~3.78 (m, 4H), 3.52 (s, 2H), 2.72 (brs, 4H), 1.35 (s, 9H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.2, 150.5, 147.2, 137.5, 130.7, 129.9, 128.0, 126.3, 66.9, 53.6, 52.8, 34.1, 29.7, 23.8. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{26}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 328.1907, found 328.1918.

(*Z*)-5-(4-Isopropylbenzylidene)-3-(morpholinomethyl)-furan-2(5*H*)-one (**9m**): Light yellow solid, yield 47%. m.p. 98~99 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.73 (d, $J=8.3$ Hz, 2H), 7.37 (s, 1H), 7.28 (d, $J=8.4$ Hz, 2H), 6.01 (s,

1H), 3.75 (overlap, 4H), 3.39 (s, 2H), 3.03~2.85 (m, 1H), 2.54 (overlap, 4H), 1.26 (d, $J=8.3$ Hz, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.4, 150.5, 146.8, 141.5, 130.7, 128.0, 127.0, 113.7, 53.6, 52.8, 34.1, 29.7, 23.8. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 314.1751, found 314.1755.

(*Z*)-5-(4-(4-methylpiperazin-1-yl)benzylidene)-3-(morpholinomethyl)furan-2(5*H*)-one (**9n**): Yellow solid, yield 16%. ^1H NMR (400 MHz, CDCl_3) δ : 7.73~7.67 (d, $J=9.0$ Hz, 2H), 7.31 (s, 1H), 6.88 (d, $J=9.0$ Hz, 2H), 5.93 (s, 1H), 3.77~3.70 (overlap, $J=4.1$ Hz, 4H), 3.36 (s, 2H), 3.31~3.34 (overlap, $J=11.8$, 6.7 Hz, 4H), 2.57 (brd, 6.7 Hz, 4H), 2.52 (brd, 4H), 2.47 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.3, 141.5, 132.2, 130.4, 126.1, 123.7, 115.4, 114.9, 66.9, 54.8, 53.6, 53.6, 48.0, 47.7, 46.1. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 370.2125, found 370.2134.

(*Z*)-5-(3-Fluoro-4-(4-methylpiperazin-1-yl)benzylidene)-3-(morpholinomethyl)furan-2(5*H*)-one (**9o**): Yellow solid, yield 19%. ^1H NMR (400 MHz, CDCl_3) δ : 7.57 (dd, $J=14.4$, 1.9 Hz, 1H), 7.39 (dd, $J=8.5$, 1.7 Hz, 1H), 7.31 (s, 1H), 6.92 (t, $J=20.5$, 8.9 Hz, 1H), 5.90 (s, 1H), 3.76~3.70 (overlap, 4H), 3.37 (s, 2H), 3.24~3.15 (overlap, 4H), 2.64~2.57 (m, 4H), 2.55~2.49 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 170.2, 146.5, 141.2, 136.2, 133.4, 127.4, 118.4, 112.5, 66.9, 55.0, 53.7, 53.6, 52.8, 49.9, 46.1. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{27}\text{FN}_3\text{O}_3$ $[\text{M}+\text{H}]^+$ 388.2031, found 388.2035.

5-Cyclohexylidene-3-(morpholinomethyl)furan-2(5*H*)-one (**9p**): Light yellow solid, yield 42%. m.p. 109~110 °C; ^1H NMR (400 MHz, DMSO) δ : 7.96 (s, 1H), 3.61~3.55 (overlap, 4H), 3.22 (s, 2H), 2.41 (overlap, 8H), 1.59 (s, 6H); ^{13}C NMR (101 MHz, DMSO) δ : 170.3, 142.3, 137.8, 130.4, 127.5, 66.6, 53.5, 52.5, 28.7, 28.1, 27.3, 26.1. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{22}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 264.1594, found 264.1591.

4.1.7 Synthesis of (*Z*)-3-(4-(*tert*-butyl)benzylidene)-indolin-2-one (**11**)

To a solution of indolin-2-one (200 mg, 1.5 mmol) in CH_3OH (1 mL) was added 4-(*tert*-butyl)benzaldehyde (365.6 mg, 2.25 mmol), and the mixture was stirred with ethylenediamine (10 mg) for 10 h at room temperature. Then, the system was concentrated to remove CH_3OH and purified by silica gel column chromatography (ethyl acetate/petroleum ether, $V:V=1:7$) to give 3-(4-(*tert*-butyl)benzylidene)indolin-2-one (**11**), 68 mg, yield 16% (yellow solid). ^1H NMR (400 MHz, CDCl_3) δ : 10.59 (s, 1H), 7.66 (overlap, 3H), 7.61 (s, 1H), 7.55 (d, $J=8.3$ Hz, 2H), 7.24 (d, $J=7.8$ Hz, 1H), 6.88 (t, $J=7.9$ Hz, 2H), 1.33 (s, 9H); ^{13}C NMR (101 MHz, CDCl_3) δ : 174.0, 158.0, 148.1, 141.0, 136.8, 135.2, 134.6, 132.1, 130.8, 127.6, 126.3, 126.2, 115.3, 39.9, 36. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{NO}$ $[\text{M}+\text{H}]^+$ 278.1539, found 278.1544.

4.1.8 Synthesis of (*Z*)-3-(4-(*tert*-butyl)benzylidene)-1-((tetra-hydro-2*H*-pyran-4-yl)methyl)indolin-2-one (**12**)

Under ice bath conditions, to a solution of 3-(4-(*tert*-

butyl)benzylidene)indolin-2-one (25 mg, 0.09 mmol) in DMF (1 mL) was added NaH (7.22 mg, 0.3 mmol), and the mixture was stirred for 30 min. Then, 4-(bromomethyl)-tetrahydro-2H-pyran (24 μ L, 0.16 mmol) was added to the system. After reacting for 9 h, the mixture was diluted into water (10 mL) and extracted with ethyl acetate (10 mL $\times$ 4). The combined organic layers were dried over MgSO_4 , concentrated, and purified by silica gel column chromatography [$V(\text{acetone}) : V(\text{petroleum ether}) = 1 : 5$] to give 3-(4-(*tert*-butyl)-benzylidene)-1-(tetrahydro-2H-pyran-4-yl)-indolin-2-one (**12**), 14 mg, yield 44% (yellow oil). ^1H NMR (400 MHz, CDCl_3) δ : 7.43 (d, $J = 7.9$ Hz, 2H), 7.38 (d, $J = 8.2$ Hz, 2H), 7.25 (t, $J = 7.2$ Hz, 1H), 6.88 (d, $J = 7.8$ Hz, 1H), 6.78 (t, $J = 7.8$ Hz, 1H), 6.55 (d, $J = 7.5$ Hz, 1H), 4.80 (s, 1H), 3.99 (d, $J = 10.2$ Hz, 2H), 3.67 (d, $J = 7.2$ Hz, 2H), 3.35 (t, $J = 11.5$ Hz, 2H), 2.12~2.08 (m, 1H), 1.63 (d, $J = 12.8$ Hz, 2H), 1.53~1.44 (m, 2H), 1.33 (s, 9H). HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{30}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 376.2271, found 376.2278.

4.2 Biological evaluation

4.2.1 Material

Fetal bovine serum (FBS), Roswell Park Memorial Institute 1640 (RPMI-1640) medium, MEM, DMEM, and penicillin-streptomycin were purchased from HyClone (Victoria, Australia). MTT, NAC, and Z-VAD-fmk were purchased from Selleck Chemicals (Houston, TX). The FITC/Annexin V Apoptosis Detection Kit was purchased from BestBio (Shanghai, China). DCFH-DA and diaminobenzidine (DAB) were purchased from Beyotime Biotechnology (Shanghai, China). Antibodies specific for actin, caspase-3, and PARP were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Antibodies specific for caspase-9 was purchased from Cell Signaling Technology (Danvers, MA). The antibody specific for survivin, cleaved caspase-3, and cleaved caspase-9 were purchased from Abcam (Cambridge, MA). Peroxidase labeled anti-rabbit, anti-mouse, and anti-goat polyclonal immunoglobulins were purchased from Bioss (Shanghai, China). The enhanced chemiluminescence (ECL) kit was purchased from Thermo Fisher Scientific (Waltham, MA).

All the tested compounds were dissolved in dimethyl sulfoxide (DMSO) to 10 mmol/L stock solutions and stored at $-20\text{ }^\circ\text{C}$. For each experiment, the stock solutions were diluted in the culture medium to obtain the desired concentration. The final DMSO content in cell culture was $\leq 0.5\%$ (volume fraction), which was found to be nontoxic to cells.

4.2.2 Cell lines and cultures

Human gastric cancer cell lines (MGC803, HGC27, and SGC7901) and human gastric epithelial cells (GES1) were obtained from the American Type Culture Collection (ATCC, Manassas, VA) and cultured at $37\text{ }^\circ\text{C}$ in an atmosphere containing $5\% \text{CO}_2$, with 10% heat-inactivated fetal bovine serum supplemented with RPMI1640 medium, 100 U/mL penicillin, and 0.1 mg/mL streptomycin. All cell lines were from ATCC and were cultured under standard

conditions as recommended by the supplier.

4.2.3 Analysis of cell proliferation

For MGC803 cells, cell survival rates were determined by MTT assay. The cells were seeded in a 96-well plate (4000 well) for 24 h, followed by adding solution of compounds ($200\text{ }\mu\text{mol/L}$) to the respective wells in the indicated final concentrations and incubated for 24 h. Then, $20\text{ }\mu\text{L}$ of 5 mg/mL MTT was added to the medium, and the cells were incubated for 4 h at $37\text{ }^\circ\text{C}$ and $5\% \text{CO}_2$. After removal of the culture medium, $150\text{ }\mu\text{L}$ of DMSO was added to dissolve the formazan crystals. The absorbance was read by an enzyme labeling instrument with a 490-nm wavelength resolution. The viability of untreated cells was set as 100% , and the viability of the other groups was calculated by comparing the optical density reading of the control cells. The IC_{50} values were calculated by nonlinear regression analysis.

4.2.4 Apoptosis analysis

The cells were treated with different concentrations of compound **91** for 24 h. Cells were collected and washed with PBS twice, and incubated with fluorescein isothiocyanate (FITC) conjugated annexin V by use of an annexin V/FITC apoptosis kit following the step-by-step protocol provided by the manufacturer. Annexin V+/PI- cells were considered as early apoptotic and Annexin V+/PI+ cells as late apoptotic/necrotic.

4.2.5 Western blot analysis

MGC803 cells were seeded in a 100-mm tissue culture plate, cultivated for 24 h, and then treated with different concentrations of compound **91** for 24 h. After treatment, cells were collected and lysed with ice-cold lysis buffer (Beyotime, Shanghai, China). Protein concentrations of the lysates were determined by a micro-BCA protein assay kit. After centrifugation at 12000 r/min for 30 min, the total cellular protein extracts were boiled with $5\times$ loading buffer, separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a nitrocellulose membrane. After blocking with 5% skimmed milk in PBST for 2 h, membranes were incubated with appropriate antibodies overnight at $4\text{ }^\circ\text{C}$, followed by horseradish peroxidase (HRP) conjugated anti-mouse, anti-goat, or anti-rabbit secondary antibodies. The detection of specific proteins was carried out with an enhanced chemiluminescence (ECL) Western blot kit according to the recommended procedure.

Images were captured on a Canon image scanner (LIDE110, Japan). The relative protein levels were calculated by use of β -actin protein as an endogenous control and quantified by Image J software.

4.2.6 Statistical analysis

The data are expressed as means $\pm$ SD. Significant differences between the groups were determined by the unpaired Student's *t*-test. $*P < 0.05$ and $**P < 0.01$ were accepted as indications of statistical significance.

Supporting Information The NMR spectra for synthetic compounds. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Manchoju, A.; Annadate, R. A.; Desquien, L.; Pansare, S. V. *Org. Biomol. Chem.* **2018**, *16*, 6224.
- [2] FÁKOVÁ, H.; Pour, M.; Kuneš, J.; Šenel, P. *Tetrahedron Lett.* **2005**, *46*, 8137.
- [3] David, S.; Peter, B.; Andreas, R.; Wilhelm, S. *Toxicol. Lett.* **2018**, *295*, 369.
- [4] Dasmahapatra, G.; Lembersky, D.; Son, M. P.; Patel, H.; Peterson, D.; Attkisson, E.; Fisher, R. I.; Friedberg, J. W.; Dent, P.; Grant, S. *Mol. Cancer Ther.* **2019**, *18*, 1180.
- [5] Germain, A. D. S.; Clavé, G.; Badetdenisot, M. A.; Pillot, J. P.; Cornu, D. *Nat. Chem. Biol.* **2016**, *12*, 1552.
- [6] Xu, H. W.; Zhao, L. J.; Liu, H. F.; Zhao, D.; Luo, J.; Xie, X. P.; Liu, W. S.; Zheng, J. X.; Dai, G. F.; Liu, H. M.; Liu, L. H.; Liang, Y. B. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 2388.
- [7] Xu, H. W.; Jia, S. L.; Xie, X. P.; Luo, J.; Wang, S. H. *Chin. J. Org. Chem.* **2017**, *37*, 902 (in Chinese). (徐海伟, 贾世龙, 谢晓平, 罗姣, 王树, 有机化学, **2017**, *37*, 902.)
- [8] Necula, L.; Matei, L.; Dragu, D.; Neagu, A. I.; Mambet, C.; Nedeianu, S.; Bleotu, C.; Diaconu, C. C.; Chivu-Economescu, M. *World J. Gastroenterol.* **2019**, *17*, 2029.
- [9] Xu, H. W.; Liu, G. Z.; Zhu, S. L.; Hong, G. F.; Liu, H. M.; Wu, Q. *Steroids* **2010**, *75*, 419.
- [10] Xu, H. W.; Wang, J. F.; Liu, G. Z.; Hong, G. F.; Liu, H. M. *Org. Biomol. Chem.* **2007**, *5*, 1247.
- [11] Caroppi, P.; Sinibaldi, F.; Fiorucci, L.; Santucci, R. *Curr. Med. Chem.* **2009**, *16*, 4058.
- [12] Fulda, S.; Debatin, K. M. *Oncogene* **2006**, *25*, 4798.
- [13] Danial, N. N.; Korsmeyer, S. J. *Cell* **2004**, *116*, 205.
- [14] Fulda, S.; Debatin, K. M. *Eur. J. Cancer* **2005**, *41*, 786.
- [15] Cui, Q.; Wang, J. A.; Yang, H. S.; Xie, L. G. *Chin. J. Org. Chem.* **2010**, *30*, 1705 (in Chinese). (崔庆, 王娇, 杨海申, 谢龙观, 徐效华, 有机化学, **2010**, *30*, 1705.)

(Lu, Y.)