NN 型双齿半夹心 Ru(II)化合物: 无受体脱氢环化合成喹啉和吡咯衍生物的催化剂

胡博文* 张宇哲 尹鸽平 陈大发

(哈尔滨工业大学化工与化学学院 哈尔滨 150001)

摘要 四个含 NN 型双齿配体的半夹心(η^6 -*p*-cymene)Ru(II)化合物被成功制备. 这四个化合物分别为(η^6 -*p*-cymene)-Ru(C₅H₄N-C₅H₃N-OH) (**1**), (η^6 -*p*-cymene)Ru(C₅H₄N-CH₂-C₅H₄N) (**2**), (η^6 -*p*-cymene)Ru(C₅H₄N-CH₂-C₅H₃N-OH) (**3**)和(η^6 -*p*-cymene)Ru(C₅H₄N-CH₂-C₅H₃N-OCH₃) (**4**). 这些化合物通过核磁氢谱、碳谱和元素分析等手段表征, 化合物 **2** 的结构被 X 射线单晶衍射证实. 将这些化合物应用于催化氨醇与酮的环化反应, 其中 **3** 的催化效率最高. 在 0.5 mol% 化合物 **3** 的存在下, 制备了一系列喹啉和吡啶衍生物.

关键词 半夹心; 钌; 无受体脱氢环化; 双齿配体; 2-羟基吡啶

Half-Sandwich Ruthenium(II) Complexes with Bidentate NN Ligands: Active Catalysts for the Synthesis of Quinolines and Pyrroles by Acceptorless Dehydrogenative Cyclization

Hu, Bowen* Zhang, Yuzhe Yin, Geping Chen, Dafa*

(School of Chemistry and Chemical Engineering, Harbin Institute of Technology, Harbin 150001)

Abstract Four (η^6 -*p*-cymene)Ru(II) complexes with bidentate NN ligands, (η^6 -*p*-cymene)Ru(C₅H₄N-C₅H₃N-OH) (**1**), (η^6 -*p*-cymene)Ru(C₅H₄N-CH₂-C₅H₄N) (**2**), (η^6 -*p*-cymene)Ru(C₅H₄N-CH₂-C₅H₃N-OH) (**3**) and (η^6 -*p*-cymene)Ru(C₅H₄N-CH₂-C₅H₃N-OCH₃) (**4**) were prepared. These complexes were all characterized by ¹H NMR, ¹³C NMR and elemental analysis, and complex **2** was further determined by single crystal crystallography. Complexes **1**~**4** were treated as catalysts for cyclizations of amino alcohols with ketones, and complex **3** exhibited the highest activity. The cyclization reactions proceeded in toluene with 0.5 mol% catalyst loading, and a series of quinolines and pyrroles were synthesized.

Keywords half-sandwich; ruthenium; acceptorless dehydrogenative cyclization; bidentate ligand; 2-hydroxypyridine

1 Introduction

The synthesis of *N*-heterocycles has drawn much attention, because *N*-heterocycles exist widely in natural products and pharmaceuticals, and they can be used for the preparation of functional materials with improved physical properties.^[1-9] However, traditional methods to *N*-heterocycles always need multiple steps and harsh reaction conditions, and sometimes the raw materials are necessary.^[10-13] Thus, there is an increasing demand for developing atom-economical, sustainable and environmental friendly

methodologies for the preparation of *N*-heterocycles.

Acceptorless dehydrogenation condensation (ADC) of amino alcohols with ketones or secondary alcohols has become an alternative method for the sustainable synthesis of *N*-heterocycles, and the only by-products are H₂ and H₂O.^[14-20] Recently, many transition-metal catalysts, such as Ir,^[21-24] Ru,^[25-36] Mn,^[37,38] Co,^[39,40] Ni,^[41-43] Cu,^[44] and Fe^[45] complexes, have been applied for such transformation. Among these reported catalysts, the ruthenium complexes, especially those with a proton-responsive site, are highly effective.^[29-36] For example, the groups of Mil-

* Corresponding author. E-mail: bowenhu@hit.edu.cn

Received August 11, 2019; revised September 16, 2019; published online September 25, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 21672045, 21871068), the China Postdoctoral Science Foundation (No. 2016M591519), the Heilongjiang Provincial Postdoctoral Foundation (No. LBH-Z16069), and the National Natural Science Foundation of Heilongjiang Province (Nos. B2018005, LC2018005).

国家自然科学基金(Nos. 21672045, 21871068)、中国博士后基金(No. 2016M591519)、黑龙江省博士后基金(No. LBH-Z16069)及黑龙江省自然科学基金(Nos. B2018005, LC2018005)资助项目.

stein,^[29,30] Sun,^[31] Yu,^[32] Kundu^[33] and Lang^[34] have independently reported Ru(II) complexes for the synthesis of pyridines, quinolines and pyrroles (Figure 1).

Figure 1 Typical Ru(II) catalysts with a proton-responsive site for the synthesis of *N*-heterocycles

Recently, our group also developed an effective Ru(II) complex with a 2-hydroxypyridyl fragment (Scheme 1).^[34] As an update of our continuous interest in H_2 -relating reactions,^[46–52] herein we present four (η^6 -*p*-cymene)Ru(II) complexes **1**~**4** with bidentate NN ligands for the catalytic reactions of amino alcohols with ketones for the preparation of quinolines and pyrroles, and the most efficient catalyst is complex **3** bearing a 2-hydroxypyridyl fragment.

2 Results and discussion

2.1 Synthesis and characterization of the Ru(II) complexes

Complex **1** was synthesized by the reaction of $[RuCl_2(p\text{-cymene})]_2$ with **L**₁ in CH_3OH , followed by the addition of NH_4PF_6 (Scheme 1). The ^1H NMR spectrum of **1** in $CDCl_3$ shows several signals between δ 9.40 and 5.76 for its aromatic protons, one multiplet at δ 2.45 and two doublets at δ 0.90 and 0.89 for its isopropyl group, and one singlet at δ 2.19 for the methyl group. The isopropyl group exhibits as two doublets and one multiplet, indicating the different chemical environments of its two methyl groups, which is caused by the asymmetry of the ligand **L**₁.

Complexes **2**~**4** were synthesized following a similar procedure as described for **1** (Scheme 1). When **L**₂~**L**₄ were treated with $[RuCl_2(p\text{-cymene})]_2$ in refluxing methanol for 3 h, followed by the addition of NH_4PF_6 , complexes **2**~**4** were isolated in 75%, 80% and 82% yields, respectively. Similar to complex **1**, each of the isopropyl groups in complexes **3** and **4** reveals three signals in the ^1H NMR spectrum. While differently, the isopropyl signals of complex **2** are presented as one multiplet and one doublet, due to

Scheme 1 Synthesis of complexes **1**~**4**

the symmetrical **L**₂ ligand.

The molecular structure of **2** (Figure 1) was further confirmed by X-ray crystallography. In the solid state, it is a cationic complex with the central Ru atom coordinating with one *p*-cymene group, one Cl atom and two pyridyl groups. Ru(1)—Cl(1), Ru(1)—N(1) and Ru(1)—N(2) bond distances are 0.2397(10), 0.2106(3) and 0.2110(3) nm, respectively, comparable to those in Ru-PPh₃ complexes.^[42–44] The two pyridine rings of the dipyriddy ligands are not coplanar, and C(5)—C(6)—C(7) angle is 113.2(3)°.

Figure 1 Molecular structure of complex **2** with all atoms labeled

All hydrogens and the hexafluorophosphate anion are omitted for the sake of clarity. Selected bond lengths (nm) and angles (°): Ru(1)—N(1) 0.2106(3), Ru(1)—N(2) 0.2110(3), Ru(1)—Cl(1) 0.2397(10), Ru(1)—C(12) 0.2216(4), Ru(1)—C(13) 0.2180(4), Ru(1)—C(14) 0.2190(4), Ru(1)—C(15) 0.2241(4), Ru(1)—C(16) 0.2213(4), Ru(1)—C(17) 0.2171(4), N(1)—Ru(1)—N(2) 84.86(13), N(1)—Ru(1)—Cl(1) 85.19(10), N(2)—Ru(1)—Cl(1) 85.65(10), C(5)—C(6)—C(7) 113.2(3)

2.2 Catalysis

To find the optimal conditions for the synthesis of 2-phenylquinoline, the reaction of *o*-aminobenzyl alcohol with acetophenone was selected as the model reaction (Table 1). In the beginning, KOH was chosen as the base. When complex **1** was selected as the catalyst, after 2 h at 120 °C in toluene, 2-phenylquinoline was isolated in 68% yield (Table 1, Entry 1). Similarly, in the same conditions, complexes **2**~**4** afforded 2-phenylquinoline in 76%, 81% and 71% yields, respectively, suggesting that **3** is the most efficient catalyst (Table 1, Entries 2~4). Weaker bases, such as K_2CO_3 and CS_2CO_3 , gave much worse results (Table 1, Entries 5 and 6). KO^tBu showed similar activity as KOH.

Table 1 Optimization of reaction conditions for the synthesis of 2-phenylquinoline^a

Entry	Cat.	Base	Yield ^b /%
1	1	KOH	68
2	2	KOH	76
3	3	KOH	81
4	4	KOH	71
5	3	K ₂ CO ₃	13
6	3	Cs ₂ CO ₃	9
7	3	KO ^t Bu	81
8	—	KOH	Trace
9 ^c	3	KOH	41

^a Conditions: *o*-aminobenzyl alcohol (2 mmol), acetophenone (2.4 mmol), base (2 mmol), toluene (2 mL), catalyst (0.5 mol%), 120 °C, 2 h. ^b Isolated yield. ^c Catalyst **3** (0.3 mol%).

(Table 1, Entry 7). In the absence of catalyst, only trace of the product was detected (Table 1, Entry 8). Thus, the con-

ditions of Entry 3 in Table 1 were selected as the optimal conditions (2 mmol of *o*-aminobenzyl alcohol, 2.4 mmol of acetophenone, 2 mmol of KOH, 2 mL of toluene, and 0.5 mol% complex **3** for 2 h under nitrogen). When the amount catalyst **3** was decreased to 0.3 mol%, the isolated yield of the product was only 41% (Table 1, Entry 9).

Having established the optimal conditions, the scope of the reaction with respect to ketones was examined, and the results are shown in Table 2. The electronic influence on the phenyl ring of acetophenone is not remarkable. When *o*-methyl, *m*-methyl and *p*-methoxy substituted acetophenones reacted with *o*-aminobenzyl alcohol, the corresponding 2-phenylquinoline derivatives were isolated in 83%, 73% and 83% yields, respectively (Table 2, Entries 2~4). Similarly, when *m*-fluoro, *m*-bromo, *p*-fluoro and *p*-chloro substituted acetophenones reacted with *o*-aminobenzyl alcohol, the corresponding products were in 70%~82% isolated yields (Table 2, Entries 5~8). Furthermore, 1-(2-chlorophenyl)ethan-1-one and propiophenone were also tested for the reaction, while the yield decreased dramatically owing to the steric hindrance (Table 2,

Table 2 Cyclization of *o*-aminobenzyl alcohol with a range of ketones^a

Entry	<i>o</i> -Aminobenzyl alcohol	Ketone	Product	Yield ^b /%
1				81
2				83
3				73
4				83
5				75
6				82
7				70

Continued

Entry	<i>o</i> -Aminobenzyl alcohol	Ketone	Product	Yield ^b /%
8				73
9				47
10				45

^a Conditions: *o*-aminobenzyl alcohol (2 mmol), ketone (2.4 mmol), KOH (2 mmol), toluene (2 mL), catalyst **3** (0.5 mol %), 120 °C, 2 h. ^b Isolated yield.

Table 3 Cyclization of β -amino alcohol with a series of ketones^a

Entry	β -Amino alcohol	Ketone	Product	Yield ^b /%
1				50
2				49
3				40
4				41
5				41
6				53
7				36

^a Conditions: β -amino alcohol (2 mmol), ketone (2.4 mmol), KOH (2 mmol), toluene (2 mL), catalyst **3** (0.5 mol %), 120 °C, 2 h. ^b Isolated yield.

Entries 9, 10).

In addition, the reactions of β -amino alcohols with acetophenone derivatives were also conducted, and a series of substituted pyrroles were obtained (Table 3). Compared to

the synthesis of 2-phenylquinoline, the yields of pyrrole derivatives were much lower, and there were always some unidentified by-products formed. The yields were not improved even with longer reaction time. 2-Amino-1-butanol

reacted with acetophenone, *p*-methoxyacetophenone, and propiophenone, giving corresponding pyrroles in 50%, 49%, and 40% isolated yields, respectively (Table 3, Entries 1~3). Similarly, the reactions of 2-amino-3-methyl-1-pentanol with acetophenone, *p*-methoxyacetophenone, *p*-chloroacetophenone and propiophenone, generated the desired products in 36%~53% yields (Table 3, Entries 4~7). Unexpectedly, when *p*-chloroacetophenone was treated with 2-amino-1-butanol, although the expected product was generated, it could not be purified because there was an unidentified species with close polarity.

3 Conclusions

In summary, four (η^6 -*p*-cymene)Ru(II) complexes **1**~**4** bearing NN bidentate ligands were synthesized and applied to the synthesis of quinolines and pyrroles. These ruthenium complexes are all active, especially for the preparation of quinolines, and complex **3** with 2-hydroxypyridyl fragment appeared the best activity. The research exhibits the potential for the construction of heterocycles via acceptorless dehydrogenative reactions catalyzed by bidentate (η^6 -*p*-cymene)Ru(II) complexes.

4 Experimental

4.1 General considerations

All manipulations were carried out under an inert nitrogen atmosphere using a Schlenk line. Solvents were distilled from appropriate drying agents under N₂ before use. All reagents were purchased from commercial sources. Liquid compounds were degassed by standard freeze-pump-thaw procedures prior to use. [RuCl₂(*p*-cymene)]₂,^[53] 6-hydroxy-2,2'-bipyridine (**L**₁),^[54] 2-(2-pyridylmethyl)pyridine (**L**₂),^[55] 6-(pyridin-2-ylmethyl)pyridin-2-ol (**L**₃),^[51] and 2-methoxy-6-(pyridin-2-ylmethyl)pyridine (**L**₄),^[51] were prepared as previously, respectively.

The ¹H NMR and ¹³C NMR spectra were recorded on a Bruker Avance 400 spectrometer. The ¹H NMR chemical shifts were referenced to residual solvent as determined relative to Me₄Si (δ =0). The ³¹P NMR chemical shifts were reported relative to external 85% H₃PO₄. The ¹³C NMR chemical shifts were reported relative to the carbon resonance of CDCl₃ (δ 77.0) or DMSO-*d*₆ (δ 39.5). Elemental analyses were performed on a PerkinElmer 240C analyzer. X-ray diffraction studies were carried out in a SuperNova X-ray single crystal diffractometer or a Bruker D8 Quest X-ray diffractometer. Data collections were performed using four-circle kappa diffractometers equipped with CCD detectors. Solution, refinement, and geometrical calculations for all crystal structures were performed by SHELXTL.

4.1.1 Synthesis of **1**

A solution of **L**₁ (0.1 g, 0.58 mmol) and RuCl₂(*p*-cymene)]₂ (0.17 g, 0.29 mmol) was heated in refluxing methanol (10 mL) for 3 h. After cooling to room temperature, the solvent was evaporated under vacuum. Water (2.7 mL) and NH₄PF₆ aqueous solution (0.26 g in 10 mL H₂O)

were then added to the residue with vigorous stirring, respectively. The crude product was precipitated and washed by water, giving complex **1** as a yellow powder (0.29 g, 86%). m.p. 162~164 °C (dec.); ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.40 (d, *J*=6.5 Hz, 1H), 8.33 (d, *J*=10.0 Hz, 1H), 8.11 (t, *J*=9.0 Hz, 1H), 7.68~7.55 (m, 3H), 6.75 (d, *J*=10.5 Hz, 1H), 6.28 (d, *J*=7.5 Hz, 1H), 6.04 (d, *J*=7.5 Hz, 1H), 5.83 (d, *J*=7.5 Hz, 1H), 5.76 (d, *J*=7.5 Hz, 1H), 2.46~2.43 (m, 1H), 2.19 (s, 3H), 0.90 (d, *J*=7.5 Hz, 3H), 0.89 (d, *J*=7.5 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 167.7, 157.0, 155.7, 153.1, 140.1, 139.7, 126.7, 123.1, 115.3, 111.2, 105.2, 102.1, 88.3, 86.5, 82.6, 82.1, 30.9, 22.3, 22.0, 19.2; ³¹P NMR (162 MHz, DMSO-*d*₆) δ : 144.2. Anal. calcd for C₂₀H₂₂ClF₆N₂OPRu: C 40.86, H 3.77, N 4.77; found C 40.92, H 3.75, N 4.71.

4.1.2 Synthesis of **2**

By using a similar procedure as described above, a solution of **L**₂ (0.1 g, 0.59 mmol) and RuCl₂(*p*-cymene)]₂ (0.18 g, 0.30 mmol) was heated in refluxing methanol (10 mL) for 3 h. After cooling to temperature, the solvent was evaporated under vacuum. Water (2.8 mL) and NH₄PF₆ aqueous solution (0.28 g in 10 mL H₂O) were then added to the residue with vigorous stirring, respectively. The crude product was precipitated and washed by water, giving complex **2** as a golden powder (0.26 g, 75%). Single crystals suitable for X-ray crystallographic determination were grown with CH₃OH/ether at ambient temperature. m.p. 181~184 °C (dec.). ¹H NMR (400 MHz, DMSO-*d*₆) δ : 8.93 (d, *J*=6.0 Hz, 2H), 8.04 (t, *J*=7.6 Hz, 2H), 7.76 (d, *J*=7.6 Hz, 2H), 7.54 (t, *J*=6.0 Hz, 2H), 6.02 (d, *J*=6.4 Hz, 2H), 5.88 (d, *J*=6.4 Hz, 2H), 4.80 (d, *J*=15.6 Hz, 1H), 4.40 (d, *J*=15.6 Hz, 1H), 2.81~2.75 (m, 1H), 1.94 (s, 3H), 1.16 (d, *J*=6.4 Hz, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 157.6, 157.2, 140.5, 125.8, 124.6, 105.9, 105.9, 100.4, 85.2, 84.7, 46.3, 30.6, 22.4, 17.8. ³¹P NMR (162 MHz, DMSO-*d*₆) δ : 144.2. Anal. calcd for C₂₁H₂₄ClF₆N₂PRu: C 43.05, H 4.13, N 4.78; found C 42.76, H 4.16, N 4.71.

4.1.3 Synthesis of **3**

By using a similar procedure as described above, a solution of **L**₃ (0.1 g, 0.53 mmol) and RuCl₂(*p*-cymene)]₂ (0.16 g, 0.27 mmol) was heated in refluxing methanol (10 mL) for 3 h. After cooling to room temperature, the solvent was evaporated under vacuum. Water (2.8 mL) and NH₄PF₆ aqueous solution (0.28 g in 10 mL H₂O) was then added to the residue with vigorous stirring, respectively. The crude product was precipitated and washed by water, giving complex **3** as a pale yellow powder (0.25 g, 80%). m.p. 185~187 °C (dec.). ¹H NMR (400 MHz, CDCl₃) δ : 8.84 (d, *J*=5.6 Hz, 1H), 7.87 (t, *J*=7.6 Hz, 1H), 7.73 (t, *J*=8.0 Hz, 1H), 7.65 (d, *J*=7.6 Hz, 1H), 7.38 (t, *J*=6.8 Hz, 1H), 7.26 (t, *J*=3.6 Hz, 1H), 6.90 (d, *J*=8.4 Hz, 1H), 5.89 (d, *J*=6.0 Hz, 1H), 5.84 (d, *J*=6.0 Hz, 1H), 5.75 (t, *J*=6.0 Hz, 2H), 4.54 (d, *J*=16 Hz, 1H), 4.46 (d, *J*=16 Hz, 1H), 2.85~2.75 (m, 1H), 2.02 (s, 3H), 1.24 (d, *J*=6.8 Hz, 3H), 1.19 (d, *J*=6.8 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 166.7, 158.1, 157.7, 154.4, 142.0, 140.3, 125.2, 124.1, 116.9,

110.8, 106.3, 99.3, 85.3, 84.5, 83.4, 82.9, 46.9, 31.2, 22.7, 22.3, 18.2; ^{31}P NMR (162 MHz, $\text{DMSO}-d_6$) δ : 144.2. Anal. calcd for $\text{C}_{21}\text{H}_{24}\text{ClF}_6\text{N}_2\text{OPRu}$: C 41.90, H 4.02, N 4.65; found C 41.63, H 3.82, N 4.60.

4.1.4 Synthesis of 4

By using a similar procedure as described above, a solution of **L**₄ (0.1 g, 0.5 mmol) and $\text{RuCl}_2(p\text{-cymene})_2$ (0.15 g, 0.25 mmol) was heated in refluxing methanol (10 mL) for 3 h. After cooling to room temperature, the solvent was evaporated under vacuum. Water (2.5 mL) and NH_4PF_6 aqueous solution (0.26 g in 10 mL H_2O) were then added to the residue with vigorous stirring, respectively. The crude product was precipitated and washed by water, giving complex **4** as a yellow powder (0.25 g, 82%). m.p. 175~177 °C (dec.); ^1H NMR (400 MHz, CDCl_3) δ : 8.93 (d, $J=5.2$ Hz, 1H), 7.82~7.76 (m, 2H), 7.60 (d, $J=7.6$ Hz, 1H), 7.30 (d, $J=6.8$ Hz, 1H), 7.23 (d, $J=7.2$ Hz, 1H), 6.87 (d, $J=8.8$ Hz, 1H), 5.76~5.65 (m, 4H), 4.52 (d, $J=15.6$ Hz, 1H), 4.43 (d, $J=15.6$ Hz, 1H), 4.06 (s, 3H), 2.89~2.82 (m, 1H), 2.14 (s, 3H), 1.25 (d, $J=6.8$ Hz, 3H), 1.18 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.9, 157.9, 156.3, 155.2, 142.5, 139.8, 125.4, 123.7, 119.1, 107.1, 106.8, 99.6, 85.3, 84.3, 83.6, 82.5, 57.9, 48.2, 31.2, 22.4, 18.5; ^{31}P NMR (162 MHz, CDCl_3) δ : 144.4. Anal. calcd for $\text{C}_{22}\text{H}_{26}\text{ClF}_6\text{N}_2\text{OPRu}$: C 42.90, H 4.25, N 4.55; found C 43.13, H 4.28, N 4.48.

4.1.5 General procedure for the synthesis of quinolines and pyrroles

Under a nitrogen atmosphere, the amino alcohol (2.0 mmol), ketone (2.4 mmol), complex **3** (6.1 mg, 0.01 mmol), KOH (0.11 g, 2.0 mmol), and 2 mL of toluene were mixed and stirred at 110 °C. After 2 h, the reaction mixture was cooled to room temperature. The mixture was then condensed under reduced pressure and subjected to purification by flash silica gel column chromatography to afford the target product, which was identified by NMR analyses. The melting points of the known compounds are consistent with those reported in literatures

2-Phenylquinoline: White solid. m.p. 84~86 °C; $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.25 (d, $J=8.4$ Hz, 2H), 8.18 (d, $J=7.2$ Hz, 2H), 7.90~7.83 (m, 2H), 7.75 (t, $J=8.0$ Hz, 1H), 7.56~7.46 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ : 157.3, 148.1, 139.5, 136.9, 129.7, 129.6, 129.3, 128.7, 127.6, 127.4, 127.2, 126.3, 119.0.

2-(*o*-tolyl)quinolone: Yellow solid. m.p. 73~77 °C; $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.17 (t, $J=8.4$ Hz, 2H), 7.84 (d, $J=7.2$ Hz, 1H), 7.73 (t, $J=6.8$ Hz, 1H), 7.56~7.48 (m, 3H), 7.32 (m, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.3, 147.9, 140.8, 136.1, 136.0, 130.9, 129.8, 129.7, 129.6, 128.5, 127.5, 126.7, 126.4, 126.1, 122.4, 20.5.

2-(*m*-tolyl)quinolone: Yellow solid. m.p. 38~40 °C; $^{[56]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.20 (t, $J=8.8$ Hz, 2H), 8.02 (s, 1H), 7.94~7.82 (m, 3H), 7.73 (t, $J=8.4$ Hz, 1H), 7.52 (t, $J=7.6$ Hz, 1H), 7.42 (t, $J=7.6$ Hz, 1H), 7.28 (d, $J=7.6$ Hz, 1H), 2.49 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 157.6, 148.3, 139.6, 138.5, 136.7, 130.1, 129.7, 129.6, 128.7,

128.2, 127.4, 127.1, 126.2, 124.7, 119.2, 21.6.

2-(4-Methoxyphenyl)quinoline: Yellow solid. m.p. 121~124 °C; $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.19~8.17 (m, 4H), 7.86~7.81 (m, 2H), 7.74 (t, $J=8.4$ Hz, 1H), 7.52 (t, $J=8.0$ Hz, 1H), 7.08 (d, $J=8.8$ Hz, 2H), 3.91 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.8, 156.8, 148.2, 136.6, 132.2, 129.6, 129.5, 128.9, 127.4, 126.9, 125.9, 118.5, 114.2, 55.3.

2-(3-Fluorophenyl)quinolone: Yellow solid. m.p. 47~48 °C; $^{[57]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.14~8.07 (m, 2H), 7.84 (d, $J=9.6$ Hz, 2H), 7.76~7.73 (m, 2H), 7.65 (d, $J=8.4$ Hz, 1H), 7.47~7.36 (m, 2H), 7.06 (t, $J=5.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 160.9, 154.4, 147.0, 140.7, 135.7, 129.1, 128.7, 126.3, 126.2, 125.4, 121.9, 117.4, 115.0 (d, $J=22.6$ Hz), 113.3 (d, $J=22.6$ Hz).

2-(3-Bromophenyl)quinoline: Yellow solid. m.p. 73~76 °C; $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.24 (s, 1H), 8.06 (d, $J=8.4$ Hz, 2H), 7.94 (d, $J=7.6$ Hz, 1H), 7.69~7.59 (m, 3H), 7.47~7.39 (m, 2H), 7.25 (t, $J=7.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 154.3, 147.0, 140.5, 135.8, 131.0, 129.4, 129.1, 128.7, 128.6, 126.3, 126.1, 125.5, 124.8, 122.0, 117.4.

2-(4-Fluorophenyl)quinoline: White solid. m.p. 91~93 °C; $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.14~8.07 (m, 4H), 7.75 (d, $J=8.8$ Hz, 2H), 7.65 (t, $J=6.8$ Hz, 1H), 7.45 (t, $J=6.8$ Hz, 1H), 7.13 (t, $J=8.4$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.9, 161.5, 155.1, 147.1, 135.8, 134.7, 128.7, 128.6, 128.3, 126.4, 126.1, 125.2, 117.5, 114.8, 114.6.

2-(4-Chlorophenyl)quinoline: White solid. m.p. 112~114 °C; $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.25 (d, $J=8.8$ Hz, 1H), 8.20~8.14 (m, 3H), 7.87~7.84 (m, 2H), 7.76 (t, $J=6.8$ Hz, 1H), 7.58~7.51 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 155.9, 148.1, 137.9, 137.1, 135.6, 129.9, 129.6, 129.0, 128.8, 127.4, 127.2, 126.5, 118.5.

2-(2-Chlorophenyl)quinoline: White solid. m.p. 73~75 °C; $^{[58]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.11 (t, $J=8.8$ Hz, 2H), 7.78 (d, $J=8.0$ Hz, 1H), 7.67~7.61 (m, 3H), 7.49 (t, $J=7.6$ Hz, 1H), 7.42 (d, $J=6.8$ Hz, 1H), 7.34~7.27 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 157.4, 148.1, 139.6, 135.7, 132.3, 131.7, 130.1, 129.9, 129.7, 127.5, 127.2, 127.1, 126.8, 122.8.

3-Methyl-2-phenylquinoline: Yellow oil. $^{[45]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.19 (d, $J=8.0$ Hz, 1H), 8.06 (s, 1H), 7.82 (d, $J=8.0$ Hz, 1H), 7.70 (t, $J=8.0$ Hz, 1H), 7.62 (d, $J=8.0$ Hz, 2H), 7.57~7.45 (m, 4H), 2.50 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.3, 137.1, 129.3, 129.0, 128.9, 128.3, 127.6, 126.7, 126.5, 20.6.

2-Ethyl-5-phenyl-1H-pyrrole: White solid. m.p. 49~50 °C; $^{[58]}$ ^1H NMR (400 MHz, CDCl_3) δ : 8.16 (s, 1H), 7.46 (d, $J=7.2$ Hz, 2H), 7.36 (t, $J=7.6$ Hz, 2H), 7.19 (t, $J=7.2$ Hz, 1H), 6.45 (t, $J=2.8$ Hz, 1H), 6.01 (t, $J=2.8$ Hz, 1H), 2.71 (q, $J=7.6$ Hz, 2H), 1.32 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 135.6, 133.0, 130.6, 128.8, 125.6, 123.4, 106.2, 106.0, 21.0, 13.6.

2-Ethyl-5-(4-methoxyphenyl)-1H-pyrrole: White solid. m.p. 69~71 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.07 (s,

1H), 7.40 (d, $J=8.8$ Hz, 2H), 6.93 (d, $J=12.8$ Hz, 2H), 6.33 (t, $J=3.0$ Hz, 1H), 6.00 (t, $J=3.0$ Hz, 1H), 3.85 (s, 3H), 2.71 (q, $J=7.6$ Hz, 2H), 1.32 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 157.9, 134.9, 130.6, 126.2, 124.8, 114.3, 105.9, 104.8, 55.3, 21.0, 13.6.

5-Ethyl-3-methyl-2-phenyl-1H-pyrrole: Colorless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.87 (s, 1H), 7.45~7.42 (m, 4H), 7.28~7.25 (m, 1H), 5.93 (s, 1H), 2.71 (q, $J=7.6$ Hz, 2H), 2.32 (s, 3H), 1.34 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 134.0, 133.9, 128.6, 126.5, 125.9, 125.4, 116.2, 108.6, 20.8, 13.6, 12.5.

2-(*sec*-Butyl)-5-phenyl-1H-pyrrole: Colorless oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.22 (s, 1H), 7.55 (d, $J=8.0$ Hz, 2H), 7.44 (t, $J=7.6$ Hz, 2H), 7.27 (t, $J=7.0$ Hz, 1H), 6.56 (t, $J=3.2$ Hz, 1H), 6.11 (t, $J=3.2$ Hz, 1H), 2.87~2.80 (m, 1H), 1.85~1.65 (m, 2H), 1.40 (d, $J=7.2$ Hz, 3H), 1.05 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 139.3, 133.2, 130.3, 128.9, 125.7, 123.5, 106.0, 105.7, 34.3, 30.2, 20.0, 12.6, 11.9.

2-(*sec*-Butyl)-5-(4-methoxyphenyl)-1H-pyrrole: Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.05 (s, 1H), 7.41 (d, $J=8.8$ Hz, 2H), 6.94 (d, $J=8.8$ Hz, 2H), 6.35 (t, $J=3.2$ Hz, 1H), 6.00 (t, $J=2.8$ Hz, 1H), 3.86 (s, 3H), 2.78~2.73 (m, 1H), 1.76~1.59 (m, 2H), 1.33 (d, $J=7.2$ Hz, 3H), 0.97 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 157.9, 138.5, 130.3, 126.3, 124.8, 114.3, 105.3, 104.7, 55.3, 34.4, 30.3, 20.1, 11.9.

2-(*sec*-Butyl)-5-(4-chlorophenyl)-1H-pyrrole: Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.09 (s, 1H), 7.41~7.32 (m, 4H), 6.45 (t, $J=3.2$ Hz, 1H), 6.02 (t, $J=3.2$ Hz, 1H), 2.80~2.71 (m, 1H), 1.78~1.58 (m, 2H), 1.33 (d, $J=7.2$ Hz, 3H), 0.97 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 139.7, 131.6, 131.0, 129.1, 128.9, 124.5, 106.4, 105.9, 34.4, 30.2, 20.1, 11.8.

5-(*sec*-Butyl)-3-methyl-2-phenyl-1H-pyrrole: Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.83 (s, 1H), 7.44~7.38 (m, 4H), 7.24~7.20 (m, 1H), 5.88 (s, 1H), 2.74~2.66 (m, 1H), 2.28 (s, 3H), 1.75~1.54 (m, 2H), 1.30 (d, $J=7.2$ Hz, 3H), 0.96 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 137.6, 134.0, 128.6, 125.8, 125.3, 116.0, 107.9, 34.2, 30.2, 20.0, 12.6, 11.9.

Supporting Information Crystallographic details of complex **2** and NMR spectra of new complexes and catalytic products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Abass, M. *Heterocycles* **2005**, 65, 901.
- [2] Michael, J. P. *Nat. Prod. Rep.* **2008**, 25, 166.
- [3] Moorthy, N. S. H. N.; Manivannan, E.; Karthikeyan, C.; Piyush, T. *Mini-Rev. Med. Chem.* **2013**, 13, 1415.
- [4] Gopaul, K.; Shintre, S. A.; Koorbanally, N. A. *Anti-Cancer Agents Med. Chem.* **2015**, 15, 631.
- [5] Zhou, Y.; Li, W.; Yu, L.; Liu, Y.; Wang, X.; Zhou, M. *Dalton Trans.* **2015**, 44, 1858.
- [6] Musiol, R. *Expert Opin. Drug Discovery* **2017**, 12, 583.
- [7] Wang, Y.; Liu, L.; Wang, Y. *Chin. J. Org. Chem.* **2018**, 38, 1009 (in Chinese).
- [8] Yan, J.; Ji, X.-Y.; Hua, S.-G.; Wang, J. *Chin. J. Org. Chem.* **2018**, 38, 791 (in Chinese).
- [9] Chen, Y.; He, B.; Qin, A.; Tang, B. Z. *Sci. China Chem.* **2019**, 62, 1017.
- [10] Varela, J.; Saá, C. *Chem. Rev.* **2003**, 103, 3787.
- [11] Zeni, Z.; Larock, R. C. *Chem. Rev.* **2006**, 106, 4644.
- [12] Marco-Contelles, J.; Pérez-Mayoral, E.; Samadi, A. *Chem. Rev.* **2009**, 109, 2652.
- [13] Chelucci, G.; Porcheddu, A. *Chem. Rev.* **2017**, 17, 200.
- [14] Schranck, J.; Tlili, A.; Beller, M. *Angew. Chem., Int. Ed.* **2013**, 52, 7642.
- [15] Nandakumar, A.; Midya, S. P.; Landge, V. G.; Balaraman, E. *Angew. Chem., Int. Ed.* **2015**, 54, 11022.
- [16] Shimizu, K. *Catal. Sci. Technol.* **2015**, 5, 1412.
- [17] Yang, Q.; Wang, Q.; Yu, Z. *Chem. Soc. Rev.* **2015**, 44, 2305.
- [18] Mukherjee, A.; Milstein, D. *ACS Catal.* **2018**, 8, 11435.
- [19] Mastalir, M.; Glatz, M.; Pittenauer, E.; Allmaier, G.; Kirchner, K. *Org. Lett.* **2019**, 21, 1116.
- [20] Irrgang, T.; Kempe, R. *Chem. Rev.* **2019**, 119, 2524.
- [21] Michlik, S.; Kempe, R. *Nat. Chem.* **2013**, 5, 140.
- [22] Hille, T.; Irrgang, T.; Kempe, R. *Angew. Chem., Int. Ed.* **2017**, 56, 371.
- [23] Forberg, D.; Schwob, T.; Kempe, R. *Nat. Commun.* **2018**, 9, 1751.
- [24] Maji, M.; Chakrabarti, K.; Panja, D.; Kundu, S. J. *Catal.* **2019**, 373, 93.
- [25] Vander Mierde, H.; Van Der Voort, P.; De Vos, D.; Verpoort, F. *Eur. J. Org. Chem.* **2008**, 1625.
- [26] Zhang, M.; Fang, X.; Neumann, H.; Beller, M. *J. Am. Chem. Soc.* **2013**, 135, 11384.
- [27] Iida, K.; Miura, T.; Ando, J.; Saito, S. *Org. Lett.* **2013**, 15, 1436.
- [28] Subramanian, M.; Sundar, S.; Rengan, R. *Appl. Organomet. Chem.* **2018**, e4582.
- [29] Srimani, D.; Ben-David, Y.; Milstein, D. *Chem. Commun.* **2013**, 49, 6632.
- [30] Srimani, D.; Ben-David, Y.; Milstein, D. *Angew. Chem., Int. Ed.* **2013**, 52, 4012.
- [31] Pan, B.; Liu, B.; Yue, E.; Liu, Q.; Yang, X.; Wang, Z.; Sun, W.-H. *ACS Catal.* **2016**, 6, 1247.
- [32] Chai, H.; Wang, L.; Liu, T.; Yu, Z. *Organometallics* **2017**, 36, 4936.
- [33] Maji, M.; Chakrabarti, K.; Paul, B.; Roy, B. C.; Kundu, S. *Adv. Synth. Catal.* **2018**, 360, 722.
- [34] Deng, D.; Hu, B.; Yang, M.; Chen, D. *Organometallics* **2018**, 37, 2386.
- [35] Guo, B.; Yu, T.-Q.; Li, H.-X.; Zhang, S.-Q.; Braunstein, P.; Young, D. J.; Li, H.-Y.; Lang, J.-P. *ChemCatChem* **2019**, 11, 2500.
- [36] Wan, X.-M.; Liu, Z.-L.; Liu, W.-Q.; Cao, X.-N.; Zhu, X.; Zhao, X.-M.; Song, B.; Hao, X.-Q.; Liu, G. *Tetrahedron* **2019**, 75, 2697.
- [37] Mastalir, M.; Glatz, M.; Pittenauer, E.; Allmaier, G.; Kirchner, K. *J. Am. Chem. Soc.* **2016**, 138, 15543.
- [38] Kallmeier, F.; Dudziec, B.; Irrgang, T.; Kempe, R. *Angew. Chem., Int. Ed.* **2017**, 56, 7261.
- [39] Midya, S. P.; Landge, V. G.; Sahoo, M. K.; Rana, J.; Balaraman, E. *Chem. Commun.* **2018**, 54, 90.
- [40] Zhang, G.; Wu, J.; Zeng, H.; Zhang, S.; Yin, Z.; Zheng, S. *Org. Lett.* **2017**, 19, 1080.
- [41] Das, S.; Maiti, D.; Sarkar, S. D. *J. Org. Chem.* **2018**, 83, 2309.
- [42] Parua, S.; Sikari, R.; Sinha, S.; Das, S.; Chakraborty, G.; Paul, N. D. *Org. Biomol. Chem.* **2018**, 16, 274.
- [43] Singh, K.; Vellakkaran, M.; Banerjee, D. *Green Chem.* **2018**, 20, 2250.
- [44] Tan, D.-W.; Li, H.-X.; Zhu, D.-L.; Li, H.-Y.; Young, D. J.; Yao, J.-L.; Lang, J.-P. *Org. Lett.* **2018**, 20, 608.
- [45] Elangova, S.; Sortais, J.-B.; Beller, M.; Darcel, C. *Angew. Chem., Int. Ed.* **2015**, 54, 14483.
- [46] Shi, J.; Hu, B.; Gong, D.; Shang, S.; Hou, G.; Chen, D. *Dalton*

- Trans.* **2016**, *45*, 4828.
- [47] Shi, J.; Hu, B.; Chen, X.; Shang, S.; Deng, D.; Sun, Y.; Shi, W.; Yang, X.; Chen, D. *ACS Omega* **2017**, *2*, 3406.
- [48] Shi, J.; Hu, B.; Ren, P.; Shang, S.; Yang, X.; Chen, D. *Organometallics* **2018**, *37*, 2795.
- [49] Deng, D.; Hu, B.; Yang, M.; Chen, D. *Organometallics* **2018**, *37*, 3353.
- [50] Zhang, C.; Zhao, J.; Hu, B.; Shi, J.; Chen, D. *Organometallics* **2019**, *38*, 654.
- [51] Deng, D.; Hu, B.; Zhang, Z.; Mo, S.; Yang, M.; Chen, D. *Organometallics* **2019**, *38*, 2218.
- [52] Gong, D.; Hu, B.; Chen, D. *Dalton Trans.* **2019**, *48*, 8826.
- [53] Bennett, M. A.; Smith, A.K. *J. Chem. Soc. Dalton. Trans.* **1974**, 233.
- [54] Salamanca, V.; Toledo, A.; Albeniz, A. C. *J. Am. Chem. Soc.* **2018**, *140*, 17851.
- [55] Dyker, G.; Muth, O. *Eur. J. Org. Chem.* **2004**, *21*, 4319.
- [56] Cacchi, S.; Fabrizi, G.; Marinelli, F. *Synlett* **1999**, 401.
- [57] Wang, R.; Fan, H.; Zhao, W.; Li, F. *Org. Lett.* **2016**, *18*, 3558.
- [58] Mo, S.; Zhu, Y.; Shen, Z. *Org. Biomol. Chem.* **2013**, *11*, 2756.
- [59] Krasniqi, B.; Geerts, K.; Dehaen, W. *J. Org. Chem.* **2019**, *84*, 5027.

(Lu, Y.)