含苯并噻唑砌块的 2,4,6-三取代嘧啶衍生物的合成及抗肿瘤活性评价

李二冬^{a,d} 孟娅琪^{a,d} 张路野^{a,b} 张洋^{a,b} 周蕊^a
 刘丽敏^a 栗娜^{b,c} 辛景超^{b,c} 郑甲信^{a,d}
 单丽红^{*,a,b,c,d} 刘宏民^{*,a,b,c,d} 张秋荣^{*,a,b,c,d}

(^a 郑州大学药学院 郑州 450001)

(^b 新药创制与药物安全性评价河南省协同创新中心 郑州 450001)

(^c 河南省药物质量评价重点实验室 郑州 450001)

(^d 教育部药物制备关键技术重点实验室 郑州 450001)

摘要 为了寻找高效的抗肿瘤药物,设计并合成了一系列含苯并噻唑砌块的 2,4,6-三取代嘧啶衍生物。采用噻唑蓝(MTT)法对目标化合物在人类四种癌细胞[EC-109(人食管癌细胞)、MGC-803(人胃癌细胞)、PC-3(人前列腺癌细胞)、HepG-2(人肝癌细胞)]、GES-1(人正常胃黏膜上皮细胞)和 HEEC(人正常食管细胞)中进行抗肿瘤活性评价,结果显示部分化合物对 MGC-803 和 PC-3 细胞表现出中度至强效的抗肿瘤活性。其中 2-(((4-(4-(吡啶-2-基)哌嗪-1-基)-6-(三氟甲基)嘧啶-2-基)硫基)甲基)苯并[d]噻唑(**13h**)和 2-(((4-(4-(嘧啶-2-基)哌嗪-1-基)-6-(三氟甲基)嘧啶-2-基)硫代)甲基)苯并[d]噻唑(**13i**)对 PC-3 表现出比较好的抗肿瘤活性,IC₅₀ 值分别 3.82 和 2.29 μmol/L,且化合物 **13h** 和 **13i** 对 GES-1 的细胞增殖毒性明显小于阳性对照 5-氟尿嘧啶。

关键词 嘧啶; 苯并噻唑; 合成; 抗肿瘤活性

Synthesis and Antitumor Evaluation of 2,4,6-Trisubstituted Pyrimidine Derivatives Containing Benzothiazole Moiety

Li, Erdong^{a,d} Meng, Yaqi^{a,d} Zhang, Luyue^{a,b} Zhang, Yang^{a,b} Zhou, Rui^a
 Liu, Limin^a Li, Na^{b,c} Xin, Jingchao^{b,c} Zheng, Jiaxin^{a,d} Shan, Lihong^{*,a,b,c,d}
 Liu, Hongmin^{*,a,b,c,d} Zhang, Qiurong^{*,a,b,c,d}

(^a School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001)

(^b Collaborative Innovation Center of New Drug Research and Safety Evaluation, Zhengzhou 450001)

(^c Key Laboratory of Henan Province for Drug Quality and Evaluation, Zhengzhou 450001)

(^d Key Laboratory of Technology of Drug Preparation, Ministry of Education, Zhengzhou 450001)

Abstract In order to find high-efficiency antitumor drugs, a series of 2,4,6-trisubstituted pyrimidine derivatives containing benzothiazole moiety were designed, synthesized and evaluated for antitumor activities against four cancer cells (EC-109, human esophageal cancer cells; MGC-803, human gastric cancer cells; PC-3, human prostate cancer cells; HepG-2, human liver cancer cells), GES-1 (human normal gastric mucosal epithelial cells), and HEEC (human normal esophagus) using thiazolyl blue (MTT) method. The results showed that some compounds exhibited moderate to strong antitumor activities against MGC-803 and PC-3 cells. Among them, 2-(((4-(4-(pyridin-2-yl)piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)-thio)methyl)benzo[d]thiazole (**13h**) and 2-(((4-(4-(pyrimidin-2-yl)piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13i**) showed the most potent antitumor activities against PC-3 cells with IC₅₀ values of 3.82 and 2.29 μmol/L, respectively. The toxicities of compounds **13h** and **13i** to GES-1 cells were significantly lower than the positive control 5-fluorouracil.

* Corresponding authors. E-mail: zqr409@yeah.net; liuhm@zzu.edu.cn; 343785851@qq.com

Received July 8, 2019; revised October 14, 2019; published online October 25, 2019.

Project supported by the National Natural Science Foundation of China (No. U1904163) and the National Key Research Program of Proteins (No. 2018YFE0195100).

国家自然科学基金(No. U1904163)和蛋白关键研究(No. 2018YFE0195100)资助项目。

Keywords pyrimidine; benzothiazole; synthesis; antitumor activity

1 Introduction

Cancer is the second most common cause of death in humans with a high incidence.^[1] In 2015, there were 1658370 new cancer cases and 589430 cancer deaths in the United States.^[2] The World Health Organization (WHO) has estimated that the incidence of cancer will reach 22 million every year in the next two years.^[3] Therefore, it is urgent to develop effective antitumor drugs.

Mercaptopyrimidine derivatives have excellent coordination ability because they contain N and S atoms and they can form complexes with many metal salts. It has been reported that mercaptopyrimidines have antileukemia, antitumor, anti-inflammatory, anti-bacterial and anti-convulsant effects.^[4~10] At the same time, Noolvi *et al.*^[11] reported that compound **1** has a good antitumor activity against non-small cell lung cancer cells with IC₅₀ value of 71.8 nmol/L (Figure 1). In 2013, compound **2** was synthesized and evaluated its cytotoxicity against six human cancer cell lines. The results showed that compound **2** exerted significant inhibitory activity compared to dasatinib.^[12] It was also found that compound **3** had an anti-bacterial activity comparable to chloramphenicol, and compound **4** showed strong cytotoxic activity against a variety of cancer cells (MDA-MB-231, Caki-1, PANC-1, A549, MKN-45, HT29 and PC-3 cells) with an IC₅₀ value of thiazole and piperazine are two pharmacophores with anti-

11~20 nmol/L.^[13,14] The above data indicated that benzothiazole moiety. According to the combination principles of pharmacodynamic groups, it is desirable to synthesize higher efficiency antitumor inhibitors if they are introduced into the pyrimidine core.

Based on previous studies, 2-mercapto-4-hydroxy-6-trifluoromethylpyrimidine was selected as the scaffold.^[15,16] Then, benzothiazole and the substituted piperazines were introduced into the mercaptopyrimidine to synthesize a series of compounds with multiple active moieties. Finally, the antitumor activities of the target compounds were determined using thiazolyl blue (MTT) method.

1 Results and discussion

1.1 Chemistry

The synthesis of the target compounds was depicted in Scheme 1. Pyrimidine core (**6**) was firstly synthesized by a

Figure 1 Antitumor agents and target compounds

Reagents and conditions: (i) H₂NCSNH₂, NaH, MeCN, 70 °C, 3 h; (ii) BF₃·OEt₂, CH₂Cl₂, r.t., 2.5 h; (iii) KOH, H₂O, dioxane, 60 °C, 6 h; (iv) POCl₃, dioxane, 90 °C, 5 h; (v) 1-Boc-piperazine, DMF, 90 °C, 2 h; (vi) CH₂Cl₂, CF₃COOH, 0 °C, 8 h; (vii) DMF, 90 °C, 3 h

Scheme 1 Synthesis of 2,4-disubstituted pyrimidine derivatives containing benzothiazole moiety

ring-forming reaction of commercially available thiourea with ethyl 4,4,4-trifluoroacetoacetate (**5**) under basic condition in ethanol. At the same time, 2-aminobenzenethiol (**7**) reacted with chloroacetyl chloride to get 2-chloromethylbenzothiazole (**8**). Next, compound **9** was synthesized from pyrimidine and 2-chloromethyl benzothiazole by nucleophilic reaction, and the chlorination reaction was followed to obtain the key intermediate compound **10**. Then, the N-Boc substituted piperazine was introduced to give compound **11**. Finally, the piperazine substitution reaction was carried out to obtain a series of target compounds **13a~13i** after the removal of *tert*-butoxycarbonyl (Boc) protecting group.

1.2 Antitumor activity

In order to explore the antitumor activities of the prepared compounds, compounds **9**, **10**, **12** and **13a~13i** were evaluated against EC-109 (human esophageal cancer cells), MGC-803 (human gastric cancer cells), PC-3 (human prostate cancer cells) and HepG-2 (human liver cancer cells) using MTT assay.^[17,18] 5-Fluorouracil (5-Fu) was used as a positive control. The results of evaluation of antitumor activity are shown in Table 1.

As shown in Table 1, the antitumor activities of compounds **9** and **10** against the four tumor cells were very weak, and their IC₅₀ values were greater than 50 μmol/L.

The activity of compound **12** against MGC-803 and PC-3 cell lines was significantly increased compared to compounds **9** and **10**. Among the target compounds **13a~13i**, compounds **13h** and **13i** exerted excellent antitumor activities against PC-3 with IC₅₀ values of 3.82 and 2.29 μmol/L, respectively. It could also find that compound **13i** was the most active compound against HepG-2 and MGC-803 cells with IC₅₀ values of 15.08 and 9.39 μmol/L, respectively, which were comparable to the positive control 5-Fu. Compounds **13a~13g** have poor antitumor activities against four tumor cells compared to the positive control 5-Fu. Compared the activity data of compounds **13a** and **13b** with **13c**, we concluded that as the carbon chain of the aliphatic substituent group grows, the antitumor activities of the compounds against the selected four cancer cells were also increased. When R was an aromatic substituent, the antitumor activity of compound was also significantly increased when the number of heteroatoms increased by comparing compounds **13e** and **13h** with **13i**.

To investigate the toxicity of the target compounds to normal cells, compounds **13h** and **13i** were selected to further explore their cytotoxicity against GES-1 (normal human gastric epithelial cell line) and HEEC (human normal esophageal cells). As shown in Table 2, the results indicated that compounds **13h** and **13i** had weak or no cytotoxicity

Table 1 *In vitro* antitumor activities evaluation results for compounds **9**, **10**, **12** and **13a~13i**

Compd.	R	IC ₅₀ ^a /(μmol·L ⁻¹)			
		EC-109	MGC-803	PC-3	HepG-2
9	—	>50	>50	>50	>50
10	—	>50	>50	>50	>50
12	—	>50	31.93 ± 1.50	15.77 ± 1.19	>50
13a	CH ₃	>50	>50	21.58 ± 1.33	>50
13b	CH ₂ CH ₃	>50	>50	13.33 ± 1.12	36.71 ± 1.58
13c	CH ₂ CH ₂ CH ₂ CH ₃	40.01 ± 1.61	12.38 ± 1.10	7.89 ± 0.46	19.35 ± 1.32
13d	CH ₂ CH ₂ OH	>50	25.89 ± 1.35	12.83 ± 1.10	>50
13e		>50	>50	>50	>50
13f		>50	44.30 ± 1.64	28.57 ± 1.45	>50
13g		>50	48.25 ± 1.46	38.09 ± 1.88	>50
13h		35.68 ± 1.40	11.32 ± 1.00	3.82 ± 0.58	23.45 ± 1.31
13i		27.35 ± 1.20	9.39 ± 0.61	2.293 ± 0.360	15.08 ± 1.14
5-Fu ^b	—	12.23 ± 1.08	8.24 ± 0.75	5.54 ± 0.74	13.12 ± 1.34

^a Antitumor activity was assayed by exposure for 72 h to substances and expressed as concentration required to inhibit tumor cells proliferation by 50% (IC₅₀). ^b Used as a positive control.

against GES-1 and HEEC, which was significantly lower than the positive control 5-Fu.

Table 2 *In vitro* antiproliferative activities of compounds **13h** and **13i** against human normal cells

Compd.	IC ₅₀ /(μmol·L ⁻¹)	
	GES-1	HEEC
13h	89.68	>100
13i	>100	>100
5-Fu	22.60	>100

2 Conclusion

In conclusion, a series of 2,4,6-trisubstituted pyrimidine derivatives containing benzothiazole moiety were designed, synthesized and evaluated for antitumor activities against EC-109, MGC-803, PC-3 and HepG-2 cells using MTT method. The antitumor activity results showed that some compounds exhibited moderate to strong antitumor activities against MGC-803 and PC-3 cells. Among them, compounds **13h** and **13i** displayed the most potent antitumor activities against PC-3 with IC₅₀ values of 3.82 and 2.29 μmol/L, respectively. The toxicities of compounds **13h** and **13i** to GES-1 cells were significantly lower than the positive control 5-Fu.

3 Experimental

3.1 Materials

¹H NMR and ¹³C NMR spectra were measured on a Bruker 400 and 101 MHz spectrometer, respectively, and TMS as an internal standard. Melting points were determined on an X-5 micromelting apparatus. High-resolution mass spectra were measured using a Waters-Micromass Q-ToF Micro High Resolution Determination of tetragonal-flight time tandem mass spectrometer. The names of the compounds were from chembiooffice 14.0. All commercially available reagents and solvents were used without further purification.

3.2 Chemistry

3.2.1 Preparation of compounds **6** and **8**

Compounds **6** and **8** were synthesized according to the literature. Compound **6**: m.p. 247~249 °C^[19], compound **8**: m.p. 34 °C^[20].

3.2.2 Preparation of 2-((benzo[d]thiazol-2-yl-methyl)-thio)-6-(trifluoromethyl)pyrimidin-4-ol (**9**)

Compound **6** (5.00 g, 25.49 mmol) was dissolved in KOH aqueous solution (33.14 mmol, 30 mL), and a solution of compound **8** (4.68 g, 25.49 mmol) in dioxane (18 mL) was added dropwise at room temperature. The reaction mixture was heated at 60 °C for 6 h, while the progress of the reaction was monitored by thin-layer chromatography (TLC). After the reaction was completed, the reaction mixture was cooled to ambient temperature, and then poured into ice-water (300 mL). Large amount of yellow solid was precipitated after the resulting aqueous mixture was adjusted to pH=7 by adding diluted hydrochloric acid (1 mL).

The suspension was filtered off via suction filtration to give compound **9** 6.56 g, 74.9% yield. Yellow solid, m.p. 191.0~191.4 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 13.60 (s, 1H), 8.06 (d, *J*=8.0 Hz, 1H), 7.96 (d, *J*=8.1 Hz, 1H), 7.51~7.47 (m, 1H), 7.43~7.36 (m, 1H), 6.70 (s, 1H), 4.91 (d, *J*=11.2 Hz, 2H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 167.5, 164.6, 152.3, 151.2, 135.1, 126.2, 125.3, 122.4, 122.1, 121.7, 119.0, 107.5, 31.8. HRMS (ESI) calcd for C₁₃H₉F₃N₃OS₂ [M+H]⁺ 344.0139, found 344.0137.

3.2.3 Preparation of 2-(((4-chloro-6-(trifluoromethyl)-pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**10**)

To a 100 mL round bottom flask was added compound **9** (6.00 g, 17.48 mmol) in dioxane (30 mL) followed by the addition of phosphorus oxychloride (8.15 mL, 87.4 mmol). The reaction mixture was heated at 90 °C for 5 h. After the reaction was completed, the reaction mixture was poured into an ice-water (300 mL) after cooling to room temperature. The mixture was stirred for 3 min, filtered under reduced pressure and dried *in vacuo* to obtain compound **10** 4.85 g, yield 46.7%. Purple red solid, m.p. 93.1~93.5 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.06 (d, *J*=8.0 Hz, 2H), 7.97 (d, *J*=8.1 Hz, 1H), 7.51 (t, *J*=7.6 Hz, 1H), 7.43 (t, *J*=7.6 Hz, 1H), 4.95 (s, 2H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 171.7, 167.4, 162.9, 155.8, 155.4, 152.4, 135.0, 126.2, 125.3, 122.4, 122.1, 114.6, 32.7. HRMS (ESI) calcd for C₁₃H₈ClF₃N₃S₂ [M+H]⁺ 361.9800, found 361.9804.

3.2.4 Preparation of *tert*-butyl 4-(2-((benzo[d]thiazol-2-ylmethyl)thio)-6-(trifluoromethyl)pyrimidin-4-yl)pyrrolidine-1-carboxylate (**11**)

To a solution of compound **10** (2.00 g 5.53 mmol) in *N,N*-dimethylformamide (DMF) (15 mL) was added an excess of 1-Boc-piperazine (3.09 g, 16.58 mmol). The reaction mixture was heated at 90 °C for 2 h. After the reaction was completed, the reaction mixture was concentrated to give the crude compound **11**, which was used in the next step without further characterization. LC-MS (ES⁺) [M+H]⁺ *m/z* 512.1.

3.2.5 Preparation of 2-(((4-(piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**12**)

A 100 mL round-bottomed flask was charged with compound **11** (1.38 g, 2.70 mmol) and dichloromethane (15 mL). Then, an excess trifluoroacetic acid (10 mL) was added and the reaction was carried out at 0 °C for 8 h. After the reaction was completed, the reaction mixture was transferred to a 100 mL beaker. The resulting mixture was adjusted to pH 6~7 by addition of saturated sodium carbonate solution. It was extracted with dichloromethane for 2~3 times and the organic layer was concentrated to get compound **12** 0.82 g, yield 73.8%. White solid, m.p. 79.4~80.1 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.03 (d, *J*=7.9 Hz, 1H), 7.96 (d, *J*=8.1 Hz, 1H), 7.57~7.46 (m, 1H), 7.46~7.35 (m, 1H), 7.06 (s, 1H), 4.82 (s, 2H), 3.87 (s, 4H), 3.07~2.82 (m, 4H), 1.19 (d, *J*=7.3 Hz, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 170.0, 169.3, 160.9, 153.3, 152.9, 152.4, 134.9, 126.1, 125.1, 122.2, 119.4, 96.6, 45.2,

43.0, 32.7. HRMS (ESI) calcd for $C_{17}H_{17}F_3N_5S_2$ $[M+H]^+$ 412.0877, found 412.0869.

3.2.6 Preparation of compounds **13a**~**13i**

A 50 mL round bottom flask was charged with compound **12** (1.00 mmol) and *N,N*-dimethylformamide (DMF) (5 mL), and different chlorine-substituted compounds (1.20 mmol) were added with stirring, respectively. The mixture was heated to 90 °C for 3 h. After the reaction was completed (monitored by TLC), the reaction mixture was cooled to room temperature and poured into the ice-water (200 mL). The suspension was filtered off via suction filtration to give compounds **13a**~**13i**, which were purified by chromatography (petroleum ether/ethyl acetate, $V:V=2\sim3:1$).

2-(((4-(4-Methylpiperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13a**): Light brown solid, yield 82.3%. m.p. 83.2~83.7 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.01 (d, $J=7.9$ Hz, 1H), 7.94 (d, $J=8.1$ Hz, 1H), 7.49 (dd, $J=11.3, 4.0$ Hz, 1H), 7.40 (t, $J=7.3$ Hz, 1H), 7.00 (s, 1H), 4.80 (s, 2H), 3.66 (s, 4H), 2.23 (s, 4H), 2.13 (s, 3H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.0, 169.3, 160.9, 153.2, 152.9, 152.5, 135.0, 126.0, 125.0, 122.2, 122.0, 96.3, 53.9, 45.3, 43.7, 32.7.

2-(((4-(4-Ethylpiperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13b**): Light brown solid, yield 77.6%. m.p. 90.1~91.5 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.02 (d, $J=7.8$ Hz, 1H), 7.94 (d, $J=8.0$ Hz, 1H), 7.52~7.47 (m, 1H), 7.40 (t, $J=7.2$ Hz, 1H), 7.00 (s, 1H), 4.79 (s, 2H), 3.65 (d, $J=28.6$ Hz, 4H), 2.54~2.49 (m, 2H), 2.25 (q, $J=7.2$ Hz, 4H), 0.95 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.2, 169.3, 160.8, 153.2, 152.8, 152.5, 134.9, 126.0, 125.0, 122.2, 122.1, 96.2, 51.6, 51.2, 32.7, 28.9, 11.7.

2-(((4-(4-Butylpiperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13c**): Light brown solid, yield 80.4%. m.p. 84.2~84.8 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.01 (d, $J=7.8$ Hz, 1H), 7.94 (d, $J=8.1$ Hz, 1H), 7.49 (t, $J=7.2$ Hz, 1H), 7.40 (t, $J=7.2$ Hz, 1H), 6.99 (s, 1H), 4.79 (s, 2H), 3.81~3.50 (m, 4H), 2.54~2.21 (m, 4H), 2.17 (d, $J=7.3$ Hz, 2H), 1.35 (dq, $J=14.3, 6.8$ Hz, 2H), 1.29~1.23 (m, 2H), 0.88 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.2, 169.3, 160.8, 153.2, 152.8, 152.5, 134.9, 126.0, 125.0, 122.2, 122.1, 96.2, 57.1, 52.1, 43.3, 32.7, 28.1, 19.9, 13.8. HRMS (ESI) calcd for $C_{21}H_{25}F_3N_5S_2$ $[M+H]^+$ 468.1503, found 468.1494.

2-(4-(2-((Benzo[d]thiazol-2-ylmethyl)thio)-6-(trifluoromethyl)pyrimidin-4-yl)piperazin-1-yl)ethan-1-ol (**13d**): Light brown solid, yield 73.2%. m.p. 81.3~82.4 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.02 (d, $J=7.9$ Hz, 1H), 7.95 (d, $J=8.1$ Hz, 1H), 7.54~7.46 (m, 1H), 7.46~7.36 (m, 1H), 7.00 (s, 1H), 4.79 (s, 2H), 4.45 (s, 1H), 3.63 (s, 4H), 3.49 (s, 2H), 2.51 (s, 2H), 2.36 (t, $J=6.0$ Hz, 4H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.1, 169.3, 160.8, 152.8, 152.5, 134.9, 126.1, 125.0, 122.2, 122.1, 119.4, 96.2, 59.8, 58.3, 52.4, 43.4, 32.7.

2-(((4-(4-Phenylpiperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13e**): Light

brown solid, yield 61.6%. m.p. 79.1~80.5 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.01 (t, $J=7.1$ Hz, 2H), 7.52 (t, $J=7.7$ Hz, 1H), 7.41 (t, $J=7.7$ Hz, 1H), 7.22 (t, $J=7.9$ Hz, 2H), 7.07 (s, 1H), 6.87~6.79 (m, 3H), 4.82 (s, 2H), 3.83 (s, 4H), 3.04 (s, 4H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.4, 169.4, 160.9, 153.3, 152.9, 152.5, 150.4, 135.0, 128.9, 126.1, 125.0, 122.3, 122.1, 119.4, 119.2, 115.6, 96.4, 47.7, 32.8.

2-(((4-(4-(4-Methoxyphenyl)piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13f**): Light brown solid, yield 67.0%. m.p. 82.6~83.1 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.03~7.99 (m, 2H), 7.54~7.50 (m, 1H), 7.44~7.40 (m, 1H), 7.07 (s, 1H), 6.85~6.79 (m, 4H), 4.83 (s, 2H), 3.86 (d, $J=21.7$ Hz, 4H), 3.71 (s, 3H), 2.89 (d, $J=31.5$ Hz, 4H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.3, 169.4, 160.8, 153.3, 152.9, 152.5, 144.7, 135.0, 126.1, 125.0, 122.3, 122.1, 119.4, 117.8, 114.2, 96.3, 55.1, 49.2, 43.3, 32.8.

2-(((4-(4-(4-Methoxybenzyl)piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13g**): White solid, yield 82.9%. m.p. 96.1~96.8 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.02 (d, $J=7.7$ Hz, 1H), 7.90 (d, $J=8.0$ Hz, 1H), 7.53~7.46 (m, 1H), 7.45~7.38 (m, 1H), 7.15 (d, $J=8.6$ Hz, 2H), 6.97 (s, 1H), 6.89 (d, $J=8.6$ Hz, 2H), 4.77 (s, 2H), 3.75 (s, 2H), 3.73~3.42 (m, 4H), 3.33 (s, 3H), 2.40~2.05 (m, 4H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.2, 169.3, 160.7, 158.3, 153.2, 152.8, 152.4, 134.9, 130.1, 129.1, 127.8, 126.1, 125.0, 122.2, 122.1, 113.5, 96.1, 60.9, 54.9, 51.6, 32.7.

2-(((4-(4-(Pyridin-2-yl)piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13h**): Light brown solid, yield 65.2%. m.p. 97.1~98.4 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.15~8.12 (m, 1H), 8.01 (t, $J=8.6$ Hz, 2H), 7.58~7.49 (m, 2H), 7.43~7.38 (m, 1H), 7.05 (s, 1H), 6.77 (d, $J=8.6$ Hz, 1H), 6.67 (dd, $J=6.9, 5.0$ Hz, 1H), 4.83 (s, 2H), 3.80 (d, $J=30.7$ Hz, 4H), 3.61~3.37 (m, 4H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 170.15, 169.35, 160.98, 158.43, 153.20, 152.87, 152.50, 147.54, 137.58, 135.00, 126.15, 125.09, 122.34, 122.16, 113.25, 107.07, 96.41, 59.71, 43.81, 32.74. HRMS (ESI) calcd for $C_{22}H_{20}F_3N_6S_2$ $[M+H]^+$ 489.1143, found 489.1128.

2-(((4-(4-(Pyrimidin-2-yl)piperazin-1-yl)-6-(trifluoromethyl)pyrimidin-2-yl)thio)methyl)benzo[d]thiazole (**13i**): Light brown solid, yield 68.1%. m.p. 96.7~97.5 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.37 (d, $J=4.7$ Hz, 2H), 7.96 (dd, $J=12.3, 8.0$ Hz, 2H), 7.50~7.43 (m, 1H), 7.41~7.34 (m, 1H), 7.03 (s, 1H), 6.63 (t, $J=4.7$ Hz, 1H), 4.82 (s, 2H), 3.76 (s, 8H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 169.8, 169.4, 161.0, 160.9, 157.8, 152.9, 152.4, 135.0, 126.0, 125.0, 122.2, 122.0, 119.4, 110.4, 96.4, 59.7, 42.5, 32.6. HRMS (ESI) calcd for $C_{21}H_{19}F_3N_7S_2$ $[M+H]^+$ 490.1095, found 49.1080.

Supporting Information The 1H NMR and ^{13}C NMR spectra of compounds **12** and **13a**~**13i**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Samadder, N. J.; Giridhar, K. V.; Baffy, N.; Riegert, J. D.; Couch, F. *J. Mayo Clin. Proc.* **2019**, *94*, 1084.
- [2] Siegel, R. L.; Miller, K. D.; Jemal, A. *CA-Cancer J. Clin.* **2015**, *65*, 5.
- [3] Ferlay, J.; Colombet, M.; Soerjomataram, I.; Mathers, C.; Parkin, D. M.; Piñeros, M.; Znaor, A.; Bray, F. *Int. J. Cancer.* **2019**, *144*, 1941.
- [4] Banerjee, S.; Arnst, K. E.; Wang, Y.; Kumar, G.; Deng, S.; Yang, L.; Miller, D. D. *J. Med. Chem.* **2018**, *61*, 1704.
- [5] Abdel, M.; Heba, T.; Ragab, F. A.; Ramla, M. M. *Eur. J. Med. Chem.* **2010**, *45*, 2336.
- [6] Bekhit, A. A.; Fahmy, H. T.; Rostom, S. A.; Baraka, A. M. *Eur. J. Med. Chem.* **2003**, *38*, 27.
- [7] Rostamizadeh, S.; Nojavan, M.; Aryan, R.; Sadeghian, H.; Davoodnejad, M. *Chin. Chem. Lett.* **2013**, *24*, 629.
- [8] Tozkoparan, B.; Ertan, M.; Kelicen, P. *Farmaco* **1999**, *54*, 588.
- [9] Deyanov, A. B.; Niyazov, R. K.; Syropyatov, B. Y.; Kolla, V. E.; Konshin, M. E. *J. Pharm. Chem.* **1991**, *25*, 248.
- [10] Nalini, C. N.; Amuthalakshmi, S.; Shanthanakrishnan, V. S. *Curr. Microwave Chem.* **2018**, *5*, 189.
- [11] Noolvi, M. N.; Harun, M. P.; Manpreet, K. *Eur. J. Med. Chem.* **2012**, *54*, 447.
- [12] Cai, J.; Sun, M.; Wu, X.; Chen, J.; Wang, P.; Zong, X.; Ji, M. *Eur. J. Med. Chem.* **2013**, *63*, 702.
- [13] Christopher, J.; Bungard, P. D.; Williams, J. S. *Med. Chem. Lett.* **2017**, *8*, 1292.
- [14] Zhang, Q. W.; Zhang, B. Y.; Zhou, A. N.; Zhang, Y.; Li, J. Q. *Chin. J. Med. Chem.* **2015**, *4*, 255.
- [15] Meng, Y. Q.; Li, E. D.; Zhang, Y.; Liu, S.; Bao, C. N.; Yang, P.; Zhang, L. Y.; Zhang, D. Q.; Wang, J. K.; Chen, Y. X.; Li, N.; Xin, J. C.; Zhao, P. R.; Ke, Y.; Zhang, Q. R.; Liu, H. M. *Chin. J. Org. Chem.* **2019**, *39*, 2541 (in Chinese).
(孟娅琪, 李二冬, 张洋, 刘栓, 包崇男, 杨鹏, 张路野, 张丹青, 王继宽, 陈雅欣, 栗娜, 辛景超, 赵培, 可钰, 张秋荣, 刘宏民, 有机化学, **2019**, *39*, 2541.)
- [16] Li, N.; Xin, J. C.; Ma, Q. S.; Li, E. D.; Meng, Y. Q.; Bao, C. N.; Yang, P.; Song, P. P.; Cui, F.; Cheng, P. J.; Gu, Y. F.; Zhao, P. R.; Ke, Y.; Liu, H. M.; Zhang, Q. R. *Chin. J. Org. Chem.* **2018**, *38*, 665 (in Chinese).
(栗娜, 辛景超, 马启胜, 李二冬, 孟娅琪, 包崇男, 杨鹏, 宋攀攀, 崔飞, 陈鹏举, 顾一飞, 赵培, 可钰, 刘宏民, 张秋荣, 有机化学, **2018**, *38*, 665.)
- [17] Xin, J. C.; Li, N.; Ma, Q. S.; Li, E. D.; Meng, X. C.; Ke, Y.; Liu, H. M.; Zhang, Q. R. *Chin. J. Org. Chem.* **2018**, *38*, 451 (in Chinese).
(辛景超, 栗娜, 马启胜, 李二冬, 孟祥川, 可钰, 刘宏民, 张秋荣, 有机化学, **2018**, *38*, 451.)
- [18] Song, P. P.; Li, N.; Cui, F.; Xin, J. C.; Zhang, X. S.; Cao, Q. P.; Wang, C. J.; Dai, W. J.; Meng, X. C.; Liu, M.; Chang, T. H.; Liu, Q. Y.; Sun, Y. H.; Ke, Y.; Zhang, Q. R.; Liu, H. M. *Chin. J. Org. Chem.* **2017**, *37*, 2725 (in Chinese).
(宋攀攀, 栗娜, 崔飞, 辛景超, 张孝松, 曹钦坡, 王超杰, 戴文杰, 孟祥川, 刘梦, 常通航, 柳晴怡, 孙月红, 可钰, 张秋荣, 刘宏民, 有机化学, **2017**, *37*, 2725.)
- [19] Miller, W. H.; Dessert, A. M.; Anderson, G. W. *J. Am. Chem. Soc.* **1948**, *70*, 500.
- [20] Puranik, N. V.; Hemalata, M. P.; Srivastava, P. *Med. Chem. Res.* **2016**, *4*, 805.

(Lu, Y.)