锰介导的 Reformatsky 反应: 高效合成多样化 β -羟基链烷酸酯

夏艳萍 欧阳露 廖建华 魏一飞 罗人仕*

(赣南医学院药学院 江西赣州 341000)

摘要 发展了一种由锰介导的简易高效的 Reformatsky 催化体系. 在便宜易得锰粉的介导下, 实现了镍催化剂高效催化碘代乙酸乙酯与醛酮的 Reformatsky 反应. 以乙腈为反应溶剂, 温和条件下可获得高收率的 Reformatsky 加成产物(up to 98%). 此外, 该催化方法学还可放大至克级规模.

关键词 锰; 镍催化剂; Reformatsky; 醛酮; β -羟基链烷酸酯

Manganese-Mediated Reformatsky Reaction: Highly Divergent Synthesis of β -Hydroxyalkanoates

Xia, Yanping Ouyang, Lu Liao, Jianhua Wei, Yifei Luo, Renshi*

(School of Pharmacy, Gannan Medical University, Ganzhou, Jiangxi 341000)

Abstract A practical, flexible, and efficient manganese-mediated catalytic system of Reformatsky reaction has been described. The cheap and readily available manganese powder acts as the reaction mediator, which conducts effectively for the preparation of useful β -hydroxyalkanoate compounds with the nickel catalyst in excellent yields (up to 98%) and mild condition. In addition, the catalytic methodology can be scaled up to the gram scale.

Keywords manganese; nickel catalyst; reformatsky; aldehydes and ketones; β -hydroxyalkanoates

1 Introduction

The β -hydroxyl carbonyl scaffold is an important pharmacophore and valuable synthon.^[1] In the past few decades, efficient assembly of this type of structure has attracted wide attention on the pharmaceutical industry and medicinal chemistry based on the fact that this structure widely exists in bioactive molecules, pharmaceuticals agents and natural products.^[2] Thus, the development of efficient synthesis of β -hydroxyl carbonyl derivatives is extremely significant.

The classical Reformatsky reaction, reported for the first time in 1887,^[3] has been proven to be the most direct and effective strategy for the construction of β -hydroxyl carbonyl compounds.^[4] Therefore, The Reformatsky reaction has become an efficient synthetic method for the construction of C—C bond, which has been widely applied in organic chemistry.^[5] Traditional method divides this transformation into two steps. The first step is the preparation of organic metallic reagent. Then the second step is the addition

of different kinds of electrophiles such as aldehydes, ketones and imines with organic metallic reagent, which is characterized by complex operation and lower efficiency.^[6] For example, a wide variety of metals or low-valent metallic salts, such as Zn powder^[7] or R_2Zn ,^[8] Mg,^[9] Sm(II) salt,^[10] are served as mediators in the Reformatsky reaction (Scheme 1a), which needs long reaction time in low yield. Therefore, it becomes essential to develop a more simple, inexpensive, and practical strategy for the Reformatsky reaction.

Manganese is one of the most abundant elements on the earth, which has the advantages of abundant source, cheap price and harmlessness to environment. Recently, more and more attention has been focused on green chemistry along with people's awareness of the importance of environment.^[11] Hence, we present a new nickel-catalyzed and manganese-mediated efficient Reformatsky reaction of aldehydes and ketones with ethyl iodoacetate, which is undoubtedly eco-friendly with cheap raw material. This strategy relies on simple and readily available materials to

* Corresponding author. E-mail: luorensi2010@163.com

Received June 29, 2020; revised July 30, 2020; published online August 26, 2020.

Project supported by the the National Natural Science Foundation of China (Nos. 21962004, 21562004), Jiangxi provincial department of science and technology (No. 20192BAB203004); Gannan Medical University (No. QD201810, YJ202027).

国家自然科学基金(Nos. 21962004, 21562004)、江西省自然科学基金(No. 20192BAB203004)、赣南医学院(Nos. QD201810, YJ202027)资助项目.

Scheme 1 Reformatsky reaction for the construction of β -hydroxyl carbonyl scaffold

generate useful β -hydroxyl esters with the advantages of high efficiency, simple operation and mild condition.

2 Results and discussion

Initially, we began our studies by using benzaldehyde (**1a**) and ethyl bromoacetate (**2b**) as the template reaction, and examined the reaction parameters including solvent and additive to identify the optimal reactions conditions (Table 1). In the absence of Ni catalyst, only 12% yield was obtained (Entry 1). In addition, the higher activity ethyl iodoacetate (**2a**) was also chosen as nucleophile, giving **3a** in the yield of 25% (Entry 2). To improve the activity, NiBr_2 was added as catalyst, but which also gave **3a** in low yield with the nucleophile of **2b** (Entry 3). However, when 1.0 equiv. of NaI was added to produce **2a** *in situ* under standard condition, the yield increased rapidly and reached to 90% (Entry 4). The evaluation of a number of common Ni catalysts, including NiBr_2 , NiCl_2 and NiI_2 (Table 1, Entries 5~7) were screened. In the presence of NiCl_2 , only 23% yield of desired product was obtained. To our delight, the reaction catalyzed by NiI_2 gave the best results with the yield of 93%. Even more important, a lower amount of catalyst (10 mol%) and a shorter reaction time (12 h) can also realize the transformation efficiently (Table 1, Entry 8). The screening of the solvents, such as Et_2O , dioxane, toluene, DCM, hexane, MeCN, confirmed that solvent had a strong influence on the reaction. The use of Et_2O and dioxane as the solvent could give the expected product **3a** in good yield (Entries 9 and 10). However, a lower conversion of **3a** was observed by using toluene, hexane and dichloromethane (DCM) as the solvents (Entries 11~13). MeCN was the best solvent to give the target

product in excellent yield under standard conditions.

Table 1 Optimization of reaction conditions^a

Entry	X	[Ni]	Solvent	Time/h	Yield ^b / % of 3a
1	Br	—	THF	36	12
2	I	—	THF	36	25
3	Br	NiBr_2	THF	36	27
4 ^c	Br	NiBr_2	THF	36	90
5	I	NiBr_2	THF	36	94
6	I	NiCl_2	THF	36	23
7	I	NiI_2	THF	36	92
8 ^d	I	NiI_2	THF	12	92
9 ^d	I	NiI_2	Et_2O	12	93
10 ^d	I	NiI_2	Dioxane	12	90
11 ^d	I	NiI_2	Toluene	12	78
12 ^d	I	NiI_2	DCM	12	80
13 ^d	I	NiI_2	Hexane	12	85
14 ^d	I	NiI_2	MeCN	12	98

^a Reaction conditions: All reactions were performed with **1a** (0.5 mmol), **2a** (1.5 equiv.), nickel catalyst (20 mol%), Mn powder (3 equiv.), CF_3COOH (20 mol%), solvent (2 mL), at 0 °C under N_2 atmosphere. ^b Yield was determined by GC with dodecane as internal standard based on **1a**. ^c 1.0 equiv. of NaI was added. ^d NiI_2 (10 mol%) was added.

With the optimized reaction condition in hand, a broad range of aldehydes were investigated to afford the β -hydroxyalkanoate compounds under the optimal conditions. As shown in Table 2, the nickel-catalyzed and manganese-mediated Reformatsky reaction proceeded smoothly over a wide range of aldehydes in good to excellent yields. Firstly, substrates with different halogen on aryl were surveyed, which obtained the desired products in the yield of 90%~96% (**3c**, **3d**, **3g**, **3h**, and **3r**~**3t**). Besides, the halogen atom with different position on the aromatic ring can also suit for this reaction condition (**3c**, **3d**, **3f** and **3g**). In addition, using the electron-donating groups, such as methyl and methoxyl, as the substrates, the reaction also proceeded well (**3b**, **3e**, **3f** and **3i**). It is worth noting that the aldehyde with higher conjugation degree (2-naphthaldehyde) performed well to access the desired products (**3j**). Gratifyingly, this protocol could be applied to aromatic heterocyclic aldehydes. For example, 2-thienaldehyde and furfural gave the desired products **3k** and **3l** in 92% and 95% yields. Notably, polysubstituted aromatic aldehydes, with the strong steric hindrance, still smoothly took place under the standard conditions (**3m**, **3o**, **3p**). Meanwhile, the substrate 2,4,6-trimethylbenzaldehyde could also be employed in the catalytic system to afford the product **3q** in 86% yield.

The scope of nickel-catalyzed and manganese-mediated Reformatsky reaction was further investigated to a variety of aliphatic aldehydes (Table 3). Gratifyingly, under the standard reaction conditions, the unsaturated aldehydes

Table 2 Screening the viable aromatic aldehydes^a

^a Reaction conditions: aldehyde (**1**) as the limiting reagent (0.5 mmol), **2a** (1.5 equiv.), NiI₂ (10 mol%), Mn powder (3 equiv.), CF₃COOH (20 mol%), MeCN (2 mL), 0 °C under N₂ atmosphere for 12 h.

Table 3 Screening the viable aliphatic aldehydes^a

^a Reaction conditions: aldehyde (**1**) as the limiting reagent (0.5 mmol), **2a** (1.5 equiv.), NiI₂ (10 mol%), Mn powder (3 equiv.), CF₃COOH (20 mol%), MeCN (2 mL), 0 °C under N₂ atmosphere for 12 h.

such as cinnamaldehydes were also tolerated in this process, which gave the corresponding products **3u**, **3v** and **3w** in good yields, especially for **3v** with large steric hindrance. Other aliphatic aldehyde was also conducted to this reaction to give the corresponding product ethyl-3-cyclohexyl-3-hydroxypropanoate efficiently (**3x**). In view of the great importance of β-hydroxyalkanoates scaffold in bio-

medicine, it is important to develop late-stage modification of β-hydroxyalkanoates. For example, natural aldehydes, such as citral (**1y**), could participate in the present transformation to generate the corresponding β-hydroxyalkanoate **3y**, respectively.

Apart from aldehydes, ketones could also be used as electrophiles in the presence of nickel-catalyzed Reformatsky reaction. Fortunately, when the above optimal reaction conditions were applied to acetophenone, 1-(*p*-tolyl)ethan-1-one, ethyl 3-hydroxy-3-(4-(trifluoromethyl)phenyl)butanoate and 1-(naphthalen-2-yl)ethan-1-one, the desired products were also obtained in lower yield (Table 4).

Besides broad functional group tolerance, the practicability of the developed protocol was tested. A scale-up experiment (**1a**, 10 mmol) was smoothly conducted under the standard conditions, which β-hydroxyalkanoate **3aa** was obtained in 91% yield (Scheme 2).

On the basis of results, a proposed mechanism was proposed for the Ni-catalyzed Reformatsky reaction in Scheme 3. The reaction was initiated with Ni⁰. Therefore, the Ni^{II} was reduced to Ni⁰ by manganese powder firstly. Then oxidation-addition of Ni⁰ was occurred and achieved **int-I**.^[12] While, the electronegativity of nickel is low, and low initial activity of **int-I** is exhibited in the cycle, which is unable to react with the carbonyl compounds directly.^[13] In order to enhance the nucleophiles, anion exchange was occurred with more acidic Bronsted acid to produced **int-II**. Then, **Int-III** was obtained by the nucleophilic ad-

Table 4 Screening the viable ketones^a

^a Reaction conditions: ketones (**4**) as the limiting reagent (0.5 mmol), **2a** (1.5 equiv.), NiI_2 (10 mol%), Mn powder (3 equiv.), CF_3COOH (20 mol%), MeCN (2 mL), 0 °C under N_2 atmosphere for 36 h.

Scheme 2 Gram-scale experiment

dition of aldehyde and **Int-II**. Hydrolysis of **Int-III** gave the desired products, and Ni^{II} complex was reduced to Ni^0 for the next catalytic cycles (Path A). Besides, the alkyl- $\text{Ni}(\text{II})$ may also be reduced to alkyl- $\text{Ni}(\text{I})$ by $\text{Mn}(0)$ firstly, then the alkyl- $\text{Ni}(\text{I})$ is able to attack the $\text{C}=\text{O}$ bond (Path B). The similar process was reported by Ruben Martin.^[14]

3 Conclusion

In conclusion, a versatile and convenient strategy for the

preparation of functionalized β -hydroxyalkanoates via nickel-catalyzed Reformatsky reaction of aldehydes and ketones with ethyl iodacetate is developed. Moreover, the simple and readily available starting materials, operating procedure and good yields make this protocol particularly practical, convenient and mild. In short, this protocol may open up a new viewpoint on classical Reformatsky reaction. Ongoing studies are focused on exploring the asymmetric catalytic Reformatsky reaction and the application.

4 Experimental section

4.1 General information

^1H NMR and ^{13}C NMR spectra were recorded by using a Bruker DRX-400 spectrometer (400 MHz for ^1H NMR; 100 MHz for ^{13}C NMR), using CDCl_3 as solvent and TMS as an internal standard. The chemical shifts are referenced to signals at δ 7.26 and 77.0, respectively. Chemical shifts (δ) are reported and quoted to the nearest δ 0.01 relative to the residual protons in CDCl_3 (δ 7.26 for ^1H NMR) or TMS (δ 0 for ^1H NMR) and CDCl_3 (δ 77.0 for ^{13}C NMR). GC analyses were performed on a GC-7900 chromatograph with an FID and equipped with an AT-SE-30 capillary column (internal diameter: 0.32 mm, length: 30 m).

4.2 Procedure for the preparation of **3** and **5**

To a 25 mL dried Shrek tube was added the mixture of anilines **1** or **4** (0.5 mmol), Mn powder (3 equiv.), NiI_2 (10 mol%), ethyl iodacetate (**2a**) (0.75 mmol), CF_3COOH (25 mol%) in MeCN (2.0 mL) successively. The mixture was stirred at room temperature for 12 h under an N_2 atmosphere. After the reaction was completed, the mixture was diluted with H_2O (15 mL), neutralized with NH_4Cl , and extracted with EtOAc (10 mL $\times$ 3). The organic extract was washed with brine (10 mL $\times$ 3) and dried over anhydrous MgSO_4 . After removal of the EtOAc in vacuum, the crude product was purified by column chromatography on silica gel with hexanes or petroleum ether/ethyl acetate (V : V = 5 : 1 to 20 : 1) to give the desired products **3** and **5**.

Scheme 3 Proposed reaction mechanism

Ethyl 3-hydroxy-3-phenylpropanoate (**3a**):^[5d] Yield: 98% (95.1 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.41~7.28 (m, 5H), 5.19~5.11 (m, 1H), 4.20 (q, *J*=7.1 Hz, 2H), 3.39 (d, *J*=3.4 Hz, 1H), 2.77~2.73 (m, 2H), 1.28 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.4, 142.6, 128.6, 127.8, 125.7, 70.3, 60.9, 43.4, 14.2.

Ethyl 3-hydroxy-3-(p-tolyl)propanoate (**3b**):^[5d] Yield: 95% (98.8 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.31~7.27 (m, 2H), 7.18 (d, *J*=8.0 Hz, 2H), 5.11 (dt, *J*=7.4, 3.4 Hz, 1H), 4.20 (q, *J*=7.1 Hz, 2H), 3.32 (d, *J*=3.4 Hz, 1H), 2.80~2.67 (m, 2H), 2.36 (s, 3H), 1.28 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.4, 139.7, 137.5, 129.2, 125.7, 70.2, 60.8, 43.4, 21.1, 14.2.

Ethyl 3-(4-chlorophenyl)-3-hydroxypropanoate (**3c**):^[8a] Yield: 93% (106.0 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.34~7.27 (m, 4H), 5.09 (dd, *J*=8.2, 4.5 Hz, 1H), 4.17 (q, *J*=7.1 Hz, 2H), 3.59 (s, 1H), 2.75~2.63 (m, 2H), 1.26 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.2, 141.1, 133.4, 128.7, 127.1, 69.6, 61.0, 43.3, 14.1.

Ethyl 3-(4-fluorophenyl)-3-hydroxypropanoate (**3d**):^[8a] Yield: 90% (95.4 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.41~7.33 (m, 2H), 7.09~7.01 (m, 2H), 5.17~5.09 (m, 1H), 4.20 (q, *J*=7.1 Hz, 2H), 3.41 (d, *J*=2.6 Hz, 1H), 2.78~2.65 (m, 2H), 1.28 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.3, 162.2 (d, *J*=244 Hz), 138.3 (d, *J*=4 Hz), 127.3 (d, *J*=8 Hz), 115.3 (d, *J*=11 Hz), 69.7, 61.0, 43.3, 14.1.

Ethyl 3-hydroxy-3-(4-methoxyphenyl)propanoate (**3e**):^[8a] Yield: 92% (103.0 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.32 (d, *J*=8.6 Hz, 2H), 6.91 (d, *J*=8.7 Hz, 2H), 5.11 (dt, *J*=8.8, 3.4 Hz, 1H), 4.20 (q, *J*=7.1 Hz, 2H), 3.82 (s, 3H), 3.22 (d, *J*=3.3 Hz, 1H), 2.73 (m, 2H), 1.34 (dt, *J*=14.3, 13.6 Hz, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.5, 159.2, 134.7, 127.0, 113.9, 70.0, 60.9, 55.3, 43.3, 14.2.

Ethyl 3-hydroxy-3-(o-tolyl)propanoate (**3f**):^[15] Yield: 90% (93.6 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.52 (d, *J*=7.5 Hz, 1H), 7.28~7.15 (m, 3H), 5.37 (dd, *J*=5.9, 2.9 Hz, 1H), 4.22 (q, *J*=7.1 Hz, 2H), 3.29 (d, *J*=2.4 Hz, 1H), 2.76~2.64 (m, 2H), 2.37 (s, 3H), 1.30 (t, *J*=7.1 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 172.6, 140.5, 134.3, 130.46, 127.6, 126.4, 125.3, 67.0, 60.9, 42.2, 19.0, 14.2.

Ethyl 3-(2-fluorophenyl)-3-hydroxypropanoate (**3g**):^[15] Yield: 91% (103.7 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.63 (d, *J*=7.6 Hz, 1H), 7.31 (dd, *J*=14.1, 7.6 Hz, 2H), 7.21 (t, *J*=7.5 Hz, 1H), 5.49 (d, *J*=9.6 Hz, 1H), 4.19 (q, *J*=7.0 Hz, 2H), 3.60 (s, 1H), 2.84 (d, *J*=16.5 Hz, 1H), 2.58 (dd, *J*=16.4, 9.7 Hz, 1H), 1.27 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.5, 140.0, 131.4, 129.3, 128.7, 127.2, 127.1, 67.0, 61.0, 41.5, 14.1.

Ethyl 3-(2-fluorophenyl)-3-hydroxypropanoate (**3h**):^[16] Yield: 93% (98.6 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.55 (t, *J*=7.5 Hz, 1H), 7.31~7.23 (m, 1H), 7.17 (t, *J*=7.5 Hz, 1H), 7.07~6.99 (m, 1H), 5.46~5.37

(m, 1H), 4.19 (q, *J*=7.2 Hz, 2H), 3.65 (d, *J*=4.0 Hz, 1H), 2.83~2.69 (m, 2H), 1.27 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.40, 159.4 (d, *J*=254 Hz), 129.4 (d, *J*=13 Hz), 129.1 (d, *J*=9 Hz), 127.2 (d, *J*=4 Hz), 124.4 (d, *J*=4 Hz), 115.2 (d, *J*=11 Hz), 64.56, 64.54, 60.96, 41.94, 41.93, 14.12.

Ethyl 3-hydroxy-3-(2-methoxyphenyl)propanoate (**3i**):^[15] Yield: 90% (100.8 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.45 (dd, *J*=7.5, 1.1 Hz, 1H), 7.31~7.26 (m, 1H), 7.00 (t, *J*=7.4 Hz, 1H), 6.90 (d, *J*=8.2 Hz, 1H), 5.38 (dd, *J*=9.2, 3.6 Hz, 1H), 4.20 (q, *J*=7.1 Hz, 2H), 3.87 (s, 3H), 2.85 (dd, *J*=16.1, 3.6 Hz, 1H), 2.73 (dd, *J*=16.1, 9.2 Hz, 1H), 1.29 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.7, 156.1, 130.5, 128.6, 126.6, 120.8, 110.3, 66.6, 60.7, 55.3, 41.7, 14.2.

Ethyl 3-hydroxy-3-(naphthalen-2-yl)propanoate (**3j**):^[5d] Yield: 90% (109.8 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 8.09 (d, *J*=8.2 Hz, 1H), 7.92~7.88 (m, 1H), 7.81 (d, *J*=8.2 Hz, 1H), 7.73 (d, *J*=7.1 Hz, 1H), 7.57~7.47 (m, 3H), 5.95 (dd, *J*=9.2, 2.7 Hz, 1H), 4.24 (q, *J*=7.1 Hz, 2H), 3.63 (s, 1H), 2.89 (qd, *J*=16.4, 6.3 Hz, 2H), 1.30 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.7, 138.2, 133.8, 130.0, 129.0, 128.3, 126.3, 125.6, 125.5, 123.0, 122.9, 67.4, 61.0, 42.8, 14.2.

Ethyl 3-hydroxy-3-(thiophen-2-yl)propanoate (**3k**):^[5d] Yield: 92% (92.0 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.27 (dd, *J*=4.7, 1.5 Hz, 1H), 6.98 (q, *J*=3.5 Hz, 2H), 5.42~5.32 (m, 1H), 4.20 (q, *J*=7.1 Hz, 2H), 3.59 (s, 1H), 2.92~2.81 (m, 2H), 1.28 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 171.9, 146.4, 126.7, 124.9, 123.7, 66.6, 61.0, 43.2, 14.2.

Ethyl 3-(furan-2-yl)-3-hydroxypropanoate (**3l**):^[8a] Yield: 95% (87.4 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.40 (dd, *J*=1.8, 0.8 Hz, 1H), 6.36 (dd, *J*=3.2, 1.8 Hz, 1H), 6.30 (d, *J*=3.3 Hz, 1H), 5.16 (dt, *J*=8.6, 4.4 Hz, 1H), 4.25~4.19 (m, 2H), 3.25 (d, *J*=5.0 Hz, 1H), 2.96~2.83 (m, 2H), 1.32~1.30 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.0, 154.7, 142.3, 110.3, 106.3, 64.2, 61.0, 39.8, 14.2.

Ethyl 3-(2,3-dimethylphenyl)-3-hydroxypropanoate (**3m**): Yield: 89% (98.8 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.40 (d, *J*=7.5 Hz, 1H), 7.19~7.10 (m, 2H), 5.44 (dd, *J*=8.2, 4.5 Hz, 1H), 4.23 (q, *J*=7.2 Hz, 2H), 2.75~2.65 (m, 2H), 2.32 (s, 3H), 2.26 (s, 3H), 1.31 (t, *J*=7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.7, 140.5, 137.0, 132.9, 129.2, 125.9, 123.0, 67.4, 60.9, 42.4, 20.6, 14.6, 14.2.

Ethyl 3-(2,5-dimethylphenyl)-3-hydroxypropanoate (**3o**):^[17] Yield: 94% (104.3 mg), colourless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.35 (s, 1H), 7.03 (q, *J*=7.7 Hz, 2H), 5.34 (d, *J*=9.1 Hz, 1H), 4.21 (q, *J*=7.1 Hz, 2H), 3.46 (s, 1H), 2.74~2.62 (m, 2H), 2.34 (d, *J*=11.4 Hz, 6H), 1.30 (td, *J*=7.1, 1.3 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.6, 140.5, 140.5, 135.77, 131.1, 130.4, 128.2, 125.9, 67.0, 60.8, 42.4, 21.1, 18.5, 14.2.

Ethyl 3-(2,6-dimethylphenyl)-3-hydroxypropanoate (**3p**):^[17] Yield: 87% (96.6 mg), colourless oil. ¹H NMR

(400 MHz, CDCl_3) δ : 7.09 (dd, $J=8.4$, 6.4 Hz, 1H), 7.01 (d, $J=7.5$ Hz, 2H), 5.66 (d, $J=10.6$ Hz, 1H), 4.25~4.20 (m, 2H), 3.08 (dd, $J=16.5$, 10.6 Hz, 1H), 3.00 (s, 1H), 2.57 (dd, $J=16.5$, 3.1 Hz, 1H), 2.48 (s, 6H), 1.31 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.7, 137.6, 136.2, 129.4, 127.5, 67.7, 60.9, 39.9, 20.8, 14.2.

Ethyl 3-hydroxy-3-mesitylpropanoate (**3q**):^[8a] Yield: 86% (101.5 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 6.85 (s, 2H), 5.63 (dd, $J=10.6$, 2.1 Hz, 1H), 4.23 (q, $J=7.1$ Hz, 2H), 3.08 (dd, $J=16.5$, 10.7 Hz, 1H), 2.59~2.50 (m, 1H), 2.44 (d, $J=10.2$ Hz, 6H), 2.27 (s, 3H), 1.32 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.7, 137.0, 136.1, 134.6, 130.2, 67.5, 60.9, 40.1, 20.8, 20.7, 14.2.

Ethyl 3-(2,4-difluorophenyl)-3-hydroxypropanoate (**3r**): Yield: 90% (103.5 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.50 (td, $J=8.5$, 6.7 Hz, 1H), 6.92~6.85 (m, 1H), 6.80~6.75 (m, 1H), 5.35 (dd, $J=8.8$, 3.4 Hz, 1H), 4.17 (q, $J=7.1$ Hz, 2H), 3.76 (s, 1H), 2.78~2.65 (m, 2H), 1.26 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.2, 162.3 (dd, $J=246$, 12 Hz), 159.4 (dd, $J=247$, 12 Hz), 128.2 (dd, $J=8.5$, 6.0 Hz), 125.6 (dd, $J=13.5$, 3.0 Hz), 111.4 (dd, $J=21$, 4 Hz), 103.6 (t, $J=25$ Hz), 64.1 (d, $J=2$ Hz), 61.0, 41.9, 14.1.

Ethyl 3-(2,5-difluorophenyl)-3-hydroxypropanoate (**3s**): Yield: 96% (110.4 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.26 (m, 1H), 7.02~6.91 (m, 2H), 5.38 (d, $J=9.3$ Hz, 1H), 4.21 (q, $J=7.1$ Hz, 2H), 3.71 (s, 1H), 2.81 (dd, $J=16.6$, 3.0 Hz, 1H), 2.66 (dd, $J=16.6$, 9.4 Hz, 1H), 1.29 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.3, 159.0 (dd, $J=240.5$, 2.0 Hz), 155.1 (dd, $J=239.5$, 2.0 Hz), 131.4 (dd, $J=16$, 7 Hz), 116.2 (dd, $J=24.5$, 9 Hz), 115.3 (dd, $J=24.5$, 9 Hz), 113.9 (dd, $J=25$, 5 Hz), 64.17 (d, $J=1.0$ Hz), 61.1, 41.7, 14.1.

Ethyl 3-(2,6-difluorophenyl)-3-hydroxypropanoate (**3t**): Yield: 91% (104.6 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.26 (td, $J=8.2$, 4.3 Hz, 1H), 6.95~6.83 (m, 2H), 5.55 (dt, $J=9.4$, 4.9 Hz, 1H), 4.18 (q, $J=7.2$ Hz, 2H), 3.26 (d, $J=6.1$ Hz, 1H), 3.15 (dd, $J=16.3$, 9.5 Hz, 1H), 2.75 (dd, $J=16.4$, 4.2 Hz, 1H), 1.26 (td, $J=7.2$, 1.6 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.6, 161.0 (dd, $J=244$, 8.5 Hz), 129.7 (t, $J=11$ Hz), 117.5 (t, $J=5$ Hz), 111.8 (dd, $J=19.5$, 3.5 Hz), 62.2 (t, $J=3$ Hz), 60.9, 40.8 (t, $J=3$ Hz), 14.0.

Ethyl (*E*)-3-hydroxy-5-phenylpent-4-enoate (**3u**):^[5d] Yield: 93% (102.3 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.40 (d, $J=7.4$ Hz, 2H), 7.34 (t, $J=7.3$ Hz, 2H), 7.29~7.25 (m, 1H), 6.68 (d, $J=15.9$ Hz, 1H), 6.25 (dd, $J=15.9$, 6.0 Hz, 1H), 4.75 (dd, $J=11.4$, 6.0 Hz, 1H), 4.21 (q, $J=7.1$ Hz, 2H), 2.71~2.60 (m, 2H), 1.30 (t, $J=7.1$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ : 172.3, 136.4, 130.8, 129.9, 128.6, 127.8, 126.6, 68.9, 60.9, 41.5, 14.2.

Ethyl (*E*)-3-hydroxy-4-methyl-5-phenylpent-4-enoate (**3v**): Yield: 90% (105.3 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.37~7.22 (m, 5H), 6.61 (s, 1H), 4.63 (dd, $J=11.4$, 5.0 Hz, 1H), 4.26~4.15 (m, 2H), 3.26 (s, 1H), 2.67 (dd, $J=8.5$, 3.7 Hz, 2H), 1.90 (d, $J=3.9$ Hz,

3H), 1.29 (td, $J=7.1$, 5.3 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.6, 138.3, 137.4, 129.0, 128.1, 126.6, 126.0, 73.6, 60.9, 40.4, 14.2, 13.8.

Ethyl (*E*)-5-(4-chlorophenyl)-3-hydroxypent-4-enoate (**3w**): Yield: 92% (116.8 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.29 (d, $J=7.8$ Hz, 2H), 7.14 (d, $J=7.6$ Hz, 2H), 6.64 (d, $J=15.9$ Hz, 1H), 6.19 (dd, $J=15.9$, 6.1 Hz, 1H), 4.73 (d, $J=5.0$ Hz, 1H), 4.21 (dd, $J=14.0$, 6.9 Hz, 2H), 2.65 (d, $J=7.4$ Hz, 2H), 2.36 (s, 3H), 1.30 (t, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.3, 137.7, 133.6, 130.7, 129.3, 128.9, 126.5, 69.0, 60.9, 41.6, 21.2, 14.2.

Ethyl 3-cyclohexyl-3-hydroxypropanoate (**3x**):^[11] Yield: 96% (96.0 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 4.16 (q, $J=7.1$ Hz, 2H), 3.76 (dd, $J=10.3$, 4.5 Hz, 1H), 2.81 (s, 1H), 2.43 (dt, $J=16.2$, 12.8 Hz, 2H), 1.86 (d, $J=12.6$ Hz, 1H), 1.75 (d, $J=9.3$ Hz, 2H), 1.66 (d, $J=12.0$ Hz, 2H), 1.41~1.32 (m, 1H), 1.29~1.11 (m, 6H), 1.08~0.96 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 173.5, 77.4, 77.1, 76.8, 72.1, 60.6, 43.1, 38.6, 28.8, 28.2, 26.4, 26.1, 26.0, 14.2.

Ethyl (*E*)-3-hydroxy-5,9-dimethyldeca-4,8-dienoate (**3y**): Yield: 82% (98.4 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 5.21 (d, $J=8.6$ Hz, 1H), 5.13~5.03 (m, 1H), 4.77 (qd, $J=9.0$, 4.1 Hz, 1H), 4.16 (q, $J=7.1$ Hz, 2H), 2.94~2.59 (m, 1H), 2.56~2.41 (m, 2H), 2.09 (dt, $J=13.6$, 6.0 Hz, 3H), 2.02~1.96 (m, 1H), 1.73~1.66 (m, 6H), 1.59 (d, $J=3.5$ Hz, 3H), 1.26 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.5, 172.5, 139.5, 139.3, 132.37, 131.7, 126.5, 125.5, 123.8, 123.8, 65.1, 64.8, 60.7, 60.6, 41.9, 41.8, 39.4, 32.4, 26.4, 26.3, 25.7, 23.3, 17.7, 17.7, 16.6, 14.2.

Ethyl 3-hydroxy-3-phenylbutanoate (**5a**):^[5d] Yield: 72% (74.9 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.48 (dd, $J=5.2$, 3.4 Hz, 2H), 7.35 (dd, $J=10.4$, 4.9 Hz, 2H), 7.28~7.23 (m, 1H), 4.44 (s, 1H), 4.07 (q, $J=7.1$ Hz, 2H), 3.00 (d, $J=15.8$ Hz, 1H), 2.81 (d, $J=15.8$ Hz, 1H), 1.57 (s, 3H), 1.15 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.7, 146.9, 128.2, 126.9, 124.5, 72.8, 60.7, 46.5, 30.7, 14.0.

Ethyl 3-hydroxy-3-(*p*-tolyl)butanoate (**5b**):^[5d] Yield: 66% (73.3 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.36 (d, $J=8.2$ Hz, 2H), 7.17 (d, $J=8.1$ Hz, 2H), 4.38 (s, 1H), 4.09 (qt, $J=4.9$, 2.6 Hz, 2H), 2.99 (d, $J=15.9$ Hz, 1H), 2.80 (d, $J=15.9$ Hz, 1H), 2.35 (s, 3H), 1.55 (s, 3H), 1.17 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.8, 144.0, 136.4, 128.9, 124.4, 72.7, 60.7, 46.5, 30.8, 21.0, 14.0.

Ethyl 3-hydroxy-3-(4-(trifluoromethyl)phenyl)butanoate (**5c**): Yield: 60% (82.8 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.64~7.54 (m, 4H), 4.56 (s, 1H), 4.08 (q, $J=7.1$ Hz, 2H), 2.99 (d, $J=16.1$ Hz, 1H), 2.83 (d, $J=16.1$ Hz, 1H), 1.55 (s, 3H), 1.15 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.5, 151.0, 129.1 (q, $J=32$ Hz), 125.2 (q, $J=41$ Hz), 125.0, 123.2 (q, $J=270$ Hz), 72.65, 60.97, 46.05, 30.54, 13.92.

Ethyl 3-hydroxy-3-(naphthalen-2-yl)butanoate (**5d**):^[5d]

Yield: 62% (80.0 mg), colourless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.98 (d, $J=1.3$ Hz, 1H), 7.86 (dt, $J=5.7$, 2.6 Hz, 3H), 7.59 (dd, $J=8.6$, 1.9 Hz, 1H), 7.53~7.46 (m, 2H), 4.61 (s, 1H), 4.07 (qd, $J=7.1$, 3.5 Hz, 2H), 3.13 (d, $J=15.9$ Hz, 1H), 2.91 (d, $J=15.9$ Hz, 1H), 1.66 (s, 3H), 1.13 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.8, 144.3, 133.2, 132.4, 128.2, 128.0, 127.5, 126.1, 125.8, 123.19, 123.1, 73.0, 60.8, 46.3, 30.7, 14.0.

Supporting Information ^1H NMR and ^{13}C NMR spectra of compounds **3a**~**3y** and **5a**~**5d**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Paterson, I.; Chen, D. Y.-K.; Coster, M. J.; Aceña, J. L.; Bach, nJ.; Gibson, K. R. L. E.; Oballa, R. M.; Trieselmann, T.; Wallace, D. J.; Hodgson, A. P.; Norcross, R. D. *Angew. Chem., Int. Ed.* **2001**, *40*, 4055.
(b) Calter, M. A.; W. Liao, J. *Am. Chem. Soc.* **2002**, *124*, 13127.
(c) Schetter, B.; R. Mahrwald, *Angew. Chem., Int. Ed.* **2006**, *45*, 7506.
(d) List, B.; Lerner, R. A.; Barbas, C. F. *J. Am. Chem. Soc.* **2000**, *122*, 2395.
(e) Cheng-Sánchez, I.; Carrillo, P.; Sánchez-Ruiz, A.; Martínez-Poveda, B.; A. Quesada, R.; M. Medina, A.; J. López-Romero, M.; Sarabia, F. *J. Org. Chem.* **2018**, *83*, 5365.
(f) Kang, T.; Han, D. Jo, S. *J. Org. Chem.* **2017**, *82*, 9335.
(g) Gomez-Palomino, A.; Romea, P.; Urpi, F. *Synthesis* **2018**, *50*, 2721.
- [2] (a) Mandavid, H.; Rodrigues, A. M. S.; Espindola, L. S.; Eparvier, V.; Stien, D. J. *Nat. Prod.* **2015**, *78*, 1735.
(b) Nakamukai, S.; Takada, K.; Furihata, K.; Ise, Y.; Okada, S.; Morii, Y.; Yamawaki, N.; Takatani, T.; Arakawa, O.; Gustafson, K. R.; Stellatolide H Matsunaga, S. *Tetrahedron Lett.* **2018**, *59*, 2532.
(c) Shin, H. J.; Rashid, M. A.; Cartner, L. K.; Bokesch, H. R.; Wilson, J. A.; McMahon, J. B.; Gustafson, K. R. *Tetrahedron Lett.* **2015**, *56*, 4215.
(d) Bae, M.; Moon, K.; Kim, J.; Park, H. J.; Lee, S. K.; Shin, J.; Oh, D.-C. *J. Nat. Prod.* **2016**, *79*, 332.
- [3] Reformatsky, S. *Ber. Dtsch. Chem. Ges.* **1887**, *20*, 1210.
- [4] (a) Choppin, S.; Ferreiro-Medeiros, L.; Barbarottoa, M.; Colobert, F. *Chem. Soc. Rev.* **2013**, *42*, 937.
(b) Kim, J. H.; Ko, Y. O.; Bouffard, J.; Lee, S. *Chem. Soc. Rev.* **2015**, *44*, 2489.
(c) Shen, Z.-L.; Wang, S.-Y.; Chok, Y.-K.; Xu, Y.-H.; Loh, T.-P. *Chem. Rev.* **2013**, *113*, 271.
(d) Cozzi, P. G. *Angew. Chem. Int. Ed.* **2007**, *46*, 2568.
- [5] (a) Yamakoshi, S.; Kawanishi, E. *Tetrahedron Lett.* **2014**, *55*, 1175.
(b) Wu, P.; Cai, W.; Chen, Q.-Y.; Xu, S.; Yin, R.; Li, Y.; Zhang, W.; Luesch, H. *Org. Lett.* **2016**, *18*, 5400.
(c) Yun, J.-J.; Zhi, M.-L.; Shi, W.-X.; Chu, X.-Q.; Shen, Z.-L.; Loh, T.-P. *Adv. Synth. Catal.* **2018**, *360*, 2632.
(d) Liu, X.-Y.; Li, X.-R.; Zhang, C.; Chu, X.-Q.; Rao, W.; Loh, T.-P.; Shen, Z.-L. *Org. Lett.* **2019**, *21*, 5873.
(e) Sinast, M.; Zuccolo, M.; Wischnat, J.; Sube, T.; Hasnik, F.; Baro, A.; Dallavalle, S.; Laschat, S. *J. Org. Chem.* **2019**, *84*, 10050.
(f) Cao, Q.; Stark, R. T.; Fallis, I. A.; Browne, D. L. *ChemSusChem* **2019**, *12*, 2554.
- [6] (a) Frstner, A. *Synthesis* **1989**, 571.
(b) Marshall, J. A. *Chemtracts* **2000**, *13*, 705.
(c) Podlech, J.; Maier, T. C. *Synthesis* **2003**, 633.
(d) Orsini, F.; Sello, G. *Curr. Org. Synth.* **2004**, *1*, 111.
(e) Nakamura, E. In *Organometallic in Synthesis*, A Manual, Ed.: Schlosser, M., Wiley, New York, **2002**, p. 579.
(f) Ocampo, R.; Dolbier Jr, W. R. *Tetrahedron* **2004**, *60*, 9325.
- [7] (a) Peng, Y.-Y.; Liu, P.; Liu, Z.-J.; Liu, J.-T.; Mao, H.-F.; Yao, Y.-L. *Tetrahedron* **2018**, *74*, 3074.
(b) Cao, C.-R.; Jiang, M.; Liu, J.-T. *Eur. J. Org. Chem.* **2015**, 1144.
(c) Huck, L.; Berton, M.; de la Hoz, A.; Díaz-Ortizb, A.; Alcázar, J. *Green Chem.* **2017**, *19*, 1420.
(d) Fernández-Sánchez, L.; Fernández-Salas, J. A.; Maestro, M. C.; García Ruano, J. L. *J. Org. Chem.* **2018**, *83*, 12903.
(e) De Munck, L.; Sukowski, V.; Vila, C.; Muñozb, M. C.; Pedro, J. R. *Org. Chem. Front.* **2017**, *4*, 1624.
- [8] (a) Fernandez-Ibanez, M. A.; Macia, B.; Minnaard, A. J.; Feringa, B. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 1317.
(b) Fornalczyk, M.; Singh, K.; Stuart, A. M. *Chem. Commun.* **2012**, *48*, 3500.
(c) Maestro, A.; de Marigorta, E. M.; Palacios, F.; Vicario, J. *Org. Lett.* **2019**, *21*, 9473.
(d) De Munck, L.; Vila, C.; Muñoz, M. C.; Pedro, J. R. *Chem. Eur. J.* **2016**, *22*, 17590.
- [9] (a) Chattopadhyay, A.; Kr Dubey, A. *J. Org. Chem.* **2007**, *72*, 9357.
(b) Moriwake, T. *J. Org. Chem.* **1966**, *31*, 983.
- [10] (a) Orsini, F.; Lucci, E. M. *Tetrahedron Lett.* **2005**, *46*, 1909.
(b) Segade, Y.; Montaos, M. A.; Rodríguez, J.; Jimenez, C. *Org. Lett.* **2014**, *16*, 5820.
(c) Nelson, C. G.; Jr. Burke, T. R. *J. Org. Chem.* **2012**, *77*, 733.
(d) Sinast, M.; Zuccolo, M.; Wischnat, J.; Sube, T.; Hasnik, F.; Baro, A.; Dallavalle, S.; Laschat, S. *J. Org. Chem.* **2019**, *84*, 10050.
- [11] (a) Hu, Y.; Zhou, B.; Wang, C. *Acc. Chem. Res.* **2018**, *51*, 816.
(b) Hammarback, L. A.; Clark, I. P.; Sazanovich, I. V.; Towrie, M.; Robinson, A.; Clarke, F.; Meyer, S.; Fairlamb, I. J. S.; Lynam, J. M. *Nat. Catal.* **2018**, *1*, 830.
(c) Zhou, B.; Hu, Y.; Liu, T.; Wang, C. *Nat. Commun.* **2017**, *8*, 1169.
(d) Yadav, V.; Landge, V. G.; Subaramanian, M.; Balaraman, E. *ACS Catal.* **2020**, *10*, 947.
(e) Wang, Y.; Zhu, L.; Shao, Z.; Li, G.; Lan, Y.; Liu, Q. *J. Am. Chem. Soc.* **2019**, *141*, 17337.
- [12] (a) Cozzi, P. G.; Rivalta, E. *Angew. Chem. Int. Ed.* **2005**, *44*, 3600.
(b) Adrian, J. C.; Snapper, J. M. L. *J. Org. Chem.* **2003**, *68*, 2143.
(c) Dondoni, A.; Massi, A.; Sabbatini, S. *Chem.-Eur. J.* **2005**, *11*, 7110.
(d) Qin, X.; Wen, M.; Lee, Y.; Zhou, J. S. *Org. Lett.* **2019**, *21*, 5990.
- [13] (a) Cozzi, P. G. *Angew. Chem. Int. Ed.* **2007**, *46*, 2568.
(b) Li, R.; Xu, H.; Zhao, N.; Jin, X.; Dang, Y. *J. Org. Chem.* **2020**, *85*, 833.
- [14] (a) Börjesson, M.; Moragas, T.; Gallego, D.; Martin, R. *ACS Catal.* **2016**, *6*, 6739.
(b) Börjesson, M.; Moragas, T.; Gallego, D.; Martin, R. *J. Am. Chem. Soc.* **2016**, *138*, 7504.
- [15] Zoua, X.-L.; Dua, G.-F.; Sun, W.-F.; He, L.; Ma, X.-W.; Gu, C.-Z.; Dai, B. *Tetrahedron* **2013**, *69*, 607.
- [16] Baudoux, J.; Lefebvre, P.; Legay, R.; Lasne, M.-C.; Rouden, J. *Green Chem.* **2010**, *12*, 252.
- [17] Wadhwa, K.; Verkade, J. G. *J. Org. Chem.* **2009**, *74*, 4368.

(Fan, Y.)