对甲苯亚磺酸钠/KI 介导端炔的需氧氧化碘代反应合成 1-碘代炔和 1,3-二炔

周 鹏* 冯尚伟 邱会华 张建涛*

(广东石油化工学院化学学院 广东茂名 525000)

摘要 基于对甲苯亚磺酸钠/KI 介导端炔的需氧氧化碘代反应, 发展了一种高效合成 1-碘代炔的绿色环保方法. 该方法以最绿色环保的空气作为氧化剂, 无需其他有毒的氧化剂, 具有效率高、反应操作简单、底物适用性广、试剂绿色环保和反应条件温和的特点. 此外, 基于对甲苯亚磺酸钠/KI 介导端炔的需氧氧化碘代反应, 发展了一种一锅法无金属催化端炔合成对称 1,3-二炔的方法.

关键词 端炔; 1-碘代炔; 空气; 亚磺酸钠; 1,3-二炔

Sodium *p*-Toluenesulfinate/KI-Mediated Aerobic Oxidative Iodination of Terminal Alkynes for Synthesis of 1-Iodoalkynes and 1,3-Diynes

Zhou, Peng* Feng, Shangwei Qiu, Huihua Zhang, Jiantao*

(College of Chemistry, Guangdong University of Petrochemical Technology, Maoming, Guangdong 525000)

Abstract A practical and environmentally friendly protocol for the synthesis of 1-iodoalkynes was developed via sodium sulfinate-KI mediated aerobic oxidative iodination of terminal alkynes. This transformation features simple operation, high efficiency, broad functional-group tolerance and mild reaction conditions. Especially, dioxygen was used in these reactions which avoided the addition of hazardous or toxic oxidants. Moreover, for the purpose of increasing the synthetic utility of this method, an efficient transition-metal-free synthetic approach to symmetrical 1,3-diynes was developed via the iodination/ homocoupling of terminal alkynes in a one-pot manner.

Keywords terminal alkyne; 1-iodoalkyne; air; sodium sulfinate; 1,3-diyne

1 Introduction

Due to the synthetic adaptability, 1-iodoalkynes have attracted considerable attention as key precursors and building blocks in the field of organic chemistry and material chemistry.^[1] What's more, some 1-iodoalkynes have potential applications in the preparation of radio-iodinated probes for imaging applications.^[2] In view of their broad significance, many studies have been directed toward the development of efficient methods for their preparation.^[3] Among the known synthetic methods, the direct oxidative iodination of terminal alkynes, as a most straightforward approach, has been extensively explored. As classical iodinating reagents, molecular iodine can react with terminal alkynes with the assistance of Ag(I) salts,^[4a] Cu(I) salts^[4b] and bases^[4c-4f] to access the corresponding 1-iodoalkynes

(Scheme 1, path a). Indeed, taking the advantages of high reactivity, mild conditions and excellent selectivity, hypervalent iodonium salts such as PhI(OAc)₂,^[5] *N*-iodosuccinimide,^[6] bis(sym-collidine)iodine(I) hexafluorophosphate,^[7] HMBMIBDCI (an ionic liquid iodinating reagent)^[8] and Me₃SI(OAc)₂,^[9] as iodinating reagents and oxidants, can also be used for the direct oxidative iodination of terminal alkynes (path b). To develop economic and environmentally benign oxidative reactions, the use of iodide salts like KI, NaI, ZnI₂ and (*n*-Bu)₄NI as iodo-sources is highly attractive, owing to the low cost and low toxicity of these compounds. In this regard, several procedures have been documented for the similar transformation with iodide salts as the iodo-source by anodic oxidation^[10] or in the presence of oxidants, such as bis(trimethylsilyl)

* Corresponding author. E-mail: zzd83@aliyun.com; 1351248781@qq.com

Received July 9, 2020; revised September 11, 2020; published online October 10, 2020.

Project supported by the National Natural Science Foundation of China (No. 21602035), the Natural Science Foundation of Guangdong Province (No. 2016A030307030) and the Program for Innovative Research Team of Guangdong University of Petrochemical Technology (No. 519124).

国家自然科学基金(No. 21602035)、广东省自然科学基金(No. 2016A030307030)和广东石油化工学院创新研究团队(No. 519124)资助项目.

peroxide (BTMSPO),^[11a] *t*-butyl hydroperoxide (TBHP),^[11b] PhI(OAc)₂,^[11c] oxone,^[11d] chloramine salt,^[11e] *tert*-butyl nitrite^[11f] and copper(II) sulfate^[2b] (path c). Though significant progresses have been made, some crucial issues have remained fairly unsolved, including the harsh reaction conditions and the use of large amounts of hazardous oxidants and iodo-sources which makes them environmentally unfavourable. Thus, any practical and environmentally friendly access to provide 1-iodoalkynes with good functional group compatibility in high yields would be extraordinarily valuable. Recently, some iodine or iodide salt/green oxidant systems were proved to be effective in the construction of C—X (X=O, N, S, *etc.*) bonds.^[12] Inspired by these works, we sought to develop a more efficient and straightforward route to 1-iodoalkynes with low cost and low toxicity iodo-sources and oxidants. There is no doubt that dioxygen is the cheapest and most environmentally friendly oxidant. In line with our interests in alkynes,^[13] we herein disclose a sodium sulfinate/ KI-mediated oxidative iodination of terminal alkynes to 1-iodoalkynes with dioxygen.

Scheme 1 Direct oxidative iodination of terminal alkynes to 1-iodoalkynes

2 Results and discussion

At the outset of our studies, phenylacetylene (**1a**, 0.5 mmol) was employed as a model substrate to react with KI (1.2 equiv.) to identify the optimal reaction conditions, and the results were summarized in Table 1. Initially, the reaction was explored by examining the iodination of **1a** with KI in methanol under air condition at room temperature. Not surprisingly, no desired product (**2a**) was obtained under such conditions (Table 1, Entry 1). It was delighted to find that 67% yield of **2a** could be afforded when the reaction was carried out with sodium *p*-toluenesulfinate (TolSO₂Na, 1 equiv.) as an additive (Entry 2). Encouraged by the promising result, the amount of TolSO₂Na was subsequently examined, and the results showed that 1.5 equiv. of TolSO₂Na gave the best results (Entries 3–5). Then, various iodide salts such as NaI, ZnI₂ and NH₄I were screened, but none of them could give comparable results as the run with KI (Entry 4 vs Entries

6–8). As the next optimization step, the solvent effect was examined by screening a variety of reaction media (Entries 9–15). It could be found that the solvent played a significant role in the process of this reaction. Among all the tested solvents [C₂H₅OH, CH₃CN, CH₂Cl₂, *p*-xylene, *N,N'*-dimethylformamide (DMF), dimethyl sulfoxide (DMSO) and H₂O], C₂H₅OH showed the highest efficiency for the present transformation (Entry 9). In addition, a control experiment confirmed that the target product was not observed when the reaction was carried out under N₂ atmosphere (Entry 16), suggesting that O₂ is essential for the reaction. Thus, the optimized reaction conditions were affirmed as follows: 1.2 equiv. of KI as iodo-source, 1.5 equiv. of TolSO₂Na as the additive in C₂H₅OH at room temperature under air atmosphere.

Table 1 Screening of reaction conditions^a

$\text{Ph}-\text{C}\equiv\text{C}-\text{H} \xrightarrow[\text{Solvent, air, r.t., 5 h}]{\text{iodide salt, TolSO}_2\text{Na}} \text{Ph}-\text{C}\equiv\text{C}-\text{I}$ <div style="display: flex; justify-content: space-around; width: 100%;"> 1a 2a </div>				
Entry	Iodide salt	<i>n</i> (TolSO ₂ Na)/mmol	Solvent	Yield ^b /%
1 ^c	KI	—	CH ₃ OH	NR
2	KI	0.5	CH ₃ OH	67
3	KI	0.6	CH ₃ OH	80
4	KI	0.75	CH ₃ OH	89
5	KI	0.85	CH ₃ OH	88
6	NaI	0.75	CH ₃ OH	85
7	ZnI ₂	0.75	CH ₃ OH	68
8	NH ₄ I	0.75	CH ₃ OH	64
9	KI	0.75	C₂H₅OH	93
10	KI	0.75	CH ₃ CN	87
11	KI	0.75	CH ₂ Cl ₂	84
12	KI	0.75	<i>p</i> -Xylene	83
13	KI	0.75	DMF	Trace
14	KI	0.75	DMSO	Trace
15	KI	0.75	H ₂ O	Trace
16 ^d	KI	0.75	C ₂ H ₅ OH	Trace

^a Reaction conditions: **1a** (0.5 mmol), iodide salt (0.6 mmol), TolSO₂Na in solvent (4 mL) at room temperature under air condition. ^b Isolated yield. ^c NR is no reaction. ^d Under N₂ (with a balloon, 101 kPa).

Under the optimized reaction conditions, a range of terminal alkynes **1** were applied in the transformation to establish the scope and generality of this protocol (Table 2). It was noticed that various aryl- and alkyl-substituted terminal alkynes were found to be suitable substrates for the TolSO₂Na-mediated aerobic oxidative iodination. A series of substituted phenylacetylenes bearing electron-rich groups (Me, Et, *n*-Pr, *i*-Pr, OMe, *t*-Bu, OCF₃ and N(CH₃)₂) and electron-poor groups (F, Cl, Br, CN and CF₃) were found to be suitable substrates for this reaction, and the corresponding 1-iodoalkyne products **2a** ~ **2q** were achieved in good yields (80%–92%). Notably, halo substituents (F, Cl and Br), trifluoromethoxy, methoxy and cyano groups were also compatible with this transformation which might offer the opportunity for further syn-

thetic modification. It was noteworthy that the reaction efficiency wasn't significantly affected by the steric effect. Sterically hindered substituted phenylacetylenes (2-F, 2-Cl and 2-Br) proceeded smoothly to furnish the desired products in good yields (**2o**~**2q**, 87%~89%). Furthermore, heteroarylethynes like 3-ethynylpyridine (**1r**) and 3-ethynylthiophene (**1s**) were also employed as the substrates and the corresponding 1-iodoalkyne products (**2r** and **2s**) were produced with no detriment in yields (90% and 87%, respectively). We turned our attention to alkyl-substituted terminal acetylenes, pleasingly, alkyl-substituted terminal acetylenes could also be transformed into the desired 1-iodoalkyne products **2t**~**2v** successfully in good yields (81%~88%). When ethynyltriisopropylsilane (**1w**), an important reagent for the construction of carbon-carbon triple bonds, was applied to this reaction, the expected product (**2w**) was formed in excellent yield (90%), which indicated good availability of this protocol.

Table 2 Synthesis of 1-iodoalkynes^{a,b}

$\text{R}-\text{C}\equiv\text{C}-\text{H} \xrightarrow[\text{C}_2\text{H}_5\text{OH (4 mL), air, r.t., 5~12 h}]{\text{KI (0.6 mmol), TolSO}_2\text{Na (0.75 mmol)}} \text{R}-\text{C}\equiv\text{C}-\text{I}$	
1	2
2a , R = H, 93% 2b , R = 4-CH ₃ , 86% 2c , R = 4-C ₂ H ₅ , 82% 2d , R = 4- <i>n</i> -Pr, 83% 2e , R = 4- <i>t</i> -Bu, 80% 2f , R = 4- <i>i</i> -Pr, 80% 2g , R = 4-OCH ₃ , 81% 2h , R = 3-CH ₃ , 86% 2i , R = 4-OCF ₃ , 88% 2j , R = 4-N(CH ₃) ₂ , 89% ^c 2k , R = 4-Cl, 92% 2l , R = 4-CF ₃ , 89% 2m , R = 4-CN, 85% 2n , R = 3-F, 90% 2o , R = 2-F, 89% 2p , R = 2-Cl, 89% 2q , R = 2-Br, 87%	2r , 90% 2s , 87% 2t , 87% 2u , 81% 2v , 88% 2w , 90%

^a Reaction conditions: alkyne (0.5 mmol), KI (1.2 equiv.), TolSO₂Na (1.5 equiv.), in C₂H₅OH (4 mL) at room temperature under air condition. ^b Isolated yield. ^c At room temperature for 24 h.

As an effective and mild method, the scale-up of the sodium sulfinate-mediated aerobic oxidative iodination might have great practical significance for the large-scale preparation of 1-iodoalkynes in the laboratory, and even in the industry. As demonstrated in Scheme 2, gram-scale experiments were successfully conducted by employing **1a** and **1w** as the substrates, and the corresponding 1-iodoalkynes **2a** and **2w** was obtained in 86% and 84% yields, respectively, demonstrating its great potential for large-scale synthesis in the industry.

Scheme 2 Large-scale experiment

In 2010, Jiang's group^[14] developed an efficient synthetic approach to symmetrical 1,3-diynes from 1-haloalkynes under transition-metal-free conditions in DMF. Based on Jiang's work, we examined the *in situ* transition-metal-free coupling of 1-iodoalkynes to symmetrical 1,3-diynes in one-pot for the purpose of increasing the synthetic utility of our methods. To our delight, It turned out that treatment of crude 1-iodoalkynes in DMF at 120 °C delivered corresponding 1,3-diynes (Scheme 3, **3a**, **3e**, **3n** and **3u**) in moderate yields (53%~79%) for the one-pot process.

Scheme 3 Transition-metal-free approach for symmetrical 1,3-diynes from terminal alkynes in one pot

To gain insight into the transformation, some control experiments were carried out (Scheme 4). The radical trapping experiments were conducted by adding 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) into the model reaction under the standard conditions. Obviously, the formation of product **2a** was completely suppressed, indicating that a free radical pathway might be involved in this process (Scheme 4, a). Gratifyingly, the reaction between **1a** and *p*-toluenesulfonyl iodide (Tsl) was examined instead

Scheme 4 Mechanism experiments

of TolSO₂Na and KI, yielding the corresponding product **2a** in moderate yield (43%) (Scheme 4, b). And the result indicated that sulfonyl iodide was likely to be a key intermediate to react with **1a** to lead to the iodoalkyne **2a**. Interestingly, trace TsI (**4**) could be detected under standard reaction conditions by thin layer chromatography (TLC) analysis, but disappeared after being stirred overnight which was in accordance with the literature (Scheme 4, c).^[15]

On the basis of the results above and previous reports,^[16-17] a plausible mechanism for this aerobic oxidative iodination was proposed as outlined in Scheme 5. Initially, TolSO₂Na was converted into sulfonyl iodide (**4**) by a combination of air/KI. Subsequently, sulfur-centered sulfonyl radical **A**, oxygen-centered sulfonyl radical **B**, together with iodine radical were generated via the homolytic cleavage of sulfonyl iodide (**4**). The iodine radical underwent direct addition to the alkyne **1** to generate the vinyl radical intermediate **C**. Finally, assisted by dioxygen and **B**, intermediate **C** underwent elimination of hydrogen to provide the iodoalkyne **2** with the release of TolSO₃H.

Scheme 5 Possible reaction mechanism

3 Conclusions

In conclusion, a practical and environmentally friendly protocol for the synthesis of 1-iodoalkynes via sodium sulfonate mediated aerobic oxidative iodination of terminal alkynes without metal catalysts was developed. This strategy was not only featured with simple operation, remarkable efficiency, but also was highlighted with broad functional-group tolerance. Especially, air, the cheapest and most environmentally benign oxidant, was used in our reactions which avoided the addition of hazardous or toxic oxidants. Besides, an efficient and transition-metal-free synthetic approach to symmetrical 1,3-diynes was developed via the iodination/homocoupling of terminal alkynes in a one-pot manner.

4 Experimental section

4.1 General information.

¹H NMR and ¹³C NMR spectra were recorded on a 400 MHz (100 MHz for ¹³C NMR) spectrometer taking CDCl₃ as solvent. And the chemical shifts were referenced to tet-

ramethylsilane (TMS). Melting points were determined on a micromelting point apparatus and uncorrected. MS was obtained using electron ionization (EI). All products were isolated by thin-layer chromatography (TLC) prepared with 600 mesh silica gel plates (GF254). All known products and their spectra were in agreement with authentic samples. All chemicals were purchased from commercial suppliers and were used without further purification.

4.2 General Procedure for the Synthesis of 1-iodoalkynes

A 25-mL round-bottom flask with a magnetic stir bar was charged with KI (0.6 mmol), TolSO₂Na (0.75 mmol) and C₂H₅OH (4 mL). After stirring of the mixture for 5 min, the terminal alkyne was added and the resulting mixture was stirred at room temperature for 5~12 h. After the reaction was complete as monitored by TLC, the solution was diluted with ethyl acetate (20 mL), washed with brine (10 mL × 3), dried over MgSO₄, filtered, and evaporated under vacuum. The crude product was purified by TLC on silica gel (taking ethyl acetate and petroleum ether as eluent) to afford the products **2**.

1-Iodo-phenylacetylene (**2a**).^[6g-6h] Pale yellow oil, 93% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.42~7.44 (m, 2H), 7.35~7.28 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 132.34, 128.79, 128.24, 123.39, 94.14, 6.04; MS (ESI) *m/z*: 229 [M+H⁺].

1-(Iodoethynyl)-4-methylbenzene (**2b**).^[6g-6h] Pale yellow oil, 86% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.32 (d, *J*=8.1 Hz, 2H), 7.11 (d, *J*=7.9 Hz, 2H), 2.35 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 139.02, 132.20, 128.98, 120.37, 94.25, 21.52, 4.84; MS (ESI) *m/z*: 243 [M+H⁺].

1-Ethyl-4-(iodoethynyl)benzene (**2c**).^[5a] Pale yellow oil, 82% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.34 (d, *J*=8.1 Hz, 2H), 7.13 (d, *J*=8.0 Hz, 2H), 2.64 (q, *J*=7.6 Hz, 2H), 1.21 (t, *J*=7.6 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 145.29, 132.31, 127.80, 120.61, 94.30, 28.85, 15.29, 4.90; MS(ESI) *m/z*: 257 [M+H⁺].

1-(Iodoethynyl)-4-propylbenzene (**2d**).^[5a] Pale yellow oil, 83% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.34 (d, *J*=8.0 Hz, 2H), 7.11 (d, *J*=8.4 Hz, 2H), 2.57 (t, *J*=7.2 Hz, 2H), 1.57~1.67 (m, 2H), 0.92 (t, *J*=7.3 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 143.78, 132.20, 128.39, 120.60, 94.30, 37.96, 24.29, 13.75, 4.86; MS (ESI) *m/z*: 271 [M+H⁺].

1-(*tert*-Butyl)-4-(iodoethynyl)benzene (**2e**).^[11e] Pale yellow solid, 80% yield. m.p. 89~91 °C (lit.^[9] 90~92 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.35 (d, *J*=8.5 Hz, 2H), 7.31 (d, *J*=8.5 Hz, 2H), 1.29 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 152.14, 131.93, 125.28, 120.45, 94.30, 34.85, 31.20, 5.15; MS (ESI) *m/z*: 285 [M+H⁺].

1-(Iodoethynyl)-4-isopropylbenzene (**2f**): Pale yellow oil, 80% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.33~7.36 (m, 2H), 7.15~7.19 (m, 2H), 2.85~2.92 (m, 1H), 1.23 (d, *J*=6.9 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 149.88, 132.32, 127.42, 126.60, 126.46, 126.38, 120.73, 94.29, 34.09, 23.77, 4.84; HRMS calcd for C₁₁H₁₂I [M+H]⁺

270.9984, found 270.9980.

1-(Iodoethynyl)-4-methoxybenzene (**2g**):^[6g-6h] Pale yellow oil, 80% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.34~7.38 (m, 2H), 6.80~6.83 (m, 2H), 3.79 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 160.13, 133.96, 115.75, 114.04, 94.17, 55.49, 4.09; MS (ESI) *m/z*: 259 [M+H⁺].

1-(Iodoethynyl)-3-methylbenzene (**2h**):^[11e] Pale yellow oil, 86% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.22~7.26 (m, 2H), 7.19 (t, *J*=7.5 Hz, 1H), 7.12~7.14 (m, 1H), 2.31 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 137.95, 132.89, 129.71, 129.38, 128.11, 123.19, 94.32, 21.19, 5.44; MS (ESI) *m/z*: 243 [M+H⁺].

1-(Iodoethynyl)-4-(trifluoromethoxy)benzene (**2i**): Pale yellow semisolid, yield 88%. ¹H NMR (400 MHz, CDCl₃) δ: 7.44~7.47 (m, 2H), 7.15~7.17 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 149.27 (d, *J*=1.9 Hz), 133.88, 122.09, 121.62, 120.70 (d, *J*=1.0 Hz), 119.06, 92.68, 7.73; HRMS calcd for C₉H₅F₃IO [M+H]⁺ 312.9337, found 312.9331.

4-(Iodoethynyl)-*N,N*-dimethylaniline (**2j**):^[4e] Brown solid, 89% yield. m.p. >120 °C (Dec.) [lit.^[4e] 120~125 °C (Dec.)]; ¹H NMR (400 MHz, CDCl₃) δ: 7.29~7.31 (m, 2H), 6.58~6.60 (m, 2H), 2.96 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 150.28, 133.34, 111.47, 110.20, 95.01, 40.11, 1.78; MS (ESI) *m/z*: 272 [M+H⁺].

1-Chloro-4-(iodoethynyl)benzene (**2k**):^[6g-6h] Pale yellow solid, 92% yield. m.p. 84~86 °C (lit.^[4e] 83~84 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.37~7.37 (m, 2H), 7.27~7.30 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 134.94, 133.55, 128.62, 121.87, 93.00, 7.67; MS (ESI) *m/z*: 263 [M+H⁺].

1-(Iodoethynyl)-4-(trifluoromethyl)benzene (**2l**):^[6g-6h] Pale yellow viscous liquid, 89% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.52~7.58 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 132.61, 130.49 (d, *J*=32.8 Hz), 127.05 (d, *J*=1.5 Hz), 125.21 (q, *J*=3.9 Hz), 122.44, 92.82, 10.13; MS (ESI) *m/z*: 297 [M+H⁺].

4-(Iodoethynyl)benzonitrile (**2m**):^[6g-6h] Pale yellow solid, 85% yield. m.p. 170~172 °C (lit.^[4e] 169~170 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.59~7.62 (m, 2H), 7.49~7.52 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 132.89, 131.97, 128.08, 118.26, 112.13, 92.57, 13.21; MS (ESI) *m/z*: 254 [M+H⁺].

1-Fluoro-3-(iodoethynyl)benzene (**2n**):^[6g-6h] Pale yellow oil, 90% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.19~7.30 (m, 2H), 7.12 (ddd, *J*=9.3, 2.4, 1.4 Hz, 1H), 7.03 (tdd, *J*=8.5, 2.6, 1.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.16 (d, *J*=246.9 Hz), 129.82 (d, *J*=8.6 Hz), 128.24 (d, *J*=3.2 Hz), 125.11 (d, *J*=9.4 Hz), 119.16 (d, *J*=22.9 Hz), 116.29 (d, *J*=21.2 Hz), 92.87 (d, *J*=3.4 Hz), 8.07; MS (ESI) *m/z*: 247 [M+H⁺].

1-Fluoro-2-(iodoethynyl)benzene (**2o**):^[6g-6h] Pale yellow oil, 89% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.41~7.45 (m, 1H), 7.27~7.31 (m, 1H), 7.04~7.11 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.74 (d, *J*=252.3 Hz), 134.20 (d, *J*=1.1 Hz), 130.48 (d, *J*=8.0 Hz), 123.85 (d, *J*=3.8 Hz), 115.50 (d, *J*=20.8 Hz), 111.94 (d, *J*=15.7 Hz), 87.36, 11.73 (d, *J*=3.4 Hz); MS (ESI) *m/z*: 247 [M+

H⁺].

1-Chloro-2-(iodoethynyl)benzene (**2p**):^[6g-6h] Pale yellow oil, 89% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.46 (dd, *J*=7.5, 1.9 Hz, 1H), 7.38 (dd, *J*=7.5, 1.9 Hz, 1H), 7.18~7.30 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 136.71, 134.17, 129.76, 129.32, 126.36, 123.24, 90.90, 12.30; MS (ESI) *m/z*: 263 [M+H⁺].

Bromo-2-(iodoethynyl)benzene (**2q**):^[6g-6h] Pale yellow oil, 87% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.56 (dd, *J*=8.0, 1.1 Hz, 1H), 7.45 (dd, *J*=7.7, 1.7 Hz, 1H), 7.24 (dt, *J*=7.6, 1.2 Hz, 1H), 7.16 (dt, *J*=7.8, 1.7 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 134.28, 132.36, 129.93, 126.96, 126.08, 125.46, 92.69, 12.09; MS (ESI) *m/z*: 308 [M+H⁺].

3-(Iodoethynyl)pyridine (**2r**):^[6g-6h] Pale yellow solid, 90% yield. m.p. 107~109 °C (lit.^[9] 104~106 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.68 (s, 1H), 8.54 (d, *J*=3.8 Hz, 1H), 7.71~7.74 (m, 1H), 7.24~7.27 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 152.94, 148.86, 139.31, 122.97, 90.71, 11.39; MS (ESI) *m/z*: 230 [M+H⁺].

3-(Iodoethynyl)thiophene (**2s**):^[6g-6h] Pale yellow oil, 87% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.47 (dd, *J*=3.0, 1.2 Hz, 1H), 7.27~7.24 (m, 1H), 7.11 (dd, *J*=5.0, 1.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 130.30, 130.00, 125.16, 122.52, 89.15, 5.67; MS (ESI) *m/z*: 235 [M+H⁺].

((3-Iodoprop-2-yn-1-yl)oxy)benzene (**2t**):^[4d] Pale yellow oil, 87% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.33~7.27 (m, 2H), 7.02~6.94 (m, 3H), 4.82 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 157.56, 129.54, 121.63, 114.87, 89.22, 57.31, 4.65; MS (ESI) *m/z*: 259 [M+H⁺].

4-Iodo-2-methylbut-3-yn-2-ol (**2u**):^[6b] Clear oil, 81% yield. ¹H NMR (400 MHz, CDCl₃) δ: 1.52 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 99.11, 66.89, 31.34, 29.70, -0.79; MS (ESI) *m/z*: 211 [M+H⁺].

(4-Iodobut-3-yn-1-yl)benzene (**2v**):^[6g-6h] Clear oil, 88% yield. ¹H NMR (400 MHz, CDCl₃) δ: 7.27~7.32 (m, 2H), 7.19~7.23 (m, 3H), 2.84 (t, *J*=7.6 Hz, 2H), 2.64 (t, *J*=7.6 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 140.26, 128.44, 128.40, 126.40, 93.90, 34.92, 23.02, -6.09; MS (ESI) *m/z*: 257 [M+H⁺].

(Iodoethynyl)triisopropylsilane (**2w**):^[18] Clear oil, 90% yield. ¹H NMR (400 MHz, CDCl₃) δ: 1.07 (s, 21H); ¹³C NMR (100 MHz, CDCl₃) δ: 100.84, 18.50, 11.41; MS (ESI) *m/z*: 309 [M+H⁺].

Supporting Information ¹H NMR and ¹³C NMR spectra of compounds **2a**~**2w**, **3a**, **3e**, **3n**, **3u** and **4**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Selected reviews on the application of 1-iodoalkynes: (a) Brand, J. P.; Waser, J. *Chem. Soc. Rev.* **2012**, *41*, 4165.
(b) Wu, W.; Jiang, H. *Acc. Chem. Res.* **2014**, *47*, 2483.
(c) Shi, W.; Lei, A. *Tetrahedron Lett.* **2014**, *55*, 2763.
(d) Danilkina, N. A.; Govdi, A. I.; Balova, I. A. *Synthesis* **2020**, *52*, 1874.

- [2] (a) Kabalka, G. W.; Mereddy, A. R. *J. Labelled Compd. Radiopharm.* **2005**, *48*, 359.
 (b) Ferris, T.; Carroll, L.; Mease, R. C.; Spivey, A. C.; Aboagye, E. O. *Tetrahedron Lett.* **2019**, *60*, 936.
 (c) Dubost, E.; McLernain, H.; Babin, V.; Sutherland, A.; Cailly, T. *J. Org. Chem.* **2020**, *85*, 8300.
- [3] (a) Mandai, T.; Matsumoto, T.; Kawada, M.; Tsuji, J. *J. Org. Chem.* **1992**, *57*, 6090.
 (b) Nishikawa, T.; Shibuya, S.; Hosokawa, S.; Isobe, M. *Synlett* **1994**, 485.
 (c) Naskar, D.; Roy, S. *J. Org. Chem.* **1999**, *64*, 6896.
 (d) Blackmore, I. J.; Boa, A. N.; Murray, E. J.; Dennis, M.; Woodward, S. *Tetrahedron Lett.* **1999**, *40*, 6671.
 (e) Luithle, J. E.; Pietruszka, J. *Eur. J. Org. Chem.* **2000**, 2557.
 (f) Das, J. P.; Roy, S. *J. Org. Chem.* **2002**, *67*, 7861.
 (g) Kabalka, G. W.; Mereddy, A. R. *Tetrahedron* **2004**, *45*, 1417.
 (h) Starkov, P.; Rota, F.; D'Oyley, J. M.; Sheppard, T. D. *Adv. Synth. Catal.* **2012**, *354*, 3217.
 (i) Pelletier, G.; Lie, S.; Mousseau, J. J.; Charette, A. B. *Org. Lett.* **2012**, *14*, 5464.
- [4] (a) Heasley, V. L.; Shellhamer, D. F.; Heasley, L. E.; Yaeger, D. B. *J. Org. Chem.* **1980**, *45*, 4649.
 (b) Chen, S.-N.; Hung, T.-T.; Lin, T.-C.; Tsai, F.-Y. *J. Chin. Chem. Soc.* **2009**, *56*, 1078.
 (c) Rao, M. L. N.; Periasamy, M. *Synth. Commun.* **1995**, *25*, 2295.
 (d) He, G.-W.; Liu, F.-W.; Xu, X.-H. *Chin. J. Org. Chem.* **2007**, *27*, 663 (in Chinese).
 (贺干武, 刘凤伟, 许新华, 有机化学, **2007**, *27*, 663.)
 (e) Meng, L.-G.; Cai, P.-J.; Guo, Q.-X.; Xue, S. *Synth. Commun.* **2008**, *38*, 225.
 (f) Xia, Z.; Corcé, V.; Zhao, F.; Przybylski, C.; Espagne, A.; Jullien, L.; Saux, T. L.; Gimbert, Y.; Dossmann, H.; Mouriès-Mansuy, V.; Ollivier, C.; Louis Fensterbank, L. *Nat. Chem.* **2019**, *11*, 797.
- [5] (a) Yan, J.; Li, J.; Cheng, D. *Synlett* **2007**, *15*, 2442.
 (b) Li, Y.; Huang, D.; Wang, D.; Huang, J.; Maruoka, K. *CN 106831283*, **2020**.
- [6] (a) Naskar, D.; Roy, S. *J. Org. Chem.* **1999**, *64*, 6896.
 (b) Hein, J. E.; Tripp, J. C.; Krasnova, L. B.; Sharpless, B.; Fokin, V. V. *Angew. Chem., Int. Ed.* **2009**, *48*, 8018.
 (c) Nçsel, P.; Lauterbach, T.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. *Chem.-Eur. J.* **2013**, *19*, 8634.
 (d) Wang, B.; Zhang, J.; Wang, X.; Liu, N.; Chen, W.; Hu, Y. *J. Org. Chem.* **2013**, *78*, 10519.
 (e) Li, M.; Li, Y.; Zhao, B.; Liang, F.; Jin, L.-Y. *RSC Adv.* **2014**, *4*, 30046.
 (f) Gómez-Herrera, A.; Nagra, F.; Brill, M.; Nolan, S. P.; Cazin, C. S. J. *ChemCatChem* **2016**, *8*, 3381.
 (g) Shi, W.; Guan, Z.; Cai, P.; Chen, H. *J. Catal.* **2017**, *353*, 199.
 (h) Yao, M.; Zhang, J.; Yang, S.; Xiong, H.; Li, L.; Liu, E.; Shi, H. *RSC Adv.* **2020**, *10*, 3946.
 (i) Yao, M.; Zhang, J.; Yang, S.; Liu, E.; Xiong, H. *Synlett* **2020**, *31*, 1102.
- [7] Brunel Y.; Rousseau, G. *Tetrahedron Lett.* **1995**, *36*, 2619.
- [8] Nouzarian, M.; Hosseinzadeh, R.; Golchoubian, H. *Synth. Commun.* **2013**, *43*, 2913.
- [9] Rao, D. S.; Reddy, T. P.; Kashyap, S. *Org. Biomol. Chem.* **2018**, *16*, 1508.
- [10] Nishiguchi, I.; Kanbe, O.; Itoh, K.; Maekawa, H. *Synlett* **2000**, 89.
- [11] (a) Casarini, A.; Dembech, P.; Reginato, G.; Ricci, A.; Seconi, G. *Tetrahedron Lett.* **1991**, *32*, 2169.
 (b) Reddy, K. R.; Venkateshwar, M.; Maheswari, C. U.; Kumar, P. S. *Tetrahedron Lett.* **2010**, *51*, 2170.
 (c) Liu, Y.; Huang, D.; Huang, J.; Maruoka, K. *J. Org. Chem.* **2017**, *82*, 11865.
 (d) Srujana, K.; Swamy, P.; Naresh, M.; Durgaiiah, C.; Rammurthy, B.; Sai, G. K.; Sony, T.; Narender, N. *ChemistrySelect* **2017**, *2*, 748.
 (e) Liu, X.; Chen, G.; Li, C.; Liu, P. *Synlett* **2018**, *29*, 2051.
 (f) Chen, S.; Zhang, X.; Zhao, H.; Guo, X.; Hu, X. *Chin. J. Org. Chem.* **2018**, *38*, 1172 (in Chinese).
 (陈锁, 章晓炜, 赵辉, 郭晓红, 胡祥国, 有机化学, **2018**, *38*, 1172.)
- [12] Selected examples: (a) Lv, Y.; Li, Y.; Xiong, T.; Lu, Y.; Liu, Q.; Zhang, Q. *Chem. Commun.* **2014**, *50*, 2367.
 (b) Tan, C.; Liu, Y.; Liu, X.; Jia, H.; Xu, K.; Huang, S.; Wang, J.; Tan, J. *Org. Chem. Front.* **2020**, *7*, 780.
 (c) Quan, J.; He, X.; Yan, X.; Li, X.; Xu, X. *Chin. J. Org. Chem.* **2020**, *40*, 1033 (in Chinese).
 (全积宁, 何小雪, 严新焕, 李小青, 许响生, 有机化学, **2020**, *40*, 1033.)
 (d) Lv, Y.; Sun, K.; Wang, T.; Li, G.; Pu, W.; Chai, N.; Shen, H.; Wu, Y. *RSC Adv.* **2015**, *5*, 72142.
 (e) Huang, L.; Yin, W.; Wang, J.; Gan, C.; Huang, Y.; Huang, C.; He, Y. *RSC Adv.* **2019**, *9*, 2381.
 (f) Yuan, J.; Ma, X.; Yi, H.; Liu, C.; Lei, A. *Chem. Commun.* **2014**, *50*, 14386.
 (g) Wu, X.-F.; Gong, J.-L.; Qi, X. *Org. Biomol. Chem.* **2014**, *12*, 5807.
- [13] (a) Zhou, P.; Qiu, H.; Shi, J. *Chin. J. Org. Chem.* **2016**, *36*, 425 (in Chinese).
 (周鹏, 邱会华, 施继成, 有机化学, **2016**, *36*, 425.)
 (b) Qiu, H.; Cheng, B.; Huang, Y.; Chen, C.; Zhou, P. *Chin. J. Org. Chem.* **2018**, *38*, 1817 (in Chinese).
 (邱会华, 成博睿, 黄颖思, 陈翠, 周鹏, 有机化学, **2018**, *38*, 1817.)
 (c) Zhou, P.; Pan, Y.; Tan, H.; Liu, W. *J. Org. Chem.* **2019**, *84*, 15662.
 (d) Qiu, H.; Zhou, P.; Liu, W.; Zhang, J.; Chen, B. *ChemistrySelect* **2020**, *5*, 2935.
- [14] Chen, Z.; Jiang, H.; Wang, A.; Yang, S. *J. Org. Chem.* **2010**, *75*, 6700.
- [15] Meesin, J.; Katrun, P.; Pareseecharoen, C.; Pohmakotr, M.; Reutrakul, V.; Soorukram, D.; Kuhakarn, C. *J. Org. Chem.* **2016**, *81*, 2744.
- [16] Recent reviews on radical transformations of alkynes: (a) Xu, J.; Song, Q. *Chin. J. Org. Chem.* **2016**, *36*, 1151 (in Chinese).
 (许健, 宋秋玲, 有机化学, **2016**, *36*, 1151.)
 (b) Huang, M.-H.; Hao, W.-J.; Li, G.; Tu, S.-J.; Jiang, B. *Chem. Commun.* **2018**, *54*, 10791.
 (c) Guerrero-Robles, M. A.; Vilchis-Reyes, M. A.; Ramos-Rivera, E. M.; Alvarado, C. *ChemistrySelect* **2019**, *4*, 13698.
- [17] Selected examples on the transformation via TolSI: (a) Katrun, P.; Chiampanichayakul, S.; Korworapan, K.; Pohmakotr, M.; Reutrakul, V.; Jaipetch, T.; Kuhakarn, C. *Eur. J. Org. Chem.* **2010**, 5633.
 (b) Buathongjan, C.; Beukeaw, D.; Yotphan, S. *Eur. J. Org. Chem.* **2015**, 1575.
 (c) Wei, W.; Liu, C.; Yang, D.; Wen, J.; You, J.; Wang, H. *Adv. Synth. Catal.* **2015**, *357*, 987.
 (d) Jiang, Y.-Y.; Wang, Q.-Q.; Liang, S.; Hu, L.-M.; Little, R. D.; Zeng, C.-C. *J. Org. Chem.* **2016**, *81*, 4713.
 (e) Lai, J.; Chang, L.; Yuan, G. *Org. Lett.* **2016**, *18*, 3194.
 (f) Kumar, R.; Dwivedi, V.; Reddy, M. S. *Adv. Synth. Catal.* **2017**, *359*, 2847.
 (g) Guo, Y.-J.; Lu, S.; Tian, L.-L.; Huang, E.-L.; Hao, X.-Q.; Zhu, X.; Shao, T.; Song, M.-P. *J. Org. Chem.* **2018**, *83*, 338.
 (h) Zhang, J.; Liang, Z.; Wang, J.; Guo, Z.; Liu, C.; Xie, M. *ACS Omega* **2018**, *3*, 18002.
 (i) Xie, F.; Lu, G.-P.; Xie, R.; Chen, Q.-H.; Jiang, H.-F.; Zhang, M. *ACS Catal.* **2019**, *9*, 2718.
- [18] Li, Y.; Brand, J. P.; Waser, J. *Angew. Chem., Int. Ed.* **2013**, *52*, 6743.

(Zhao, C.)