无过渡金属催化条件下合成 3,3'-二吲哚甲烷衍生物的最新进展

张振国 刘笑笑 宗鑫龙 苑亚林 刘双磊

张 婷 吴子尚 杨静莹* 贾振华*,^a

(南京工业大学化学与分子工程学院 先进化学制造研究院 南京 211816)

摘要 3,3'-二吲哚甲烷类化合物是一类重要的吲哚生物碱,其结构单元广泛存在于天然产物、功能性材料以及合成药物分子中.因其具有多样的生物活性和功能,如抗氧化、抗炎症、抗血管生成、抗菌以及抗癌活性等,构筑 3,3'-二吲哚甲烷类杂环化合物备受关注.传统的合成方法,尤其是对称结构的 3,3'-二吲哚甲烷类化合物的合成主要在化学量的 Brønsted 酸或 Lewis 酸存在下,吲哚衍生物与羰基化合物经傅-克反应缩合得到.而过渡金属的使用可引起化合物中金属残留以及环境污染.总结和探讨了从 2010 年至今 3,3'-二吲哚甲烷类化合物的合成方法,尤其是对无过渡金属参与条件下,对称结构的 3,3'-二吲哚甲烷类化合物以及非对称结构 3,3'-二吲哚甲烷类化合物制备的最新进展以及相应的反应机理,旨在为该类药物生物活性测试提供重要的理论依据和技术支持.

关键词 无过渡金属; 3,3'-二吲哚甲烷; 生物活性; 多样性

Recent Advance on the Synthesis of 3,3'-Bisindolylmethane Derivatives under Transition-Metal-Free Catalytic Conditions

Zhang, Zhenguo Liu, Xiaoxiao Zong, Xinlong Yuan, Yalin Liu, Shuanglei

Zhang, Ting Wu, Zishang Yang, Jingying* Jia, Zhenhua*

(Institute of Advanced Synthesis, School of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816)

Abstract 3,3'-Bisindolylmethanes (3,3'-BIMs) compounds are important indole alkyloads and their units are widely found in various natural products, functional materials and synthetic pharmaceutical compounds. Due to diverse biological activities and functionalities, for instance, antioxidant, anti-inflammatory, antiangiogenic, anti-bacterial and anti-cancer *etc.*, the construction of 3,3'-BIMs is raised considerable concerns. Conventional methods especially focused on symmetrical 3,3'-BIMs were the condensation of indoles with carbonyl compounds via Friedel-Crafts pathway in the presence of Brønsted acids or Lewis acids. However, the utilization of transition metals led the residue into the products and environmental contamination. The recent advance on the synthesis of 3,3'-BIMs since 2010, mainly concerned on the approaches and corresponding mechanism to prepare symmetrical and unsymmetrical 3,3'-BIMs under transition-metal-free conditions is summarized and discussed, aiming to provide important theoretical evidence and technical support for further biological evaluations on desired compounds.

Keywords transition-metal-free; 3,3'-bisindolylmethane; biological activity; diversity

1 Introduction

Indole scaffolds are important structural constitutions of numerous natural products, pharmaceutical drugs, fine chemicals and agrochemicals.^[1] Among skeletons of indole alkaloids, 3,3'-bisindolylmethanes (BIMs) units are ubiquitous, which are widely found in various bioactive natural

products, synthetic compounds, pharmaceuticals and functional materials (Scheme 1). From a variety of Terrestrial and Marine natural resources, 3,3'-BIMs are separated and identified the common structure consist of two indole rings at 3- and 3'-positions connected with a single carbon atom. These compounds exhibit a series of important biological

* Corresponding authors. E-mail: iasjyyang@njtech.edu.cn; iaszhjia@njtech.edu.cn

Received August 5, 2020; revised October 13, 2020; published online October 21, 2020.

Dedicated to Professor Henry N. C. Wong on the occasion of his 70th birthday.

Project supported by the Start-up Grant of Nanjing Tech University (No. 38037037), the Synergetic Innovation Center for Advanced Materials Fellowship and Postgraduate Research & Practice Innovation Program of Jiangsu Province (No. KYCX20_1031).

南京工业大学启动基金(No. 38037037)、江苏省协同联合创新中心和江苏省研究生科研与实践创新计划(No. KYCX20_1031)资助项目.

activities,^[2-12] such as antioxidant,^[13-14] anti-inflammatory,^[15-17] antiangiogenic,^[18] anti-leishmanial activity,^[19] anti-staphylococcus activity,^[20] anti-MRSA (Methicillin-resistant staphylococcus aureus), anti-cancer activity^[21] and antifungal activity.^[22]

In 1886, 3,3'-BIMs was synthesized by Fischer *et al.* for the first time.^[23-24] The general method for the synthesis of 3,3'-BIMs was the reaction between indoles and carbonyl compounds via Friedel-Crafts pathway in the presence of stoichiometric acids or bases. Accompanying with the exploring the potential applications of BIMs in the field of pharmaceutical industry, diverse strategies were developed for the construction of 3,3'-BIMs. In particular, the catalysis involving transition metals promoted the efficiencies and decreased the price of the preparation of 3,3'-BIMs. However, the utilization of transition metals could result in the contamination of target products, which were essentially considered during the further evaluation of their bio-activities. In addition, the generation of the stoichiometric chemical and metallic wastes could cause serious environmental issues, which was not suitable with the development of modern sustainable chemistry.^[25-33] Bisindolylmethanes (BIMs) and trisindolylmethanes (TIMs) were well reviewed by Shiri *et al.*^[34] in 2010, in which they summarized the various approaches to BIMs and TIMs,

including 3,3'-BIMs, 2,2'-BIMs, 2,3'-BIMs and TIMs.

Recently, Xu,^[35] Xie,^[36] He,^[37] Yin,^[38] Yang,^[39] Zhong,^[40] Mei,^[41] He,^[42] Xia,^[43] Shi^[44] and others have developed various methods for preparing BIMs. This mini-review will focus on the recent advance on the synthesis of 3,3'-bisindolylmethanes (BIMs), especially, it will summarize the synthetic protocols of the symmetric and asymmetric 3,3'-BIMs under transition-metal-free conditions since 2010. The transformations catalyzed by transition metals for the synthesis of BIMs will be excluded in this review.

2 Synthetic approaches to symmetrical 3,3'-BIMs

The general mechanism for the formation of 3,3'-BIMs in the presence of acids is depicted in Scheme 2. During this Friedel-Crafts pathway, the key intermediate is enamine (**III**), tautomerized with carbon cationic intermediate (**IV**) that undergo further addition with the other indole molecular to achieve symmetrical 3,3'-BIMs (Scheme 2).

In 2013, Huo *et al.*^[45] developed a sustainable strategy to prepare symmetrical 3,3'-BIMs with a recyclable system. They utilized the complex of sodium triphenylphosphine-sulfonate (TPPMS) with CBr₄ as a highly efficient cata-

(A) Representative nature products containing 3,3'-bisindolylmethanes (BIMs) units

(B) Synthetic (BIMs) compounds and materials with diverse biological activities and functional applications

Scheme 1 Representative nature products and synthetic compounds containing 3,3'-BIMs units

Scheme 2 General Friedel-Crafts pathway to achieve symmetrical 3,3'-BIMs

lyst for the synthesis of 3,3'-BIMs from a range of carbonyl compounds, such as aldehydes, ketones and acetals, in excellent yields. The process underwent smoothly at room temperature without considering the effects of atmosphere and water. In addition, the catalyst was stable and easy-stored in air. Notably, the complex could be separated

from the reaction mixture and easily recovered to use for 10 times without obvious loss of the catalytic activities (Scheme 3).

In 2014, Taha *et al.*^[46] identified bisindolylmethane-hydrazone hybrids as novel inhibitors of β -glucuronidase. To obtain various target products to evaluate their corresponding inhibitory potential, a sequential route was designed. Bisindolylmethane ester (**23**) was prepared up to 93% yield in refluxing conditions, nevertheless other conditions gave **23** in low yields, for instance, replacing acetic acid by $\text{NaBrO}_3/\text{NaHSO}_3$, ethylammonium nitrate or $\text{BF}_3 \cdot \text{Et}_2\text{O}$. The ester was transformed into hydrazone intermediate **24** followed by condensation with diverse aldehydes to generate the desired products (Scheme 4).

Incorporation of a trifluoromethyl group (CF_3) in pharmaceutical molecules affected their bioactivities and physico-chemical properties, compared to parent molecules. Anbarsasan *et al.*^[47] developed an efficient and general one-pot trifluoromethylative functionalization of amides for the synthesis of various trifluoromethylated bis(indolyl)-arylmethanes (**28**). They successfully applied this method to synthesize trifluoromethylated analog of hypolipidemic and anti-obesity agent (**29**). During the trifluoromethylative process, the essential trifluoromethylated iminium ion was generated *in-situ* from trifluoromethyltrimethylsilane and trapped by substituted indoles for achieving the final products (Scheme 5).

The C—C triple bond of the alkynes could be activated

Scheme 3 Huo's synthesis of symmetrical 3,3'-BIMs by recyclable process

Scheme 4 Taha's synthesis sequential route of bisindolylmethane-hydrazone hybrids

Selected examples

Scheme 5 Anbarsasan's synthesis of trifluoromethylated bis(indolyl)arylmethanes

via π -coordination to the typical π -Lewis acid, $B(C_6F_5)_3$. In 2018, Zhong *et al.*^[48] reported $B(C_6F_5)_3$ -catalyzed dihydroindolization of aryl alkynes by double Markovnikov addition to indoles under neat conditions, providing a viable access to versatile symmetrical 3,3'-BIMs. The mechanism was also described in Scheme 6. Firstly, coordination of $B(C_6F_5)_3$ with aryl alkyne **30** afforded the intermediate **I**. Subsequently, nucleophilic attack of **26** on the internal carbon of the C—C triple bond occurred to generate zwitterionic boron species **II**. The first proton transfer and aromatization was followed to give the intermediate **III** and release $B(C_6F_5)_3$ at the same time. $B(C_6F_5)_3$ coordinated with intermediate **III** again to form the intermediate **IV**, which was attacked by another **26** to involve the second proton transfer and aromatization, producing the double

Markovnikov addition products **31** and regeneration of the borane catalyst.

In the same year, He *et al.*^[49] described similar metal-free protocol of $B(C_6F_5)_3$ -catalyzed synthesis of the symmetrical 3,3'-BIMs. Instead of aldehydes, more readily available, abundant, stable and easy-to-handle carboxylic acids were treated as alkylating reagents. Mechanistic studies disclosed that the carboxylic acids **33** initially reacted with silane to obtain the silyl ester **I** with release of hydrogen gas. Meanwhile, the Si—H bond of silane was coordinated to $B(C_6F_5)_3$ reversibly, generating the adduct *in-situ*, which donated the silica cation to the oxygen of carboxylic acids to form ion pair **II**. Subsequently, the hydride transfer occurred to generate disilyl acetals or aldehydes under acidic conditions. Double nucleophilic attacks of the complex of

Scheme 6 Zhong's $B(C_6F_5)_3$ -catalyzed synthesis of the symmetrical 3,3'-BIMs

$B(C_6F_5)_3$ with indole derivatives towards disilyl acetals or aldehydes proceeded to achieve the symmetrical 3,3'-BIMs and regenerate $B(C_6F_5)_3$ (Scheme 7).

Huo *et al.*^[50] discovered that the cheap, commercially available and stable radical cation triarylaminium salt could be applied as an initiator for Friedel-Crafts reaction of indoles with enamides at low catalyst loading (2 mol%) to construct the complex of 3,3'-BIMs derivatives. They proposed that the pathway of this protocol would involve the process of single electron transfer (SET) and radical intermediates from vinyl ethers, which were essential to furnish the alkylation process efficiently (Scheme 8).

Toy *et al.*^[51] synthesized a bidentate halogen bond donor catalyst for the synthesis of symmetrical 3,3'-BIMs. Compared with mono-dentate analogue and thiourea organocatalyst, the bidentate catalyst was proved to be more effective in the halogen bond catalyzed Friedel-Crafts reactions of diverse aldehydes and ketones with indoles. The natural products, such as arundine, trisindoline and vibrindole A, were successfully accomplished in one-pot from good to excellent yields (Scheme 9).

Halimehjani *et al.*^[52] synthesized novel tetra-cationic acidic organic salts (TCAOS) containing both sulfonic acid and hydrogen sulfate groups, based on DABCO (1,4-diazabicyclo[2.2.2]octane). Subsequently, TCAOS

was applied to synthesize the symmetrical 3,3'-BIMs in water with excellent turnover frequency. This process was efficient, environment friendly and the catalyst was recycled and reused in the reaction more than five times without significant loss of its catalytic ability (Scheme 10).

Recently, photo-catalysis was introduced for organic transformations, due to its sustainability and efficient capture and utilization of solar energy. Itoh *et al.*^[53] developed a one-pot synthesis of the symmetrical 3,3'-BIMs from benzyl alcohols as alkylation reagents, using catalytic iodine in oxygen atmosphere under visible light irradiation. Under optimized conditions, indole and its derivatives were coupled with various benzyl alcohols, but pyrrole was alkylated in only 18% yield. Mechanistic studies disclosed that benzyl alcohols were transferred to corresponding aldehydes irradiated by visible light in the presence of catalytic amount of modular iodine (Scheme 11).

Singh *et al.*^[54] reported a visible light mediated strategy to construct bisindolyl acetate derivatives with Eosin Y as the photo-sensitizer. Activating C—F and C—Br bonds of ethyl bromofluoroacetate (**44**) generated corresponding radicals, which were subsequently trapped by indoles to form the target products. The postulated mechanism was provided based on several control experiments, but the

Scheme 7 He's $B(C_6F_5)_3$ -catalyzed reductive synthesis of the symmetrical 3,3'-BIMs

Scheme 8 Huo's Friedel-Crafts alkylation initiated by triarylammonium salt

Selected examples:

Scheme 9 Toy's synthesis of the symmetrical 3,3'-BIMs by halogen bond catalyzed strategy

Scheme 10 Halimehjeni's tetra-cationic acidic organic salts (TCAOS)

Scheme 11 Itoh's photo-induced I_2 -catalyzed synthesis of symmetrical 3,3'-BIMs in O_2 atmosphere

pathway was still not clear (Scheme 12).

In 2007, Xia *et al.*^[55] developed a photo-redox protocol of 1,3-dipolar cycloaddition of nitrones with alkenes to

obtain isoxazolidine derivatives. When indole was subjected to the reaction conditions, the symmetrical 3,3'-BIMs were achieved in moderate to good yield. To the respect of mechanism, the catalyst was excited by visible light, which accepted one electron from indole to generate the radical cation intermediate **I**. Afterwards, this radical cation was trapped by nitron to form the intermediate **III**. After elimination of phenyl-hydroxylamine radical, cationic intermediate **IV** was formed, which could react with another indole molecular indole. Deprotonation of intermediate **IV** by phenyl-hydroxylamine radical led to the target product generated (Scheme 13).

In 2020, Qiu *et al.*^[56] reported a concise CF_3SO_2Na -induced synthesis of symmetrical 3,3'-BIMs under UV irradiation and oxygen atmosphere. Cyclic or acyclic ketones and various aldehydes were coupled with indoles at molar ratio of 1 : 2. Especially, selected target products showed anticancer activities in human bladder cancer cell lines EJ and T24. From the insight of mechanism, the key radical species was formed *in-situ* from CF_3SO_2Na under irradiation of UV light (Scheme 14).

In 2014, Ning *et al.*^[57] reported an unexpected Br_2 -

Scheme 12 Singh's organo-photo-catalyst mediated synthesis of symmetrical 3,3'-BIMs

Scheme 13 Xia's photo-redox synthesis of symmetrical 3,3'-BIMs

Selected examples:

Scheme 14 Qiu's $\text{CF}_3\text{SO}_2\text{Na}$ -induced synthesis of symmetrical 3,3'-BIMs

catalyzed synthesis of 3,3'-BIMs from indoles and carbonyl compounds with low catalyst loading (down to 2 mol%), high efficiency and broad substrate scope, and the reaction could be conducted in either organic solvents or water. This robust catalytic activity of bromine was treated as an underrated Lewis acid, which might inspire more novel Br_2 -catalyzed organic transformations and provide the potential for more focus on the possible function of bromine

(Scheme 15).

Zhang *et al.*^[58] described a facile synthesis of 3,3'-BIMs using fiber-supported ionic liquids (ILs) in water at room temperature. During the process of Friedel-Crafts alkylation, the relevant characterizations of ILs were investigated to confirm that ILs were immobilized on the surface layer of polyacrylonitrile fibers (PANF) and could be recycled with excellent stability. The mechanism was proposed as

Scheme 15 Ning's Br_2 -catalyzed synthesis of symmetrical 3,3'-BIMs

the simple “release and catch” catalytic process in detail. The mild reaction conditions and the simple procedure combined with the easy recycle of the catalyst and solvent proved that this method was environmentally friendly and economical for the synthesis of bioactive BIMs. As far as reported, this is the first example of utilizing the inexpensive and commercially available PANF as the support for novel ILs (Scheme 16).

Scheme 16 Zhang's fiber-supported ILs-catalyzed synthesis of symmetrical 3,3'-BIMs

In 2018, Arai *et al.*^[59] developed the protocol of 2-iodoimidazolium salt-catalyzed Friedel-Crafts reaction to synthesize bis(indolyl)methane alkaloids using indoles and aldehydes. Under mild conditions, various 3,3'-BIMs derivatives were obtained in good to high yields (Scheme 17).

Yamajala *et al.*^[60] reported that thiamine hydrochloride (VB_1) was an efficient and reusable catalyst to synthesize 3,3'-BIMs under water, solvent-free and grinding conditions with low catalyst loadings (down to 1 mol%). The desired products were isolated without column chromatography. Green matrix calculations showed high atom-economy

Scheme 17 Arai's 2-iodoimidazolium salt-catalyzed synthesis of symmetrical 3,3'-BIMs

and small E-factor. The gram-scale reaction was proven to have potential industrial application. Additionally, the catalyst could also be utilized effectively to prepare a varieties of alkaloids derived BIMs and biscoumarins (Scheme 18).

Scheme 18 Yamajala's VB_1 -catalyzed synthesis of symmetrical 3,3'-BIMs

More recently, Wang *et al.*^[61] described a simple and transition-metal-free method for the synthesis of 3,3'-BIMs via a new graphene oxide-induced alkylation strategy. 3,3'-BIMs derivatives were generated under mild reaction conditions, with a broad indole scope and excellent regioselectivities, in moderate to excellent yields. The dichotomous catalytic behavior of graphene oxide was efficient to promote the process of C—C bond oxidative coupling and C—O bond cleavage (Scheme 19).

3 Synthetic approaches to unsymmetrical 3,3'-BIMs

The retro-synthesis analysis disclosed that two different indolyl substrates were coupled to construct unsymmetrical 3,3'-BIMs. Conventional strategy was utilizing a leaving group pre-installed on one indole substrate. When the leaving groups were removed under various conditions, the corresponding cationic intermediates were generated *in-situ* trapped by another different indole substrate to obtain unsymmetrical 3,3'-BIMs via the subsequent process of deprotonation (Scheme 20).

Selected examples

Scheme 19 Wang's graphene oxide-catalyzed synthesis of symmetrical 3,3'-BIMs **Scheme 20** General strategy for the synthesis of unsymmetrical 3,3'-BIMs

In 2012, Sigman *et al.*^[62] discovered that *p*-TsOH catalyzed hydroarylation of vinyl indoles (**56**) to form the putative cationic intermediate with any additional stabilization, which was trapped by indole derivatives to form the diverse unsymmetrical 3,3'-BIMs in good to excellent yields. Further evaluation of the biological activities of several prepared compounds, especially product **57** exhibited promising cytotoxic activity against and selectivity for MCF-7 breast cancer cell line (Scheme 21).

Sasaki *et al.*^[63] developed a trifluoroacetic acid catalyzed method to introduce trifluoromethyl group into 3,3'-BIMs. They screened reaction conditions to evaluate the impact of different Brønsted acids in diverse solvents. Moreover, the selective synthesis of unsymmetrical 3,3'-BIMs mainly depended on the stability of the α -carbocation (Scheme 22).

In 2016, Xiao *et al.*^[64] described a green and efficient process, in which indolyl alcohols in water reacted with diverse nucleophiles, including indole derivatives, to obtain unsymmetrical 3,3'-BIMs in the absence of any catalyst. The mechanism was proposed that water might be treated as

Selected examples:

Scheme 21 Sigman's *p*-TsOH catalyzed synthesis of unsymmetrical 3,3'-BIMs

Selected examples

Scheme 22 Sasaki's synthesis of trifluoromethyl-functionalized unsymmetrical 3,3'-BIMs

a weak acid, target products were obtained in good yields when the reaction temperature was elevated to higher. The

putative mechanism of this dehydration process was $\text{S}_{\text{N}}1$ -type pathway catalyzed by water (Scheme 23).

Scheme 23 Xiao's synthesis of unsymmetrical 3,3'-BIMs in water

Müller *et al.*^[65] described another efficient approach to unsymmetrical 3,3'-BIMs in water without any catalyst and additives involved. (3-Indolylmethyl)trimethylammonium iodides were utilized as the starting materials to couple with a broad range of substituted indole derivatives. Interestingly, 7-azaindole led to 3,3'-BIMs, while 4- and 5-azaindoles reacted at the N(1)-nitrogen atom to afford novel aza-BIMs derivatives (Scheme 24).

Selected examples:

Scheme 24 Müller's synthesis of unsymmetrical 3,3'-BIMs in water

4 Synthetic approaches to asymmetrical 3,3'-BIMs

Only a few examples focused on the enantioselective synthesis of 3,3'-bisindolylmethane derivatives. In 2010, You *et al.*^[66] pioneered the synthesis of asymmetrical

3,3'-bisindolylmethanes. Their method, using chiral phosphoric acid derived phosphoric acid as the chiral catalyst, mediated asymmetric Friedel-Crafts alkylation of 1-methyl indole with 3-indolylmethanamines, to generate optically pure 3,3'-bisindolylmethanes derivatives up to 80% yield and 68% *ee* (Scheme 25).

Scheme 25 You's the first enantioselective synthesis of chiral 3,3'-BIMs

In 2014, Zhang *et al.*^[67] achieved enantioselective synthesis of unsymmetrical 3,3'-BIMs and pyrrolylsubstituted triarylmethanes in good to excellent yields with *ee* up to 95%, using a novel imidodiphosphoric acid catalyst derived from (*R*)-BINOL framework. The process of Friedel-Crafts alkylation exhibited low catalyst loading, broad functional group tolerance and provided potentially practical methods for the synthesis of enantioselective bioactive 3,3'-BIMs (Scheme 26).

Kim *et al.*^[68] provided a similar route for the synthesis of bisindolylmethane sulfamate derivatives in good yields and moderate to good enantioselectivities (up to 89% yield, 94 : 6 *er*) with similar chiral phosphoric acid as the catalyst. This protocol was well-tolerated to various 3-indolylsulfamates and indoles. In addition, the biological activities of these products were evaluated and confirmed to prevent peripheral nerve degeneration efficiently (Scheme 27).

Selected examples:

Scheme 26 Zhang's enantioselective synthesis of chiral 3,3'-BIMs

5 Conclusions and perspectives

Although the chemical synthetic principles of 3,3'-bis-indolylmethanes (3,3'-BIMs) have been described more than a century ago, the emerging of catalysis in the absence of transition metals was blooming as increasing concerns on environmental pollution. In recent years, 3,3'-BIMs have demonstrated various biological activities and functionalities, therefore, approaches under transition-metal-free conditions to 3,3'-BIMs were explored to meet the further biological evaluations. This review summarized useful and practical methods for the synthesis of different types of 3,3'-BIMs reported in last decade, and illustrated the versatilities of indole derivatives coupled with other partners bearing diverse functional groups in reasonable yields.

Considering the sustainability of chemical synthesis in the field of pharmaceutical chemistry, we envision remarkable concerns on the development of novel protocols to-

Selected examples

Scheme 27 Kim's chiral phosphoric acid catalytic synthesis of chiral 3,3'-BIMs sulfamate derivatives

wards the practical synthesis of 3,3'-BIMs. On account of increasing requirements of screening biological activities, the diversities of 3,3'-BIMs are expected, especially stereoselective synthesis of 3,3'-BIMs, in future.

References

- [1] Katritzky, A. R.; Ramsden, C. A.; Scriven, E. F. V.; Taylor, R. J. K. *Comprehensive Heterocyclic Chemistry III*, Elsevier, Oxford, **2008**.
- [2] Irie, T.; Kubushira, K.; Suzuki, K.; Tsukazaki, K.; Umezawa, K.; Nozawa, S. *Anticancer Res.* **1999**, *31*, 3061.
- [3] Hong, C.; Firestone, G. L.; Bjeldanes, L. F. *Biochem. Pharmacol.* **2002**, *63*, 1085.
- [4] Porszasz, J.; Gibiszer-Porszasz, K.; Foldeak, S.; Matkovics, B. *Experientia* **1965**, *21*, 93.
- [5] (a) Kuethe, J. T. *Chimia* **2006**, *60*, 543.
(b) Foldeak, S.; Czombas, J.; Matkovics, B. *Acta Phys. Chem.* **1965**, *11*, 115.
- [6] Bifulco, G.; Bruno, I.; Minale, L.; Riccio, R.; Calignano, A.; Debitus, C. *J. Nat. Prod.* **1994**, *57*, 1294.
- [7] Bifulco, G.; Bruno, I.; Riccio, R.; Lavayre, J.; Bourdy, G. *J. Nat. Prod.* **1995**, *58*, 1254.
- [8] Parmeggiani, F.; Rué Casamajo, A.; Walton, C. J. W.; Galman, J. L.; Turner, N. J.; Chica, R. A. *ACS Catal.* **2019**, *9*, 3482.
- [9] Wan, Y. C.; Li, Y. H.; Yan, C. X.; Yan, M.; Tang, Z. L. *Eur. J. Med. Chem.* **2019**, *183*, 111691.
- [10] Bandari, C.; Scull, E. M.; Bavineni, T.; Nimmo, S. L.; Gardner, E.

- D.; Bensen, R. C.; Singh, S. *MedChemComm* **2019**, *10*, 1465.
- [11] Wang, S.; Fang, K.; Dong, G.; Chen, S.; Liu, N.; Miao, Z.; Yao, J.; Li, J.; Zhang, W.; Sheng, C. *J. Med. Chem.* **2015**, *58*, 6678.
- [12] Zhang, M.-Z.; Chen, Q.; Yang, G.-F. *Eur. J. Med. Chem.* **2015**, *89*, 421.
- [13] Tran, P. H.; Nguyen, X.-T. T.; Chau, D.-K. N. *Asian J. Org. Chem.* **2018**, *7*, 232.
- [14] Benabadi, S. H.; Wen, R.; Zheng, J.; Dong, X.; Yuan, S. *Acta Pharmacol. Sin.* **2004**, *25*, 666.
- [15] Bharate, S. B.; Bharate, J. B.; Khan, S. I.; Tekwani, B. L.; Khan, I. A.; Vishwakarma, R. A. *Eur. J. Med. Chem.* **2013**, *63*, 435.
- [16] Zhou, J. M.; Yuan, K. Y.; Lin, W. Z.; Hu, X. C.; Jin, Q. Q.; Niu, C. G. *J. Shanghai JiaoTong Univ. (Med. Sci.)* **2018**, *38*, 138.
- [17] Yang, W. J.; Jiao, R.; Liu, Y.; Sun, X. D.; Sang, M. *Herb. Med.* **2019**, *38*, 1146.
- [18] Dong, Y. X.; Lushnikova, T.; Golla, R. M.; Wang, X. F.; Wang, G. S. *Bioorg. Med. Chem.* **2017**, *25*, 864.
- [19] Cho, H. J.; Seon, M. R.; Lee, Y. M.; Kim, J.; Kim, J. K.; Kim, S. G.; Park, J. H. *J. Nutr.* **2008**, *138*, 17.
- [20] Kunimasa, K.; Kobayashi, T.; Kaji, K.; Ohta, T. *J. Nutr.* **2010**, *140*, 1.
- [21] Xue, L.; Firestone, G. L.; Bjeldanes, L. F. *Oncogene* **2005**, *24*, 2343.
- [22] Wang, X. L.; Liu, F.; Wang, Q. Z.; Xu, S.; Luo, J. Y. *J. For. Eng.* **2017**, *2*, 0.
- [23] Fischer, H. E. *Chem. Ber.* **1886**, *19*, 2988.
- [24] Fischer, H. E. *Justus Liebigs Ann. Chem.* **1887**, *242*, 372.
- [25] Yadav, J. S.; Reddy, B. V. S.; Padmavani, B.; Gupta, M. K. *Tetrahedron Lett.* **2004**, *45*, 7577.
- [26] Ferrer, C.; Amijs, C. H. M.; Echavarren, A. M. *Chem.-Eur. J.* **2007**, *13*, 1358.
- [27] Gao, R.; Yi, C. S. *J. Org. Chem.* **2010**, *75*, 3144.
- [28] Tsuchimoto, T.; Kanbara, M. *Org. Lett.* **2011**, *13*, 912.
- [29] Xia, D.; Wang, Y.; Du, Z. T.; Zheng, Q. Y.; Wang, C. Y. *Org. Lett.* **2012**, *14*, 588.
- [30] Vicenzi, D.; Sgarbossa, P.; Biffis, A.; Tubaro, C.; Basato, M.; Bogi-
alli, S.; Pastore, P.; Venzo, A. *Organometallics* **2013**, *32*, 7153.
- [31] Chen, S. P.; Li, Y. X.; Ni, P. H.; Huang, H. W.; Deng, G. J. *Org. Lett.* **2016**, *18*, 5384.
- [32] Cooper, L.; Alonso, J. M.; Eagling, L.; Newson, H.; Herath, S.; Cox, B.; Muoz, M. P. *Chem.-Eur. J.* **2018**, *24*, 6105.
- [33] Abe, T.; Nakamura, S.; Yanada, R.; Choshi, T.; Hibino, S.; Ishikura, M. *Org. Lett.* **2013**, *15*, 3622.
- [34] Shiri, M.; Zolfigol, M. A.; Kruger, H. G.; Tanbakouchian, Z. *Chem. Rev.* **2010**, *110*, 2250.
- [35] Yi, W. G.; Jia, Z. Y.; Li, N. B.; Qiu, R. H.; Chen, J. Y.; Xu, X. H. *Chin. J. Org. Chem.* **2012**, *32*, 2390 (in Chinese).
(易卫国, 贾振永, 李宁波, 邱仁华, 陈锦杨, 许新华, 有机化学, **2012**, *32*, 2390.)
- [36] Gong, H.-W.; Xie, Z.-F. *Chin. J. Org. Chem.* **2012**, *32*, 1195 (in Chinese).
(宫海伟, 解正峰, 有机化学, **2012**, *32*, 1195.)
- [37] He, L.; Wang, X. B.; Du, G. F.; Dai, B.; Jian, T. Y. *Chin. J. Org. Chem.* **2013**, *33*, 988 (in Chinese).
(何林, 王湘波, 杜广芬, 代斌, 简腾跃, 有机化学, **2013**, *33*, 988.)
- [38] Li, L. L.; Ban, D. M.; Fu, H.; Gong, W.; Chen, Z.; Yin, X. G. *Chin. J. Synth. Chem.* **2018**, *26*, 757.
- [39] Yang, Y. S.; Cao, B. X.; Li, S. B.; Zhang, Y. P. *J. Lanzhou Univ. Technol.* **2018**, *44*, 72.
- [40] Zhang, L. Y.; Wu, B. Q.; Chen, Z. T.; Hu, J. J.; Zeng, X. F.; Zhong, G. F. *Chin. J. Org. Chem.* **2018**, *38*, 2028 (in Chinese).
(章吕烨, 吴彬强, 陈张涛, 胡锦锦, 曾晓飞, 钟国富, 有机化学, **2018**, *38*, 2028.)
- [41] Wu, P.; Wu, J. L.; Wang, J. Y.; Mei, G. J. *Chin. J. Org. Chem.* **2018**, *38*, 1251 (in Chinese).
(伍平, 吴迦勒, 王静怡, 梅光建, 有机化学, **2018**, *38*, 1251.)
- [42] Liu, T. W.; Zhang, T. W.; Zhang, S. T.; He, J. H.; Zhang, Y. T. *Chem. J. Chin. Univ.* **2019**, *40*, 719 (in Chinese).
(刘天伟, 张苏韬, 何江华, 张越涛, 高等学校化学学报, **2019**, *40*, 719.)
- [43] Zhang, R. Z.; Wang, G. D.; Li, H. S.; Duan, G. Y.; Wang, K.; Xia, C. C. *Chin. J. Org. Chem.* **2019**, *39*, 1429 (in Chinese).
(张瑞泽, 王国栋, 李洪爽, 段桂运, 王凯, 夏成才, 有机化学, **2019**, *39*, 1429.)
- [44] Mao, Y. J.; Lu, Y. N.; Li, T. Z.; Wu, Q.; Tan, W.; Shi, F. *Chin. J. Org. Chem.* **2020**, *40*, 3895 (in Chinese).
(毛雨佳, 陆一楠, 李天真, 吴琼, 谭伟, 石枫, 有机化学, **2020**, *40*, 3895.)
- [45] Huo, C. D.; Sun, C. G.; Wang, C.; Jia, X. D.; Chang, W. J. *ACS Sustainable Chem. Eng.* **2013**, *1*, 549.
- [46] Taha, M.; Ismail, N. H.; Imran, S.; Anouar, E. H.; Ali, M.; Jamil, W.; Uddin, N.; Kashif, S. M. *RSC Adv.* **2016**, *6*, 3276.
- [47] Vinay, K. P.; Pazhamalai, A. J. *Org. Chem.* **2017**, *82*, 12328.
- [48] Ling, F.; Xiao, L.; Zhong, W. H. *Org. Biomol. Chem.* **2018**, *16*, 9274.
- [49] Qiao, C.; Liu, X. F.; He, L. N. *Chem.-Asian J.* **2018**, *13*, 2664.
- [50] Huo, C. D.; Kang, L. S.; Xu, X. L.; Jia, X. D.; Wang, X. C.; Xie, H. S.; Yuan, Y. *Tetrahedron Lett.* **2014**, *55*, 954.
- [51] Liu, X. L.; Ma, S.; Toy, P. H. *Org. Lett.* **2019**, *21*, 9212.
- [52] Halimehjani, A. Z.; Barati, V. *ChemistrySelect* **2018**, *3*, 3024.
- [53] Nobuta, T.; Fujiya, A.; Tada, N.; Miura, T.; Itoh, A. *Synlett* **2012**, *23*, 2975.
- [54] Jadhav, S. D.; Bakshi, D.; Singh, A. J. *Org. Chem.* **2015**, *80*, 10187.
- [55] Zheng, L. W.; Gao, F.; Yang, C.; Gao, G. L.; Zhao, Y. T.; Gao, Y.; Xia, W. J. *Org. Lett.* **2017**, *19*, 5086.
- [56] Yang, T. B.; Lu, H. A.; Shu, Y. X.; Ou, Y. F.; Hong, L.; Au, C. T.; Qiu, R. H. *Org. Lett.* **2020**, *22*, 827.
- [57] Liang, D. Q.; Huang, W. Z.; Yuan, L.; Ma, Y. H.; Ma, J. M.; Ning, D. M. *Catal. Commun.* **2014**, *55*, 11.
- [58] Shi, X. L.; Lin, H. K.; Li, P. Y.; Zhang, W. Q. *ChemCatChem* **2014**, *6*, 2947.
- [59] Kuwano, S.; Suzuki, T.; Arai, T. *Heterocycles* **2018**, *97*, 163.
- [60] Mathavan, S.; Kannan, K.; Yamajala, B. R. D. *Org. Biomol. Chem.* **2019**, *17*, 9620.
- [61] Peng, X. J.; Zen, Y.; Liu, Q.; Liu, L. X.; Wang, H. S. *Org. Chem. Front.* **2019**, *6*, 3615.
- [62] Pathak, T. P.; Osiak, J. G.; Vaden, R. M.; Welm, B. E.; Sigman, M. S. *Tetrahedron* **2012**, *26*, 5203.
- [63] Sasaki, S.; Ikekame, Y.; Tanayama, M.; Yamauchi, T.; Higashiyama, K. *Synlett* **2012**, *23*, 2699.
- [64] Xiao, J.; Wen, H.; Wang, L.; Xu, L. B.; Hao, Z. H.; Shao, C. L.; Wang, C. Y. *Green Chem.* **2016**, *18*, 1032.
- [65] Pillaiyar, T.; Gorska, E.; Schnakenburg, G.; Müller, C. E. *J. Org. Chem.* **2018**, *83*, 9902.
- [66] Sun, F. L.; Zheng, X. J.; Gu, Q.; He, Q. L.; You, S. L. *Eur. J. Org. Chem.* **2010**, *47*.
- [67] Zhuo, M. H.; Jiang, Y. J.; Fan, Y. S.; Gao, Y.; Liu, S.; Zhang, S. Q. *Org. Lett.* **2014**, *16*, 1096.
- [68] Kim, Y.; Lee, J.; Jung, J. Y.; Kim, S. G. *Tetrahedron Lett.* **2019**, *60*, 1625.

(Lu, Y.)