以氰甲基亚磷酸酯为含磷试剂的铜催化磷酰化 异喹啉酮类化合物的高效合成

赵苏艳^{a,b} 官雪芹^c 甘子玉^c 颜秋莉^b
刘学良^{*,d} 杨道山^{*,b,c}

(^a 吉林警察学院实验技术中心 长春 130117)

(^b 青岛科技大学化学与分子工程学院 山东青岛 266042)

(^c 曲阜师范大学化学与化工学院 山东曲阜 273165)

(^d 新乡医学院基础医学院 河南新乡 453003)

摘要 报道了一种铜催化“一锅法”高效合成磷酰化异喹啉酮类化合物的新方法。该反应是第一例基于铜催化 Ullmann 偶联策略以氰甲基亚磷酸酯为含磷试剂制备含磷取代氮杂环化合物的新方法。

关键词 铜催化; 异喹啉酮; 串联反应; 有机磷化合物

Efficient Copper-Catalyzed Domino Synthesis of Phosphonated Isoquinolin-1(2*H*)-ones Using Cyanomethylphosphonates as Building Blocks

Zhao, Suyan^{a,b} Gong, Xueqin^c Gan, Ziyu^c Yan, Qiuli^b
Liu, Xueliang^{*,d} Yang, Daoshan^{*,b,c}

(^a Experimental Technology Center, Jilin Police College, Changchun 130117)

(^b College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, Qingdao, Shandong 266042)

(^c School of Chemistry and Chemical Engineering, Qufu Normal University, Qufu, Shandong 273165)

(^d Basic Medical College, Xinxiang Medical University, Xinxiang, Henan 453003)

Abstract An efficient and convenient copper-catalyzed cascade synthesis of C-4 phosphonated isoquinolin-1(2*H*)-ones has been initially proposed. This is the first example for the construction of phosphine-containing heterocycles through copper-catalyzed Ullmann-type coupling reactions using cyanomethylphosphonates as the building blocks, and it will broaden the strategies of organophosphorus synthesis in the field of organic and pharmaceutical chemistry.

Keywords copper-catalysis; isoquinolin-1(2*H*)-one; tandem reaction; organophosphorus

1 Introduction

Isoquinolin-1(2*H*)-one framework represents an important heterocyclic unit that exists in numerous natural products, pharmaceuticals and biologically active compounds (Scheme 1, a).^[1] These molecules possess diverse biological and pharmacological properties as well as structural complexity and have attracted considerable attention from organic and medicinal chemists.^[2] The exten-

sive medicinal and biological activities of isoquinolin-1(2*H*)-ones make them an attractive and challenging synthetic target in organic and medical chemistry.

On the other hand, phosphine-containing motifs are of great importance in pharmaceuticals, agrochemicals, bioactive products and materials chemistry.^[3] Importantly, they are also widely utilized as building blocks of functional materials and ligands for a variety of transition metals.^[4]

* Corresponding authors. E-mail: liuxueliang@xxmu.edu.cn; yangdaoshan@tsinghua.org.cn

Received August 25, 2020; revised October 18, 2020; published online October 21, 2020.

Project supported by the Natural Science Foundation of Shandong Province (No. ZR2016JL012) and the Scientific Research Foundation of Qingdao University of Science and Technology.

山东省自然科学基金(No. ZR2016JL012)、青岛科技大学高科技人才资助项目。

Therefore, the creation of new organophosphorus compounds and development of efficient and practical methods for organophosphines synthesis is a significant goal in modern organic synthesis. Organic phosphate compounds have attracted considerable attention in recent years because of their wide biological applications in medical chemistry (Scheme 1, c).^[5] Furthermore, they are also intermediates in the synthesis of potentially significant enzyme inhibitors, antibacterial agents and phosphonic acid-based antibiotics.^[6] Therefore, it is still a challenging and attractive subject to develop more efficient and practical synthesis methods for introducing phosphate groups into organic skeletons.

The combined molecules of phosphate and isoquinolin-1(2*H*)-one frameworks might be valuable substrates for

the formation of biologically active molecules with different structural features from the two units (Scheme 1, b). However, the methods for their preparation are very rare.^[7] Therefore, it is necessary to develop an efficient and convenient approach to phosphonated isoquinolin-1(2*H*)-one derivatives.

Cyanomethylphosphonates are useful and easily prepared chemical materials, which have been used in the phosphine-containing compounds and cyanides synthesis.^[8] However, a literature survey showed that using cyanomethylphosphonates as building blocks in transition-metal-catalyzed organic reactions is rather limited, especially in the synthesis of phosphonated heterocycles.^[9] Consequently, it is of great significance to explore the transition-metal-catalyzed transformations for the synthesis

Scheme 1 Bioactive molecules containing isoquinolin-1(2*H*)-one and phosphonate moieties and our design of phosphonated isoquinolin-1(2*H*)-ones

(2 equiv.) at 100 °C (Entries 1~4), and *L*-proline provided the highest yield (Entry 1). Notably, the target product (**3b**) was obtained only in 15% yield in the absence of ligand (Entry 10), which indicated the crucial acceleration of the ligand in the current transformation. Next, various bases were investigated (Entries 1, 11~14), and K_2CO_3 was found to be the most effective base (Entry 1). We attempted to use different solvents including *N,N*-dimethylformamide (DMF), DMSO, toluene, and *N*-methyl pyrrolidone (NMP), and DMSO gave the highest yield (Entries 11, 15~17). Subse-

quently, five copper catalysts were examined, and CuCl showed the highest reaction activity (Entries 11, 18~21). Finally, reaction temperatures were also screened and the yield of the target product maximized at 100 °C (Entries 11, 22 and 23).

The scope of copper-catalyzed domino synthesis of C-4 phosphonated isoquinolin-1(2*H*)-one derivatives from reactions of substituted 2-bromobenzamides with cyanomethylphosphonates was investigated under the optimal conditions (Table 2). We were pleased to find that various

Table 2 Reaction scope^{a,b}

^a Reaction conditions: N_2 atmosphere, **1** (0.2 mmol), **2** (0.3 mmol), CuCl (0.02 mmol), *L*-proline (0.04 mmol), K_2CO_3 (0.4 mmol), DMSO (2 mL), 6 h. ^b Isolated yield.

2-bromobenzamides derivatives could be transferred into the C-4 phosphonated isoquinolin-1(2*H*)-one derivatives in moderate to good yields. The structure of the product was unambiguously confirmed by the single crystal X-ray diffraction study of compound **3b**. Electron-donating groups such as methoxy, methyl and electron-withdrawing groups such as chloro-, bromo- and trifluoromethyl groups were well-tolerated under the copper-catalyzed reaction conditions. Different cyanomethylphosphonates were well compatible in the present domino transformation, and they did not display evidently difference of reactivity.

Furthermore, the gram-scale reaction was also performed in the usual laboratory setup with a one-neck round-bottomed flask fitted with a nitrogen balloon, and the reaction afforded **3a** in 51% yield (Scheme 2, a). In addition, the application of our present protocol for the synthesis of C-6 phosphonated benzo[4,5]imidazo[1,2-*b*]isoquinolin-11(5*H*)-one was explored. Satisfactorily, when 2-bromo-*N*-(2-bromophenyl)benzamide was used as the substrate, the corresponding conjugate containing isoquinolin-1(2*H*)-one and benzo[*d*]imidazole frameworks was obtained in 58% yield

(Scheme 2, b).

To understand the mechanism further, some control experiments were performed as shown in Scheme 3. Treatment of 2-bromo-*N*-phenylbenzamide (**1a**) with diethyl (cyanomethyl)phosphonate (**2a**) in the absence of CuCl did not afford the desired product **3a** (Scheme 3a), indicating that copper catalyst is essential for this transformation. Coupling of *N*-phenylbenzamide (**1m**) with (**2a**) did not work under our standard condition (Scheme 3, b).

On the basis of the literature reports,^[12] a possible mechanism for the copper-catalyzed domino method for the synthesis of C-4 phosphonated isoquinolin-1(2*H*)-one derivatives is suggested in Scheme 4. Reaction of CuCl with *L*-proline gives the copper complex CuL. Oxidative addition of CuL with 2-bromobenzamides affords the intermediate **I**. Treatment of cyanomethylphosphonates **2** with **I** in the presence of base K₂CO₃ leads to the complex **II**, and then reductive elimination of **II** produces the intermediate **III**. Intramolecular nucleophilic of NH amide to CN with the assistance of CuL affords **IV**, and isomerization of **IV** forms **3**.

Scheme 2 Synthesis of **3a** on a gram scale and copper-catalyzed domino synthesis of C-6 phosphonated benzimidazo-[1,2-*b*]-isoquinolin-11-one

Scheme 3 Control experiments

Scheme 3 Probable reaction mechanism

3 Conclusions

In summary, a protocol for copper-catalyzed cross-coupling of a variety of 2-bromobenzamides with cyanomethylphosphonates to afford C-4 phosphonated isoquinolin-1(2H)-one derivatives was developed. This is the first example for the construction of phosphine-containing heterocycles through transition-metal-catalyzed Ullmann-type coupling reactions using cyanomethylphosphonates as the phosphine sources. The developed method will broaden the strategies of organophosphorus synthesis in the field of pharmaceutical and organic chemistry.

4 Experimental section

4.1 General information

^1H NMR and ^{13}C NMR spectra were recorded at 25 °C at 500 and 125 MHz on a Bruker 500 spectrometer. ^1H NMR and ^{13}C NMR were recorded using tetramethylsilane (TMS) as the internal standard in CDCl_3 (^1H NMR: TMS at δ 0.00, CDCl_3 at δ 7.28; ^{13}C NMR: CDCl_3 at δ 77.0), or in $\text{DMSO}-d_6$ (^1H NMR: TMS at δ 0.00, $\text{DMSO}-d_6$ at δ 2.50; ^{13}C NMR: $\text{DMSO}-d_6$ at δ 44.0). Mass analyses and HRMS were obtained on a Finnigan-LCQDECA mass spectrometer and a Bruker Daltonics Bio-TOF-Q mass spectrometer by the ESI method, respectively. Column chromatography was performed on silica gel (200~300 mesh). All commercially available reagent-grade chemicals were purchased from chemical suppliers and used as received without further purification.

4.2 General experimental procedures for the synthesis of phosphonated isoquinolin-1(2H)-ones (3)

A 25 mL Schlenk tube equipped with a magnetic stirring

bar was charged with the substituted 2-bromo-*N*-phenylbenzamides (0.2 mmol), cyanomethylphosphonates (0.3 mmol), CuCl (0.02 mmol) and K_2CO_3 (0.4 mmol). The tube was evacuated twice and backfilled with nitrogen, and 2 mL of DMSO and *L*-proline (0.04 mmol) were added to the tube under nitrogen atmosphere by an injection syringe. Then the mixture was allowed to stir under nitrogen atmosphere at 100 °C for 6 h. After being cooled to room temperature, the mixture was poured to 10 mL of water, and the resulting solution was extracted with ethyl acetate (10 mL $\times$ 4). The combined organic layer was concentrated and the residue was purified by column chromatography on silica gel using petroleum ether/ethyl acetate as eluent to provide the desired product.

Diethyl (3-amino-1-oxo-2-phenyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3a**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). Yellow solid, 46.9 mg, 63% yield. m.p. 132~133 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.24 (d, $J=7.8$ Hz, 1H), 7.82 (d, $J=8.4$ Hz, 1H), 7.60~7.51 (m, 4H), 7.31 (d, $J=7.8$ Hz, 2H), 7.18 (t, $J=7.5$ Hz, 1H), 4.21~4.14 (m, 2H), 4.08~4.00 (m, 2H), 1.32 (t, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.5, 154.5 (d, $J=21.3$ Hz), 137.5 (d, $J=7.4$ Hz), 135.1, 133.1, 130.4, 129.7, 128.8, 128.2, 123.3 (d, $J=3.3$ Hz), 122.7, 120.0 (d, $J=12.5$ Hz), 73.2 (d, $J=199.9$ Hz), 61.7 (d, $J=4.6$ Hz), 16.1 (d, $J=7.0$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 24.4. HRMS calcd for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_4\text{P}$ [$\text{M}+\text{H}$] $^+$ 373.1312, found 373.1312.

Diethyl (3-amino-1-oxo-2-(*p*-tolyl)-1,2-dihydroisoquinolin-4-yl)phosphonate (**3b**): Eluent: petroleum ether/ethyl acetate ($V:V=4:1$). Yellow solid, 61.0 mg, 79% yield. m.p. 136~137 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.25 (d, $J=7.8$ Hz, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.55 (t, $J=7.7$ Hz, 1H), 7.39 (d, $J=7.9$ Hz, 2H), 7.18 (t, $J=7.9$ Hz, 3H), 4.26~4.14 (m, 2H), 4.14~3.90 (m, 2H), 2.44 (s, 3H), 1.33 (t, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.7, 154.3 (d, $J=20.9$ Hz), 139.9, 137.5 (d, $J=7.4$ Hz), 133.1, 132.3, 131.1, 128.5, 128.3, 123.3 (d, $J=3.2$ Hz), 122.7, 120.1 (d, $J=12.6$ Hz), 73.1 (d, $J=199.8$ Hz), 61.7 (d, $J=4.5$ Hz), 21.2, 16.2 (d, $J=7.1$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 24.5; HRMS calcd for $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_4\text{P}$ [$\text{M}+\text{H}$] $^+$ 387.1468, found 387.1462.

Diethyl(3-amino-2-(4-methoxyphenyl)-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3c**): Eluent: petroleum ether/ethyl acetate ($V:V=4:1$). White solid, 62.7 mg, 78% yield. m.p. 167~168 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.25 (d, $J=8.0$ Hz, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.57~7.53 (m, 1H), 7.24~7.21 (m, 2H), 7.18 (t, $J=7.5$ Hz, 1H), 7.10~7.08 (m, 2H), 4.22~4.14 (m, 2H), 4.11~4.02 (m, 2H), 3.87 (s, 3H), 1.33 (t, $J=7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.8, 160.3, 154.4 (d, $J=20.9$ Hz), 137.5 (d, $J=7.4$ Hz), 133.1, 129.9, 128.3, 127.3, 123.3 (d, $J=3.2$ Hz), 122.6, 120.1 (d, $J=12.6$ Hz), 115.7, 73.1 (d, $J=200$ Hz), 61.7 (d, $J=4.5$ Hz), 55.5, 16.2 (d, $J=7.1$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 24.5; HRMS calcd for $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_5\text{P}$ [$\text{M}+\text{H}$] $^+$ 403.1417, found 403.1420.

Diethyl(3-amino-2-(4-chlorophenyl)-1-oxo-1,2-dihydro-

isoquinolin-4-yl)phosphonate (**3d**): Eluent: petroleum ether/ethyl acetate ($V:V=4:1$). Yellow oil, 59.3 mg, 73% yield. m.p. 142~144 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.24 (d, $J=8.0$ Hz, 1H), 7.82 (d, $J=8.4$ Hz, 1H), 7.58~7.55 (m, 3H), 7.28 (d, 2H, $J=8.8$ Hz), 7.20 (t, $J=7.5$ Hz, 1H), 4.22~4.15 (m, 2H), 4.09~4.01 (m, 2H), 1.33 (t, $J=7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.5, 153.7 (d, $J=21.3$ Hz), 137.5 (d, $J=7.4$ Hz), 135.9, 133.6, 133.4, 130.8, 130.4, 128.3, 123.5 (d, $J=3.2$ Hz), 122.9, 120.0 (d, $J=12.5$ Hz), 73.7 (d, $J=199.7$ Hz), 61.8 (d, $J=4.6$ Hz), 16.2 (d, $J=7.1$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 24.1; HRMS calcd for $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 407.0922, found 407.0920.

Diethyl(3-amino-2-(3-chlorophenyl)-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3e**): Eluent: petroleum ether/ethyl acetate ($V:V=5:1$). Yellow oil, 54.4 mg, 67% yield. m.p. 112~113 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.21 (d, $J=7.9$ Hz, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.56~7.49 (m, 3H), 7.34 (s, 1H), 7.22 (d, $J=7.0$ Hz, 1H), 7.18 (t, $J=7.5$ Hz, 1H), 4.20~4.13 (m, 2H), 4.07~3.99 (m, 2H), 1.31 (t, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.3, 153.5 (d, $J=21.2$ Hz), 137.4, 136.2, 135.8, 133.3, 131.3, 130.0, 129.3, 128.2, 127.2, 123.4, 122.8, 119.8 (d, $J=12.4$ Hz), 73.6 (d, $J=199.2$ Hz), 61.7, 16.1 (d, $J=6.8$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 24.1; HRMS calcd for $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 407.0922, found 407.0920.

Diethyl(3-amino-2-(4-bromophenyl)-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3f**): Eluent: petroleum ether/ethyl acetate ($V:V=5:1$). Yellow solid, 54.9 mg, 61% yield. m.p. 171~172 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.22 (d, $J=8.0$ Hz, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.71 (d, $J=8.5$ Hz, 2H), 7.55 (t, $J=7.7$ Hz, 1H), 7.19 (t, $J=8.9$ Hz, 3H), 4.21~4.14 (m, 2H), 4.11~4.00 (m, 2H), 1.32 (t, $J=7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.4, 153.6 (d, $J=21.2$ Hz), 137.5 (d, $J=7.3$ Hz), 134.1, 133.7, 133.3, 130.7, 128.3, 124.0, 123.4 (d, $J=3.2$ Hz), 122.9, 119.9 (d, $J=12.5$ Hz), 73.7 (d, $J=200$ Hz), 61.8 (d, $J=4.6$ Hz), 16.2 (d, $J=7.0$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 24.1; HRMS calcd for $\text{C}_{19}\text{H}_{21}\text{BrN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 451.0417, found 451.0412.

Diethyl(3-amino-6-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3g**): Eluent: petroleum ether/ethyl acetate ($V:V=2:1$). Yellow solid, 50.4 mg, 62% yield. m.p. 170~171 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.15 (d, $J=8.5$ Hz, 1H), 7.81 (s, 1H), 7.59 (t, $J=7.3$ Hz, 2H), 7.56~7.49 (m, 1H), 7.30 (d, $J=7.6$ Hz, 2H), 7.12 (d, $J=8.5$ Hz, 1H), 4.22~4.17 (m, 2H), 4.09~4.04 (m, 2H), 1.35 (t, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.0, 154.6 (d, $J=21.3$ Hz), 139.9, 139.0 (d, $J=7.0$ Hz), 134.8, 130.5, 130.0, 128.8, 123.2, 122.8 (d, $J=3.2$ Hz), 118.4, 118.3, 73.0 (d, $J=201.3$ Hz), 62.0 (d, $J=4.9$ Hz), 16.2 (d, $J=6.8$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 23.3. HRMS calcd for $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 407.0922; found 407.0927.

Diethyl(3-amino-2-butyl-6-chloro-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3h**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). Yellow oil, 44.0 mg,

57% yield. m.p. 125~126 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.17 (d, $J=8.6$ Hz, 1H), 7.75 (d, $J=1.6$ Hz, 1H), 7.11 (d, $J=8.6$ Hz, 1H), 6.95 (br s, 2H), 4.20~4.15 (m, 2H), 4.14~4.08 (m, 2H), 4.05~3.97 (m, 2H), 1.72 (s, 2H), 1.46 (d, $J=7.6$ Hz, 2H), 1.32 (t, $J=7.1$ Hz, 6H), 0.99 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 161.9, 154.7 (d, $J=20$ Hz), 139.4, 138.4, 129.9, 123.2, 122.7 (d, $J=3.4$ Hz), 117.9, 73.7 (d, $J=200$ Hz), 61.9 (d, $J=4.7$ Hz), 41.8, 29.6, 20.3, 16.2 (d, $J=6.9$ Hz), 13.7; ^{31}P NMR (202 MHz, CDCl_3) δ : 23.8; HRMS calcd for $\text{C}_{17}\text{H}_{25}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 387.1235; found 387.1240.

Diethyl(3-amino-7-methyl-1-oxo-2-phenethyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3i**): Eluent: petroleum ether/ethyl acetate ($V:V=4:1$). Yellow solid, 43.9 mg, 53% yield. m.p. 139~141 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.03 (s, 1H), 7.62 (d, $J=8.5$ Hz, 1H), 7.28 (d, $J=7.6$ Hz, 1H), 7.25~7.20 (m, 2H), 7.18 (d, $J=6.4$ Hz, 3H), 6.40 (br s, 2H), 4.25 (t, $J=7.4$ Hz, 2H), 4.06~3.98 (m, 2H), 3.90~3.82 (m, 2H), 3.01 (t, $J=7.5$ Hz, 2H), 2.33 (s, 3H), 1.20 (t, $J=7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.5, 153.9 (d, $J=21.1$ Hz), 138.1, 134.4, 132.7, 128.9, 128.7, 127.6, 127.0, 123.4, 119.8, 119.7, 74.7 (d, $J=197.5$ Hz), 61.6 (d, $J=4.2$ Hz), 44.1, 33.9, 20.8, 16.1 (d, $J=7.1$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 19.7; HRMS calcd for $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 415.1751, found 415.1754.

Diethyl(3-amino-7-chloro-1-oxo-2-phenethyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3j**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). Yellow solid, 56.4 mg, 65% yield. m.p. 147~148 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.17 (s, 1H), 7.65 (d, $J=9.0$ Hz, 1H), 7.38 (d, $J=8.9$ Hz, 1H), 7.25~7.22 (m, 2H), 7.20~7.17 (m, 3H), 6.52 (br s, 2H), 4.24 (t, $J=7.4$ Hz, 2H), 4.06~3.99 (m, 2H), 3.91~3.83 (m, 2H), 3.01 (t, $J=7.5$ Hz, 2H), 1.20 (t, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 161.5, 154.4 (d, $J=21.3$ Hz), 137.9, 135.4, 135.4, 133.1, 129.0, 128.6 (d, $J=10$ Hz), 127.3, 127.2, 125.2, 120.9 (d, $J=12.8$ Hz), 74.6 (d, $J=200.1$ Hz), 61.8 (d, $J=4.3$ Hz), 44.2, 33.7, 16.1 (d, $J=7.0$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 23.9; HRMS calcd for $\text{C}_{21}\text{H}_{25}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 435.1235; found 435.1163.

Diethyl(3-amino-2-butyl-1-oxo-7-(trifluoromethyl)-1,2-dihydroisoquinolin-4-yl)phosphonate (**3k**): Eluent: petroleum ether/ethyl acetate ($V:V=7:1$). Yellow solid, 63.9 mg, 76% yield. m.p. 111~113 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.52 (s, 1H), 7.84 (d, $J=8.8$ Hz, 1H), 7.66 (d, $J=8.8$ Hz, 1H), 7.11 (br s, 2H), 4.19~4.08 (m, 4H), 4.03~3.95 (m, 2H), 1.76~1.70 (m, 2H), 1.49~1.44 (m, 2H), 1.29 (t, $J=7.1$ Hz, 6H), 0.99 (t, $J=7.3$ Hz, 3H); ^{31}P NMR (202 MHz, CDCl_3) δ : 23.7 (d, $J=12.1$ Hz); HRMS calcd for $\text{C}_{18}\text{H}_{24}\text{F}_3\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 421.1499, found 421.1493.

Diisopropyl (3-amino-1-oxo-2-(*p*-tolyl)-1,2-dihydroisoquinolin-4-yl)phosphonate (**3l**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). Yellow solid, 71.2 mg, 86% yield. m.p. 124~125 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.23 (d, $J=8.0$ Hz, 1H), 7.82 (d, $J=8.5$ Hz, 1H), 7.53~7.50 (m, 1H), 7.37 (d, $J=8.0$ Hz, 2H), 7.19~7.13 (m, 3H), 4.69~4.60 (m, 2H), 2.42 (s, 3H), 1.41 (d, $J=6.2$ Hz, 6H),

1.18 (d, $J=6.2$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.7, 153.7 (d, $J=20$ Hz), 139.7, 137.6 (d, $J=7.4$ Hz), 132.6, 132.4, 131.0, 128.5, 128.1, 123.8 (d, $J=3.2$ Hz), 122.4, 120.0 (d, $J=12.5$ Hz), 74.5 (d, $J=200$ Hz), 70.7 (d, $J=5$ Hz), 24.1 (d, $J=3.8$ Hz), 23.8 (d, $J=6.3$ Hz), 21.1; ^{31}P NMR (202 MHz, CDCl_3) δ : 21.9; HRMS calcd for $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 415.1781, found 415.1776.

Diisopropyl (3-amino-2-butyl-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3m**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). Yellow solid, 52.5 mg, 69% yield. m.p. 119~121 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.18 (d, $J=8.0$ Hz, 1H), 7.71 (d, $J=8.5$ Hz, 1H), 7.41 (t, $J=7.7$ Hz, 1H), 7.09 (t, $J=7.5$ Hz, 1H), 6.91 (br s, 2H), 4.56~4.50 (m, 2H), 4.09~4.06 (m, 2H), 1.69~1.66 (m, 2H), 1.42~1.36 (m, 2H), 1.34 (t, $J=6.5$ Hz, 6H), 1.07 (d, $J=6.2$ Hz, 6H), 0.92 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.5, 153.7 (d, $J=21.3$ Hz), 137.1 (d, $J=7.2$ Hz), 132.2, 128.0, 123.8 (d, $J=3.2$ Hz), 122.6, 119.6 (d, $J=12.5$ Hz), 75.5 (d, $J=197.5$ Hz), 70.6 (d, $J=4.6$ Hz), 41.8, 29.6, 23.9 (d, $J=2.5$ Hz), 23.6 (d, $J=5$ Hz), 20.2, 13.6; ^{31}P NMR (202 MHz, CDCl_3) δ : 22.3; HRMS calcd for $\text{C}_{19}\text{H}_{30}\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 381.1938, found 381.1941.

Diisopropyl(3-amino-2-(3-chlorophenyl)-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3n**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). White solid, 61.2 mg, 73% yield. m.p. 166~167 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.21 (d, $J=8.0$ Hz, 1H), 7.83 (d, $J=8.5$ Hz, 1H), 7.55~7.48 (m, 3H), 7.35 (s, 1H), 7.24~7.22 (m, 1H), 7.17 (t, $J=7.5$ Hz, 1H), 4.68~4.61 (m, 2H), 1.41 (d, $J=6.2$ Hz, 6H), 1.18 (dd, $J=6.1$, 3.0 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.4, 153.1 (d, $J=21.4$ Hz), 137.6 (d, $J=7.2$ Hz), 136.3, 135.9, 132.9, 131.3, 130.0, 129.3, 128.1, 127.3, 124.0 (d, $J=3.3$ Hz), 122.7, 119.9 (d, $J=12.5$ Hz), 75.1 (d, $J=199.4$ Hz), 70.8 (d, $J=2.5$ Hz), 70.7 (d, $J=3.0$ Hz), 23.8 (d, $J=33.8$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 21.5. HRMS calcd for $\text{C}_{21}\text{H}_{25}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 435.1235, found 435.1163.

Diisopropyl(3-amino-2-(4-methoxyphenyl)-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3o**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). Yellow solid, 67.1 mg, 78% yield. m.p. 132~133 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.22 (d, $J=7.9$ Hz, 1H), 7.81 (d, $J=8.5$ Hz, 1H), 7.51 (t, $J=7.7$ Hz, 1H), 7.22 (d, $J=8.7$ Hz, 2H), 7.15 (t, $J=7.5$ Hz, 1H), 7.06 (d, $J=8.7$ Hz, 2H), 4.67~4.61 (m, 2H), 3.84 (s, 3H), 1.41 (d, $J=6.2$ Hz, 6H), 1.18 (d, $J=6.2$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.9, 160.2, 153.9 (d, $J=21.2$ Hz), 137.6 (d, $J=7.3$ Hz), 132.7, 129.9, 128.1, 127.3, 123.8, 122.5, 120.0 (d, $J=12.6$ Hz), 115.6, 74.5 (d, $J=200.1$ Hz), 70.6 (d, $J=3.8$ Hz), 55.5, 24.0 (d, $J=3.8$ Hz), 23.7 (d, $J=6.3$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 21.9; HRMS calcd for $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_5\text{P}$ $[\text{M}+\text{H}]^+$ 431.1730, found 431.1736.

Diisopropyl(3-amino-6-chloro-1-oxo-2-phenyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3p**): Eluent: petroleum ether/ethyl acetate ($V:V=3:1$). White solid, 66.0 mg, 76% yield. m.p. 163~164 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.15 (d, $J=8.5$ Hz, 1H), 7.86 (d, $J=1.7$ Hz, 1H), 7.58 (t,

$J=7.6$ Hz, 2H), 7.52 (t, $J=7.4$ Hz, 1H), 7.32~7.29 (m, 2H), 7.11 (d, $J=8.5$ Hz, 1H), 4.72~4.65 (m, 2H), 1.45 (d, $J=6.2$ Hz, 6H), 1.25 (d, $J=6.2$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 161.8, 154.1 (d, $J=20.1$ Hz), 139.3, 139.0 (d, $J=6.8$ Hz), 134.7, 130.3, 129.7, 128.7, 123.2 (d, $J=3.3$ Hz), 122.8, 118.3, 118.2, 74.1 (d, $J=201.3$ Hz), 70.8 (d, $J=5.1$ Hz), 23.8 (d, $J=3.8$ Hz), 23.6 (d, $J=5.0$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 20.6. HRMS calcd for $\text{C}_{21}\text{H}_{25}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 435.1235, found 435.1230.

Diisopropyl(3-amino-2-butyl-6-chloro-1-oxo-1,2-dihydroisoquinolin-4-yl)phosphonate (**3q**): Eluent: petroleum ether/ethyl acetate ($V:V=4:1$). Yellow solid, 37.3 mg, 45% yield. m.p. 110~112 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.15 (d, $J=8.6$ Hz, 1H), 7.77 (s, 1H), 7.08 (d, $J=8.6$ Hz, 1H), 7.00 (br s, 2H), 4.64~4.58 (m, 2H), 4.11~4.08 (m, 2H), 1.73~1.70 (m, 2H), 1.47~1.44 (m, 2H), 1.41 (d, $J=6.0$ Hz, 6H), 1.17 (d, $J=6.1$ Hz, 6H), 0.97 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 161.9, 154.2 (d, $J=20.1$ Hz), 139.0, 138.5 (d, $J=6.8$ Hz), 129.8, 123.2 (d, $J=2.5$ Hz), 123.0, 117.9 (d, $J=12.3$ Hz), 75.1 (d, $J=200.1$ Hz), 71.0 (d, $J=5.1$ Hz), 41.8, 29.6 (d, $J=6.5$ Hz), 23.9 (d, $J=3.8$ Hz), 23.7 (d, $J=5.2$ Hz), 20.3, 13.7; ^{31}P NMR (202 MHz, CDCl_3) δ : 21.1; HRMS calcd for $\text{C}_{19}\text{H}_{29}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 415.1548, found 415.1534.

Diisopropyl(3-amino-7-methyl-1-oxo-2-phenethyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3r**): Eluent: petroleum ether/ethyl acetate ($V:V=5:1$). Yellow solid, 63.7 mg, 72% yield. m.p. 135~137 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.09 (s, 1H), 7.71 (d, $J=8.5$ Hz, 1H), 7.35~7.28 (m, 4H), 7.25 (s, 2H), 6.48 (br s, 2H), 4.57~4.50 (m, 2H), 4.33 (t, $J=7.4$ Hz, 2H), 3.09 (t, $J=7.5$ Hz, 2H), 2.39 (s, 3H), 1.38 (d, $J=6.2$ Hz, 6H), 1.12 (d, $J=6.2$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.6, 153.4 (d, $J=21.3$ Hz), 138.2, 134.6 (d, $J=7.3$ Hz), 134.0, 132.5, 128.9, 128.7, 127.4, 127.0, 124.1 (d, $J=3.1$ Hz), 119.8 (d, $J=12.5$ Hz), 75.6, 70.6 (d, $J=4.4$ Hz), 44.0, 33.9, 24.0 (d, $J=3.8$ Hz), 23.6 (d, $J=5$ Hz), 20.8; ^{31}P NMR (202 MHz, CDCl_3) δ : 22.1; HRMS calcd for $\text{C}_{24}\text{H}_{32}\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 443.4951, found 443.4946.

Diisopropyl(3-amino-7-chloro-1-oxo-2-phenethyl-1,2-dihydroisoquinolin-4-yl)phosphonate (**3s**): Eluent: petroleum ether/ethyl acetate ($V:V=5:1$). Yellow solid, 59.2 mg, 64% yield. m.p. 146~147 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.23 (s, 1H), 7.74 (d, $J=9.0$ Hz, 1H), 7.43 (d, $J=9.0$ Hz, 1H), 7.30~7.28 (m, 2H), 7.20 (d, $J=7.9$ Hz, 3H), 6.61 (br s, 2H), 4.57~4.51 (m, 2H), 4.33~4.28 (m, 2H), 3.08 (t, $J=7.5$ Hz, 2H), 1.38 (d, $J=6.2$ Hz, 6H), 1.12 (d, $J=6.2$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 161.5, 153.9 (d, $J=25$ Hz), 137.9, 135.6 (d, $J=7.2$ Hz), 132.7, 128.9, 128.7, 128.5, 127.1 (d, $J=6.3$ Hz), 125.7 (d, $J=3.0$ Hz), 120.9, 120.8, 75.4, 70.9 (d, $J=4.6$ Hz), 44.2, 33.7, 24.0 (d, $J=3.7$ Hz), 23.6 (d, $J=5.4$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 21.2; HRMS calcd for $\text{C}_{23}\text{H}_{29}\text{ClN}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 463.1548, found 463.1550.

Diisopropyl(3-amino-2-butyl-1-oxo-7-(trifluoromethyl)-1,2-dihydroisoquinolin-4-yl)phosphonate (**3t**): Eluent: petroleum ether/ethyl acetate ($V:V=4:1$). Yellow solid,

59.2 mg, 66% yield. m.p. 86~88 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.49 (s, 1H), 7.85 (d, $J=8.8$ Hz, 1H), 7.63 (d, $J=8.8$ Hz, 1H), 7.18 (br s, 2H), 4.64~4.56 (m, 2H), 4.14~4.11 (m, 2H), 1.74~1.71 (m, 2H), 1.47~1.38 (m, 2H), 1.38 (s, 6H), 1.13 (d, $J=6.2$ Hz, 6H), 0.97 (t, $J=7.3$ Hz, 3H); ^{31}P NMR (202 MHz, CDCl_3) δ : 23.1; HRMS calcd for $\text{C}_{20}\text{H}_{29}\text{F}_3\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 449.1812, found 449.1819.

Diethyl(11-oxo-5,11-dihydrobenzo[4,5]imidazo[1,2-*b*]-isoquinolin-6-yl)phosphonate (**3u**): Eluent: petroleum ether/ethyl acetate ($V:V=7:1$). Yellow solid, 42.9 mg, 58% yield. m.p. 204~205 °C; ^1H NMR (500 MHz, CDCl_3) δ : 11.18 (br s, 1H), 8.76 (d, $J=8.0$ Hz, 1H), 8.55 (d, $J=8.1$ Hz, 1H), 7.92 (d, $J=8.4$ Hz, 1H), 7.67 (t, $J=7.6$ Hz, 1H), 7.43 (t, $J=7.7$ Hz, 1H), 7.34~7.29 (m, 3H), 4.23~4.12 (m, 2H), 4.06~3.98 (m, 2H), 1.32 (t, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 160.3, 147.8 (d, $J=19.6$ Hz), 136.6 (d, $J=7.0$ Hz), 133.0, 131.2, 128.2, 128.1, 126.4, 123.5 (d, $J=3.5$ Hz), 123.0, 121.9, 119.4, 116.9, 109.7, 71.8 (d, $J=216.1$ Hz), 62.1 (d, $J=4.3$ Hz), 16.2 (d, $J=7.0$ Hz); ^{31}P NMR (202 MHz, CDCl_3) δ : 22.4; HRMS calcd for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}_4\text{P}$ $[\text{M}+\text{H}]^+$ 371.1155, found 371.1162.

Supporting Information ^1H NMR and ^{13}C NMR spectra for all compounds and ORTEP structures and spectroscopic data of compound **3b**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Ratnayake, R.; Lacey, E.; Tennant, S.; Gill, J. H.; Capon, R. J. *Chem.-Eur. J.* **2007**, *13*, 1610.
(b) Billamboz, M.; Bailly, F.; Lion, C.; Touati, N.; Vezin, H.; Calmels, C.; Andreola, M.-L.; Christ, F.; Debyser, Z.; Cotellet, P. J. *Med. Chem.* **2011**, *54*, 1812.
(c) Dickinson, R. P.; Bell, A. W.; Hitchcock, C. A.; Narayana-Swami, S.; Ray, S. J.; Richardson, K.; Troke, P. F. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 2031.
(d) Jatav, V.; Mishra, P.; Kashaw, S.; Stables, J. P. *Eur. J. Med. Chem.* **2008**, *43*, 135.
(e) Glushkov, V. A.; Shklyayev, Y. V. *Chem. Heterocycl. Compd.* **2001**, *37*, 66.
- [2] (a) Kashaw, S. K.; Gupta, V.; Kashaw, V.; Mishra, P.; Stables, J. P.; Jain, N. K. *Med. Chem. Res.* **2010**, *19*, 250.
(b) Wang, Z.; He, W.-M. *Chin. J. Org. Chem.* **2019**, *39*, 3594 (in Chinese).
(王峥, 何卫民, 有机化学, **2019**, *39*, 3594.)
(c) Li, X.-L.; Wang, J.-Q.; Li, L.; Yin, Y.-W.; Ye, L.-W. *Acta Chim. Sinica* **2016**, *74*, 49 (in Chinese).
(李新玲, 王佳琪, 李龙, 尹应武, 叶龙武, 化学学报, **2016**, *74*, 49.)
- [3] (a) Horsman, G. P.; Zechel, D. L. *Chem. Rev.* **2017**, *117*, 5704.
(b) Gahungu, M.; Arguelles-Arias, A.; Fickers, P.; Zervosen, A.; Joris, B.; Damblon, C.; Luxen, A. *Bioorg. Med. Chem.* **2013**, *21*, 4958.
(c) Tang, W.; Zhang, X. *Chem. Rev.* **2003**, *103*, 3029.
(d) Bhattacharya, A. K.; Thyagarajan, G. *Chem. Rev.* **1981**, *81*, 415.
(e) Kirumakki, S.; Huang, J.; Subbiah, A.; Yao, J.; Rowland, A.; Smith, B.; Mukherjee, A.; Samarajeewa, S.; Clearfield, A. J. *Mater. Chem.* **2009**, *19*, 2593.
(f) Krylov, S.; Kashemirov, B. A.; Hilfinger, J. M.; McKenna, C. E. *Mol. Pharmaceutics* **2013**, *10*, 445.
(g) Qiao, B.; Cao, H.-Q.; Huang, Y.-J.; Zhang, Y.; Nie, J.; Zhang, F.-G.; Ma, J.-A. *Chin. J. Chem.* **2018**, *36*, 809.
- [4] (a) Schlummer, B.; Scholz, U. *Adv. Synth. Catal.* **2004**, *346*, 1599.
(b) Hayashi, T. *Acc. Chem. Res.* **2000**, *33*, 354.
- [5] (a) Zhuang, R.; Xu, J.; Cai, Z.; Tang, G.; Fang, M.; Zhao, Y. *Org. Lett.* **2011**, *13*, 2110.
(b) Lee, H.; Yun, J. *Org. Lett.* **2018**, *20*, 7961.
(c) Mehellou, Y.; Rattan, H. S.; Balzarini, J. J. *Med. Chem.* **2018**, *61*, 2211.
- [6] Demmer, C. S.; Krogsgaard-Larsen, N.; Bunch, L. *Chem. Rev.* **2011**, *111*, 7981.
- [7] Li, B.; Yang, J.; Xu, H.; Song, H.; Wang, B. *J. Org. Chem.* **2015**, *80*, 12397.
- [8] (a) Shi, W.; Li, J.; Su, D.; Wang, X.; Zhang, L.; Pan, L.; Wu, X.; Wu, H. *Angew. Chem., Int. Ed.* **2019**, *58*, 1106.
(b) White, J. D.; Laura, Q.; Wang, G. J. *Org. Chem.* **2007**, *72*, 1717.
- [9] Zhu, W.; Zhang, D.; Yang, N.; Liu, H. *Chem. Commun.* **2014**, *50*, 10634.
- [10] (a) Yang, D.; An, B.; Wei, W.; Tian, L.; Huang, B.; Wang, H. *ACS Comb. Sci.* **2015**, *17*, 113.
(b) Yan, K.; Yang, D.; Wei, W.; Lu, S.; Li, G.; Zhao, C.; Zhang, Q.; Wang, H. *Org. Chem. Front.* **2016**, *3*, 66.
(c) Gan, Z.; Yan, Q.; Li, G.; Li, Q.; Dou, X.; Li, G.-Y.; Yang, D. *Adv. Synth. Catal.* **2019**, *361*, 4558.
- [11] (a) Bhunia, S.; Pawar, G. G.; Kumar, S. V.; Jiang, Y.; Ma, D. *Angew. Chem., Int. Ed.* **2017**, *56*, 16136.
(b) Xu, S.; Lu, J.; Fu, H. *Chem. Commun.* **2011**, *47*, 5596.
(c) Liu, X.; Fu, H.; Jiang, Y.; Zhao, Y. *Angew. Chem., Int. Ed.* **2009**, *48*, 348.
(d) Xu, L.; Jiang, Y.; Ma, D. *Org. Lett.* **2012**, *14*, 1150.
(e) Ma, D.; Cai, Q. *Acc. Chem. Res.* **2008**, *41*, 1450.
(f) Lv, X.; Bao, W. L. *J. Org. Chem.* **2009**, *74*, 5618.
(g) Martin, R.; Rivero, R.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2006**, *45*, 7079.
(h) Liu, Y.; Wan, J.-P. *Org. Biomol. Chem.* **2011**, *9*, 6873.
(i) Verma, A. K.; Kesharwani, T.; Singh, J.; Tandon, V.; Larock, R. C. *Angew. Chem., Int. Ed.* **2009**, *48*, 1138.
(j) Moessner, C.; Bolm, C. *Org. Lett.* **2005**, *7*, 2667.
(k) Cheng, L.; Ge, X.; Liu, X.; Feng, Y. *Chin. J. Org. Chem.* **2020**, *40*, 2008 (in Chinese).
(成琳, 葛新, 刘学民, 冯云辉, 有机化学, **2020**, *40*, 2008.)
(l) Xie, J.; Wang, X.; Wu, F.; Zhang, J. *Chin. J. Org. Chem.* **2019**, *39*, 3026 (in Chinese).
(谢建伟, 汪小创, 吴丰田, 张洁, 有机化学, **2019**, *39*, 3026.)
- [12] (a) Lu, B.; Ma, D. *Org. Lett.* **2006**, *8*, 6115.
(b) Xie, X.; Cai, G.; Ma, D. *Org. Lett.* **2005**, *7*, 4693.
(c) Liu, T.; Wang, R.; Yang, H.; Fu, H. *Chem.-Eur. J.* **2011**, *17*, 6765.
(d) Li, M.; Ning, J.; Yu, L.; Wen, L. *Chin. J. Org. Chem.* **2016**, *36*, 2715 (in Chinese).
(李明, 宁加彬, 于乐, 文丽荣, 有机化学, **2016**, *36*, 2715.)
(e) Zhao, S.; Wang, Z.-L. *Chin. J. Org. Chem.* **2016**, *36*, 862 (in Chinese).
(赵苏艳, 王祖利, 有机化学, **2016**, *36*, 862.)

(Zhao, C.)