铜催化羧酸与芳氨基甲酰氯的脱羧交叉偶联

周敦 樊爱红 李翔 陈春霞* 孙鹏 彭进松*

(东北林业大学化学化工与资源利用学院 哈尔滨 150040)

摘要 报道了在不加碱的条件下, 无配体铜盐能有效催化羧酸和氨基甲酰氯脱羧交叉偶联反应, 以 CuCl_2 为催化剂、苯为溶剂, 于 $120\text{ }^\circ\text{C}$ 条件下反应 48 h 即可实现脱羧交叉偶联过程. 在此标准条件下, 该催化体系对官能团普适性较好, 以较高收率获得了各种酰胺, 产物经 ^1H NMR、 ^{13}C NMR 和 HRMS 表征

关键词 铜催化; 脱羧交叉偶联; 碳-碳成键; 芳氨基甲酰氯; 酰胺

Copper-Catalyzed Decarboxylative Cross-Coupling of Carboxylic Acids and Arylcarbamoyl Chlorides

Zhou, Dun Fan, Aihong Li, Xiang Chen, Chunxia* Sun, Peng Peng, Jinsong*

(College of Chemistry, Chemical Engineering and Resource Utilization, Northeast Forestry University, Harbin 150040)

Abstract A ligand-free copper-catalyzed decarboxylative cross-coupling reaction of carboxylic acids and carbamoyl chlorides in the absence of base was developed. With CuCl_2 as the catalyst, the decarboxylative cross-coupling process could be realized in benzene at $120\text{ }^\circ\text{C}$ in 48 h. Under the standard condition, the catalytic system had good functional group tolerance, and diverse amides were obtained in good to high yields. The structures of products were elucidated by ^1H NMR, ^{13}C NMR and HRMS spectra.

Keywords copper-catalysis; decarboxylative cross-coupling; C—C bond formation; arylcarbamoyl chloride; amide

1 Introduction

Amides are a highly important class of compounds with a wide variety of applications in biochemistry, pharmacy and material science. They are usually prepared through the condensation of carboxylic acids or derivatives and amines.^[1] An alternative and less used approach to amides is the cross-coupling of organometallic reagents and carbamoyl chlorides (Scheme 1, a, b).^[2] For example, Jousseume *et al.*^[3a] reported the first palladium-catalyzed cross-coupling reaction between aryl- and vinyl-organotin and carbamoyl chlorides to give amide derivatives. Ni-catalyzed cross-coupling reactions of carbamoyl chlorides and Grignard reagents were developed for the synthesis of tertiary amides by Lemoucheux *et al.*^[4] The palladium-catalyzed cross-coupling reactions between organozinc and carbamoyl chlorides also provided the corresponding amides.^[5] Using organoboronic reagents (aryl boronic esters^[6] or acids,^[7] alkylboranes,^[8] and potassium trifluoro-

(a) Transition metal-catalyzed cross-coupling using organometallic reagents

(b) Transition metal-free cross-coupling of organometallic reagents

(c) This work: copper-catalyzed decarboxylative coupling

Scheme 1 Preparation of amides using carbamoyl chlorides borates^[9], Pd-catalyzed Suzuki-type reactions have been widely employed for the coupling reaction with carbamoyl

* Corresponding authors. E-mail: ccx0109@nefu.edu.cn; jspeng1998@nefu.edu.cn

Received July 30, 2020; revised October 20, 2020; published online November 18, 2020.

Project supported by the Fundamental Research Funds for the Central Universities (Nos. 2572019CG06, 2572020DR07), the Natural Science Foundation of Heilongjiang Province (Nos. LC2018003, B2017002), and the Programme of Introducing Talents of Discipline to Universities (111 Project, No. B20088).

中央高校基本科研业务费专项资金 (Nos. 2572019CG06, 2572020DR07)、黑龙江省自然科学基金 (Nos. LC2018003, B2017002) 和高等学校学科创新引智计划 (111 计划, No. B20088) 资助项目.

chlorides in the presence of an appropriate base. Without organometallic reagents, Kakiuchi *et al.*^[10] described ruthenium-catalyzed regioselective C—H aminocarbonylation using the carbamoyl chlorides to give tertiary amides. Metal catalyst-free direct coupling of organolithiums^[11] or organocuprates^[12] with carbamoyl chlorides is now frequently used to synthesize secondary and tertiary amides (Scheme 1, b). For example, Rouden *et al.*^[12] described the preparation of tertiary amides via the coupling reactions of organocuprates with carbamoyl chlorides.

Organolithium, magnesium and zinc reagents were used as precursors to make the organocuprates in this reaction. Over the past decades, the carboxylic acids are promising alternatives to organic halides and organometallic reagents in organic synthesis. Transition-metal-catalyzed decarboxylative coupling reactions of carboxylic acids have emerged as powerful tools for carbon-carbon bond formation.^[13] This synthetic strategy does not use expensive and sensitive organometallic reagents, and generates CO₂ instead of metal salts. Palladium,^[14] copper,^[15] silver,^[16] gold,^[17] and rhodium^[18] are capable of inducing the decarboxylation of carboxylic acids to generate organometallic species *in situ*, which can be utilized in various cross-coupling reactions. Inspired by Liu's^[19] report on the decarboxylative cross-coupling of potassium polyfluorobenzoates with aryl halides catalyzed by a CuI/Phen system, herein, we are intrigued by ligand-free copper-catalyzed direct C—C coupling of carboxylic acids with carbamoyl chlorides in the absence of base and set out to explore this possibility (Scheme 1, c).

2 Results and discussion

Using carbamoyl chloride **1a** as the coupling partner, the coupling of acetic acid **2a** instead of acetate salts was examined (Table 1). The blank experiment (without the catalyst) was examined in benzene at 80 °C for 24 h, and no desired product **3aa** was obtained. It was found that **3aa** was formed in 32% yield in the presence of 15 mol% Cu(OAc)₂ in benzene at 80 °C for 24 h (Table 1, Entry 1).

Thus, to optimize reaction conditions, different catalysts, solvents, reaction time and temperature were examined (Table 1). The effect of copper sources [such as Cu(OAc)₂, CuCl₂, CuI, CuBr, CuCl, CuO, and Cu(OTf)₂] on the reaction was first investigated, divalent CuCl₂ was found to give the best result (53% yield, Entry 2), and CuI gave no product (Entry 3). A higher ratio of CuCl₂ to **1a** is preferable to afford a better yield [25% (5 mol%), 30% (10 mol%), and 53% (15 mol%); Entries 2, 6 and 7]. Secondly, a survey of reaction media showed that non-polar solvents (benzene, *o*-xylene, *m*-xylene and xylene, Entries 8, 9, 11, and 13) generally provided better yields than weak or strong polar solvents [tetrahydrofuran (THF), acetonitrile, *N,N*-dimethylformamide (DMF), and dimethyl sulfoxide (DMSO), Entries 10, and 14~16). Finally, a series of experiments were conducted to reveal the role of reaction time (Entries 17 and 18) and temperature (Entries 2 and

13) on the reaction. Generally, higher yields were obtained with the reaction time and temperature increasing. In general, CuCl₂ (15 mol%) was found to be the best catalyst for decarboxylative cross-coupling reaction of carboxylic acids and carbamoyl chlorides in benzene at 120 °C for 48 h.

Table 1 Optimization of Cu-catalyzed decarboxylative coupling^a

Entry	Cu	Solvent	Time/h	Yield ^b /%
1 ^c	Cu(OAc) ₂	Benzene	24	32
2 ^c	CuCl ₂	Benzene	24	53
3 ^c	CuI/CuBr/CuCl	Benzene	24	0
4 ^c	CuO	Benzene	24	0
5 ^c	Cu(OTf) ₂	Benzene	24	0
6 ^{c,d}	CuCl ₂	Benzene	24	25
7 ^{c,e}	CuCl ₂	Benzene	24	30
8	CuCl ₂	<i>o</i> -Xylene	24	38
9	CuCl ₂	<i>m</i> -Xylene	24	11
10	CuCl ₂	THF	24	Trace
11	CuCl ₂	Xylene	24	60
12	CuCl ₂	Toluene	24/48	Trace
13	CuCl ₂	Benzene	24	69
14	CuCl ₂	CH ₃ CN	24	Trace
15	CuCl ₂	DMF	24	0
16	CuCl ₂	DMSO	24	Trace
17	CuCl ₂	Benzene	36	79
18	CuCl ₂	Benzene	48	81

^a Conditions: **1a** (0.2 mmol), **2a** (0.4 mmol), catalyst (15 mol%), and solvent (1.0 mL), 120 °C, 24 h. ^b Isolated yield. ^c 80 °C. ^d 5 mol% of catalyst. ^e 10 mol% of catalyst.

With the optimized reaction conditions in hand, the scope and limitations of this decarboxylative C—C coupling reaction was then explored, a variety of diversely substituted amides **3** were obtained (Tables 2 and 3). Firstly, the generality of carboxylic acids was investigated in the process (Table 2). It was found that aliphatic carboxylic acids (**2a**~**2c**), 2-arylacetic acids (**2d**~**2j**), 3-phenylpropionic acid (**2k**), benzoic acids (**2l**~**2n**) and heteroaromatic acid (**2o**) could be converted into the amides **3** in moderate to good yields (38%~81%). However, when isonicotinic acid **2p** was used, the reaction could not proceed to give the corresponding product **3ap**. For aliphatic carboxylic acids, the length of the carbon chain (from C3 to C14, **3ab** and **3ac**) seemed not have any effect on the yield. The standard conditions were compatible with various groups such as methyl, trifluoromethyl, fluorine, chlorine, and nitro functionality at the *ortho*-, *meta*-, and *para*-position of the aromatic ring. In general, both the *para*- and *meta*-substituted aromatic carboxylic acids afforded a superior yield than *ortho*-substituted analogues

Table 2 Variation of the carboxylic acids^{a,b}

^a Reaction conditions: **1a** (0.2 mmol), carboxylic acid **2** (0.4 mmol), 15 mol% of CuCl₂, benzene (1.0 mL), 120 °C, 48 h. ^b Isolated yield.

 Table 3 Variation of the arylcarbamoyl chlorides^{a,b}

^a Reaction conditions: arylcarbamoyl chlorides **1** (0.2 mmol), **2d** (0.4 mmol), 15 mol% of CuCl₂, benzene (1.0 mL), 120 °C, 48 h. ^b Isolated yield. ^c Acetic acid was used.

(for example, **3af**, 38% yield) as the result of the steric hindrance of substituents.

Secondly, the scope of arylcarbamoyl chlorides was then explored using *p*-tolyl acetic acid **2d** as the coupling partner (Table 3). A variety of substituents on the aryl moiety were applicable, affording the corresponding products **3bd**~**3md** in 35%~89% yields (Table 3). Both electron-donating (Me, OMe) and -withdrawing groups (CF₃, Cl, and CN) were tolerant under the reaction conditions. The electronic nature of aniline skeleton seemed to affect the efficiency to some extent, arylcarbamoyl chlorides con-

taining electron-donating groups underwent the reaction smoothly and afforded good yields (**3bd**~**3ed**, 50%~89%). *N*-Methyl arylcarbamoyl chlorides containing groups at the *para*-position of aniline (**1d**, **1g**) were generally more reactive than those bearing substituents at the *ortho*- or *meta*-position (**1b**, **1c**, and **1f**) and provided higher yields. In addition, β -naphthylcarbamoyl chloride could also be transformed into the desired products **3jd** in 75% yield. Phenylcarbamoyl chlorides **1k**~**1n** were finally investigated to elucidate the influence of different *N*-substituted groups (methyl, benzyl, cyclohexyl, and

phenyl) on the reaction. *N*-Aliphatic substituted substrates generally provided high yields (68%~85%), however, the introduction of the aromatic substituent on the N atom (for example, diphenylcarbamoyl chloride **1m**) decreased reactivity and the product **3md** was obtained in 43% yield. Two alkyl substituted carbamoyl chloride has been examined, for example, the decarboxylative C—C coupling reaction of morpholine-4-carbonyl chloride and benzoic acid gave the corresponding product **3ol** in 76% yield.

To gain insight into the mechanism of the reaction, 2,2,6,6-tetramethylpiperidinoxy (TEMPO) as a radical probe was added to the reaction of morpholine-4-carbonyl chloride and benzoic acid, the decarboxylation step was not blocked and gave the product **3ol** in 72% yield. On the basis of the above results, a plausible pathway was proposed for this Cu(II)-catalyzed decarboxylative C—C coupling reaction. Scheme 2 shows a simplified sequence of events beginning with the active Cu(II) catalyst. The extrusion of HCl took place between carboxylic acid **2** and CuCl₂ to give copper(II) carboxylate **4**, which would undergo the decarboxylative reaction to afford organo-copper species **5** with nucleophilic reactivity via the extrusion of carbon dioxide. The carbon dioxide and HCl generated from the above process were detected by headspace gas chromatography (HS-GC). The subsequent step involved the coordination of arylcarbamoyl chloride **1** to the organo-Cu(II) center to form **6**, which was followed by a nucleophilic addition of organo-copper species across the C=O bond to provide an intermediate complex **7**. β -Chlorine elimination afforded the amide **3** and regenerated the Cu(II) catalyst.

Scheme 2 Proposed catalytic pathway

3 Conclusions

In summary, a ligand-free Cu(II)-catalyzed decarboxylation reaction that can use directly different types of carboxylic acids has been developed. Under the mild condition, a broad range of functional groups is compatible with this C—C coupling reaction, and diverse amides can be

obtained in good to high yields.

4 Experimental section

4.1 General experimental methods

Melting points were determined on a melting point apparatus in open capillaries and uncorrected. Infrared spectra of samples were recorded from 4000 to 500 cm⁻¹ in ATR (attenuated total reflectance) mode using an FT-IR instrument. ¹H NMR spectra were recorded on a 500 MHz spectrometer, and ¹³C NMR spectra were recorded at 126 MHz. Unless otherwise stated, deuteriochloroform (CDCl₃) was used as a solvent. Chemical shifts (δ) are given relative to tetramethylsilane (TMS, δ_{H} 0) or CHCl₃ (δ_{C} 77.16). High-resolution mass spectra were recorded on a BIO TOF Q mass spectrometer equipped with an electrospray ion source (ESI), and operated in the positive mode.

Chemicals were all purchased from commercial supplies and used without further purification unless otherwise stated. Solvents were dried and purified according to the standard procedures before use. Reactions were monitored by analytical thin-layer chromatography (TLC). All reactions were conducted in dried glassware. Purification of reaction products was done by flash chromatography with 230~400 mesh silica gel. All the arylcarbamoyl chloride substrates were prepared according to literature methods.^[20]

4.2 General procedure for Copper-catalyzed decarboxylative coupling

A 10 mL vial equipped with a magnetic stirring bar was charged with arylcarbamoyl chloride **1** (0.2 mmol, 1.0 equiv.), carboxylic acid **2** (0.4 mmol, 2.0 equiv.), and then CuCl₂ (0.03 mmol, 4.0 mg) were added. Finally, benzene (1.0 mL) was added to the mixture via syringe at room temperature under air. The tube was sealed and put into a preheated oil bath at 120 °C for 48 h. The mixture was cooled to room temperature, quenched with water (5 mL), and diluted with ethyl acetate (5 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (5 mL $\times$ 2). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated in vacuo. The crude product was then purified by flash chromatography on silica gel (H) with ethyl acetate/petroleum ether (*V* : *V* = 1 : 3) as eluent. *N*-Ethyl-*N*-phenylacetamide (**3aa**): Yield 81% (26.1 mg), yellow solid. m.p. 55~56 °C (lit.^[21] 55 °C); ¹H NMR (500 MHz, CDCl₃) δ : 7.40 (t, *J*=7.8 Hz, 2H), 7.33 (t, *J*=7.4 Hz, 1H), 7.16~7.13 (m, 2H), 3.73 (q, *J*=7.2 Hz, 2H), 1.80 (s, 3H), 1.09 (t, *J*=7.1 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 169.9, 142.9, 129.7, 128.2, 127.8, 43.8, 22.8, 13.1; IR (KBr) ν : 2980, 1642, 1494, 1401, 1297, 768, 709 cm⁻¹.

N-Ethyl-*N*-phenylpropionamide (**3ab**):^[22] Yield 68% (24.1 mg), yellow liquid. ¹H NMR (500 MHz, CDCl₃) δ : 7.40 (t, *J*=7.5 Hz, 2H), 7.33 (t, *J*=7.2 Hz, 1H), 7.16~7.12 (m, 2H), 3.74 (q, *J*=7.1 Hz, 2H), 2.01 (q, *J*=7.5 Hz, 2H), 1.09 (t, *J*=7.1 Hz, 3H), 1.02 (t, *J*=7.3 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 172.4, 141.5, 128.6, 127.4, 126.8,

42.9, 26.9, 12.1, 8.6; IR (KBr) ν : 2970, 1658, 1496, 1404, 1261, 702 cm^{-1} .

N-Ethyl-*N*-phenyltetradecanamide (**3ac**): Yield 68% (45.0 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.39 (t, $J=7.5$ Hz, 2H), 7.32 (t, $J=7.5$ Hz, 1H), 7.12 (d, $J=7.7$ Hz, 2H), 3.72 (q, $J=7.2$ Hz, 2H), 1.98 (t, $J=7.6$ Hz, 2H), 1.54~1.51 (m, 2H), 1.30~1.11 (m, 20H), 1.08 (t, $J=7.3$ Hz, 3H), 0.85 (t, $J=6.8$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 171.8, 141.6, 128.6, 127.4, 126.8, 42.9, 33.5, 30.9, 28.7, 28.6, 28.5, 28.4, 28.34, 28.30, 28.25, 24.5, 21.7, 13.1, 12.1; IR (KBr) ν : 2925, 1662, 1496, 1261, 768 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{22}\text{H}_{37}\text{NONa}$ [$\text{M} + \text{Na}$] $^+$ 354.27674, found 354.27679.

N-Ethyl-*N*-phenyl-2-(*p*-tolyl)acetamide (**3ad**):^[23] Yield 73% (36.9 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.41~7.34 (m, 3H), 7.09 (d, $J=7.0$ Hz, 2H), 7.04 (d, $J=7.6$ Hz, 2H), 6.94 (d, $J=7.7$ Hz, 2H), 3.75 (q, $J=7.2$ Hz, 2H), 3.36 (s, 2H), 2.30 (s, 3H), 1.10 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.5, 141.3, 135.0, 131.4, 128.5, 127.9, 127.9, 127.7, 126.9, 43.2, 39.8, 20.0, 12.0; IR (KBr) ν : 3483, 2928, 1660, 1494, 1130, 769 cm^{-1} .

N-Ethyl-*N*-phenyl-2-(*m*-tolyl)acetamide (**3ae**): Yield 66% (33.4 mg), yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.44~7.32 (m, 3H), 7.19~7.01 (m, 6H), 3.78 (q, $J=7.2$ Hz, 2H), 3.40 (s, 2H), 2.02 (s, 3H), 1.14 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.4, 141.3, 135.4, 133.4, 129.0, 128.6, 128.4, 127.5, 127.0, 125.7, 124.8, 43.3, 38.2, 18.4, 12.0; IR (KBr) ν : 3490, 2971, 1660, 1494, 1131, 746 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{19}\text{NONa}$ [$\text{M} + \text{Na}$] $^+$ 276.13589, found 276.13538.

N-Ethyl-*N*-phenyl-2-(*o*-tolyl)acetamide (**3af**): Yield 38% (19.3 mg), light red liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.41~7.34 (m, 3H), 7.14~7.04 (m, 6H), 3.78 (q, $J=7.2$ Hz, 2H), 3.38 (s, 2H), 2.02 (s, 3H), 1.14 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.4, 141.3, 135.4, 133.4, 129.0, 128.6, 128.4, 127.5, 127.0, 125.7, 124.9, 43.3, 38.2, 18.4, 12.1; IR (KBr) ν : 3482, 2970, 1659, 1494, 1131, 746 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{19}\text{NONa}$ [$\text{M} + \text{Na}$] $^+$ 276.13589, found 276.13550.

N-Ethyl-*N*-phenyl-2-(4-(trifluoromethyl)phenyl)acetamide (**3ag**): Yield 46% (28.2 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.48 (d, $J=8.1$ Hz, 2H), 7.44~7.37 (m, 3H), 7.17 (d, $J=7.9$ Hz, 2H), 7.12~7.07 (m, 2H), 3.76 (q, $J=7.1$ Hz, 2H), 3.45 (s, 2H), 1.11 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 168.4, 141.0, 138.6, 128.7, 128.4, 127.8 (128.2, 128.0, 127.7, 127.5, q, $^2J_{\text{C-F}}=32$ Hz), 127.6, 127.2, 124.1 (124.17, 124.14, 124.11, 124.08, q, $^3J_{\text{C-F}}=4$ Hz), 123.2 (126.5, 124.3, 122.1, 120.0, q, $^1J_{\text{C-F}}=272$ Hz), 43.4, 40.1, 11.9; IR (KBr) ν : 2972, 1659, 1495, 1326, 1126, 1067, 701 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{16}\text{F}_3\text{NONa}$ [$\text{M} + \text{Na}$] $^+$ 330.10762, found 330.10773.

N-Ethyl-2-(4-nitrophenyl)-*N*-phenylacetamide (**3ah**):^[23] Yield 70% (39.8 mg), red oil. IR (KBr) ν : 3490, 2974, 1659, 1346, 856, 732, 560 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 8.15~8.02 (m, 2H), 7.48~7.34 (m, 3H), 7.22

(d, $J=8.6$ Hz, 2H), 7.10 (d, $J=7.6$ Hz, 2H), 3.76 (q, $J=7.1$ Hz, 2H), 3.49 (s, 2H), 1.11 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 167.8, 145.8, 142.1, 140.8, 129.1, 128.9, 127.5, 127.4, 122.4, 43.4, 40.2, 11.9.

N-Ethyl-2-(3-fluorophenyl)-*N*-phenylacetamide (**3ai**): Yield 48% (24.7 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.44~7.33 (m, 3H), 7.17 (q, $J=7.8$ Hz, 1H), 7.08 (d, $J=6.9$ Hz, 2H), 6.87 (td, $J=8.5$, 2.1 Hz, 1H), 6.81 (d, $J=7.6$ Hz, 1H), 6.77 (d, $J=9.9$ Hz, 1H), 3.75 (q, $J=7.1$ Hz, 2H), 3.39 (s, 2H), 1.11 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 168.7, 161.6 (162.6, 160.7, d, $^1J_{\text{C-F}}=246$ Hz), 141.0, 136.9 (136.91, 136.85, d, $^3J_{\text{C-F}}=8$ Hz), 128.6, 128.5, 127.6, 127.1, 123.7 (123.72, 123.70, d, $^4J_{\text{C-F}}=3$ Hz), 115.0 (115.1, 114.9, d, $^2J_{\text{C-F}}=21$ Hz), 112.4 (112.5, 112.3, d, $^2J_{\text{C-F}}=20$ Hz), 43.3, 40.0 (d, $^4J_{\text{C-F}}=1$ Hz), 12.0. IR (KBr) ν : 2973, 1657, 1593, 1267, 1132, 769, 702 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{16}\text{H}_{16}\text{FNONa}$ [$\text{M} + \text{Na}$] $^+$ 280.11081, found 280.11093.

N-Ethyl-2-(naphthalen-2-yl)-*N*-phenylacetamide (**3aj**): Yield 66% (38.2 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.72~7.66 (m, 1H), 7.62 (t, $J=7.9$ Hz, 2H), 7.38~7.31 (m, 3H), 7.30~7.22 (m, 3H), 7.20~7.07 (m, 1H), 7.02~6.94 (m, 2H), 3.68 (q, $J=7.1$ Hz, 2H), 3.49 (s, 2H), 1.03 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.3, 141.2, 132.4, 132.0, 131.2, 128.5, 127.7, 127.0, 126.8, 126.6, 126.54, 126.52, 126.4, 124.8, 124.4, 43.3, 40.6, 12.0; IR (KBr) ν : 3482, 3055, 2971, 2932, 1654, 1494, 1129, 701, 561 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{20}\text{H}_{19}\text{NONa}$ [$\text{M} + \text{Na}$] $^+$ 312.13589, found 312.13593.

N-Ethyl-*N*,3-diphenylpropionamide (**3ak**): Yield 48% (23.9 mg), white solid, m.p. 100~101 $^{\circ}\text{C}$ (lit.^[24] 100~101 $^{\circ}\text{C}$); ^1H NMR (500 MHz, CDCl_3) δ : 7.39~7.28 (m, 3H), 7.26~7.17 (m, 3H), 7.15~7.08 (m, 2H), 7.00 (d, $J=7.4$ Hz, 2H), 3.78 (q, $J=7.0$ Hz, 2H), 1.09 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 152.9, 140.7, 131.4, 128.8, 128.1, 127.7, 127.3, 127.1, 119.5, 89.7, 81.8, 42.4, 12.0; IR (KBr) ν : 2964, 2213, 1632, 1492, 1347, 1310, 1261, 1025, 759, 700, 529 cm^{-1} .

N-Ethyl-*N*-phenylbenzamide (**3al**):^[25] Yield 73% (20.3 mg), black liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.20 (t, $J=7.7$ Hz, 2H), 7.17~7.11 (m, 3H), 7.10~7.03 (m, 3H), 6.95 (d, $J=7.6$ Hz, 2H), 3.91 (q, $J=7.2$ Hz, 2H), 1.15 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.3, 142.1, 135.2, 128.4, 128.1, 127.6, 126.9, 126.7, 125.6, 44.4, 11.9; IR (KBr) ν : 3483, 2973, 1638, 1493, 1118, 699, 585 cm^{-1} .

4-Chloro-*N*-ethyl-*N*-phenylbenzamide (**3am**):^[23] Yield 73% (37.8 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.20~7.13 (m, 4H), 7.13~7.08 (m, 1H), 7.05 (d, $J=8.3$ Hz, 2H), 6.94 (d, $J=8.0$ Hz, 2H), 3.90 (q, $J=7.1$ Hz, 2H), 1.14 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 168.0, 142.0, 134.5, 133.7, 129.2, 128.3, 126.93, 126.85, 125.9, 44.5, 11.8; IR (KBr) ν : 2973, 1647, 1493, 1091, 754, 583 cm^{-1} .

N-Ethyl-4-nitro-*N*-phenylbenzamide (**3an**): Yield 54% (29.2 mg), light yellow solid. m.p. 117~119 $^{\circ}\text{C}$ (lit.^[23] 119~120 $^{\circ}\text{C}$); ^1H NMR (500 MHz, CDCl_3) δ : 7.99 (d,

$J=8.1$ Hz, 2H), 7.42 (d, $J=7.9$ Hz, 2H), 7.28~7.13 (m, 3H), 7.01 (d, $J=7.6$ Hz, 2H), 3.98 (q, $J=7.0$ Hz, 2H), 1.22 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 167.9, 147.9, 142.6, 142.1, 129.52, 129.49, 128.0, 127.5, 123.0, 45.4, 12.8; IR (KBr) ν : 2976, 1651, 1347, 846, 763, 699, 580 cm^{-1} .

N-Ethyl-*N*-phenylfuran-2-carboxamide (**3ao**): Yield 76% (32.7 mg), white solid. m.p. $174\sim 177\text{ }^\circ\text{C}$ (lit.^[26] $175\sim 176\text{ }^\circ\text{C}$); ^1H NMR (500 MHz, CDCl_3) δ : 7.44~7.32 (m, 3H), 7.30 (d, $J=1.0$ Hz, 1H), 7.21~7.14 (m, 2H), 6.16 (dd, $J=3.5, 1.7$ Hz, 1H), 5.72 (d, $J=2.7$ Hz, 1H), 3.90 (q, $J=7.1$ Hz, 2H), 1.20 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 158.8, 147.2, 144.2, 142.4, 129.5, 128.5, 128.0, 116.1, 110.9, 45.4, 12.8; IR (KBr) ν : 3127, 3107, 2984, 1628, 1469, 1412, 1309, 1120, 1028, 704, 581 cm^{-1} .

N-Methyl-*N*-(*o*-tolyl)-2-(*p*-tolyl)acetamide (**3bd**): Yield 64% (32.4 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.23~7.13 (m, 3H), 6.98 (d, $J=7.6$ Hz, 1H), 6.94 (d, $J=7.8$ Hz, 2H), 6.81 (d, $J=7.9$ Hz, 2H), 3.23 (s, 2H), 3.09 (s, 3H), 2.20 (s, 3H), 1.99 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.3, 141.5, 135.1, 134.8, 130.9, 130.4, 128.0, 128.0, 127.4, 126.3, 39.4, 35.2, 20.0, 16.2; IR (KBr) ν : 3482, 2923, 1657, 1493, 1376, 1103, 727, 569 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{20}\text{NO}$ $[\text{M} + \text{H}]^+$ 254.1539, found 254.1532.

N-Methyl-*N*-(*m*-tolyl)-2-(*p*-tolyl)acetamide (**3cd**): Yield 50% (25.3 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.19 (t, $J=7.7$ Hz, 1H), 7.07 (d, $J=7.6$ Hz, 1H), 6.96 (d, $J=7.7$ Hz, 2H), 6.91~6.82 (m, 3H), 6.80 (s, 1H), 3.32 (s, 2H), 3.16 (s, 3H), 2.26 (s, 3H), 2.21 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.2, 142.9, 138.6, 135.0, 131.5, 128.4, 127.9, 127.6, 127.3, 123.5, 39.5, 36.5, 20.2, 20.0; IR (KBr) ν : 3483, 2924, 1659, 1517, 1047, 794, 706 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{20}\text{NO}$ $[\text{M} + \text{H}]^+$ 254.1539, found 254.1534.

N-Methyl-*N*,2-di-*p*-tolylacetamide (**3dd**): Yield 81% (41.0 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.11 (d, $J=8.0$ Hz, 2H), 6.96 (d, $J=7.8$ Hz, 2H), 6.93 (d, $J=7.8$ Hz, 2H), 6.88 (d, $J=7.9$ Hz, 2H), 3.32 (s, 2H), 3.16 (s, 3H), 2.30 (s, 3H), 2.21 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.3, 140.5, 136.8, 135.0, 131.4, 129.2, 128.1, 127.9, 126.3, 124.5, 39.2, 36.6, 20.1, 20.0; IR (KBr) ν : 3483, 2924, 1659, 1514, 1434, 1377, 1121, 826, 780, 557 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{20}\text{NO}$ $[\text{M} + \text{H}]^+$ 254.1539, found 254.1533.

N-(4-Methoxyphenyl)-*N*-methyl-2-(*p*-tolyl)acetamide (**3ed**): Yield 89% (47.9 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.08~7.00 (m, 4H), 6.96 (d, $J=7.7$ Hz, 2H), 6.90 (d, $J=8.8$ Hz, 2H), 3.83 (s, 3H), 3.40 (s, 2H), 3.23 (s, 3H), 2.29 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.6, 157.9, 135.8, 135.0, 131.4, 128.0, 127.9, 127.7, 113.7, 54.5, 39.3, 36.7, 20.0; IR (KBr) ν : 3483, 2933, 2837, 1651, 1513, 1248, 916, 730, 563 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{20}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 270.1489, found 270.1481.

N-(2-Chlorophenyl)-*N*-methyl-2-(*p*-tolyl)acetamide

(**3fd**): Yield 35% (19.1 mg), light red solid. m.p. $56\sim 58\text{ }^\circ\text{C}$ (lit.^[27] $66\sim 67\text{ }^\circ\text{C}$); ^1H NMR (500 MHz, CDCl_3) δ : 7.43 (dd, $J=7.9, 1.5$ Hz, 1H), 7.30~7.17 (m, 2H), 7.08 (dd, $J=7.6, 1.7$ Hz, 1H), 6.96 (d, $J=7.9$ Hz, 2H), 6.85 (d, $J=8.0$ Hz, 2H), 3.25 (dd, $J=50.2, 14.9$ Hz, 2H), 3.13 (s, 3H), 2.22 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.2, 140.1, 135.1, 132.2, 130.8, 129.6, 129.3, 128.6, 128.0, 127.9, 127.1, 39.5, 35.0, 20.0; IR (KBr) ν : 2918, 1670, 1480, 1370, 1125, 1063, 771, 568, 466 cm^{-1} .

N-(4-Chlorophenyl)-*N*-methyl-2-(*p*-tolyl)acetamide (**3gd**): Yield 63% (34.4 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.36 (d, $J=8.5$ Hz, 2H), 7.05 (d, $J=7.8$ Hz, 4H), 6.94 (d, $J=7.2$ Hz, 2H), 3.41 (s, 2H), 3.24 (s, 3H), 2.30 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 171.1, 142.6, 136.3, 133.7, 132.0, 129.9, 129.1, 129.0, 128.8, 40.6, 37.6, 21.1; IR (KBr) ν : 2924, 1660, 1490, 1374, 1120, 1014, 722, 554 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{16}\text{H}_{17}\text{ClNO}$ $[\text{M} + \text{H}]^+$ 274.0993, found 274.0991.

N-Methyl-2-(*p*-tolyl)-*N*-(4-(trifluoromethyl)phenyl)acetamide (**3hd**): Yield 57% (35.0 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.57 (d, $J=8.3$ Hz, 2H), 7.17 (d, $J=6.4$ Hz, 2H), 6.97 (d, $J=7.4$ Hz, 2H), 6.86 (br, 2H), 3.37 (s, 2H), 3.21 (s, 3H), 2.22 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.9, 146.1, 135.4, 130.7, 128.2, 127.7, 127.0, 125.7, 122.7, 125.9, 123.8, 121.6, 119.5, q, $J_{\text{C-F}}=272\text{ Hz}$, 39.7, 36.6, 20.0; IR (KBr) ν : 2925, 1664, 1326, 1124, 852, 612 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{17}\text{F}_3\text{NO}$ $[\text{M} + \text{H}]^+$ 308.1257, found 308.1255.

N-(4-Cyanophenyl)-*N*-methyl-2-(*p*-tolyl)acetamide (**3id**): Yield 64% (33.8 mg), light yellow liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.60 (d, $J=8.4$ Hz, 2H), 7.20~7.14 (m, 2H), 6.98 (d, $J=7.7$ Hz, 2H), 6.86 (d, $J=6.3$ Hz, 2H), 3.42 (s, 2H), 3.21 (s, 3H), 2.23 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.8, 146.9, 135.5, 132.5, 130.4, 128.3, 128.2, 127.6, 127.1, 117.0, 39.9, 36.6, 20.0; IR (KBr) ν : 3507, 2924, 2228, 1667, 1505, 1118, 851, 576 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{17}\text{H}_{16}\text{N}_2\text{ONa}$ $[\text{M} + \text{Na}]^+$ 287.11548, found 287.11569.

N-Methyl-*N*-(naphthalen-2-yl)-2-(*p*-tolyl)acetamide (**3jd**): Yield 75% (43.4 mg), light yellow solid, m.p. $65\sim 68\text{ }^\circ\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 7.78 (d, $J=8.6$ Hz, 2H), 7.71~7.63 (m, 1H), 7.52~7.36 (m, 3H), 7.14 (d, $J=6.3$ Hz, 1H), 6.93 (d, $J=7.3$ Hz, 2H), 6.85 (d, $J=6.8$ Hz, 2H), 3.36 (s, 2H), 3.25 (s, 3H), 2.20 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.3, 140.3, 135.1, 132.5, 131.4, 131.3, 128.8, 128.0, 127.9, 126.8, 126.7, 125.8, 125.7, 125.1, 124.4, 39.6, 36.6, 20.0; IR (KBr) ν : 3053, 2917, 1651, 1504, 1381, 1107, 863, 781, 581, 504 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{20}\text{H}_{19}\text{NONa}$ $[\text{M} + \text{Na}]^+$ 312.13589, found 312.13574.

N-Benzyl-*N*-phenyl-2-(*p*-tolyl)acetamide (**3kd**): Yield 85% (53.6 mg), light yellow solid. m.p. $71\sim 74\text{ }^\circ\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 7.24~7.18 (m, 3H), 7.18~7.13 (m, 3H), 7.11~7.07 (m, 2H), 6.96 (d, $J=7.7$ Hz, 2H), 6.87 (d, $J=7.7$ Hz, 2H), 6.85~6.79 (m, 2H), 4.80 (s, 2H), 3.33 (s, 2H), 2.22 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.0, 141.2, 136.4, 135.1, 131.3, 128.4, 128.0,

127.93, 127.85, 127.7, 127.3, 127.0, 126.3, 52.2, 39.7, 20.0; IR (KBr) ν : 2980, 1642, 1596, 1494, 1401, 1297, 768, 709 cm^{-1} . HRMS-ESI calcd for $\text{C}_{22}\text{H}_{21}\text{NONa}$ [$\text{M} + \text{Na}$] $^{+}$ 338.15154, found 338.15109.

N-Cyclohexyl-*N*-phenyl-2-(*p*-tolyl)acetamide (**3ld**): Yield 80% (49.1 mg), yellow solid. m.p. 68~70 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 7.41~7.30 (m, 3H), 7.07~6.98 (m, 4H), 6.91 (d, $J=7.9$ Hz, 2H), 4.61 (tt, $J=12.1$, 3.5 Hz, 1H), 3.26 (s, 2H), 2.29 (s, 3H), 1.82 (d, $J=11.5$ Hz, 2H), 1.70 (d, $J=13.6$ Hz, 2H), 1.55 (d, $J=13.1$ Hz, 1H), 1.38 (qt, $J=13.2$, 3.3 Hz, 2H), 1.01 (qd, $J=12.4$, 3.5 Hz, 2H), 0.89 (qt, $J=13.1$, 3.7 Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.5, 139.1, 135.9, 132.6, 130.7, 129.0, 128.9, 128.2, 54.3, 41.5, 31.6, 25.8, 25.4, 21.1; IR (KBr) ν : 2926, 2853, 1655, 1592, 1492, 1258, 1067, 767, 713, 486 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{21}\text{H}_{26}\text{NO}$ [$\text{M} + \text{H}$] $^{+}$ 308.2009, found 308.2001.

N,N-Diphenyl-2-(*p*-tolyl)acetamide (**3md**): Yield 43% (25.9 mg), yellow solid. m.p. 72~76 $^{\circ}\text{C}$ (m.p. $^{[28]}$ 74~77 $^{\circ}\text{C}$); ^1H NMR (500 MHz, CDCl_3) δ : 7.34 (br, 5H), 7.25~7.12 (m, 5H), 7.07 (d, $J=7.9$ Hz, 2H), 7.01 (d, $J=7.9$ Hz, 2H), 3.61 (s, 2H), 2.32 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 170.2, 141.8, 135.2, 131.0, 128.4, 128.0, 127.9, 126.8, 125.3, 40.7, 20.0; IR (KBr) ν : 2935, 1671, 1589, 1352, 1308, 1279, 1148, 805, 759, 703, 581, 490 cm^{-1} .

N-Methyl-*N*-phenylacetamide (**3na**): Yield 68% (20.3 mg), light red solid. m.p. 100~102 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 7.41 (t, $J=7.5$ Hz, 2H), 7.32 (t, $J=7.2$ Hz, 1H), 7.18 (d, $J=7.6$ Hz, 2H), 3.25 (s, 3H), 1.86 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.5, 143.6, 128.7, 126.7, 126.1, 36.1, 21.4; IR (KBr) ν : 3046, 1656, 1598, 1496, 1297, 707, 559 cm^{-1} ; HRMS-ESI calcd for $\text{C}_9\text{H}_{11}\text{NONa}$ [$\text{M} + \text{Na}$] $^{+}$ 172.07329, found 172.07333.

Morpholino(phenyl)methanone (**3ol**): Yield 76% (29.0 mg), white solid, m.p. 73~74 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 7.40~7.28 (m, 5H), 4.06~3.10 (m, 8H); ^{13}C NMR (126 MHz, CDCl_3) δ : 169.41, 134.30, 128.86, 127.54, 126.07, 65.87; IR (KBr) ν : 3414, 1621, 1436, 1274, 1109, 1012, 731, 709 cm^{-1} ; HRMS-ESI calcd for $\text{C}_{11}\text{H}_{13}\text{NO}_2\text{Na}$ [$\text{M} + \text{Na}$] $^{+}$ 214.08385, found 214.08389.

Supporting Information ^1H NMR and ^{13}C NMR spectra of **3aa**~**3ao**, **3bd**~**3md**, **3na**, **3ol** and HRMS spectra of **3ac**, **3ae**~**3ag**, **3ai**, **3aj**, **3bd**~**3ed**, **3gd**~**3ld**, **3na**, **3ol** are available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- (a) Valeur, E.; Bradley, M. *Chem. Soc. Rev.* **2009**, 38, 606.
(b) de Figueiredo, R. M.; Suppo, J.-S.; Campagne, J.-M. *Chem. Rev.* **2016**, 116, 12029.
- Roy, S.; Roy, S.; Gribble, G. W. *Tetrahedron* **2012**, 68, 9867.
- Organotin reagents, see:
(a) Balas, L.; Jousseau, B.; Shin, H.; Verlhac, J.-B.; Wallian, F. *Organometallics* **1991**, 10, 366.
(b) Jousseau, B.; Kwon, H.; Verlhac, J.-B.; Denat, F.; Dubac, J. *Synlett* **1993**, 117.
- (c) Murakami, M.; Hoshino, Y.; Ito, H.; Ito, Y. *Chem. Lett.* **1998**, 163.
(d) Hu, W.; Zheng, J.; Li, M.; Wu, W.; Liu, H.; Jiang, H. *Chin. J. Chem.* **2018**, 36, 712.
- Lemoucheux, L.; Rouden, J.; Lasne, M.-C. *Tetrahedron Lett.* **2000**, 41, 9997.
- Rieke, R. D.; Kim, S.-H. *Tetrahedron Lett.* **2012**, 53, 3478.
- Lysén, M.; Kelleher, S.; Begtrup, M.; Kristensen, J. L. *J. Org. Chem.* **2005**, 70, 5342.
- Duan, Y.-Z.; Deng, M.-Z. *Synlett* **2005**, 355.
- Yasui, Y.; Tsuchida, S.; Miyabe, H.; Takemoto, Y. *J. Org. Chem.* **2007**, 72, 5898.
- Krishnamoorthy, R.; Lam, S. Q.; Manley, C. M.; Herr, R. J. *J. Org. Chem.* **2010**, 75, 1251.
- Kochi, T.; Urano, S.; Seki, H.; Mizushima, E.; Sato, M.; Kakiuchi, F. *J. Am. Chem. Soc.* **2009**, 131, 2792.
- (a) Matsuzono, M.; Fukuda, T.; Iwao, M. *Tetrahedron Lett.* **2001**, 42, 7621.
(b) Chao, W.-R.; Yean, D.; Amin, K.; Green, C.; Jong, L. *J. Med. Chem.* **2007**, 50, 3412.
- Lemoucheux, L.; Seitz, T.; Rouden, J.; Lasne, M.-C. *Org. Lett.* **2004**, 6, 3703.
- Selected reviews, see:
(a) Baudoin, O. *Angew. Chem., Int. Ed.* **2007**, 46, 1373.
(b) Gooßen, L. J.; Rodriguez, N.; Gooßen, K. *Angew. Chem., Int. Ed.* **2008**, 47, 3100.
(c) Rodriguez, N.; Gooßen, L. J. *Chem. Soc. Rev.* **2011**, 40, 5030.
(d) Dzik, W. I.; Lange, P. P.; Gooßen, L. J. *Chem. Sci.* **2012**, 3, 2671.
(e) Wei, Y.; Hu, P.; Zhang, M.; Su, W. *Chem. Rev.* **2017**, 117, 8864.
- Selected examples, see:
(a) Myers, A. G.; Tanaka, D.; Mannion, M. R. *J. Am. Chem. Soc.* **2002**, 124, 11250.
(b) Tanaka, D.; Romeril, S. P.; Myers, A. G. *J. Am. Chem. Soc.* **2005**, 127, 10323.
- Gooßen, L. J.; Deng, G.; Levy, L. M. *Science* **2006**, 313, 662.
- (a) Gooßen, L. J.; Linder, C.; Rodriguez, N.; Lange, P. P.; Fromm, A. *Chem. Commun.* **2009**, 7173.
(b) Cornella, J.; Sanchez, C.; Banawa, D.; Larrosa, I. *Chem. Commun.* **2009**, 7176.
- (a) Cornella, J.; Rosillo-Lopez, M.; Larrosa, I. *Adv. Synth. Catal.* **2011**, 353, 1359.
(b) Dupuy, S.; Lazreg, F.; Slawin, A. M. Z.; Cazin, C. S. J.; Nolan, S. P. *Chem. Commun.* **2011**, 47, 5455.
- Sun, Z.-M.; Zhao, P. *Angew. Chem., Int. Ed.* **2009**, 48, 6726.
- Shang, R.; Fu, Y.; Wang, Y.; Xu, Q.; Yu, H.-Z.; Liu, L. *Angew. Chem., Int. Ed.* **2009**, 48, 9350.
- (a) Jiao, J.; Zhang, X.-R.; Chang, N.-H.; Wang, J.; Wei, J.-F.; Shi, X.-Y.; Chen, Z.-G. *J. Org. Chem.* **2011**, 76, 1180.
(b) Iwai, T.; Fujihara, T.; Terao, J.; Tsuji, Y. *J. Am. Chem. Soc.* **2010**, 132, 9602.
- Yamasaki, R.; Morita, K.; Iizumi, H.; Ito, A.; Fukuda, K.; Okamoto, I. *Chem.-Eur. J.* **2019**, 25, 10118.
- Allah, T. N.; ne Savourey, S.; Berthet, J.-C.; Nicolas, E.; Cantat, T. *Angew. Chem., Int. Ed.* **2019**, 58, 10884.
- Song, G.; Sun, G.; Tang, Y.; Mai, W. *J. Chem. Res.* **2013**, 630.
- Zhou, Y.; Zhang, X.; Zhang, Y.; Ruan, L.; Zhang, J.; Zhang-Negrerie, D.; Du, Y. *Org. Lett.* **2017**, 19, 150.
- Zhang, Z.; Liu, Y.-H.; Zhang, X.; Wang, X.-C. *Tetrahedron* **2019**, 75, 2763.
- Bao, Y.-S.; Zhaorigetu, B.; Agula, B.; Baiyin, M.; Jia, M. *J. Org. Chem.* **2014**, 79, 803.
- Ackermann, L.; Vicente, R.; Hofmann, N. *Org. Lett.* **2009**, 11, 4274.
- Das, K. G.; Funke, P. T.; Bose, A. K. *J. Am. Chem. Soc.* **1964**, 86, 3729.

(Cheng, F.)