空气条件下温和高效催化 *N*-吡啶吲哚啉的氧化脱氢

唐 灏 张贝贝 陈卫东 骆钧飞*

(宁波大学材料科学与化学工程学院 浙江宁波 315211)

摘要 报道了空气条件下温和高效的 *N*-羟基邻苯二甲酰亚胺和铜共同催化 *N*-吡啶吲哚啉的氧化脱氢反应。已知的 *N*-吡啶吲哚啉的氧化过程只能在过量强氧化剂 2,3-二氯-5,6-二氰对苯醌存在下实现, 该反应以空气作为单一的氧化剂, 具有绿色、安全、低成本以及操作简便等优点, 为 *N*-吡啶吲哚啉类化合物氧化生成相应吲哚衍生物提供了一条新的绿色途径。

关键词 *N*-羟基邻苯二甲酰亚胺; 脱氢; *N*-吡啶吲哚啉; 空气条件; 氧化反应

A Mild and Efficient Catalytic Aerobic Oxidative Dehydrogenation of *N*-Pyridylindolines

Tang, Hao Zhang, Beibei Chen, Weidong Luo, Junfei*

(School of Materials Science and Chemical Engineering, Ningbo University, Ningbo, Zhejiang 315211)

Abstract An efficient and mild *N*-hydroxyphthalimide and copper co-catalyzed aerobic oxidative dehydrogenation of *N*-pyridylindolines is developed. The use of air as sole oxidant allows a green, safe, low-cost and simple operation. This method provides an alternative to the known method for the oxidation of *N*-pyridylindoline derivatives by the use of excess stoichiometric strong oxidant 2,3-dichloro-5,6-dicyanobenzoquinone.

Keywords *N*-hydroxyphthalimide; dehydrogenation; *N*-pyridylindolines; air condition; oxidation reaction

1 Introduction

Indole skeletons are frequently found in natural products, pharmaceuticals and biologically active compounds,^[1] thus, a variety of methodologies for the synthesis of indoles have been reported.^[2] Among those, the oxidation of indolines is an important methodology to produce the uncommonly substituted indole derivatives.^[3] To date, many methods for the oxidation of indolines to indoles by various oxidants have been discovered, while the use of stoichiometric oxidants resulted in the generation of considerable amounts of toxic by-products.^[4]

The transition-metal-catalyzed dehydrogenation provides an alternative to the traditional oxidation reactions using stoichiometric oxidants.^[5] In this context, the methods for Ru,^[6] Pd,^[7] Au,^[8] Co^[9] and others metals^[10] catalyzed oxidative dehydrogenation of indolines to the corresponding indoles. In addition, the borane-catalyzed dehydrogenation of indoline has also been reported.^[11] More recently, we^[12] have presented a *N*-hydroxyphthalimide-catalyzed aerobic

dehydrogenation of *N*-heterocycles. These methods only produce hydrogen or water as the by-product. However, high reaction temperatures are always required to proceed efficiently. Moreover, a significant disadvantage is that only *N*-free indolines or *N*-alkylated indolines were probed to undergo the dehydrogenation process.

C(7)-Substituted indoles are particularly important as they are ubiquitous in numerous biologically active molecules.^[13] The prevalent method for the synthesis of C(7) decorated indoles is through the transition-metal-catalyzed C—H functionalization of indolines, along with subsequent oxidation.^[14] In this regard, *N*-pyridylindolines or *N*-pyrimidylindolines are frequently used as the starting materials to achieve the C(7) decorated indoles via C—H activation and oxidation steps. However, to our surprise, the oxidation of *N*-pyridylindolines or *N*-pyrimidylindolines still requires the use of a considerable amount of strong oxidant 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ).^[15] Thus, the development of a “mild and green” method for the oxidation of *N*-pyridylindoline with high functional group

* Corresponding author. E-mail: luojunfei@nbu.edu.cn

Received September 27, 2020; revised November 2, 2020; published online November 18, 2020

Project supported by the Natural Science Foundation of Ningbo City (No. 2019A610027).

宁波市自然科学基金(No. 2019A610027)资助项目.

tolerance would still represent a useful contribution to this area.

(a) (Oxidative) dehydrogenation of indolines

Limitation: R is limited in H or alkyl groups

(b) Oxidation of *N*-pyridyl indolines using stoichiometric DDQ

Limitation: generation of considerable amount of toxic by-products

(c) This work: NHPI and Cu co-catalyzed oxidative dehydrogenation of *N*-pyridyl indolines

Mild conditions Cheap catalyst Air as sole oxidant

Scheme 1 Transformations of indolines to indoles

Herein, we report a new *N*-hydroxyphthalimide (NHPI) and copper co-catalyzed oxidative dehydrogenation of *N*-pyridylindolines under mild conditions. The method displays good scope, tolerating both electron-donating and electron-withdrawing substituents on the indoline. The use of air as sole oxidant is green, safe and low-cost.

2 Results and discussion

N-Pyridylindoline (**1a**) was chosen as a model substrate to optimize the reaction conditions. During the experimentation, it was found that the oxidation of **1a** proceeded efficiently under room temperature with 20 mol% NHPI and 5 mol% $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ in MeCN, with 94% of desired *N*-pyridyl indole (**2a**) being obtained under 101 kPa O_2 atmosphere (Table, Entry 1). Other solvents such as toluene, 1,4-dioxane and tetrahydrofuran (THF) were less efficient, and *N,N*-dimethylformamide (DMF) was completely unreactive (Entries 2–5). The amount of NHPI catalyst affected the oxidation process. The indole product **2a** could be obtained in 85% yield by the use of 40 mol% NHPI (Entry 6). However, the reaction completely stopped in the absence of NHPI catalyst and showed much less efficiency without adding $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (Entries 7 and 8). Attempt to oxidize **1a** under air conditions displayed high efficiency as well with 91% yield of **2a** being obtained (Entry 9), on the other hand, only trace of *N*-pyridylindole was observed when conducting the reaction under nitrogen atmosphere (Entry 10). The use of other metal salt such as FeCl_3 dramatically restrained the oxidation (Entry 11). Carrying the reaction out under air conditions was chosen as the optimal reaction

conditions to explore the substrate scope although the oxidation proceeded slightly worse than when under O_2 atmosphere. Notably, this oxidation process can be conducted in an open flask without heating in the summer days.

Table 1 Optimization of reaction conditions^a

Entry	$\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ / mol%	NHPI/ mol%	Solvent	Yield/%
1	5	20	MeCN	94
2	5	20	Toluene	60
3	5	20	1,4-Dioxane	61
4	5	20	THF	40
5	5	20	DMF	0
6	5	40	MeCN	85
7	5	—	MeCN	0
8	—	20	MeCN	30
9 ^b	5	20	MeCN	93
10 ^c	5	20	MeCN	Trace
11 ^d	—	20	MeCN	6

^a Unless otherwise noted, the reactions were performed using 0.2 mmol of **1a** in 3.0 mL of solvent at room temperature under 101 kPa O_2 atmosphere for 12 h. Yields are of pure isolated products; ^b The reaction was performed under air. ^c The reaction was performed under nitrogen. ^d The reaction was performed with 5 mol% of FeCl_3 .

With the optimized conditions in hand, the versatility of this oxidation process was then explored by probing diversely substituted indolines **1** (Table 2). Generally, the oxidation process confided high functional group tolerance. Explicitly, the electron donor groups substituted either on C(4) or C(5) position were well compatible with the reaction, undergoing the oxidation process efficiently to give the corresponding indole derivatives in good to excellent yields (Table 2, **2b**–**2d**, **2j**, **2k**). The substrates with weak electron acceptors halogen atoms, fluoro, chloro and bromo, that can further proceed various useful transformations, were all found to be well tolerated, giving the halogenated indoles in yields above 80% by slightly increasing the reaction temperature to 35 °C (Table 2, **2e**–**2g**, **2k**–**2m**). The indolines with electron-withdrawing ester group, on the other hand, required further heating to 45 °C to achieve the satisfied yields (Table 2, **2h** and **2o**). Importantly, the strong electron-withdrawing nitro group substituted indoline could also be efficiently transformed into the corresponding indole **2i** in 83% yield although the reaction was required to be heated at 100 °C. The typical methyl substituent had been chosen to test the positional effect, and without big difference being observed (Table 2, **2p** and **2q**). Of note is that the congested C(2)- or C(3)-methyl substituted indolines could still go through the oxidation efficiently, affording the cor-

responding indole products **2r** and **2s** in 70% and 83% yields, respectively. The pyrimidyl group had similar coordination ability but more electron negative in comparison with pyridyl group. Satisfyingly, *N*-pyrimidylindoline was tested and delivered *N*-pyrimidylindole in 81% yield by heating the reaction at 45 °C (Table 2, **2t**).

Table 2 Scope of *N*-pyridylindolines oxidation^a

^a Unless otherwise noted, the reactions were performed using 0.2 mmol of **1** at room temperature for 12 h. Yields are of pure isolated products; ^b The reaction was performed at 35 °C; ^c The reaction was performed at 45 °C; ^d The reaction was performed at 100 °C.

To investigate the utility of this reaction on the preparative scale, a gram-scale reaction of indoline **1a** (1.08 g, 5.5 mmol) was carried out under the standard condition in an

open flask at room temperature (Scheme 2). Satisfyingly, the desired product **2a** could be readily obtained in 91% yield.

Scheme 2 Scale-up reaction.

Taking advantage of the myriad methods available for C(7) C—H functionalization of *N*-pyridyl or pyrimidylindolines, the C(7) decorated indole derivatives could be accessed from the corresponding indolines through this aerobic catalytic method by avoiding the use of strong oxidant DDQ. The C(7) amination,^[16] acetoxylation^[17] and benzenesulfonylation^[18] products have been tested and all underwent the oxidation process to furnish the desired indole derivatives in good yields (Scheme 3).

Scheme 3 Oxidation of C(7)-decorated indolines

Control reactions showed that the oxidative dehydrogenation still worked well at elevated temperature in the absence of $\text{Cu(OAc)}_2\cdot\text{H}_2\text{O}$ (Scheme 4, a). It is suggested that the copper might promote the transformation of phthalimide-*N*-oxyl (PINO) from NHPI under mild conditions. Additionally, a radical pathway might be involved as the reaction worked poorly in the presence of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) (Scheme 4, b). On the basis of these results and the previous reports,^[12,19] the reaction may begin with the transformation of NHPI to PINO promoted by copper. A radical cation **I**

and NHPI anion could be generated by the reaction of PINO and *N*-pyridylindoline through single electron transfer (SET), then deprotonation of **I** by NHPI anion gives radical **II** and reproduces NHPI. Molecular oxygen reacts with radical **II** to afford radical **III**, which can be protonated by NHPI to generate intermediate **IV**. The final *N*-pyridylindole substrate can be obtained upon expulsion of H₂O₂ (Scheme 5).

Scheme 4 Control reactions

Scheme 5 Proposed reaction mechanism

3 Conclusions

In summary, a NHPI and copper co-catalyzed aerobic dehydrogenation of *N*-pyridyl-indolines under mild conditions has been developed. The method displays merits including being green, safe, low-cost and simple to operate. To the best of our knowledge, the existing methods for oxidation of *N*-pyridylindoline derivatives require the use of strong oxidants at elevated temperature. This is the first example of the oxidation of *N*-pyridylindolines using oxy-

gen as a sole oxidant under mild conditions. Owing to the wide utilization of *N*-pyridyl or pyrimidylindoline substrates in transition-metal-catalyzed C—H functionalization, the method would be applicable in the synthesis of indoles, especially for the C(7) decorated indole derivatives. The desirable result of the scale-up reaction demonstrates that the method has potential use in industry.

4 Experimental section

4.1 General information

¹H NMR and ¹³C NMR spectra were recorded in 400 MHz apparatus (Bruker AV-400), and the frequencies for ¹H NMR and ¹³C NMR test are 400 MHz and 100 MHz, respectively. Melting points were obtained using a Bibby Stuart Scientific apparatus. HRMS were obtained by using Q Exactive HF (analyzer type: Orbitrap MS). All reagents and solvents were purchased from commercial suppliers and used without further purification. All the reactions were carried out without any precautions to exclude air and moisture. Reactions were monitored by thin-layer chromatography (TLC) on silica gel GF-254 pre-coated glass plates. Chromatograms were visualized by fluorescence quenching with UV light at 254 nm.

4.2. General procedure for the synthesis of *N*-pyridylindolines

1-(Pyridin-2-yl)indolines (**1**) (0.2 mmol), *N*-hydroxyphthalimide (NHPI) (0.04 mmol) and Cu(OAc)₂·H₂O (0.01 mmol) were dissolved in MeCN (3.0 mL). The reaction mixture was stirred at room temperature for 12 h. After that, the reaction mixture was filtered and evaporated under vacuum. The crude residue was purified by flash column chromatography (petroleum ether/EtOAc, *V* : *V* = 15 : 1) to afford the corresponding indole derivatives **2**.

1-(Pyridin-2-yl)-1*H*-indole (**2a**).^[20] Light yellow liquid (36.1 mg, 93% yield). ¹H NMR (400 MHz, CDCl₃) δ: 8.58 (ddd, *J* = 4.9, 1.9, 0.8 Hz, 1H), 8.25~8.20 (m, 1H), 7.85~7.79 (m, 1H), 7.74 (d, *J* = 3.5 Hz, 1H), 7.70~7.67 (m, 1H), 7.50 (d, *J* = 8.2 Hz, 1H), 7.34~7.30 (m, 1H), 7.25~7.21 (m, 1H), 7.17 (ddd, *J* = 7.3, 4.9, 0.8 Hz, 1H), 6.76~6.70 (m, 1H); ¹³C NMR (101 MHz, CDCl₃) δ: 152.5, 148.9, 138.4, 135.0, 130.4, 125.9, 123.1, 121.2, 121.0, 120.0, 114.6, 112.9, 105.5.

5-Methoxy-1-(pyridin-2-yl)-1*H*-indole (**2b**).^[21] Colorless liquid (37.2 mg, 83% yield). ¹H NMR (400 MHz, CDCl₃) δ: 8.55 (d, *J* = 4.8 Hz, 1H), 8.17 (d, *J* = 9.0 Hz, 1H), 7.82~7.78 (m, 1H), 7.70 (d, *J* = 3.4 Hz, 1H), 7.46 (d, *J* = 8.2 Hz, 1H), 7.16~7.11 (m, 2H), 6.95~6.93 (m, 1H), 6.64 (d, *J* = 3.4 Hz, 1H), 3.88 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 154.9, 152.5, 148.8, 138.3, 131.1, 130.2, 126.2, 119.7, 114.1, 113.9, 112.7, 105.3, 102.9, 55.7.

5-(Benzyloxy)-1-(pyridin-2-yl)-1*H*-indole (**2c**): White solid (48.7 mg, 81% yield). m.p. 128~130 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.55~8.54 (m, 1H), 8.17 (d, *J* = 9.0 Hz, 1H), 7.82~7.79 (m, 1H), 7.69 (d, *J* = 3.5 Hz, 1H), 7.50~7.45 (m, 3H), 7.40 (t, *J* = 7.5 Hz, 2H), 7.34 (d, *J* = 7.1 Hz, 1H), 7.19 (d, *J* = 2.6 Hz, 1H), 7.16~7.14 (m, 1H),

7.04~7.01 (m, 1H), 6.63 (d, $J=3.4$ Hz, 1H), 5.14 (s, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 154.1, 152.5, 148.7, 138.4, 137.5, 131.0, 130.3, 128.5, 127.8, 127.5, 126.3, 119.8, 114.1, 114.0, 113.4, 105.3, 104.3, 70.6; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{17}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$ 301.1341, found 301.1337.

5-Methyl-1-(pyridin-2-yl)-1*H*-indole (**2d**): Yellow solid (31.7 mg, 76% yield). m.p. 58~59 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.56~8.55 (m, 1H), 8.10 (d, $J=8.5$ Hz, 1H), 7.82~7.79 (m, 1H), 7.71 (d, $J=3.4$ Hz, 1H), 7.49 (d, $J=8.2$ Hz, 1H), 7.45 (s, 1H), 7.16~7.12 (m, 2H), 6.64 (d, $J=3.4$ Hz, 1H), 2.47 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.6, 148.9, 138.3, 133.3, 130.7, 130.6, 125.9, 124.6, 120.8, 119.8, 114.2, 112.7, 105.1, 21.3; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{13}\text{N}_2$ $[\text{M}+\text{H}]^+$ 209.1079, found 209.1075.

5-Fluoro-1-(pyridin-2-yl)-1*H*-indole (**2e**): White solid (34.8 mg, 82% yield). m.p. 62~64 °C (lit.^[22] 62~64 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.57 (d, $J=3.8$ Hz, 1H), 8.23 (dd, $J=9.1, 4.6$ Hz, 1H), 7.85~7.81 (m, 1H), 7.73 (d, $J=3.5$ Hz, 1H), 7.45 (d, $J=8.2$ Hz, 1H), 7.30 (dd, $J=9.2, 2.6$ Hz, 1H), 7.18 (dd, $J=7.3, 4.9$ Hz, 1H), 7.06~7.01 (m, 1H), 6.67 (d, $J=3.5$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 158.5 (d, $J_{\text{C-F}}=236.8$ Hz), 152.3, 148.9, 138.5, 131.7, 130.9 (d, $J_{\text{C-F}}=10.1$ Hz), 127.2, 120.1, 114.2 (d, $J_{\text{C-F}}=9.2$ Hz), 114.1, 111.2 (d, $J_{\text{C-F}}=25.5$ Hz), 105.9 (d, $J_{\text{C-F}}=23.4$ Hz), 105.4 (d, $J_{\text{C-F}}=4.3$ Hz).

5-Chloro-1-(pyridin-2-yl)-1*H*-indole (**2f**): Yellow solid (37.5 mg, 82% yield). m.p. 78~80 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.57~8.56 (m, 1H), 8.20 (d, $J=8.9$ Hz, 1H), 7.83~7.81 (m, 1H), 7.71 (d, $J=3.5$ Hz, 1H), 7.62 (d, $J=2.1$ Hz, 1H), 7.44~7.43 (m, 1H), 7.25~7.23 (m, 1H), 7.20~7.17 (m, 1H), 6.65 (dd, $J=3.5, 0.7$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 148.9, 138.5, 133.5, 131.4, 127.0, 126.8, 123.3, 120.4, 120.3, 114.4, 114.3, 105.0; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{10}\text{ClN}_2$ $[\text{M}+\text{H}]^+$ 229.0533, found 229.0530.

5-Bromo-1-(pyridin-2-yl)-1*H*-indole (**2g**): White solid (45.3 mg, 83% yield). m.p. 98~100 °C (lit.^[21] 98~100 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.58~8.56 (m, 1H), 8.15 (d, $J=8.9$ Hz, 1H), 7.86~7.82 (m, 1H), 7.78 (d, $J=1.9$ Hz, 1H), 7.70 (d, $J=3.5$ Hz, 1H), 7.45 (d, $J=8.2$ Hz, 1H), 7.37 (dd, $J=8.9, 2.0$ Hz, 1H), 7.21~7.18 (m, 1H), 6.65 (d, $J=3.5$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 149.0, 138.5, 133.8, 132.0, 126.9, 125.9, 123.5, 120.4, 114.8, 114.4, 114.4, 104.9.

Methyl 1-(pyridin-2-yl)-1*H*-indole-5-carboxylate (**2h**): White solid (36.3 mg, 72% yield). m.p. 101~103 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.60 (d, $J=3.9$ Hz, 1H), 8.41 (d, $J=1.3$ Hz, 1H), 8.22 (d, $J=8.8$ Hz, 1H), 7.99 (dd, $J=8.8, 1.6$ Hz, 1H), 7.86 (t, $J=7.8$ Hz, 1H), 7.76 (d, $J=3.5$ Hz, 1H), 7.50 (d, $J=8.2$ Hz, 1H), 7.24~7.21 (m, 1H), 6.80 (d, $J=3.5$ Hz, 1H), 3.95 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 167.8, 152.0, 149.0, 138.5, 137.5, 129.9, 127.3, 124.4, 123.7, 123.1, 120.7, 114.8, 112.7, 106.4, 51.9; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{13}\text{N}_2\text{O}_2$ $[\text{M}+\text{H}]^+$ 253.0977, found 253.0974.

5-Nitro-1-(pyridin-2-yl)-1*H*-indole (**2i**): Light yellow solid (39.7 mg, 83% yield). m.p. 88~90 °C (lit.^[22] 88~90 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.62~8.59 (m, 2H),

8.32 (d, $J=9.2$ Hz, 1H), 8.19~8.16 (m, 1H), 7.90 (td, $J=8.1, 1.8$ Hz, 1H), 7.81 (d, $J=3.5$ Hz, 1H), 7.49 (d, $J=8.2$ Hz, 1H), 7.29 (dd, $J=7.1, 5.2$ Hz, 1H), 6.88 (d, $J=3.4$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.7, 149.2, 142.6, 138.9, 138.0, 129.7, 128.9, 121.4, 118.5, 117.8, 114.9, 113.4, 107.0.

4-(Benzyloxy)-1-(pyridin-2-yl)-1*H*-indole (**2j**): White solid (48.1 mg, 80% yield). m.p. 90~92 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.58 (s, 1H), 7.82 (t, $J=9.8$ Hz, 2H), 7.66 (s, 1H), 7.54 (d, $J=6.8$ Hz, 2H), 7.51 (d, $J=7.9$ Hz, 1H), 7.42 (t, $J=6.8$ Hz, 2H), 7.36 (d, $J=6.7$ Hz, 1H), 7.22 (t, $J=7.7$ Hz, 1H), 7.17 (s, 1H), 6.93 (s, 1H), 6.72 (d, $J=7.5$ Hz, 1H), 5.27 (s, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.5, 148.9, 138.3, 137.4, 136.4, 128.8, 128.5, 127.7, 127.3, 124.6, 123.8, 121.2, 120.1, 115.1, 114.7, 106.4, 102.8, 70.0; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{17}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$ 301.1341, found 301.1337.1

4-Methyl-1-(pyridin-2-yl)-1*H*-indole (**2k**): Light yellow liquid (38.3 mg, 92% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.64 (d, $J=4.8$ Hz, 1H), 8.10 (d, $J=8.2$ Hz, 1H), 7.87~7.80 (m, 2H), 7.55 (d, $J=8.2$ Hz, 1H), 7.32~7.27 (m, 1H), 7.23~7.20 (m, 1H), 7.10 (d, $J=7.1$ Hz, 1H), 6.82 (d, $J=2.9$ Hz, 1H), 2.67 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.4, 148.9, 138.2, 134.7, 130.3, 130.1, 125.4, 123.1, 121.5, 119.9, 114.6, 110.4, 103.8, 18.6; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{13}\text{N}_2$ $[\text{M}+\text{H}]^+$ 209.1079, found 209.1075.

4-Fluoro-1-(pyridin-2-yl)-1*H*-indole (**2l**): Yellow solid (36.9 mg, 87% yield). m.p. 53~55 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.55~8.54 (m, 1H), 7.97 (d, $J=8.4$ Hz, 1H), 7.80~7.76 (m, 1H), 7.64 (d, $J=3.5$ Hz, 1H), 7.43 (d, $J=8.2$ Hz, 1H), 7.23~7.14 (m, 2H), 6.87 (dd, $J=9.9, 7.9$ Hz, 1H), 6.79 (d, $J=3.5$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 156.1 (d, $J_{\text{C-F}}=246.6$ Hz), 152.2, 148.9, 138.5, 137.4 (d, $J_{\text{C-F}}=10.4$ Hz), 125.9, 123.6 (d, $J_{\text{C-F}}=7.6$ Hz), 120.5, 119.2 (d, $J_{\text{C-F}}=22.5$ Hz), 114.6, 109.1, 106.1 (d, $J_{\text{C-F}}=18.4$ Hz), 101.1; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{10}\text{FN}_2$ $[\text{M}+\text{H}]^+$ 213.0828, found 213.0825.

4-Chloro-1-(pyridin-2-yl)-1*H*-indole (**2m**): Light yellow liquid (38.9 mg, 85% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.58~8.57 (m, 1H), 8.15~8.12 (m, 1H), 7.85~7.82 (m, 1H), 7.73 (d, $J=3.5$ Hz, 1H), 7.46 (d, $J=8.2$ Hz, 1H), 7.22~7.19 (m, 3H), 6.84 (m, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 149.0, 138.5, 135.8, 129.0, 126.5, 126.1, 123.7, 121.0, 120.6, 114.7, 111.7, 103.8; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{10}\text{ClN}_2$ $[\text{M}+\text{H}]^+$ 229.0533, found 229.0530.

4-Bromo-1-(pyridin-2-yl)-1*H*-indole (**2n**):^[21] Light yellow liquid (44.8 mg, 82% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.58 (dd, $J=4.8, 1.7$ Hz, 1H), 8.18 (d, $J=8.4$ Hz, 1H), 7.86~7.74 (m, 1H), 7.75 (d, $J=3.5$ Hz, 1H), 7.47 (d, $J=8.2$ Hz, 1H), 7.38 (d, $J=7.6$ Hz, 1H), 7.21 (dd, $J=7.4, 4.9$ Hz, 1H), 7.15 (t, $J=8.0$ Hz, 1H), 6.79 (d, $J=3.5$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 149.0, 138.6, 135.4, 130.9, 126.5, 124.1, 124.0, 120.6, 114.80, 114.78, 112.3, 105.6.

Methyl 1-(pyridin-2-yl)-1*H*-indole-4-carboxylate (**2o**): Light yellow liquid (44.4 mg, 88% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.58~8.57 (m, 1H), 8.48 (d, $J=8.3$ Hz,

1H), 7.99~7.98 (m, 1H), 7.84~7.81 (m, 1H), 7.79 (d, $J=3.5$ Hz, 1H), 7.45 (d, $J=8.2$ Hz, 1H), 7.41~7.39 (m, 1H), 7.33 (t, $J=9$ Hz, 1H), 7.21~7.19 (m, 1H), 4.00 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 167.7, 152.2, 148.9, 138.6, 135.8, 130.0, 127.8, 124.5, 122.3, 121.8, 120.5, 118.0, 114.9, 106.3, 51.8; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{13}\text{N}_2\text{O}_2$ $[\text{M}+\text{H}]^+$ 253.0972, found 253.0974.

6-Methyl-1-(pyridin-2-yl)-1H-indole (**2p**):^[23] Yellow liquid (32.9 mg, 79% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.59 (dd, $J=4.9, 1.8$ Hz, 1H), 8.04 (s, 1H), 7.84~7.79 (m, 1H), 7.65 (d, $J=3.5$ Hz, 1H), 7.56 (d, $J=8.0$ Hz, 1H), 7.48 (d, $J=7.8$ Hz, 1H), 7.16 (dd, $J=7.3, 4.9$ Hz, 1H), 7.06 (d, $J=8.0$ Hz, 1H), 6.68 (d, $J=3.5$ Hz, 1H), 2.53 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.5, 148.8, 138.3, 135.4, 132.9, 128.1, 125.4, 122.9, 120.6, 119.8, 114.6, 112.9, 105.3, 22.0.

7-Methyl-1-(pyridin-2-yl)-1H-indole (**2q**):^[24] Yellow liquid (30.4 mg, 73% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.60 (d, $J=4.4$ Hz, 1H), 7.83 (td, $J=7.7, 1.9$ Hz, 1H), 7.54 (d, $J=7.8$ Hz, 1H), 7.35~7.30 (m, 3H), 7.11 (t, $J=7.5$ Hz, 1H), 7.02 (d, $J=7.2$ Hz, 1H), 6.68 (d, $J=3.3$ Hz, 1H), 2.14 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 153.4, 148.8, 137.8, 135.1, 130.4, 129.6, 125.4, 122.1 (2C), 121.0, 120.6, 118.9, 104.6, 20.6.

2-Methyl-1-(pyridin-2-yl)-1H-indole (**2r**):^[25] Yellow liquid (29.2 mg, 70% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.63 (dd, $J=4.9, 1.8$ Hz, 1H), 7.84~7.80 (m, 1H), 7.56~7.53 (m, 1H), 7.39~7.34 (m, 2H), 7.27~7.24 (m, 1H), 7.14~7.09 (m, 2H), 6.41 (s, 1H), 2.44 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.3, 149.4, 138.1, 137.0, 136.7, 128.6, 121.8, 121.4, 120.7, 120.6, 119.6, 110.2, 103.2, 13.9.

3-Methyl-1-(pyridin-2-yl)-1H-indole (**2s**):^[26] Light yellow liquid (34.6 mg, 83% yield). ^1H NMR (400 MHz, CDCl_3) δ : 8.53 (dd, $J=4.8, 1.5$ Hz, 1H), 8.22 (d, $J=8.3$ Hz, 1H), 7.79~7.74 (m, 1H), 7.59 (d, $J=7.8$ Hz, 1H), 7.51 (s, 1H), 7.43 (d, $J=8.3$ Hz, 1H), 7.30 (t, $J=7.2$ Hz, 1H), 7.24~7.20 (m, 1H), 7.09 (dd, $J=7.3, 4.9$ Hz, 1H), 2.37 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.5, 148.8, 138.2, 135.3, 131.0, 123.2, 123.1, 120.8, 119.4, 119.0, 114.7, 114.0, 113.0, 9.7.

1-(Pyrimidin-2-yl)-1H-indole (**2t**): White solid (31.6 mg, 81% yield). m.p. 86~88 °C (lit.^[21] 86~88 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.80 (d, $J=8.3$ Hz, 1H), 8.64~8.63 (m, 2H), 8.26 (s, 1H), 7.61 (d, $J=7.6$ Hz, 1H), 7.33 (t, $J=7.6$ Hz, 1H), 7.23 (t, $J=7.2$ Hz, 1H), 6.95 (s, 1H), 6.69 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 158.1, 157.8, 135.4, 131.3, 125.8, 123.6, 122.1, 120.8, 116.3, 116.1, 106.9.

N,N-diphenyl-1-(pyridin-2-yl)-1H-indol-7-amine (**2u**): White solid (54.2 mg, 75% yield). m.p. 140~142 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.21~8.14 (m, 1H), 7.56 (d, $J=7.8$ Hz, 1H), 7.45 (td, $J=7.8, 1.8$ Hz, 1H), 7.33 (d, $J=3.3$ Hz, 1H), 7.17 (t, $J=7.7$ Hz, 1H), 7.04 (q, $J=8.6, 8.0$ Hz, 7H), 6.86 (t, $J=7.3$ Hz, 2H), 6.74 (d, $J=7.8$ Hz, 4H), 6.70 (d, $J=3.3$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.7, 147.6, 146.2, 136.8, 132.8, 132.5, 129.8, 128.5, 122.8, 121.7, 121.5, 121.0, 119.6, 118.1, 104.5; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{20}\text{N}_3$ $[\text{M}+\text{H}]^+$ 362.1652, found

362.1647.

1-(Pyrimidin-2-yl)-1H-indol-7-yl acetate (**2v**): White solid (36.5 mg, 72% yield). m.p. 74~76 °C (lit.^[27] 74~76 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.70 (d, $J=4.8$ Hz, 2H), 7.99 (d, $J=3.5$ Hz, 1H), 7.51 (d, $J=7.8$ Hz, 1H), 7.23 (t, $J=7.8$ Hz, 1H), 7.12 (t, $J=4.8$ Hz, 1H), 7.03 (d, $J=7.8$ Hz, 1H), 6.73 (d, $J=3.5$ Hz, 1H), 2.19 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.1, 158.2, 157.5, 137.5, 134.7, 129.6, 126.7, 122.5, 119.0, 118.3, 117.0, 107.1, 21.4.

7-(Phenylsulfonyl)-1-(pyrimidin-2-yl)-1H-indole (**2w**): Yellow solid (47.0 mg, 70% yield). m.p. 97~99 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.41 (d, $J=4.8$ Hz, 2H), 7.89~7.84 (m, 2H), 7.82~7.77 (m, 2H), 7.67 (d, $J=7.7$ Hz, 1H), 7.56 (t, $J=7.4$ Hz, 1H), 7.49 (t, $J=7.5$ Hz, 2H), 7.29 (d, $J=7.8$ Hz, 1H), 7.10 (t, $J=4.8$ Hz, 1H), 6.82 (d, $J=3.5$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 159.0, 158.27, 143.7, 133.6, 132.30, 131.7, 131.1, 128.7, 127.7, 126.8, 126.3, 125.8, 121.3, 118.5, 106.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{N}_3\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 336.0801, found 336.0797.

Supporting Information ^1H NMR and ^{13}C NMR spectra of compounds **2a**~**2w**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Bariwal, J.; Voskressensky, L. G.; Van der Eycken, E. V. *Chem. Soc. Rev.* **2018**, 47, 3831.
(b) Xie, T.-Z.; Zhao, Y.-L.; Ma, W.-G.; Wang, Y.-F.; Yu, H.-F.; Wang, B.; Wei, X.; Huang, Z.-P.; Zhu, P.-F.; Liu, Y.-P.; Luo, X.-D. *Chin. J. Org. Chem.* **2020**, 40, 679 (in Chinese).
(解天珍, 赵云丽, 马伟光, 王易芬, 于浩飞, 王 蓓, 魏 鑫, 黄之镨, 朱培凤, 刘亚平, 罗晓东, 有机化学, **2020**, 40, 679.)
(c) Huang, J.; Luo, S.; Gong, L. *Acta Chim. Sinica* **2013**, 71, 879 (in Chinese).
(黄建洲, 罗时玮, 龚流柱, 化学学报, **2013**, 71, 879.)
(d) America, M.; Asgharia, A.; Amoozadeha, A.; Bakherad, M.; *Chin. Chem. Lett.* **2017**, 28, 1031.
(e) Liang, K. J.; Xia, C. F. *Chin. J. Chem.* **2017**, 35, 255.
- [2] (a) Bandini, M.; Eichholzer, A. *Angew. Chem., Int. Ed.* **2009**, 48, 9608.
(b) Barluenga, J.; Rodriguez, F.; Fananas, F. J. *Chem.-Asian J.* **2009**, 4, 1036.
(c) Shi, Z.; Zhang, C.; Li, S.; Pan, D.; Ding, S.; Cui, Y.; Jiao, N. *Angew. Chem., Int. Ed.* **2009**, 48, 4572.
(d) Taber, D. F.; Tirunahari, P. K. *Tetrahedron* **2011**, 67, 7195.
(e) Shi, Z.; Nie, K.; Liu, C.; Zhang, M.; Zhang, W. *Chin. J. Org. Chem.* **2020**, 40, 327 (in Chinese).
(施展, 聂克睿, 刘畅, 张明智, 章维华, 有机化学, **2020**, 40, 327.)
(f) Yu, K.; Gao, B.; Ding, H. *Acta Chim. Sinica* **2016**, 74, 410 (in Chinese).
(余宽, 高北岭, 丁寒峰, 化学学报, **2016**, 74, 410.)
(g) Huang, E.; Zhang, L.; Xiao, C.; Meng, G.; Zhang, B.; Hu, J.; Wan, D. C.-C.; Meng, Q.; Jin, Z.; Hua, C. *Chin. Chem. Lett.*, **2019**, 30, 2157.
(h) Cerveri, A.; Bandini, M. *Chin. J. Chem.* **2020**, 38, 287.
(i) Yang, P.; Xu, R.-Q.; Zheng, C.; You, S.-L. *Chin. J. Chem.* **2020**, 38, 235.
- [3] (a) Thikekar, T. U.; Selvaraju, M.; Sun, C.-M. *Org. Lett.* **2016**, 18, 316.
(b) Peng, F.; McLaughlin, M.; Liu, Y.; Mangion, I.; Tschäen, D. M.;

- Xu, Y. *J. Org. Chem.* **2016**, *81*, 10009.
- (c) Zhang, J.; Chen, S.; Chen, F.; Xu, W.; Deng, G.-J.; Gong, H. *Adv. Synth. Catal.* **2017**, *359*, 2358.
- (d) Huo, H.-R.; Tang, X.-Y.; Gong, Y.-F. *Synthesis* **2018**, *50*, 2727.
- (e) Taboonpong, P.; Chavasiri, W. *Catal. Commun.* **2018**, *104*, 9.
- (f) Zhu, R.; Zhai, M.; Liu, S.; Liu, X.; Wang, Z.; Ju, R.; Yu, X. *Chin. J. Org. Chem.* **2020**, *40*, 2045 (in Chinese).
(朱润宇, 翟民, 刘霜, 刘星彤, 王振, 居瑞军, 虞心红, 有机化学, **2020**, *40*, 2045.)
- [4] (a) Pratt, E. F.; McGovern, T. P. *J. Org. Chem.* **1964**, *29*, 1540.
(b) Kawase, M.; Miyake, Y.; Kikugawa, Y. *J. Chem. Soc., Perkin Trans. 1* **1984**, 1401.
(c) Giethlen, B.; Schaus, J. M. *Tetrahedron Lett.* **1997**, *38*, 8483.
(d) Mukaiyama, T.; Kawana, A.; Fukuda, Y.; Matsuo, J. *Chem. Lett.* **2001**, 390.
(e) Tilstam, Kawana, U.; Harre, M.; Heckrodt, T.; Weinmann, H. *Tetrahedron Lett.* **2001**, *42*, 5385.
- [5] (a) Zhang, X.; Liu, W.; Sun, D.; Huang, J.; Qiu, X.; Li, Z.; Wu, Z. *ChemSusChem* **2020**, *4774*.
(b) Chen, H.; Liu, L.; Huang, T.; Chen, J.; Chen, T. *Adv. Synth. Catal.* **2020**, 3332.
(c) Tang, H.; Luo, J.; Xie, P. *Chin. J. Org. Chem.* **2019**, *39*, 2735 (in Chinese).
(唐灏, 骆钧飞, 解攀, 有机化学, **2019**, *39*, 2735.)
(d) Zhang, Z.; Tian, Q.; Qian, J.; Liu, Q.; Liu, T.; Shi, L.; Zhang, G. *J. Org. Chem.* **2014**, *79*, 8182.
(e) Qian, J.; Zhang, Z.; Liu, Q.; Liu, T.; Zhang, G. *Adv. Synth. Catal.* **2014**, *356*, 3119.
(f) Zhang, Z.; Zhang, W.; Li, J.; Liu, Q.; Liu, T.; Zhang, G. *J. Org. Chem.* **2014**, *79*, 11226.
- [6] (a) Wang, Q.; Chai, H.; Yu, Z. *Organometallics* **2018**, *37*, 584.
(b) He, K. H.; Tan, F. F.; Zhou, C. Z.; Zhou, G.-J.; Yang, X.-L.; Li, Y. *Angew. Chem., Int. Ed.* **2017**, *56*, 3080.
- [7] (a) Lou, C.; Qin, S.; Zhang, S.; Lv, Z.; Senan, A.; Chen, Z.; Yin, G. *Catal. Commun.* **2017**, *90*, 5.
(b) Hara, T.; Mori, K.; Mizugaki, T.; Ebitani, K.; Kaneda, K. *Tetrahedron Lett.* **2003**, *44*, 6207.
- [8] (a) Elumalai, K.; Weng, K. L. *Tetrahedron Lett.* **2018**, *59*, 3958.
(b) So, M. H.; Liu, Y.; Ho, C.-M.; Che, C.-M. *Chem.-Asian J.* **2009**, *4*, 1551.
- [9] (a) Zhou, W.; Chen, D.; Sun, F.; Qian, J.; He, M.; Chen, Q. *Tetrahedron Lett.* **2018**, *59*, 949.
(b) Iosub, A. V.; Stahl, S. S. *Org. Lett.* **2015**, *17*, 4404.
- [10] (a) Jawale, D. V.; Gravel, E.; Shah, N.; Dauvois, V.; Li, H.; Namboothiri, I. N. N.; Doris, E. *Chem.-Eur. J.* **2015**, *21*, 7039.
(b) Damodara, D.; Arundhathi, R.; Likhara, P. R. *Adv. Synth. Catal.* **2014**, *356*, 189.
(c) Zhou, W.; Taboonpong, P.; Aboo, A. H.; Zhang, L.; Jiang, J.; Xiao, J. *Synlett* **2016**, *27*, 1806.
(d) Cui, X.; Li, Y.; Bachmann, S.; Scalone, M.; Surkus, A.; Junge, K.; Topf, C.; Beller, M. *J. Am. Chem. Soc.* **2015**, *137*, 10652.
(e) Wendlandt, A. E.; Stahl, S. S. *J. Am. Chem. Soc.* **2014**, *136*, 506.
(f) Karki, M.; Araujo, H. C.; Magolan, J. *Synlett* **2013**, *24*, 1675.
- [11] (a) Kojima, M.; Kanai, M. *Angew. Chem., Int. Ed.* **2016**, *55*, 12224.
(b) Maier, A. F. G.; Tussing, S.; Schneider, T.; Florke, U.; Qu, Z.-W.; Grimme, S.; Paradies, J. *Angew. Chem., Int. Ed.* **2016**, *55*, 12219.
- [12] Chen, W.; Tang, H.; Wang, W.; Fu, Q.; Luo, J. *Adv. Synth. Catal.* **2020**, *362*, 3905.
- [13] (a) Burke, J. P.; Bian, Z.; Shaw, S.; Zhao, B.; Goodwin, C. M.; Belmar, J.; Browning, C. F.; Vigil, D.; Friberg, A.; Camper, D. V.; Rossanese, O. W.; Lee, T.; Olejniczak, E. T.; Fesik, S. W. *J. Med. Chem.* **2015**, *58*, 3794.
(b) Ozawa, Y.; Kusano, K.; Owa, T.; Yokoi, A.; Asada, M.; Yoshimatsu, K. *Cancer Chemother. Pharmacol.* **2012**, *69*, 1353.
- (c) Dydio, P.; Zielinski, T.; Jurczak, J. *Chem. Commun.* **2009**, 4560.
(d) Ferandin, Y.; Bettayeb, K.; Kritsanida, M.; Lozach, O.; Polychronopoulos, P.; Magiatis, P.; Skaltsounis, A.-L.; Meijer, L. *J. Med. Chem.* **2006**, *49*, 4638.
- [14] (a) Yang, X.-F.; Hu, X.-H.; Loh, T.-P. *Org. Lett.* **2015**, *17*, 1481.
(b) Shin, Y.; Sharma, S.; Mishra, N. K.; Han, S.; Park, J.; Oh, H.; Ha, J.; Yoo, H.; Jung, Y. H.; Kim, I. S. *Adv. Synth. Catal.* **2015**, *357*, 594.
(c) Mishra, N. K.; Jeong, T.; Sharma, S.; Shin, Y.; Han, S.; Park, J.; Oh, J. S.; Kwak, J. H.; Jung, Y. H.; Kim, I. S. *Adv. Synth. Catal.* **2015**, *357*, 1293.
(d) Xie, W.; Li, B.; Wang, B. *J. Org. Chem.* **2015**, *81*, 396.
(e) Wang, X.; Tang, H.; Feng, H.; Li, Y.; Yang, Y.; Zhou, B. *J. Org. Chem.* **2015**, *80*, 6238.
(f) Hou, W.; Yang, Y.; Ai, W.; Wu, Y.; Wang, X.; Zhou, B.; Li, Y. *Eur. J. Org. Chem.* **2015**, 395.
(g) Park, J.; Mishra, N. K.; Sharma, S.; Han, S.; Shin, Y.; Jeong, T.; Oh, J. S.; Kwak, J. H.; Jung, Y. H.; Kim, I. S. *J. Org. Chem.* **2015**, *80*, 1818.
(h) Yang, D.; Mao, S.; Gao, Y.-R.; Guo, D.-D.; Guo, S.-H.; Li, B.; Wang, Y.-Q. *RSC Adv.* **2015**, *5*, 23727.
(i) Kim, M.; Mishra, N. K.; Park, J.; Han, S.; Shin, Y.; Sharma, S.; Lee, Y.; Lee, E.-K.; Kwak, J. H.; Kim, I. S. *Chem. Commun.* **2014**, *50*, 14249.
(j) Pan, S.; Wakaki, T.; Ryu, N.; Shibata, T. *Chem.-Asian J.* **2014**, *9*, 1257.
(k) Urones, B.; Arrayas, R. G.; Carretero, J. C. *Org. Lett.* **2013**, *15*, 1120.
- [15] (a) Ahmad, A.; Dutta, H. S.; Kumar, M. Khan, A. A.; Raziullah, Koley, D. *Org. Lett.* **2020**, *22*, 5870.
(b) Ahmad, A.; Dutta, H. S.; Khan, B.; Kant, R.; Koley, D. *Adv. Synth. Catal.* **2018**, *360*, 1644.
(c) Zhai, W.; Li, B.; Wang, B. *ChemistrySelect* **2018**, *3*, 8035.
(d) Ai, W.; Yang, X.; Wu, Y.; Wang, X.; Li, Y.; Yang, Y.; Zhou, B. *Chem.-Eur. J.* **2014**, *20*, 17653.
(e) Wu, Y.; Yang, Y.; Zhou, B.; Li, Y. *J. Org. Chem.* **2015**, *80*, 1946.
- [16] Kumar, M.; Sharma, R.; Raziullah, Khan, A. A.; Ahmad, A.; Dutta, H. S.; Koley, D. *Org. Lett.* **2020**, *22*, 2152.
- [17] Gao, S.; Xu, X.; Tang, H.; Wu, J.-Q.; Luo, J. *ChemistrySelect* **2019**, *4*, 5835.
- [18] Zhi, C.; Wang, Q.; Liu, S.; Xue, Y.; Shi, L.; Zhu, X.; Hao, X.-Q.; Song, M.-P. *J. Org. Chem.* **2020**, *85*, 1022.
- [19] (a) Lee, J. M.; Park, E. J.; Cho, S. H.; Chang, S. *J. Am. Chem. Soc.* **2008**, *130*, 7824.
(b) Ishii, Y.; Sakaguchi, S.; Iwahamab, T. *Adv. Synth. Catal.* **2001**, *343*, 393.
(c) Bag, R.; Sar, D.; Punniyamurthy, T. *Org. Lett.* **2015**, *17*, 2010.
(d) Zhou, W.-J.; Cao, G.-M.; Shen, G.; Zhu, X.-Y.; Gui, Y.-Y.; Ye, J.-H.; Sun, L.; Liao, L.-L.; Li, J.; Yu, D.-G. *Angew. Chem., Int. Ed.* **2017**, *56*, 15683.
(e) Parasram, M.; Chuentragool, P.; Wang, Y.; Shi, Y.; Gevorgyan, V. *J. Am. Chem. Soc.* **2017**, *139*, 14857.
- [20] Ackermann, L.; Lygin, A. V. *Org. Lett.* **2011**, *13*, 3332.
- [21] Xu, S.; Huang, X.; Hong, X.; Xu, B. *Org. Lett.* **2012**, *14*, 4614.
- [22] Liang, L.; Fu, S.; Lin, D.; Zhang, X.-Q.; Deng, Y.; Jiang, H.; Zeng, W. *J. Org. Chem.* **2014**, *79*, 9472.
- [23] Li, T.; Wang, Z.; Zhang, M.; Zhang, H.-J.; Wen, T.-B. *Chem. Commun.* **2015**, *51*, 6777.
- [24] Liu, S.-L.; Li, Y.; Guo, J.-R.; Yang, G.-C.; Li, X.-H.; Gong, J.-F.; Song, M.-P. *Org. Lett.* **2017**, *19*, 4042.
- [25] Manna, M. K.; Bairy, G.; Jana, R. *J. Org. Chem.* **2018**, *83*, 8390.
- [26] Rull, S. G.; Blandez, J. F.; Fructos, M. R.; Belderrain, T. R.; Nicasio, M. C. *Adv. Syn. Catal.* **2015**, *357*, 907.
- [27] Gao, S.; Xu, X.; Tang, H.; Wu, J.-Q.; Luo, J. *ChemistrySelect* **2019**, *4*, 5835.

(Zhao, C.)