无过渡金属及无碱参与的 *N*-苄基-*N*-叔丁基羰基酰胺的转酰胺化叶丹锋^{a,b} 陈浩^b 刘志园^b 雷川虎^{*,b}^(a) 上海大学材料科学与工程学院 上海 200444)^(b) 上海大学理学院化学系 超分子化学与催化研究中心 上海 200444)

摘要 报道了一种无过渡金属和无碱条件下 *N*-苄基-*N*-叔丁基羰基酰胺发生转酰胺化的方法。该反应从高活性的 *N*-叔丁基羰基酰亚胺出发, 历经亲核取代机理, 可以中等到优良的收率得到一系列取代酰胺产物。值得一提的是, 由于反应体系无需添加额外的碱, 使得氨基酸衍生的手性亲核试剂及具有手性碳的酰胺在反应过程中其手性均能得到保持。此外, 应用该方法还合成得到一些前药及抗抑郁症药物吗氯贝胺, 展示了该方法在有机合成中的潜在应用。

关键词 转酰胺化; *N*-苄基-*N*-叔丁基羰基酰胺; 无过渡金属; 无碱

Transamidation of *N*-Benzyl-*N*-Boc-amides under Transition Metal-Free and Base-Free ConditionsYe, Danfeng^{a,b} Chen, Hao^b Liu, Zhiyuan^b Lei, Chuanhu^{*,b}^(a) School of Materials Science and Engineering, Shanghai University, Shanghai 200444)^(b) Center for Supramolecular Chemistry & Catalysis and Department of Chemistry, College of Sciences, Shanghai University, Shanghai 200444)

Abstract A new protocol for the transamidation of *N*-benzyl-*N*-Boc-amides under transition metal-free and base-free conditions is described. The reaction features high reactivity of *N*-Boc-imides via direct acyl nucleophilic substitution mechanism, and provides access to a diverse array of substituted amides in good to excellent yields. Notably, the base-free condition preserves enantiopurity with respect to chiral amino-acid-derived nucleophiles, and *N*-Boc-amides bearing an epimerizable stereocenter. The application of this method is also demonstrated through the synthesis of pro-drugs and antidepressant moclobemide.

Keywords transamidation; *N*-benzyl-*N*-Boc-amides; transition metal-free; base-free

1 Introduction

Amide is one of the most fundamental functional groups found in a vast array of natural products, pharmaceuticals, fine chemicals, materials, and constitutes the basic linkage of proteins.^[1] The preparation of amide and their further synthetic conversions have continuously attracted considerable attention. Among many synthetic methods for amide transformations,^[2] transamidation processes are attractive due to the ready availability of amides.^[3] However, the chemical stability of amides, ascribed to C(O)—N amidic resonance, generally translates into reduced reactivity and constrained synthetic utility. Thus, tailor-made metal complex,^[4] harsh experimental conditions^[5] or highly evolved

enzymes^[6] constitute the basis for transamidation reactions. A breakthrough of the field was made by Garg and co-workers who demonstrated that nickel-catalyzed C(O)—N bond activation^[7] could affect *N*-benzyl-*N*-Boc-amide transamidation.^[8] Nearly in the same time, Szostak *et al.*^[9] reported that Pd/N-heterocyclic carbene complexes can be used to realize the transamidation of secondary carboxamides carrying *N*-*tert*-butoxycarbonyl (Boc) group (Scheme 1a). Despite these seminal advances, transition metal-free conditions are needed,^[10] especially for drug manufacturing industry, where trace metal contaminants can be problematic. Within the cadre of this latter focus, important progress has been made by Szostak and cowork-

* Corresponding author. E-mail: chlei@shu.edu.cn

Received September 22, 2020; revised October 23, 2020; published online December 9, 2020.

Project supported by the National Natural Science Foundation of China (No. 21901155), the Shanghai Rising-Star Program (No. 20QA1403700), the Eastern Scholars Program from the Shanghai Municipal Education Committee, and the Shanghai University Startup Funding (No. N.13-G210-20-263).

国家自然科学基金(No. 21901155)、上海市青年科技启明星计划(No. 20QA140370)、上海高校特聘教授(东方学者)及上海大学启动基金(No. N.13-G210-20-263)资助项目。

ers.^[11] By exploiting *N*-Boc/Ts activated amides or *N*-acyl-glutarimides as the electrophiles, they have realized transamidations under transition metal-free conditions, although stoichiometric quantities of base, *e.g.* Et₃N or LiHMDS, were required to promote the reported transformations (Scheme 1b). In 2018, Zeng group reported CsF-catalyzed transamidation of *N,N*-diBoc-amides. The key step of this reaction involves formation of an acyl fluoride intermediate that proceeds at relatively high temperatures.^[12] Meanwhile, Verho *et al.*^[13] demonstrated that *N*-Boc-*N*-(8-quinoliny)amides could be employed to accomplish metal-free transamidations in the absence of any additional reagents. Later on, Szostak^[14] and Chandrasekaran^[15] also achieved base-free transamidations by employing twisted *N*-Boc-*N*-phenyl-amides and *N*-benzyl-*N*-Ts-amides, respectively (Scheme 1c). Despite these progresses, challenges remain. For instance, the substrate scope in these transition-metal free transformations has yet to be expanded to include optically enriched materials. Moreover, the development of more general transamidation methods that proceed under base-free conditions is still in need.^[16]

(a) Garg and Szostak: transition metal-catalyzed transamidation^[8-9]

(b) Szostak: transition metal-free transamidation in the presence of base^[11]

(c) Verho, Szostak and Chandrasekaran: Transition metal-free and base-free transamidation^[13-15]

(d) This work: transition metal-free and base-free transamidation of *N*-benzyl-*N*-Boc amides

Scheme 1 Methods for the transamidation reaction

Generally, aliphatic amines are better nucleophiles than alcohols. Inspired by the success of our recent esterification of *N*-benzyl-*N*-Boc-amides promoted by catalytical quantities of Cs₂CO₃,^[17] we envisioned that the transamidation of *N*-benzyl-*N*-Boc-amides with aliphatic amines might be feasible. Realization of this transformation would complement the existing transamidation protocols, and might expand the synthetic utility of *N*-benzyl-*N*-Boc-

amides as a bench-stable, readily available acyl-transfer reagents in organic synthesis. Here we report a transamidation method using *N*-benzyl-*N*-Boc-amides under transition metal-free and base-free conditions. As detailed below, this procedure could be applied to substrates bearing chiral stereocenters, with no detectable erosion of enantiopurity in most cases (Scheme 1d).

2 Results and discussion

We started our investigations by optimizing the transamidation between *N*-benzyl-*N*-Boc-amide **1a** and pyrrolidine **2a** (Table 1). Following our previous studies involving the Cs₂CO₃-catalyzed esterification of *N*-activated amides,^[17] Cs₂CO₃ was initially selected as the putative catalyst and DMF as the solvent. To our surprise, the yield increased as the loading of Cs₂CO₃ decreased. In fact, the reaction was found to occur in the absence of base, producing the desired amide **3a** in 81% yield. It was rationalized that possible formation of Cs⁺-pyrrolidine Lewis acid-base adduct reduces the nucleophilicity of amine (Entries 1~4).^[18] Such a proposed adduct was expected to depend on the polarity of the medium. Therefore, a screening of solvents was carried out. It was found that toluene was the most effective medium (Entry 9). For example, the yields for **3a** were obtained in 91%, 83%, 78%, and 81% when the reaction was performed in toluene, THF, CH₃CN and diethyl ether, respectively (Entries 5~9). When the reaction was performed in toluene at 80 °C, the yield was only 50% as determined by GC analysis. However, increasing temperature to reflux resulted in an almost quantitative GC yield corresponding to 90% isolated yield (Entry 11). When the stoichiometry of amine was reduced to 1.0 equiv., the yield decreased to 79%. Thus, the optimized conditions involved the use of 1.5 equiv. of amine in toluene at reflux in the absence of any other reagents.

Next, a number of disubstituted benzamides (imides) including *N*-acylanilide, other *N*-Boc-amides, *N*-tosyl amides, and *N*-glutarimide were tested. These substrates were chosen since the structural differences were expected to affect the stability of the amide bond, thereby influencing the reaction outcomes. As shown in Scheme 2, *N*-methyl-*N*-phenyl benzamide (**1aa**) proved completely unreactive. Amide **1ab** bearing *N*-phenyl group displayed reactivity comparable to that of *N*-benzyl amide **1a**. When benzyl group in **1a** was replaced by methyl one, the yield of the reaction dramatically decreased to 66% (**1ac**). Intriguingly, moving from methyl to ethyl resulted in an increase of yield by *ca.* 30% (**1ac** vs **1ad**). In the case of *N*-tosyl amides (**1af** and **1ag**), the yields decreased significantly, presumably as the result involving *N*-tosyl cleavage. Additionally, it was observed that, *N*-glutarimide (**1ah**), a highly reactive substrate under transition metal-catalyzed N-C activation conditions,^[19] led to a reduced yield.

Table 1 Optimization of the transamidation of *N*-benzyl-*N*-Boc-amide (**1a**) with pyrrolidine **2a**

Entry	CS ₂ CO ₃ /equiv.	Solvent	T/°C	Conv./% of 1a ^a	Yield ^a /% of 3a
1	2.0	DMF	100	41	33
2	1.0	DMF	100	70	48
3	0.2	DMF	100	85	63
4	0	DMF	100	84	81
5	0	THF	Reflux	97	83
6	0	CH ₃ CN	Reflux	79	78
7	0	Ether	Reflux	93	81
8	0	DCM	Reflux	7	6
9	0	Toluene	100	91	91
10	0	Toluene	80	54	50
11	0	Toluene	Reflux	99	97 (90) ^b
12 ^c	0	Toluene	Reflux	86	79

^a Conversion and yield were determined by GC using *n*-dodecane as an internal standard. ^b Yield in parenthesis refers to the isolated yield. ^c 1.0 equiv. of **2a** was used.

Scheme 2 Effect of *N*-substituents on amides

Reaction conditions: amide (0.1 mmol), **2a** (0.15 mmol), and toluene (0.5 mL) were heated with stirring under reflux for 12 h. Yields of the transamidation reaction were determined by GC analysis using an internal standard (yields in parentheses refer to *N*-activated amide starting material recovered from the reaction mixtures)

Using the above optimized reaction conditions and pyrrolidine **2a** as the model nucleophile, we set out to explore the reaction scope of acyl moiety, including evaluation of the effect of different substituents on the aryl rings (Scheme 3). Aryl amide bearing *ortho*-methyl substituents showed diminished reactivity as compared to analogous *meta*- and *para*-substrates (**3b** versus **3c** and **3d**). Such a finding leads us to infer that the present transamidation protocol is sensitive to steric effects. In contrast, it proved less susceptible to electronic effects in a diverse array of *para*-substituted *N*-benzyl-*N*-Boc-arylamides, including those bearing ester, fluoro, chloro, bromo, iodo, trifluoromethyl, nitro, and cyano groups (**3g**~**3n**). The method also proved amenable for heterocycles. For instance, fu-

ran-, thiophene- and benzo-thiophene-based substrates underwent transamidation in good yields (**3p**~**3r**). Furthermore, the present transamidation protocol was not restricted to (hetero)aryl substrate; the cinnamyl pyrrolidine amide **3s** was obtained in 82% yield. Alkyl amides such as phenylpropyl- and hexyl-amides furnished products **3t** and **3u** in 77% and 39% yields, respectively. Moreover, pyrrolidine-based cyclohexyl-amide (**3v**) could be accessed, albeit in a poor yield.

Scheme 3 Amide products accessed via the transamidation of the corresponding *N*-benzyl-*N*-Boc-amides

Reaction conditions: **1** (0.5 mmol), **2a** (0.75 mmol), and toluene (2.5 mL) were heated with stirring under reflux for 12 h. Yields refer to isolated yields (the yields in parentheses refer to initial amides recovered in the reaction mixtures)

Next, we sought to examine the generality of this transformation by testing a variety of aliphatic amines using *N*-benzyl-*N*-Boc-amide (**1a**) as the substrate (Scheme 4). Reaction of various cyclic secondary aliphatic amines occurred smoothly (**3w**~**3ad**). With azetidine as the nucleophile, the reaction furnished **3w** in 94% yield. This stands in contrast to its analogous azepane, where the large ring size severely diminished the yield (**3x**).^[20] Piperidine bearing α -methyl group was found much less effective, giving

3z in a low yield. It is of interest to note that the reaction is chemoselective. In the case of 4-hydroxypiperidine, where both *N*- and *O*-nucleophilic sites exist, the only *N*-acylated product **3ab** was obtained, along with considerable recovered starting amide. The low yield (33%) may reflect the detrimental effect of the alcohol moiety in 4-hydroxypiperidine on the reaction. In contrast to the cyclic secondary amines, acyclic *N*-methyl-4-methoxybenzylamine gave the transamidation product **3ae** in 29% yield. This low yield is attributed to the relatively low basicity and nucleophilicity of acyclic secondary amines, since the pK_a of protonated *N*-methyl-4-methoxybenzylamine was found to be 9.97.^[21]

Scheme 4 Products obtained when *N*-benzyl-*N*-Boc-amide (**1a**) was subject to the present procedure using a variety of amines. Reaction conditions: **1a** (0.5 mmol), **2** (0.75 mmol), and toluene (2.5 mL) were heated with stirring under reflux for 12 h. Yields refer to isolated yield (the yields in parentheses refer to the amide **1a** recovered in the reaction mixtures).

The substrate scope with respect to the primary aliphatic amine was also investigated. It was observed that various

primary amines, including linear and branched ones were compatible with the standard reaction conditions allowing access to secondary amides in 50%~88% yields (Scheme 4b). Unfortunately, attempts to transamidate using a sterically hindered *tert*-butylamine, as well as less nucleophilic aromatic amines, *e.g.* aniline, *p*-anisidine, and indoline, were unsuccessful. In these cases, full recovery of the starting amides was observed.

The potential application of the present transamidation procedure was showcased in the synthesis of antidepressant moclobemide, the first reversible inhibitor of monoamine oxidase (MAO) marketed by Roche.^[22] This one-step method delivered the desired amide **3ar** in 79% yield (Scheme 5a). Furthermore, the post-transformation of commercial drugs, *e.g.* pregabalin^[23] and amlodipine,^[24] which were used in the treatment of peripheral neuropathic pain and high blood pressure, respectively, could likewise be achieved. The corresponding products **3as** and **3at** were obtained in good yields as pro-drugs to provide an increased duration of active compound in the plasma (Schemes 5b and 5c).^[25]

Scheme 5 Synthesis of moclobemide and post-transformation of drugs.

Yields refer to isolated yields after column chromatography.

To the best of our knowledge, transamidation reactions that involve optically enriched substrates remain elusive.^[8] A recurring issue in conventional amidation processes is the ease of racemization.^[26] As a test of our methodology, the free amino-based amino esters **4a**~**4d** were employed as nucleophiles. Pleasingly, all the reactions proceeded smoothly without erosion of the stereochemical fidelity, albeit in low yields. On the other hand, the yields based on the recovered starting materials (brsm) were excellent. As shown in Scheme 6, the yields gradually decreased when the side chain of amino ester was changed from ethyl to *tert*-butyl (products **5e**~**5g**); presumably. This reflects steric effects on the nucleophilicity of amino acid derivatives.

To further explore the viability of the present approach with regard to stereochemical integrity, an amide **1w** with an epimerizable stereocenter was treated with pyrrolidine **2a** under the standard reaction conditions. Only a slightly reduced enantioselectivity was observed for **5h** (98% *ee*),^[8b] while for the transamidation between **1w** and 2-methoxyethylamine (**2m**), no appreciable loss of enantiopurity occurred during the course of the reaction.

Aryl esters are well-established as activating groups that can undergo O—C(acyl) cleavage under conditions of either transition metal-mediated or transition metal-free amidation.^[27,11b] A competition experiment was therefore performed to gauge selectivity, as shown in Scheme 7. Here, substrate **1x** bearing *N*-benzyl-*N*-Boc-amide group and phenolic ester, was subjected to our typical reaction protocol. Under these conditions, products **3au** and **3av** were obtained in 57% and 9% yields, respectively. Such a finding is consistent with the canonical thinking that activated esters are more reactive than *N*-benzyl-*N*-Boc-amides. While beyond the scope of the present study, in principle this distinction could be exploited to access to unsymmetrical terephthalamides, a scaffold that was widely found in natural products.^[28] For instance, when **3au** was treated with *n*-hexyl amine **2k**, the corresponding product **3aw** was obtained in 82% yield.

In an effort to obtain insights into the underlying mechanism of the present protocol, we performed an intermolecular competition experiment using two representative aliphatic amines while monitoring the kinetics of amidation by GC (Figure 1). In accord with expectations, the secondary amine pyrrolidine **2a** reacted rather quickly to afford **3a** in ca. 90% yield within 6 h. In contrast, the less nucleophilic primary amine *n*-hexanamine **2k** only provided a low yield (ca. 12%) of the corresponding product (**3af**) over the course of 12 hours. This finding, considered in concert with the lack of reactivity seen for less nucleophilic aromatic amines (vide supra), supports the conclusion that the nucleophilicity of the amine substrates and the reactivity seen in the present transamidation protocol are closely correlated.^[29] Furthermore, we determined the product yields when pyrrolidine **2a** was allowed to react with three differently substituted aryl amides under our transamidation conditions. As can be seen from Figure 1,

Scheme 6 Transamidation of enantioenriched amino esters and *N*-activated amides

Reaction conditions: **1** (0.5 mmol), amine **4** or **2** (0.75 mmol), and toluene (2.5 mL) were heated with stirring under reflux for 12 h. Yields refer to isolated yields (yields in parentheses are the original *N*-activated amides recovered in the reaction mixtures). The enantiomeric excess (*ee*) was determined by HPLC analysis using a stationary phase chiral column

Scheme 7 Selectivity tests involving amide **1x** that contains phenolic ester and an *N*-benzyl-*N*-Boc

Reaction conditions: **1x** (0.5 mmol), **2a** (0.75 mmol), and toluene (2.5 mL) were heated with stirring under reflux for 12 h. Yields refer to isolated yields

Figure 1 Kinetic profile for test transamidations

Reaction of **1a** with pyrrolidine **2a** (giving product **3a**) and *n*-hexanamine **2k** (giving product **3af**) under the standard conditions of the present study (top), and reaction of **1a** (R=H), **1d** (R=4-MeC₆H₄) and **1n** (R=4-NCC₆H₄) with pyrrolidine **2a** at 90 °C (bottom)

which depicts the yield versus time for the transamidation, a higher reaction rate was seen for the electron-deficient 4-cyano-benzamide **1n** (giving product **3n**) as compared to the corresponding electron-rich 4-methyl-benzamide **1d** and non-substituted benzamide **1a** (giving products **3d** and **3a**, respectively). Taken together, the data lead us to propose a nucleophilic acyl substitution mechanism that involves direct nucleophilic attack of amine on the *N*-activated amide.^[30] Moreover, we rationalize that the driving force for the catalyst- and additive-free transamidation lies in the nucleophilic addition of amine to the weakened *N*-benzyl-*N*-Boc-amide bond, and the formation of thermodynamic more stable amide product.^[31]

3 Conclusions

In summary, a protocol that permits the transamidation of *N*-benzyl-*N*-Boc-amides under transition metal-free and base-free conditions is presented. It provides access to a range of amides in moderate to good yields. An advantage of this method is that the chiral centers that are present in both amide and amine substrates are permitted. The readily availability of *N*-benzyl-*N*-Boc-amides coupled with the

broad substrate scope and functional group tolerance of present transamidation protocol leads us to predict that it could emerge as a useful complement to related methodologies and could also be used to allow ready access to value-added amides.

4 Experimental section

4.1 General information

All NMR spectra were acquired on a Bruker 600 MHz or a JEOL ECZ 400 MHz NMR spectrometer. ¹H NMR chemical shifts were recorded relative to TMS (δ 0.00) or residual protonated solvents (CDCl₃ δ 7.26). Proof of purity of new compounds is provided through the attached copies of ¹H NMR, ¹³C NMR, and ¹⁹F NMR spectra. Mass spectrometric data consistent with the proposed structures were also obtained.

4.2 General procedure for the transamidation of *N*-benzyl-*N*-Boc-amides

A 10-mL vial containing a magnetic stir bar was charged with *N*-activated amide (0.5 mmol), aliphatic amine (0.75 mmol) and 2.5 mL of toluene. The reaction mixture was heated with stirring under reflux overnight (*ca.* 12 h). After cooling, the volatiles were evaporated off under reduced pressure and the residue purified by flash chromatography over silica gel using ethyl acetate/petroleum ether (*V* : *V* = 5 : 1~1 : 1) as eluent to give product.

Phenyl(pyrrolidin-1-yl)methanone (3a):^[11a] Colorless oil (79 mg, 90% yield). ¹H NMR (400 MHz, CDCl₃) δ : 7.53~7.50 (m, 2H), 7.42~7.38 (m, 3H), 3.65 (t, *J*=7.0 Hz, 2H), 3.42 (t, *J*=6.6 Hz, 2H), 2.00~1.95 (m, 2H), 1.90~1.84 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ : 169.8, 137.3, 129.9, 128.3, 127.2, 49.7, 46.3, 26.5, 24.6; GC-MS *m/z*: 175.0.

Pyrrolidin-1-yl(*o*-tolyl)methanone (3b):^[32] Yellow oil (21 mg, 22% yield). ¹H NMR (600 MHz, CDCl₃) δ : 7.26~7.24 (m, 1H), 7.21~7.19 (m, 3H), 3.66 (t, *J*=7.0 Hz, 2H), 3.13 (t, *J*=6.8 Hz, 2H), 2.31 (s, 3H), 1.98~1.94 (m, 2H), 1.89~1.84 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ : 170.1, 138.0, 133.9, 130.5, 128.9, 126.0, 125.7, 48.5, 45.4, 26.2, 24.8, 19.1; GC-MS *m/z*: 189.0.

Pyrrolidin-1-yl(*m*-tolyl)methanone (3c):^[32] Yellow oil (59 mg, 63% yield). ¹H NMR (600 MHz, CDCl₃) δ : 7.33 (s, 1H), 7.29~7.26 (m, 2H), 7.22~7.21 (d, *J*=4.8 Hz, 1H), 3.64 (t, *J*=6.9 Hz, 2H), 3.42 (t, *J*=6.6 Hz, 2H), 2.37 (s, 3H), 1.98~1.93 (m, 2H), 1.89~1.84 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ : 170.1, 138.2, 137.3, 130.5, 128.2, 127.8, 124.1, 49.7, 46.2, 26.5, 24.6, 21.5; GC-MS *m/z*: 189.0.

Pyrrolidin-1-yl(*p*-tolyl)methanone (3d):^[32] Yellow solid (68 mg, 72% yield). m.p. 77~78 °C (Lit. 76~78 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.44~7.41 (m, 2H), 7.19 (d, *J*=7.6 Hz, 2H), 3.64 (t, *J*=7.0 Hz, 2H), 3.44 (t, *J*=6.6 Hz, 2H), 2.37 (s, 3H), 1.99~1.92 (m, 2H), 1.89~1.83 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ : 169.9, 140.0, 134.4, 128.9, 127.3, 49.8, 46.3, 26.5, 24.6, 21.5; GC-MS *m/z*: 189.0.

(4-Methoxyphenyl)(pyrrolidin-1-yl)methanone (**3e**):^[32] White solid (85 mg, 83% yield). m.p. 76~77 °C (Lit. 74~76 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.54~7.50 (m, 2H), 6.92~6.88 (m, 2H), 3.83 (s, 3H), 3.63 (t, *J*=6.8 Hz, 2H), 3.48 (t, *J*=6.4 Hz, 2H), 1.99~1.92 (m, 2H), 1.90~1.84 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.5, 160.9, 129.3, 113.5, 55.4, 49.9, 46.4, 26.6, 24.6; GC-MS *m/z*: 205.0.

(3,5-Dimethoxyphenyl)(pyrrolidin-1-yl)methanone (**3f**): Yellow oil (99 mg, 84% yield). ¹H NMR (400 MHz, CDCl₃) δ: 6.63 (d, *J*=2.4 Hz, 2H), 6.49 (t, *J*=2.2 Hz, 1H), 3.80 (s, 6H), 3.63 (t, *J*=7.0 Hz, 2H), 3.42 (t, *J*=6.6 Hz, 2H), 2.00~1.93 (m, 2H), 1.90~1.84 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.5, 160.8, 139.3, 105.1, 102.0, 55.6, 49.7, 46.3, 26.5, 24.6; HRMS (ESI) calcd for C₁₃H₁₈NO₃ [M+H]⁺: 236.1281, found 236.1281.

Methyl 4-(pyrrolidine-1-carbonyl)benzoate (**3g**):^[33] White solid (112 mg, 96% yield). m.p. 103~104 °C (Lit. 96~98 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.09~8.06 (m, 2H), 7.59~7.56 (m, 2H), 3.94 (s, 3H), 3.66 (t, *J*=6.8 Hz, 2H), 3.39 (t, *J*=6.6 Hz, 2H), 2.01~1.95 (m, 2H), 1.93~1.86 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 168.8, 166.5, 141.5, 131.3, 129.7, 127.1, 52.4, 49.5, 46.3, 26.5, 24.5; GC-MS *m/z*: 233.0.

(4-Fluorophenyl)(pyrrolidin-1-yl)methanone (**3h**):^[33] White solid (83 mg, 86% yield). m.p. 86~87 °C (Lit. 87~90 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.56~7.52 (m, 2H), 7.10~7.05 (m, 2H), 3.64 (t, *J*=7.0 Hz, 2H), 3.43 (t, *J*=6.6 Hz, 2H), 2.00~1.93 (m, 2H), 1.91~1.85 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 168.8, 163.56 (d, *J*=249.7 Hz), 133.3, 129.54 (d, *J*=8.6 Hz), 115.38 (d, *J*=21.7 Hz), 49.8, 46.4, 26.6, 24.6; ¹⁹F NMR (376 MHz, CDCl₃) δ: -110.4; GC-MS *m/z*: 193.0.

(4-Chlorophenyl)(pyrrolidin-1-yl)methanone (**3i**):^[32] White solid (100 mg, 96% yield). m.p. 70~71 °C (Lit. 68~70 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.47 (d, *J*=5.6 Hz, 2H), 7.37 (d, *J*=5.6 Hz, 2H), 3.64 (t, *J*=4.6 Hz, 2H), 3.42 (t, *J*=4.4 Hz, 2H), 1.99~1.94 (m, 2H), 1.91~1.86 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 168.7, 136.0, 135.6, 128.8, 128.6, 49.7, 46.4, 26.6, 24.6; GC-MS *m/z*: 209.0.

(4-Bromophenyl)(pyrrolidin-1-yl)methanone (**3j**):^[32] White solid (101 mg, 80% yield). m.p. 76~77 °C (Lit. 76~78 °C); ¹H NMR (600 MHz, CDCl₃) δ: 7.54 (d, *J*=8.4 Hz, 2H), 7.40 (d, *J*=8.4 Hz, 2H), 3.63 (t, *J*=6.9 Hz, 2H), 3.41 (t, *J*=6.9 Hz, 2H), 1.99~1.94 (m, 2H), 1.91~1.86 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ: 168.7, 136.1, 131.6, 129.0, 124.2, 49.7, 46.4, 26.5, 24.5; GC-MS *m/z*: 253.0.

(4-Iodophenyl)(pyrrolidin-1-yl)methanone (**3k**):^[34] Yellow oil (138 mg, 92% yield). ¹H NMR (600 MHz, CDCl₃) δ: 7.76~7.73 (m, 2H), 7.28~7.27 (m, 1H), 7.26~7.25 (m, 3H), 3.63 (t, *J*=6.9 Hz, 2H), 3.41 (t, *J*=6.6 Hz, 2H), 2.00~1.85 (m, 4H); ¹³C NMR (150 MHz, CDCl₃) δ: 168.9, 137.6, 136.7, 129.0, 96.3, 49.7, 46.4, 26.5, 24.6; GC-MS *m/z*: 301.0.

Cyclopentyl(4-(trifluoromethyl)phenyl)methanone

(**3l**):^[32] Yellow oil (120 mg, 99% yield). ¹H NMR (600 MHz, CDCl₃) δ: 7.67 (d, *J*=8.1 Hz, 1H), 7.63 (d, *J*=8.1 Hz, 1H), 3.66 (t, *J*=6.9 Hz, 2H), 3.39 (t, *J*=6.6 Hz, 2H), 2.01~1.96 (m, 2H), 1.92~1.88 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ: 168.4, 140.8, 131.7 (q, *J*=34.5 Hz), 127.6, 125.5 (q, *J*=3.8 Hz), 123.9 (q, *J*=272.3 Hz), 49.6, 46.4, 26.5, 24.5; ¹⁹F NMR (376 MHz, CDCl₃) δ: -62.9; GC-MS *m/z*: 243.0.

(4-Nitrophenyl)(pyrrolidin-1-yl)methanone (**3m**):^[33] Yellow solid (93 mg, 84% yield). m.p. 83~84 °C (Lit. 78~80 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.29~8.28 (m, 1H), 8.27~8.26 (m, 1H), 7.70~7.69 (m, 1H), 7.68~7.67 (m, 1H), 3.67 (t, *J*=7.0 Hz, 2H), 3.39 (t, *J*=6.4 Hz, 2H), 2.04~1.89 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.5, 143.2, 128.3, 123.8, 49.6, 46.5, 26.5, 24.5; GC-MS *m/z*: 220.0.

4-(Pyrrolidine-1-carbonyl)benzonitrile (**3n**):^[32] White solid (99 mg, 99% yield). m.p. 85~86 °C (Lit. 87~88 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.73~7.70 (m, 2H), 7.63~7.60 (m, 2H), 3.66 (t, *J*=6.8 Hz, 2H), 3.38 (t, *J*=6.6 Hz, 2H), 2.03~1.96 (m, 2H), 1.95~1.88 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.8, 141.5, 132.4, 127.9, 118.3, 113.7, 49.6, 46.5, 26.5, 24.5; GC-MS *m/z*: 200.0.

Naphthalen-2-yl(pyrrolidin-1-yl)methanone (**3o**):^[33] White solid (99 mg, 88% yield). m.p. 69~70 °C (Lit. 76~77 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.01 (s, 1H), 7.88~7.85 (m, 3H), 7.62 (dd, *J*=8.4, 1.6 Hz, 1H), 7.55~7.50 (m, 2H), 3.71 (t, *J*=7.2 Hz, 2H), 3.50 (t, *J*=6.6 Hz, 2H), 2.03~1.86 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.9, 134.7, 133.9, 132.7, 128.6, 128.2, 127.9, 127.2, 127.1, 126.7, 124.6, 49.9, 46.4, 26.6, 24.7; GC-MS *m/z*: 225.0.

Furan-2-yl(pyrrolidin-1-yl)methanone (**3p**):^[32] Yellow oil (61 mg, 74% yield). ¹H NMR (600 MHz, CDCl₃) δ: 7.51 (s, 1H), 7.06 (d, *J*=3.6 Hz, 1H), 6.49~6.48 (m, 1H), 3.83 (t, *J*=6.6 Hz, 2H), 3.65 (t, *J*=6.9 Hz, 2H), 2.01~1.98 (m, 2H), 1.92~1.88 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ: 158.2, 148.8, 144.1, 115.7, 111.4, 47.9, 47.1, 26.6, 23.8; GC-MS *m/z*: 165.0.

Pyrrolidin-1-yl(thiophen-2-yl)methanone (**3q**):^[32] White solid (54 mg, 60% yield). m.p. 71~72 °C (Lit. 72~73 °C); ¹H NMR (600 MHz, CDCl₃) δ: 7.52 (d, *J*=3.6 Hz, 1H), 7.47 (d, *J*=5.4 Hz, 1H), 7.08~7.07 (m, 1H), 3.78~3.77 (m, 2H), 3.69~3.67 (m, 2H), 2.02~1.99 (m, 2H), 1.96~1.93 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ: 161.9, 139.6, 129.61, 129.60, 127.2, 49.0, 47.4, 26.8, 24.2; GC-MS *m/z*: 181.0.

Benzo[*b*]thiophen-2-yl(pyrrolidin-1-yl)methanone (**3r**):^[35] White solid (52 mg, 45% yield). m.p. 90~91 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.87~7.81 (m, 2H), 7.70 (d, *J*=0.4 Hz, 1H), 7.43~7.36 (m, 2H), 3.84 (t, *J*=6.4 Hz, 2H), 3.71 (t, *J*=6.6 Hz, 2H), 2.04~1.95 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.4, 140.7, 139.6, 139.2, 126.09, 126.06, 124.99, 124.74, 122.5, 49.2, 47.6, 26.8, 24.2; GC-MS *m/z*: 231.0.

(*E*)-3-Phenyl-1-(pyrrolidin-1-yl)prop-2-en-1-one (**3s**):^[36] Colorless oil (82 mg, 82% yield). m.p. 95~96 °C (Lit.

97~98 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.69 (d, *J*=15.5 Hz, 1H), 7.55~7.52 (m, 2H), 7.40~7.34 (m, 3H), 6.72 (d, *J*=15.4 Hz, 1H), 3.66~3.58 (m, 4H), 2.05~1.98 (m, 2H), 1.94~1.87 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 164.8, 141.8, 135.5, 129.7, 128.9, 128.0, 119.0, 46.7, 46.2, 26.3, 24.5; GC-MS *m/z*: 201.0.

3-Phenyl-1-(pyrrolidin-1-yl)propan-1-one (**3t**):^[37] Colorless oil (78 mg, 77% yield). ¹H NMR (600 MHz, CDCl₃) δ: 7.30~7.27 (m, 2H), 7.24~7.18 (m, 3H), 3.47 (t, *J*=6.8 Hz, 2H), 3.29 (t, *J*=6.8 Hz, 2H), 2.99 (t, *J*=7.8 Hz, 2H), 2.56 (t, *J*=7.8 Hz, 2H), 1.91~1.86 (m, 2H), 1.84~1.80 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ: 170.9, 141.7, 128.60, 128.57, 126.2, 47.0, 46.0, 36.9, 31.4, 26.2, 24.5; GC-MS *m/z*: 203.0.

1-(Pyrrolidin-1-yl)hexan-1-one (**3u**):^[38] Colorless oil (33 mg, 39% yield). ¹H NMR (400 MHz, CDCl₃) δ: 3.48~3.39 (m, 4H), 2.27~2.23 (m, 2H), 1.98~1.91 (m, 2H), 1.88~1.81 (m, 2H), 1.69~1.61 (m, 2H), 1.36~1.29 (m, 4H), 0.90 (t, *J*=6.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 172.0, 46.7, 45.7, 34.9, 31.8, 26.2, 24.7, 24.5, 22.6, 14.1; GC-MS *m/z*: 169.0.

Cyclohexyl(pyrrolidin-1-yl)methanone (**3v**):^[39] Colorless oil (12 mg, 13% yield). ¹H NMR (400 MHz, CDCl₃) δ: 3.48~3.43 (m, 4H), 2.37~2.29 (m, 1H), 1.98~1.91 (m, 2H), 1.87~1.66 (m, 8H), 1.58~1.48 (m, 2H), 1.32~1.19 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 174.9, 46.4, 45.8, 43.0, 29.0, 26.3, 26.0, 25.9, 24.4; GC-MS *m/z*: 181.0.

Azetidin-1-yl(phenyl)methanone (**3w**):^[11d] Colorless oil (75 mg, 94% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.64~7.61 (m, 2H), 7.47~7.38 (m, 3H), 4.32~4.22 (m, 4H), 2.38~2.30 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.4, 133.4, 131.0, 128.4, 127.9, 53.5, 49.0, 16.2; GC-MS *m/z*: 161.0.

Azepan-1-yl(phenyl)methanone (**3x**):^[40] Yellow oil (58 mg, 57% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.41~7.36 (m, 5H), 3.68 (t, *J*=6.0 Hz, 2H), 3.36 (t, *J*=5.4 Hz, 2H), 1.87~1.85 (m, 2H), 1.68~1.60 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 171.7, 137.4, 129.1, 128.5, 126.5, 49.8, 46.4, 29.6, 27.9, 27.4, 26.6; GC-MS *m/z*: 203.0.

(4-Methoxypiperidin-1-yl)(phenyl)methanone (**3y**): Yellow oil (67 mg, 61% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.42~7.38 (m, 5H), 4.05 (s, 1H), 3.61~3.46 (m, 3H), 3.37 (s, 3H), 3.21 (s, 1H), 1.98~1.68 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.4, 136.1, 129.6, 128.5, 126.8, 75.4, 55.8, 44.8, 39.4, 31.3, 30.2; HRMS (ESI) calcd for C₁₃H₁₈NO₂: 220.1332, found 220.1331.

(2-Methylpiperidin-1-yl)(phenyl)methanone (**3z**):^[41] Colorless oil (25 mg, 25% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.43~7.38 (m, 3H), 7.33~7.28 (m, 2H), 4.35 (s, 1H), 3.83 (d, *J*=12.1 Hz, 1H), 2.99~2.92 (m, 1H), 1.69~1.54 (m, 4H), 1.50~1.45 (m, 1H), 1.43~1.32 (m, 1H), 1.17 (d, *J*=6.9 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 169.8, 137.9, 129.4, 128.8, 126.7, 47.2, 30.4, 26.0, 19.0, 16.4; GC-MS *m/z*: 203.0.

(4-Methylpiperidin-1-yl)(phenyl)methanone (**3aa**):^[42] Yellow oil (69 mg, 68% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.42~7.38 (m, 5H), 4.22 (s, 1H), 4.00~3.94

(m, 1H), 3.66 (s, 1H), 3.29 (d, *J*=70.4 Hz, 2H), 2.05~1.49 (m, 7H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.4, 136.7, 129.5, 128.5, 126.9, 48.2, 42.6, 34.9, 34.0, 31.3, 21.9; GC-MS *m/z*: 203.0.

(4-Hydroxypiperidin-1-yl)(phenyl)methanone (**3ab**):^[43] White solid (34 mg, 33% yield). m.p. 89~90 °C (Lit. 84~86 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.42~7.36 (m, 5H), 4.20 (s, 1H), 3.96~7.90 (m, 1H), 3.64 (s, 1H), 3.36~3.17 (m, 2H), 2.29 (s, 1H), 2.00~1.79 (m, 2H), 1.61~1.43 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.6, 136.1, 129.7, 128.6, 126.7, 67.0, 45.0, 39.5, 34.7, 34.0; GC-MS *m/z*: 205.0.

Morpholino(phenyl)methanone (**3ac**):^[44] Yellow oil (47 mg, 49% yield); ¹H NMR (600 MHz, CDCl₃) δ: 7.41~7.39 (m, 5H), 3.78~3.46 (m, 8H); ¹³C NMR (150 MHz, CDCl₃) δ: 170.6, 135.4, 130.0, 128.7, 127.2, 67.0, 48.3, 42.6; GC-MS *m/z*: 191.0.

(4-Methylpiperazin-1-yl)(phenyl)methanone (**3ad**):^[44] Yellow oil (90 mg, 88% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.40 (s, 5H), 3.81 (s, 2H), 3.44 (s, 2H), 2.49 (s, 2H), 2.35 (s, 2H), 2.32 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.4, 135.9, 129.8, 128.6, 127.2, 55.4, 54.9, 47.8, 46.2, 42.2; GC-MS *m/z*: 204.0.

N-(4-Methoxybenzyl)-*N*-methylbenzamide (**3ae**):^[45] Colorless oil (37 mg, 29% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.46~7.37 (m, 5H), 7.30 (d, *J*=8.2 Hz, 1H), 7.08 (d, *J*=8.0 Hz, 1H), 6.89 (d, *J*=7.7 Hz, 2H), 4.69 (s, 1H), 4.44 (s, 1H), 3.81 (s, 3H), 3.01~2.84 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 159.2, 136.5, 129.7, 128.55, 128.50, 128.2, 127.1, 126.9, 114.2, 55.4, 54.7, 50.3, 36.9, 33.1; GC-MS *m/z*: 255.0.

N-Hexylbenzamide (**3af**):^[16e] White solid (87 mg, 85% yield). m.p. 42~43 °C (Lit. 43~44 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.78~7.75 (m, 2H), 7.51~7.40 (m, 3H), 6.21 (s, 1H), 3.47~3.42 (m, 2H), 1.65~1.57 (m, 2H), 1.42~1.29 (m, 6H), 0.91~0.88 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.7, 135.0, 131.4, 128.6, 127.0, 40.2, 31.6, 29.8, 26.8, 22.7, 14.1; GC-MS *m/z*: 205.0.

N-Neopentylbenzamide (**3ag**):^[46] White solid (72 mg, 75% yield). m.p. 112~113 °C (Lit. 113~114 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.78~7.76 (m, 2H), 7.52~7.48 (m, 1H), 7.46~7.42 (m, 2H), 6.23 (s, 1H), 3.28 (d, *J*=6.4 Hz, 2H), 0.99 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.8, 135.2, 131.5, 128.8, 127.0, 51.1, 32.3, 27.4; GC-MS *m/z*: 191.0.

N-(2-Methoxyethyl)benzamide (**3ah**):^[47] Colorless oil (78 mg, 88% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.80~7.77 (m, 2H), 7.52~7.41 (m, 3H), 6.58 (s, 1H), 3.68~3.64 (m, 2H), 3.58~3.56 (m, 2H), 3.39 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.6, 134.6, 131.6, 128.7, 127.1, 71.3, 59.0, 39.8; GC-MS *m/z*: 179.0.

N-(But-3-en-1-yl)benzamide (**3ai**):^[32] Yellow oil (53 mg, 61% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.77~7.74 (m, 2H), 7.51~7.46 (m, 1H), 7.43~7.38 (m, 2H), 6.47 (s, 1H), 5.88~5.77 (m, 1H), 5.17~5.09 (m, 2H), 3.54~3.49 (m, 2H), 2.40~2.34 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.7, 135.4, 134.8, 131.4, 128.6, 126.9,

117.4, 38.9, 33.8; GC-MS m/z : 175.0.

t-Butyl 4-benzamidobutanoate (**3aj**): White solid (67 mg, 51% yield). m.p. 49~50 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.81~7.78 (m, 2H), 7.51~7.47 (m, 1H), 7.45~7.40 (m, 2H), 6.69 (s, 1H), 3.52~3.48 (m, 2H), 2.37 (t, $J=6.9$ Hz, 2H), 1.96~1.89 (m, 2H), 1.44 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 173.5, 167.6, 134.6, 131.5, 128.6, 127.0, 80.9, 40.0, 33.4, 28.2, 24.4; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{21}\text{NO}_3$ $[\text{M}+\text{H}]^+$: 286.1414, found 286.1420.

N-(4-Methoxybenzyl)benzamide (**3ak**):^[3c] Yellow solid (74 mg, 61% yield). m.p. 95~96 °C (Lit. 98~99 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.79~7.76 (m, 2H), 7.51~7.47 (m, 1H), 7.44~7.39 (m, 2H), 7.30~7.27 (m, 2H), 6.90~6.86 (m, 2H), 6.40 (s, 1H), 4.58~4.57 (m, 2H), 3.80 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.4, 159.3, 134.6, 131.6, 130.4, 129.5, 128.7, 127.1, 114.3, 55.5, 43.8; GC-MS m/z : 241.0.

N-Cyclobutylbenzamide (**3al**):^[48] Yellow solid (55 mg, 63% yield). m.p. 116~117 °C (Lit. 119~121 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.77~7.74 (m, 2H), 7.51~7.40 (m, 3H), 6.25 (s, 1H), 4.63~4.57 (m, 1H), 2.48~2.41 (m, 2H), 1.99~1.94 (m, 2H), 1.81~1.77 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.7, 134.8, 131.5, 128.7, 127.0, 45.3, 31.5, 15.3; GC-MS m/z : 175.0.

N-Cyclopentylbenzamide (**3am**):^[49] White solid (54 mg, 57% yield). m.p. 161~162 °C (Lit. 158~160 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.76~7.73 (m, 2H), 7.51~7.47 (m, 1H), 7.45~7.40 (m, 2H), 6.06 (s, 1H), 4.44~4.38 (m, 1H), 2.14~2.05 (m, 2H), 1.76~1.64 (m, 2H), 1.54~1.46 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.3, 135.1, 131.4, 128.7, 1276.9, 77.5, 51.8, 33.4, 23.9; GC-MS m/z : 189.0.

N-Cyclohexylbenzamide (**3an**):^[49] White solid (65 mg, 64% yield). m.p. 147~149 °C (Lit. 145~147 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.77~7.74 (m, 2H), 7.51~7.47 (m, 1H), 7.49~7.40 (m, 2H), 5.96 (s, 1H), 4.03~3.94 (m, 1H), 2.07~2.00 (m, 2H), 1.79~1.73 (m, 2H), 1.69~1.63 (m, 1H), 1.49~1.38 (m, 2H), 1.29~1.18 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.8, 135.3, 131.4, 128.7, 127.0, 48.8, 33.4, 25.7, 25.0; GC-MS m/z : 203.0.

N-(4-Methylcyclohexyl)benzamide (**3ao**):^[50] Yellow oil (66 mg, 61% yield). ^1H NMR (400 MHz, CDCl_3) δ : 7.76~7.73 (m, 2H), 7.51~7.46 (m, 1H), 7.45~7.40 (m, 2H), 5.89 (s, 1H), 3.96~3.87 (m, 1H), 2.10~2.06 (m, 2H), 1.78~1.74 (m, 2H), 1.60~1.58 (m, 1H), 1.42~1.06 (m, 6H), 0.92 (d, $J=6.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.9, 135.2, 131.4, 128.7, 126.9, 49.1, 34.0, 33.4, 32.2, 22.3; GC-MS m/z : 217.0.

N-Cycloheptylbenzamide (**3ap**):^[49] White solid (56 mg, 52% yield). m.p. 125~126 °C (Lit. 127~128 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.76~7.73 (m, 2H), 7.51~7.46 (m, 1H), 7.44~7.40 (m, 2H), 6.03 (s, 1H), 4.21~4.13 (m, 1H), 2.07~2.01 (m, 2H), 1.71~1.63 (m, 4H), 1.61~1.51 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.5, 135.3, 131.4, 128.7, 126.9, 51.0, 35.3, 28.2, 24.3; GC-MS m/z : 217.0.

N-Cyclooctylbenzamide (**3aq**):^[49] White solid (58 mg,

50% yield). m.p. 104~105 °C (Lit. 102~104 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.76~7.73 (m, 2H), 7.50~7.46 (m, 1H), 7.44~7.40 (m, 2H), 6.06 (s, 1H), 4.24~4.16 (m, 1H), 1.97~1.89 (m, 2H), 1.72~1.60 (m, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.4, 135.3, 131.3, 128.7, 126.9, 50.0, 32.5, 27.3, 25.6, 23.9; GC-MS m/z : 231.0.

4-Chloro-*N*-(2-morpholinoethyl)benzamide (**3ar**):^[32] White solid (106 mg, 79% yield). m.p. 137~138 °C (Lit. 135~137 °C); ^1H NMR (600 MHz, CDCl_3) δ : 7.73~7.71 (m, 2H), 7.43~7.42 (m, 2H), 6.66 (s, 1H), 3.74~3.73 (m, 4H), 3.57~3.54 (m, 2H), 2.62~2.60 (m, 2H), 2.52~2.51 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 166.5, 137.8, 133.1, 129.0, 128.5, 67.1, 57.0, 53.5, 36.2; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{18}\text{ClN}_2\text{O}_2$ $[\text{M}+\text{H}]^+$: 269.1051, found 269.1054.

Methyl (*S*)-3-(benzamidomethyl)-5-methylhexanoate (**3as**): Colorless oil (48 mg, 35% yield). ^1H NMR (600 MHz, CDCl_3) δ : 7.79 (d, $J=7.3$ Hz, 2H), 7.49 (t, $J=7.1$ Hz, 1H), 7.43 (t, $J=7.3$ Hz, 2H), 6.78 (s, 1H), 3.63 (s, 3H), 3.56~3.52 (m, 1H), 3.36~3.32 (m, 1H), 2.44~2.40 (m, 1H), 2.36~2.32 (m, 1H), 1.73~1.68 (m, 1H), 1.28~1.24 (m, 2H), 1.21~1.17 (m, 1H), 0.92 (dd, $J=16.2, 6.3$ Hz, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 174.5, 167.5, 134.5, 131.5, 128.7, 127.0, 51.9, 44.4, 42.1, 38.0, 33.3, 25.4, 22.8, 22.7; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{24}\text{NO}_3$ $[\text{M}+\text{H}]^+$: 278.1751, found 278.1759.

3-Ethyl-methyl 2-((2-benzamidoethoxy)methyl)-4-(2-chlorophenyl)-6-methyl-1,4-dihydropyridine-3,5-dicarboxylate (**3at**): White solid (192 mg, 75% yield). m.p. 189~190 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.82~7.79 (m, 2H), 7.56~7.51 (m, 1H), 7.48~7.44 (m, 2H), 7.36 (dd, $J=7.7, 1.7$ Hz, 1H), 7.25 (s, 1H), 7.22 (dd, $J=7.9, 1.4$ Hz, 1H), 7.14~7.08 (m, 1H), 7.06~7.00 (m, 1H), 6.45 (s, 1H), 5.39 (s, 1H), 4.81~4.68 (m, 2H), 4.10~3.98 (m, 2H), 3.82~3.70 (m, 4H), 3.61 (s, 3H), 2.32 (s, 3H), 1.18 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 168.2, 168.1, 167.2, 145.8, 145.1, 144.6, 134.2, 132.2, 131.6, 131.4, 129.1, 128.5, 127.4, 127.0, 126.9, 103.6, 101.6, 70.5, 67.9, 59.8, 50.7, 39.7, 37.0, 19.1, 14.2; HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{29}\text{ClN}_2\text{NaO}_6$ $[\text{M}+\text{Na}]^+$: 535.1606, found 535.1608.

Ethyl benzoyl-*L*-valinate (**5a**): White solid (43 mg, 35% yield, 99% ee). m.p. 61~62 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.83~7.80 (m, 2H), 7.55~7.50 (m, 1H), 7.48~7.43 (m, 2H), 6.65 (d, $J=8.7$ Hz, 1H), 4.80~4.76 (m, 1H), 4.30~4.19 (m, 2H), 2.34~2.24 (m, 1H), 1.31 (t, $J=7.1$ Hz, 3H), 1.01~0.99 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.3, 167.4, 134.4, 131.9, 128.8, 127.2, 61.6, 57.5, 31.9, 19.1, 18.0, 14.4; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{20}\text{NO}_3$ $[\text{M}+\text{H}]^+$: 250.1438, found 250.1439; HPLC (Daicel CHIRALPAK IB, V (hexane) : V (isopropanol) = 90 : 10, Flow rate = 1 mL/min, λ = 225 nm): t_R = 11.369 min (minor enantiomer), t_R = 12.265 min (major enantiomer).

Ethyl (4-chlorobenzoyl)-*L*-valinate (**5b**): White solid (56 mg, 40% yield, 99% ee). m.p. 67~68 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.77~7.73 (m, 2H), 7.44~7.41 (m, 2H),

6.61~6.59 (m, 1H), 4.76~4.74 (m, 1H), 4.29~4.20 (m, 2H), 2.31~2.26 (m, 1H), 1.33~1.30 (m, 3H), 1.02~0.92 (m, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.3, 166.4, 138.1, 132.7, 129.0, 128.6, 61.6, 57.6, 31.8, 19.1, 18.1, 14.4; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{19}\text{ClNO}_3$ $[\text{M} + \text{H}]^+$: 284.1048, found 284.1049; HPLC (Daicel CHIRALPAK IB, $V(\text{hexane}) : V(2\text{-propanol}) = 90 : 10$, Flow rate = 0.5 mL/min, $\lambda = 206$ nm): $t_{\text{R}} = 10.236$ min (minor enantiomer), $t_{\text{R}} = 12.793$ min (major enantiomer).

Ethyl (4-bromobenzoyl)-L-valinate (**5c**): White solid (64 mg, 39% yield, 99% ee). m.p. 69~70 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.69~7.67 (m, 2H), 7.60~7.58 (m, 2H), 6.60 (d, $J = 8.6$ Hz, 1H), 4.75~4.74 (m, 1H), 4.29~4.20 (m, 2H), 2.31~2.26 (m, 1H), 1.31 (t, $J = 7.1$ Hz, 3H), 1.02~0.98 (m, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.2, 166.4, 133.2, 132.0, 128.8, 126.6, 61.6, 57.6, 31.8, 19.1, 18.1, 14.4; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{19}\text{BrNO}_3$ $[\text{M} + \text{H}]^+$: 328.0543, found 328.0550; HPLC (Daicel CHIRALPAK IB, $V(\text{hexane}) : V(2\text{-propanol}) = 90 : 10$, Flow rate = 0.5 mL/min, $\lambda = 240$ nm): $t_{\text{R}} = 10.496$ min (minor enantiomer), $t_{\text{R}} = 13.098$ min (major enantiomer).

Ethyl (4-cyanobenzoyl)-L-valinate (**5d**): White solid (57 mg, 42% yield, 99% ee). m.p. 87~88 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.92~7.89 (m, 2H), 7.77~7.73 (m, 2H), 6.75~6.69 (m, 1H), 4.76~4.73 (m, 1H), 4.30~4.21 (m, 2H), 2.32~2.26 (m, 1H), 1.33~1.28 (m, 3H), 1.03~0.98 (m, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.0, 165.7, 138.3, 132.6, 127.9, 118.1, 115.4, 61.8, 57.8, 31.8, 19.1, 18.0, 14.4; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$: 275.1390, found 275.1399; HPLC (Daicel CHIRALPAK IB, $V(\text{hexane}) : V(2\text{-propanol}) = 80 : 20$, Flow rate = 1 mL/min, $\lambda = 205$ nm): $t_{\text{R}} = 6.299$ min (minor enantiomer), $t_{\text{R}} = 8.978$ min (major enantiomer).

Methyl (S)-2-(4-cyanobenzamido)butanoate (**5e**): White solid (55 mg, 45% yield, 99% ee). m.p. 134~135 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.93~7.90 (m, 2H), 7.77~7.75 (m, 2H), 6.75 (d, $J = 7.7$ Hz, 1H), 4.81~4.77 (m, 1H), 3.81 (s, 3H), 2.08~2.01 (m, 1H), 1.90~1.83 (m, 1H), 0.97 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.9, 165.4, 138.0, 132.6, 127.9, 118.1, 115.5, 54.0, 52.8, 25.8, 9.6; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{15}\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$: 247.1077, found 247.1080; HPLC (Daicel CHIRALPAK IB, $V(\text{hexane}) : V(2\text{-propanol}) = 90 : 10$, Flow rate = 1 mL/min, $\lambda = 237$ nm): $t_{\text{R}} = 17.056$ min (minor enantiomer), $t_{\text{R}} = 23.356$ min (major enantiomer).

Methyl (S)-2-(4-cyanobenzamido)-2-cyclohexylacetate (**5f**): Yellow oil (60 mg, 40% yield, 99% ee). ^1H NMR (400 MHz, CDCl_3) δ : 7.93~7.90 (m, 2H), 7.77~7.74 (m, 2H), 6.71 (d, $J = 8.6$ Hz, 1H), 4.77~4.74 (m, 1H), 3.79 (s, 3H), 1.96~1.87 (m, 1H), 1.81~1.74 (m, 3H), 1.70~1.66 (m, 2H), 1.32~1.07 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.5, 165.5, 138.1, 132.6, 127.9, 118.1, 115.4, 57.5, 52.6, 41.4, 29.7, 28.5, 26.0; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{21}\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$: 301.1547, found 301.1548; HPLC (Daicel CHIRALPAK IB, $V(\text{hexane}) : V(2\text{-propanol}) = 80 : 20$, Flow rate = 1 mL/min, $\lambda = 237$ nm): $t_{\text{R}} = 7.519$ min (minor enantiomer), $t_{\text{R}} = 11.591$ min (major enantio-

mer).

Methyl (S)-2-(4-cyanobenzamido)-3,3-dimethylbutanoate (**5g**): Colorless oil (35 mg, 25% yield, 99% ee). ^1H NMR (600 MHz, CDCl_3) δ : 7.91~7.89 (m, 2H), 7.77~7.76 (m, 2H), 6.67 (d, $J = 9.3$ Hz, 1H), 4.69 (d, $J = 9.3$ Hz, 1H), 3.78 (s, 3H), 1.06 (s, 9H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.0, 165.4, 138.1, 132.6, 127.8, 117.9, 115.4, 60.5, 52.1, 35.3, 26.7; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$: 275.1390, found 275.1393; HPLC (Daicel CHIRALPAK IB, $V(\text{hexane}) : V(2\text{-propanol}) = 90 : 10$, Flow rate = 1 mL/min, $\lambda = 236$ nm): $t_{\text{R}} = 11.040$ min (minor enantiomer), $t_{\text{R}} = 15.379$ min (major enantiomer).

(S)-Cyclohex-3-en-1-yl(pyrrolidin-1-yl)methanone (**5h**): White solid (31 mg, 34% yield, 98% ee). m.p. 64~65 °C; ^1H NMR (600 MHz, CDCl_3) δ : 5.75~5.66 (m, 2H), 3.51~3.46 (m, 4H), 2.63~2.56 (m, 1H), 2.40~2.31 (m, 1H), 2.19~2.03 (m, 3H), 2.00~1.93 (m, 2H), 1.89~1.72 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 174.7, 126.5, 126.1, 46.5, 45.9, 38.9, 27.7, 26.3, 25.5, 25.2, 24.4; HRMS (ESI) calcd for $\text{C}_{11}\text{H}_{18}\text{NO}$ $[\text{M} + \text{H}]^+$: 180.1383, found 180.1386; HPLC (Daicel CHIRALPAK IC, $V(\text{hexane}) : V(2\text{-propanol}) = 90 : 10$, Flow rate = 1 mL/min, $\lambda = 208$ nm): $t_{\text{R}} = 37.545$ min (minor enantiomer), $t_{\text{R}} = 38.937$ min (major enantiomer).

(S)-N-(2-Methoxyethyl)cyclohex-3-ene-1-carboxamide (**5i**): Colorless oil (30 mg, 33% yield, 99% ee). ^1H NMR (400 MHz, CDCl_3) δ : 5.94 (s, 1H), 5.70 (d, $J = 2.2$ Hz, 2H), 3.47~3.45 (m, 4H), 3.36 (s, 3H), 2.40~2.30 (m, 1H), 2.27~2.21 (m, 1H), 2.13~2.07 (m, 2H), 1.95~1.89 (m, 1H), 1.77~1.67 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 176.0, 126.9, 125.5, 71.4, 58.9, 41.4, 39.2, 28.2, 25.9, 24.8; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{18}\text{NO}_2$ $[\text{M} + \text{H}]^+$: 184.1332, found 184.1339; HPLC (Daicel CHIRALPAK OJ, $V(\text{hexane}) : V(2\text{-propanol}) = 95 : 5$, Flow rate = 0.5 mL/min, $\lambda = 205$ nm): $t_{\text{R}} = 15.959$ min (minor enantiomer), $t_{\text{R}} = 16.814$ min (major enantiomer).

t-Butyl benzyl(4-(pyrrolidine-1-carbonyl)benzoyl)carbamate (**3au**): White solid (117 mg, 57% yield). m.p. 132~133 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.544~7.543 (m, 4H), 7.44~7.41 (m, 2H), 7.36~7.32 (m, 2H), 7.29~7.27 (m, 1H), 4.99 (s, 2H), 3.65 (t, $J = 6.9$ Hz, 2H), 3.39 (t, $J = 6.6$ Hz, 2H), 2.00~1.93 (m, 2H), 1.92~1.85 (m, 2H), 1.15 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.5, 168.7, 153.3, 139.7, 139.0, 137.8, 128.6, 128.3, 127.6, 127.5, 127.1, 83.7, 49.7, 49.0, 46.4, 27.5, 26.6, 24.5; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{29}\text{N}_2\text{O}_4$ $[\text{M} + \text{H}]^+$: 409.2122, found 409.2120.

Phenyl 4-(pyrrolidine-1-carbonyl)benzoate (**3av**): White solid (13 mg, 9% yield). m.p. 130~131 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.26~8.23 (m, 2H), 7.66~7.63 (m, 2H), 7.47~7.42 (m, 2H), 7.31~7.27 (m, 1H), 7.24~7.21 (m, 2H), 3.68 (t, $J = 6.9$ Hz, 2H), 3.41 (t, $J = 6.6$ Hz, 2H), 2.03~1.96 (m, 2H), 1.95~1.88 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 168.7, 164.7, 150.9, 142.2, 130.7, 130.4, 129.7, 127.4, 126.2, 121.8, 49.6, 46.4, 26.5, 24.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{18}\text{NO}_3$ $[\text{M} + \text{H}]^+$: 296.1281, found 296.1288.

N-hexyl-4-(pyrrolidine-1-carbonyl)benzamide (**3aw**): Yellow solid (49 mg, 82% yield). m.p. 123~124 °C; ¹H NMR (600 MHz, CDCl₃) δ: 7.78 (d, *J*=8.2 Hz, 2H), 7.48 (d, *J*=8.2 Hz, 2H), 6.74 (t, *J*=5.7 Hz, 1H), 3.63 (t, *J*=7.0 Hz, 2H), 3.48~3.41 (m, 2H), 3.37 (t, *J*=6.7 Hz, 2H), 2.00~1.95 (m, 2H), 1.91~1.86 (m, 2H), 1.64~1.59 (m, 2H), 1.40~1.35 (m, 2H), 1.34~1.30 (m, 4H), 0.89 (t, *J*=3.5 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 169.0, 167.0, 139.7, 136.2, 127.2, 127.1, 49.6, 46.3, 40.3, 31.6, 29.6, 26.8, 26.4, 24.5, 22.7, 14.1; HRMS (ESI) calcd for C₁₈H₂₇N₂O₂ [M+H]⁺: 303.2067; found 303.2068.

Supporting Information ¹H NMR, ¹³C NMR and HRMS spectra for compounds synthesized in the manuscript. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>

References

- [1] (a) Greenberg, A. R.; Breneman, C. M.; Liebman, J. F. *The Amide Linkage: Structural Significance in Chemistry, Biochemistry, and Materials Science*, Wiley-VCH, New York, 2003.
(b) Hua, X.; Liu, N.; Fan, Z.; Zong, G.; Ma, Y.; Lei, K.; Yin, H.; Wang, G. *Chin. J. Org. Chem.* **2019**, 39, 2581 (in Chinese).
(华学文, 刘南南, 范志金, 宗广宁, 马翼, 雷康, 殷昊, 王桂清, 有机化学, **2019**, 39, 2581.)
(c) Zhong, L.; Jiang, T.; Zhang, F.; Fu, Q.; Liu, X.; Xu, T.; Ding, C.; Chen, J.; Yuan, J.; Tan, C. *Chin. J. Org. Chem.* **2019**, 39, 2655 (in Chinese).
(钟良坤, 江涛, 张帆, 付庆, 刘幸海, 许天明, 丁成荣, 陈杰, 袁静, 谭成侠, 有机化学, **2019**, 39, 2655.)
- [2] For reviews, see:
(a) Allen, C. L.; Williams, J. M. J. *Chem. Soc. Rev.* **2011**, 40, 3405.
(b) Pattabiraman, V. R.; Bode, J. W. *Nature* **2011**, 480, 471.
(c) Lundberg, H.; Tinnis, F.; Selander, N.; Adolfsson, H. *Chem. Soc. Rev.* **2014**, 43, 2714.
(d) Wan, J.-P.; Jing, Y. *Beilstein J. Org. Chem.* **2015**, 11, 2209.
(e) de Figueiredo, R. M.; Suppo, J.-S.; Campagne, J.-M. *Chem. Rev.* **2016**, 116, 12029.
(f) Ojeda-Porras, A.; Gamba-Sánchez, D. *J. Org. Chem.* **2016**, 81, 11548.
For recent examples on the synthesis of amide, see:
(g) Wang, Z.; Yang, L.; Liu, H.; Bao, W.; Tan, Y.; Wang, M.; Tang, Z.; He, W. *Chin. J. Org. Chem.* **2018**, 38, 2639 (in Chinese).
(王峥, 杨柳, 刘慧兰, 谭英芝, 包文虎, 汪明, 唐子龙, 何卫民, 有机化学, **2018**, 38, 2639.)
(h) Xie, L. Y.; Peng, S.; Liu, F.; Yi, J. Y.; Wang, M.; Tang, Z. L.; Xu, X. H.; He, W. M. *Adv. Synth. Catal.* **2018**, 360, 4259.
(i) Dissanayake, D. M. M. M.; Melville, A. D.; Vannucci, A. K. *Green Chem.* **2019**, 21, 3165.
(j) Tu, Z.; Du, Y.; Cao, X.; Liu, Y. *Adv. Synth. Catal.* **2019**, 361, 4989.
(k) Xie, L.-Y.; Hu, J.-L.; Song, Y.-X.; Jia, G.-K.; Lin, Y.-W.; He, J.-Y.; Cao, Z.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, 7, 19993.
(l) Yue, H.; Bao, P.; Wang, L.; Lü, X.; Yang, D.; Wang, H.; Wei, W. *Chin. J. Org. Chem.* **2019**, 39, 463 (in Chinese).
(岳会兰, 鲍鹏丽, 王雷, 吕晓霞, 杨道山, 王桦, 魏伟, 有机化学, **2019**, 39, 463.)
(m) Sheng, R.; Li, P.; Zhou, Z.; Hu, G.; Zhang, X. *Chin. J. Org. Chem.* **2020**, 40, 462 (in Chinese).
(圣戎, 李萍, 周志强, 胡贵文, 张小祥, 有机化学, **2020**, 40, 462.)
- [3] (a) Lanigan, R. M.; Sheppard, T. D. *Eur. J. Org. Chem.* **2013**, 7453.
(b) Zheng, J.-F.; Jin, L.-R.; Huang, P.-Q. *Org. Lett.* **2004**, 6, 1139.
(c) Tyrrell, E.; Brawn, P.; Carew, M.; Greenwood, I. *Tetrahedron Lett.* **2011**, 52, 369.
(d) Allen, C. L.; Atkinson, B. N.; Williams, J. M. J. *Angew. Chem. Int. Ed.* **2012**, 51, 1383.
(e) Rao, S. N.; Mohan, D. C.; Adimurthy, S. *Org. Lett.* **2013**, 15, 1496.
- [4] (a) Eldred, S. E.; Stone, D. A.; Gellman, S. H.; Stahl, S. S. *J. Am. Chem. Soc.* **2003**, 125, 3422.
(b) Hoerter, J. M.; Otte, K. M.; Gellman, S. H.; Cui, Q.; Stahl, S. S. *J. Am. Chem. Soc.* **2008**, 130, 647.
- [5] Bon, E.; Bigg, D. C. H.; Bertrand, G. *J. Org. Chem.* **1994**, 59, 4035.
- [6] (a) Gotor, V.; Brieva, R.; González, C.; Rebollo, F. *Tetrahedron* **1991**, 47, 9207.
(b) Sergeeva, M. V.; Mozhaev, V. V.; Rich, J. O.; Khmel'nitsky, Y. L. *Biotechnol. Lett.* **2000**, 22, 1419.
- [7] (a) Hie, L.; Fine Nathel, N. F.; Shah, T. K.; Baker, E. L.; Hong, X.; Yang, Y.-F.; Liu, P.; Houk, K. N.; Garg, N. K. *Nature* **2015**, 524, 79.
(b) Weires, N. A.; Baker, E. L.; Garg, N. K. *Nat. Chem.* **2016**, 8, 75.
- [8] (a) Baker, E. L.; Yamano, M. M.; Zhou, Y.; Anthony, S. M.; Garg, N. K. *Nat. Commun.* **2016**, 7, 11554.
(b) Dander, J. E.; Baker, E. L.; Garg, N. K. *Chem. Sci.* **2017**, 8, 6433.
- [9] (a) Meng, G.; Lei, P.; Szostak, M. *Org. Lett.* **2017**, 19, 2158.
(b) Shi, S.; Szostak, M. *Chem. Commun.* **2017**, 53, 10584.
- [10] Li, G.; Szostak, M. *Chem. Rec.* **2020**, 20, 649.
- [11] (a) Liu, Y.; Shi, S.; Achtenhagen, M.; Liu, R.; Szostak, M. *Org. Lett.* **2017**, 19, 1614.
(b) Li, G.; Szostak, M. *Nat. Commun.* **2018**, 9, 4165.
(c) Liu, Y.; Achtenhagen, M.; Liu, R.; Szostak, M. *Org. Biomol. Chem.* **2018**, 16, 1322.
(d) Li, G.; Ji, C.-L.; Hong, X.; Szostak, M. *J. Am. Chem. Soc.* **2019**, 141, 11161.
- [12] Guo, W.; Huang, J.; Wu, H.; Liu, T.; Luo, Z.; Jian, J.; Zeng, Z. *Org. Chem. Front.* **2018**, 5, 2950.
- [13] Verho, O.; Pourghasemi Lati, M.; Oschmann, M. *J. Org. Chem.* **2018**, 83, 4464.
- [14] Rahman, M. M.; Li, G.; Szostak, M. *J. Org. Chem.* **2019**, 84, 12091.
- [15] Ramkumar, R.; Chandrasekaran, S. *Synthesis* **2018**, 51, 921.
- [16] Other recent methods referring to the transamidation of amides (a) Cheung, C. W.; Ploeger, M. L.; Hu, X. *ACS Catal.* **2017**, 7, 7092.
(b) Cheung, C. W.; Ma, J.-A.; Hu, X. *J. Am. Chem. Soc.* **2018**, 140, 6789.
(c) Mishra, A.; Singh, S.; Srivastava, V. *Asian J. Org. Chem.* **2018**, 7, 1600.
(d) Ghosh, T.; Jana, S.; Dash, J. *Org. Lett.* **2019**, 21, 6690.
(e) Sureshbabu, P.; Azeez, S.; Chaudhary, P.; Kandasamy, J. *Org. Biomol. Chem.* **2019**, 17, 845.
(f) Chen, J.; Xia, Y.; Lee, S. *Org. Lett.* **2020**, 22, 3504.
- [17] Ye, D.; Liu, Z.; Chen, H.; Sessler, J. L.; Lei, C. *Org. Lett.* **2019**, 21, 6888.
- [18] (a) Galli, C. *Org. Prep. Proced. Int.* **1992**, 24, 285.
(b) Salvatore, R. N.; Nagle, A. S.; Jung, K. W. *J. Org. Chem.* **2002**, 67, 674.
- [19] (a) Meng, G.; Szostak, M. *Angew. Chem., Int. Ed.* **2015**, 54, 14518.
(b) Meng, G.; Szostak, M. *Org. Lett.* **2015**, 17, 4364.
(c) Shi, S.; Meng, G.; Szostak, M. *Angew. Chem., Int. Ed.* **2016**, 55, 6959.
(d) Shi, S.; Szostak, M. *Chem.-Eur. J.* **2016**, 22, 10420.
- [20] Graton, J.; Berthelot, M.; Laurence, C. *J. Chem. Soc., Perkin Trans. 2* **2001**, 2130.
- [21] The pK_a of protonated *N*-methyl-4-methoxybenzylamine was 9.97 from website: <http://ibond.nankai.edu.cn/pka/>.
- [22] Bonnet, U. *CNS Drug Rev.* **2002**, 8, 283.
- [23] Dworkin, R. H.; Kirkpatrick, P. *Nat. Rev. Drug Discovery* **2005**, 4, 455.
- [24] Sheraz, M. A.; Ahsan, S. F.; Khan, M. F.; Ahmed, S.; Ahmad, I. *J. Pharm.* **2016**, 2016, 8961621.
- [25] Bryans, J. S.; Williams, S. C.; Blakemore, D. C. *EP 1178034*, **2001**.

- [26] (a) Yamada, S.; Hongo, C.; Yoshioka, R.; Chibata, I. *J. Org. Chem.* **1983**, *48*, 843.
(b) Harry, L. G.; Pugnière, M.; Castro, B.; Previero, A. *Int. J. Peptide Protein Res.* **1993**, *41*, 323.
(c) Ramachandran, U.; Kumar, S.; Chawla, H. P. S. *Org. Prep. Proced. Int.* **2003**, *35*, 616.
(d) Wang, S. M.; Zhao, C.; Zhang, X.; Qin, H. L. *Org. Biomol. Chem.* **2019**, *17*, 4087.
- [27] (a) Cheung, C. W.; Ploeger, M. L.; Hu, X. *Nat. Commun.* **2017**, *8*, 14878.
(b) Halima, T. B.; Masson-Makdisi, J.; Newman, S. G. *Angew. Chem. Int. Ed.* **2018**, *57*, 12925.
- [28] (a) D, H. A.; Hang, Y. *EP 20050713917*, **2005**.
(b) Abergel, R. J.; Raymond, K. N. *Inorg. Chem.* **2006**, *45*, 3622.
- [29] Kanzian, T.; Nigst, T. A.; Maier, A.; Pichl, S.; Mayr, H. *Eur. J. Org. Chem.* **2009**, 6379.
- [30] Fox, J. M.; Dmitrenko, O.; Liao, L.-a.; Bach, R. D. *J. Org. Chem.* **2004**, *69*, 7317.
- [31] Adachi, S.; Kumagai, N.; Shibasaki, M. *Tetrahedron Lett.* **2018**, *59*, 1147.
- [32] Kumar, V.; Connon, S. J. *Chem. Commun.* **2017**, *53*, 10212.
- [33] Wang, X.-F.; Yu, S.-S.; Wang, C.; Xue, D.; Xiao, J. *Org. Biomol. Chem.* **2016**, *14*, 7028.
- [34] Chardon, A.; Mohy El Dine, T.; Legay, R.; De Paolis, M.; Rouden, J.; Blanchet, J. *Chem.-Eur. J.* **2017**, *23*, 2005.
- [35] Ning, X.-Q.; Lou, S.-J.; Mao, Y.-J.; Xu, Z.-Y.; Xu, D.-Q. *Org. Lett.* **2018**, *20*, 2445.
- [36] Zhang, B.; Feng, P.; Cui, Y.; Jiao, N. *Chem. Commun.* **2012**, *48*, 7280.
- [37] Chaudhari, M. B.; Bisht, G. S.; Kumari, P.; Gnanaprakasam, B. *Org. Biomol. Chem.* **2016**, *14*, 9215.
- [38] Mondal, A.; Subaramanian, M.; Nandakumar, A.; Balaraman, E. *Org. Lett.* **2018**, *20*, 3381.
- [39] Qian, C.; Zhang, X.; Zhang, Y.; Shen, Q. *J. Organomet. Chem.* **2010**, *695*, 747.
- [40] Zhu, M.; Fujita, K.-i.; Yamaguchi, R. *J. Org. Chem.* **2012**, *77*, 9102.
- [41] Sawant, D. N.; Bagal, D. B.; Ogawa, S.; Selvam, K.; Saito, S. *Org. Lett.* **2018**, *20*, 4397.
- [42] Zhu, J.; Zhang, Y.; Shi, F.; Deng, Y. *Tetrahedron Lett.* **2012**, *53*, 3178.
- [43] Soulard, V.; Villa, G.; Vollmar, D. P.; Renaud, P. *J. Am. Chem. Soc.* **2018**, *140*, 155.
- [44] Zhao, Q.; Li, H.; Wang, L. *Org. Biomol. Chem.* **2013**, *11*, 6772.
- [45] Hamada, S.; Sugimoto, K.; Iida, M.; Furuta, T. *Tetrahedron Lett.* **2019**, *60*, 151277.
- [46] Rossi, S. A.; Shimkin, K. W.; Xu, Q.; Mori-Quiroz, L. M.; Watson, D. A. *Org. Lett.* **2013**, *15*, 2314.
- [47] Yuan, Y.-C.; Kamaraj, R.; Bruneau, C.; Labasque, T.; Roisnel, T.; Gramage-Doria, R. *Org. Lett.* **2017**, *19*, 6404.
- [48] Chernykh, A. V.; Melnykov, K. P.; Tolmacheva, N. A.; Kondratov, I. S.; Radchenko, D. S.; Daniliuc, C. G.; Volochnyuk, D. M.; Ryabukhin, S. V.; Kuchkovska, Y. O.; Grygorenko, O. O. *J. Org. Chem.* **2019**, *84*, 8487.
- [49] Tran, B. L.; Li, B.; Driess, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2014**, *136*, 2555.
- [50] Wang, Y.; Hu, X.; Morales-Rivera, C. A.; Li, G.-X.; Huang, X.; He, G.; Liu, P.; Chen, G. *J. Am. Chem. Soc.* **2018**, *140*, 9678.

(Li, L.; Fan, Y.)