3,7-二甲基-7-羟基-2-辛烯-6-内酯类似物的合成及杀菌活性

董宏波 王卫伟 赵宇 刘鑫磊 王明安*

(中国农业大学应用化学系 农药创新研究中心 北京 100193)

摘要 以环氧化-内酯化和 Sharpless 不对称双羟基化反应作为关键步骤,以 55%~90% 的总收率实现了消旋及光活性 3,7-二甲基-7-羟基-2-辛烯-6-内酯类似物的合成。它们的结构经过 ^1H NMR, ^{13}C NMR, HR-ESI-MS 和 X 射线衍射的表征。对它们的杀菌活性进行了评价,结果表明活性最好的化合物 3-苯基-7-甲基-7-羟基-2-辛烯-6-内酯(4)和 3-(呋喃-2-基)-7-甲基-7-羟基-2-辛烯-6-内酯(5)对 6 种植物病原菌的 EC_{50} 值为 0.5~20.0 $\mu\text{g/mL}$, 可以作为先导结构进行进一步结构优化。

关键词 3,7-二甲基-7-羟基-2-辛烯-6-内酯; 环氧化-内酯化反应; 不对称双羟基化反应; 杀菌活性

Synthesis and Antifungal Activity of 3,7-Dimethyl-7-hydroxy-2-octen-6-olide Analogues

Dong, Hongbo Wang, Weiwei Zhao, Yu Liu, Xinlei Wang, Ming'an*

(Innovation Center of Pesticide Research, Department of Applied Chemistry, China Agricultural University, Beijing 100193)

Abstract The synthesis of racemic and optical 3,7-dimethyl-7-hydroxy-2-octen-6-olide analogues has been achieved via epoxidation-lactonization procedure and Sharpless asymmetric dihydroxylation as the key steps in 55%~90% overall yields, respectively. Their structures were fully characterized by ^1H NMR, ^{13}C NMR, HRMS data, and X-ray diffraction analysis. Their antifungal activities were evaluated, and showed that the EC_{50} values of the most active compounds 3-phenyl-7-methyl-7-hydroxy-2-octen-6-olide (4) and 3-(furan-2-yl)-7-methyl-7-hydroxy-2-octen-6-olide (5) were in the range of 0.5~20.0 $\mu\text{g/mL}$ against the tested six phytopathogens, and they were the lead structures to be optimized.

Keywords 3,7-dimethyl-7-hydroxy-2-octen-6-olide; epoxidation-lactonization; asymmetric dihydroxylation; antifungal activity

1 Introduction

The lactone motif plays an important role in the chemical communication between wide varieties of organisms.^[1]

3,7-Dimethyl-7-hydroxy-2-octen-6-olide (1) is a seven-membered lactone natural product that was first isolated from the honey bee fungal entomopathogen *Ascosphaera apis*, as well as the fruit of plant *Litsea cubeba* in Tibet, and it exhibited good antifungal and antioxidant activities.^[2-3] Structurally, it belongs to α,β -unsaturated lactone family. This class of natural products displays a wide range of biological properties such as antifungal, antimicrobial, and phytotoxic activities as well as cytotoxicity against human tumour cells.^[4-7] Their interesting structures and promising biological profiles have attracted significant attention from a number of researchers, and several interesting strategies for the synthesis of these compounds have

been reported and reviewed, but most of them focused on these natural products with five- or six-membered as well as eight-, nine-, ten- and eleven-membered lactone rings.^[1,8-13] The natural products with seven-membered lactone have rarely reported and synthesized in the literatures (Figure 1).^[14-22] Because they have a unique ring strain seven-membered lactone structure, there are very practical challenge to synthesis them. To date, a few synthetic strategies of natural and non-natural seven-membered lactones were reported by oxidative lactonization of diols,^[15,23] the palladium-catalyzed carbonylative annulation of internal alkynes,^[24] Co-catalyzed intermolecular reductive coupling/lactonization,^[23] bromolactonization using a sulfur-based zwitterionic organocatalyst,^[25] isocyanide-based multicomponent reaction/Wittig sequence starting from phosphonium salt precursors,^[26] Pd- and Rh-catalyzed hydroarylation of γ -(2-methoxycarbonylphenyl)-

* Corresponding author. E-mail: wangma@cau.edu.cn

Received October 9, 2020; revised November 20, 2020; published online December 18, 2020.

Project supported by the National Natural Science Foundation of China (Nos. 21772229, 21172254).

国家自然科学基金(Nos. 21772229, 21172254)资助项目。

propargylic alcohols,^[27] Pd-catalyzed sequential carbon-carbon bond cleavage/formation producing arylated benzolactones,^[28] Baeyer-Villiger oxidation of cyclic ketones,^[29] the ring-closing metathesis (RCM) synthesis of unsaturated lactones^[16] and enzymatic synthesis.^[17] These strategies, however, are not suitable for the synthesis of lactones containing an olefin or isopropanol side chain such as **1**. Therefore, there has been great requirement for highly efficient synthetic methods for natural and non-natural seven-membered lactones. For this scenario, we paid our attention to the total synthesis of natural products with seven-membered lactone moiety and their biological activities.^[30-32] In the previous paper, the synthesis of the racemic 3,7-dimethyl-7-hydroxy-2-octen-6-olide, 3,7-dimethyl-2,6-octa-dien-1,6-olide and 7-methyl-7-hydroxy-2,3-benzo[*c*]-octa-1,6-olide was reported in our laboratory.^[30-32] In order to get insights into the structure-activity relationship of this class of compounds, the racemic analogues **2**~**5** of CF₃, CH₃O, C₆H₅, furan-2-yl replaced methyl in **1** and benzolactones **6**~**14** (Figure 2), (6*R*)- and (6*S*)-3,7-dimethyl-7-hydroxy-2-octen-6-olide (*R*-**1** and *S*-**1**), and their CF₃ (*R*-**2** and *S*-**2**), OCH₃ (*R*-**3** and *S*-**3**) and 4,5-dihydrobenzo[*c*]oxepin-1(3*H*)-one (*R*-**6** and *S*-**6**) analogues were designed based on the biological profiles of benzolactone.^[33-38] So the strategies to conduct effectively synthesis of various seven-membered lactones with the desired configurations and high enantiomeric purity are highly required. The efficient synthesis of racemic lactones **2**~**14** by epoxidation-lactonization approaches and asymmetric synthesis of *R*-**1**-*S*-**6** via Sharpless asymmetric dihydroxylation, and their antifungal activity evaluation were first reported in this paper.

Figure 1 Representing natural occurring seven-membered lactones

1 Results and discussion

The synthesis of racemic 3,7-dimethyl-7-hydroxy-2-octen-6-olide (**1**) using *cis*-geraniol as the starting material and epoxidation-lactonization as the key steps (Table 1, Entry 1) was investigated in the previous paper.^[30-31] In an effort to promote the application of this epoxidation-lactonization procedure and obtain the analogues of **1**, compounds **2**~**5** were designed and synthesized (Scheme 1). The key olefin acids **2b**~**5b** were prepared from esters **2a**~**5a** by LiOH hydrolyzation, then the olefin acids **2b**~**5b** were epoxidized by peracetic acid to give epoxides **2c**~**5c**, which were cyclized to afford the seven-membered lactone compounds **2**~**5** in the presence of catalyst (+)-camphorsulfonic acid (CSA) in 60%~90% overall yields for two steps (Table 1, Entries 2~5). After accomplishing the synthesis of compounds **1**~**5**, compounds **6**~**14** were designed and synthesized to observe if the antifungal activity would improve significantly when the double bond was replaced with benzene ring (Scheme 2). The key olefin acids **6a**~**13a** were prepared from easily available material substituted *o*-methyl benzoic acid with 1-bromo-3-methylbut-2-ene in the presence of lithium diisopropylamine (LDA) under -78°C . The olefin acids **6a**~**13a** were also epoxidized by peracetic acid to give epoxides **6b**~**13b**, which were cyclized to produce seven-membered lactone compounds **6**~**13** in the presence of catalyst (+)-CSA as before in 70%~86% overall yields for 2 steps (Table 1, Entries 6~13). In Entries 14~16, the olefin acids without 2,3-double bond were also successfully transferred to lactones in overall 72%~80% yields. Then one methyl group of compound **13** was removed by BBr₃ to afford compound **14**. The seven-membered lactones were confirmed by the crystal structure of **10**, which was depicted in Figure 3.

Figure 2 Structures of designed compounds in this paper

bered lactone compounds **2**~**5** in the presence of catalyst (+)-camphorsulfonic acid (CSA) in 60%~90% overall yields for two steps (Table 1, Entries 2~5). After accomplishing the synthesis of compounds **1**~**5**, compounds **6**~**14** were designed and synthesized to observe if the antifungal activity would improve significantly when the double bond was replaced with benzene ring (Scheme 2). The key olefin acids **6a**~**13a** were prepared from easily available material substituted *o*-methyl benzoic acid with 1-bromo-3-methylbut-2-ene in the presence of lithium diisopropylamine (LDA) under -78°C . The olefin acids **6a**~**13a** were also epoxidized by peracetic acid to give epoxides **6b**~**13b**, which were cyclized to produce seven-membered lactone compounds **6**~**13** in the presence of catalyst (+)-CSA as before in 70%~86% overall yields for 2 steps (Table 1, Entries 6~13). In Entries 14~16, the olefin acids without 2,3-double bond were also successfully transferred to lactones in overall 72%~80% yields. Then one methyl group of compound **13** was removed by BBr₃ to afford compound **14**. The seven-membered lactones were confirmed by the crystal structure of **10**, which was depicted in Figure 3.

Scheme 1 Synthesis of **2**~**5** by epoxidation-lactonization

In order to get insights into the structure-activity relationship of this kind of compounds, some representing compounds were selected to prepare their optical isomers. First, In an effort to obtain the optical isomers of **1**, Shi's asymmetric epoxidation^[39] of *cis*-geranic acid was first utilized to afford the chiral epoxide **15**, then cyclization of corresponding epoxide **15** in the presence of (+)-CSA to yield (6*R*)-3,7-dimethyl-7-hydroxy-2-octen-6-olide (*R*-**1**) in 60% yield with only 47% *ee* (Scheme 3). When the amounts of catalyst and phase-transfer catalyst (Bu₄NH-SO₄) were increased such as in the reference,^[40] the enantiomeric excess (*e.e.*) value did not improve significantly. So, the low *e.e.* value and only one isomer obtained were not satisfied.

Table 1 Synthesis of seven-membered lactone via epoxidation-lactonization strategy

Entry	R	Olefin acid	Lactone	Yield ^a /%	Entry	R	Olefin acid	Lactone	Yield ^a /%
1	CH ₃	1b	1	65	9	8-Cl	9a	9	70
2	CF ₃	2b	2	90	10	7-OCH ₃	10a	10	74
3	OCH ₃	3b	3	77	11	6-OCH ₃	11a	11	80
4	C ₆ H ₅	4b	4	62	12	6-Cl	12a	12	78
5	2-C ₄ H ₉ O	5b	5	60	13	7,9-(OCH ₃) ₂	13a	13	85
6	H	6a	6	86	14				72
7	8-CH ₃	7a	7	76	15				80
8	8-F	8a	8	75	16				73

^a Isolated yields. **Scheme 2** Synthesis of **6~14** by epoxidation-lactonization **Figure 3** X-ray crystal structure of seven-membered lactone **10** and **S-6**

Based on the drawback of the above approach, we resorted to the Sharpless asymmetric dihydroxylation for the lactone **R-1** and **S-1** (Scheme 4) due to its high *ee* values

Scheme 3 Synthesis of **R-1** by Shi's asymmetric epoxidation

and ability to produce multiply isomers.^[41–42] First *cis*-geranic acid was esterified with methanol to provide ester **16**, subsequent dihydroxylation of compound **16** with AD-mix- β gave compound **17a** in 93% yield and 90.2% *ee*.^[41] After removal of methyl group by LiOH hydrolyzation at room temperature, the carboxylic acid was treated with 2,4,6-trichlorobenzoyl chloride (2,4,6-TCBC) via Yamaguchi reaction in THF,^[43–45] then the mixture was heated in toluene to successfully produce lactone **R-1** in 60% yield with 92.5% *ee*. The optical rotation data of **R-1** was consistence with that in literatures.^[2–3] The enantiomer **S-1** was also prepared by the same procedure with AD-mix- α replacing AD-mix- β in 65% yield with 93.9% *ee*. After **R-1** and **S-1** were obtained, the 3-CF₃ and 3-OCH₃ analogues (**R-2** and **S-2**, **R-3** and **S-3**) were also synthesized in the similar approaches via diol ester intermediates **18a**, **18b**, **19a** and **19b** (Scheme 5). (*E*)-Methyl 3-trifluoromethyl-7-

methylocta-2,6-dienoate (**2a**) and (*E*)-methyl 3-methoxy-7-methylocta-2,6-dienoate (**3a**) were converted to chiral diol esters **18a** and **18b**, **19a** and **19b** under Sharpless asymmetric dihydroxylation condition. Then LiOH hydrolyzation and Yamaguchi reaction were used to afford compounds *R*-**2** and *S*-**2** from **18a** and **18b** in 55% and 56% yields with 90.7% and 93.0% *ee*, compounds *R*-**3** and *S*-**3** from **19a** and **19b** in 82% and 86% yields with 91.5% and 92.3% *ee*, respectively.

The racemic 3-(2-hydroxypropan-2-yl)-4,5-dihydrobenzo[c]oxepin-1(3*H*)-one (**6**) was synthesized in 9-step reaction in 10.3% overall yield, and it showed better antifungal activity than racemic **1** and *R*-**1** in the previous paper.^[32] But it could be synthesized in 3-step reaction (Table 1, Entry 6) in much higher overall yield from *o*-methyl benzoic acid, because alkylation of *o*-methyl benzoic acid with 1-bromo-3-methylbut-2-ene in the presence of LDA under -78°C gave the key intermediate acid **6a** in 74% yield.^[46] So *R*-**6** and *S*-**6** were synthesized in the similar approaches (Scheme 6). Compound **6a** was esterified to produce methyl ester **20**, and **20** was subjected to Sharpless asymmetric dihydroxylation of the olefins to afford chiral diol ester **21a** and **21b**, respectively. Finally, **21a** and **21b** were hydrolyzed and converted in reflux toluene via Yamaguchi reaction to afford derivatives *R*-**6** and *S*-**6** in 50% and 56% yields with 89.3% and 88.8% *ee*, respectively.

The absolute configurations of **18a**, **19a**, and **21a** were determined using Mosher ester method^[47] by the reaction products of **18a**, **19a**, and **21a** with the Mosher acid chloride having *R*- and *S*-configurations (Figure 4). So the absolute configurations of **18b**, **19b** and **21b** were also ob-

tained. Then the absolute configurations of *R*-**2**, *S*-**2**, *R*-**3**, *S*-**4**, *R*-**6** and *S*-**6** were deduced from the configurations of their starting materials. The absolute configurations were confirmed by the crystal structure of *S*-**6** in Figure 3, which showed that its absolute configuration was *S* at chiral center of C-6 and consistent with that of **21b**.

The antifungal activities of synthesized seven-membered lactone were evaluated using the mycelial growth rate test at $(25 \pm 0.5)^{\circ}\text{C}$ following the protocol in reference,^[48-50] chlorothalonil and carbendazim used as positive control, and EC_{50} values were determined from the inhibition rates of five different concentrations for the compounds having more than 70% inhibition rates, which were given in Table 2. The data showed that most of the synthesized seven-membered lactones have significant antifungal activities against these phytopathogens, especially against *R. solani* and *A. solani* with EC_{50} values of 2.8~134.2 $\mu\text{g/mL}$. From these data, it was found that both non-benzolactones **1**, **3**~**5** were more active than benzolactones **6**~**13**, lactones **4** and **5** with 3-aromatic groups were more active than compounds **1**~**3** against *R. solani* and *A. solani*. The *R*- and *S*-isomers of **1** and **3** had the same level of activities as that of racemic **1** and **3** against *A. solani* as well as **3** against *R. solani*, and the racemic **1** and lactone *R*-**1** had much better activities than that of *S*-**1** against *R. solani*. However, the *R*- and *S*-isomers of **6** exhibited much better activities than that of racemic benzolactone **6** against *A. solani*. It was deduced that compound **6** may partially lose one molecule of water at the *tert*-alcohol position to reduce its antifungal activity. These results implied that 3-aromatic group and *tert*-alcohol moiety played a crucial role for the antifungal

Scheme 4 Synthesis of *R*-**1** and *S*-**1** by Sharpless asymmetric dihydroxylation

Scheme 5 Synthesis of *R*-**2** and *S*-**2**, *R*-**3** and *S*-**3** by Sharpless asymmetric dihydroxylation

Scheme 6 Synthesis of *R*-6 and *S*-6 by Sharpless asymmetric dihydroxylation

Figure 4 Absolute configurations of secondary alcohols **18a**, **19a** and **21a** by modified Mosher's method. $\Delta\delta_{\text{H}}$ ($\delta_{\text{S}} - \delta_{\text{R}}$) values are given

Table 2 EC_{50} ($\mu\text{g/mL}$) values for antifungal activity of synthesized seven-membered lactones **1**~**14**

Compd.	<i>R. solani</i>	<i>A. solani</i>	<i>S. sclerotiorum</i>	<i>B. cinerea</i>	<i>F. graminearum</i>	<i>P. capsici</i>
1	65.7	42.2	— ^a	—	—	—
<i>R</i> - 1	56.5	44.9	—	—	—	—
<i>S</i> - 1	231.6	52.0	—	—	—	—
2	—	—	—	—	—	—
3	75.6	23.5	—	—	—	—
<i>R</i> - 3	87.1	32.3	—	—	—	—
<i>S</i> - 3	73.0	40.5	—	—	—	—
4	10.7	3.7	5.1	0.5	19.4	18.6
5	2.8	4.6	9.8	—	10.7	5.1
6	—	115.1	—	—	—	—
<i>R</i> - 6	—	27.4	—	—	—	—
<i>S</i> - 6	—	18.4	—	—	—	—
7	48.9	—	—	—	323.7	—
8	134.2	8.1	186.8	—	132.9	—
9	—	—	—	—	—	—
10	—	—	—	—	—	—
11	—	—	—	—	—	—
12	—	—	—	—	—	—
13	41.4	6.6	—	—	—	—
14	—	—	—	—	—	—
Carbendazim	0.05	0.05	7.9	528.2	0.07	77.8

^a “—” indicated that the EC_{50} values were not determined due to weak inhibitory activities.

activities. Among these compounds, compounds **4** and **5** were broad spectrum against six fungus and EC_{50} values were in the range of 0.5~20.0 $\mu\text{g/mL}$, which showed that they were the most active compounds and could be used as the lead structures of modification to further improve the activity, as well as the *R*- and *S*-isomers of **4** and **5** would be synthesized to explore their activity differences. These

works were in progress in our laboratory.

2 Conclusion

In conclusion, a concise synthetic strategy has been developed for the synthesis of seven-membered lactones with α -hydroxyl side chains through epoxidation-lactonization procedure, and stereoselective synthesis of optical isomers

of natural product 3,7-dimethyl-7-hydroxy-2-octen-6-olide and their analogues with Sharpless asymmetric dihydroxylation as the key steps. The absolute configurations of these optical isomers were deduced from Mosher esters of chiral diol esters, and absolute configuration of compound **S-6** determined by *x*-ray diffraction analysis. The EC₅₀ values of **4** and **5** were in the range of 0.5~20.0 µg/mL against six phytopathogens, which showed that they were the most active compounds and could be used as the lead structures.

3 Experimental section

3.1 General information

All reactions were performed with magnetic stirring. Unless otherwise stated, all reagents were purchased from commercial suppliers and used without further purification. Organic solutions were concentrated under reduced pressure using a rotary evaporator or oil pump. Flash column chromatography was performed using a Qingdao Haiyang silica gel (200~300 mesh). Melting points were measured on a Yanagimoto apparatus (Yanagimoto MFG Co., Kyoto, Japan) and are uncorrected. ¹H NMR and ¹³C NMR spectra were obtained on a Bruker DPX 300 spectrometer (Bruker Biospin Co., Stuttgart, Germany) with CDCl₃ as a solvent and TMS as an internal standard; HR-ESI-MS spectra were analyzed on a Bruker Apex II mass spectrometer (Bruker Co., Bremen, Germany). The crystal structure was analyzed with a Thermo Fisher ESCALAB 250 four-circle X-ray diffractometry. The solvents were analytical grade and newly distilled before usage. The *e.e.* values were analyzed by an Agilent LC 1100 HPLC instrument equipped with a chiral Chiralpak AD column (250×4.6 mm), eluent: hexane/isopropanol (*V*:*V*=95:5, 90:10, 85:15), flow rate: 1.0 mL/min, UV detection wavelength: 230 nm.

3.2 Synthesis of seven-membered lactones **1**~**14**

3.2.1 Synthesis of compounds **1b**~**5b**

Compounds **1b**~**5b** were synthesized following the procedure of reference [30]. The NMR and ESI-MS data of **1b** were identical with that reported in literature.^[30]

(*E*)-7-Methyl-3-(trifluoromethyl)octa-2,6-dienoic acid (**2b**): Yellow oil, yield 97%. ¹H NMR (300 MHz, CDCl₃) δ: 9.62 (br.s, 1H), 6.35 (s, 1H), 5.16~5.08 (m, 1H), 2.71~2.66 (m, 2H), 2.32~2.21 (m, 2H), 1.71 (s, 3H), 1.61 (s, 3H); ESI-MS *m/z*: 223 [M+H]⁺. HR-ESI-MS calcd for C₁₀H₁₄F₃O₂ [M+H]⁺ 223.0940, found 223.0942.

(*E*)-7-Methyl-3-methoxyocta-2,6-dienoic acid (**3b**): White solid, yield 76%. m.p. 102~104 °C; ¹H NMR (300 MHz, CDCl₃) δ: 11.48 (br, 1H), 5.16~5.11 (m, 1H), 4.99 (s, 1H), 3.65 (s, 3H), 2.75 (dd, *J*=9.3, 7.6 Hz, 2H), 2.23 (q, *J*=7.6 Hz, 2H), 1.68 (s, 1H), 1.61 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ: 178.34, 173.38, 132.43, 123.08, 90.24, 55.55, 32.37, 26.07, 25.61, 17.45; HR-ESI-MS calcd for C₁₀H₁₇O₃ [M+H]⁺ 185.1172, found 185.1172.

(*E*)-7-Methyl-3-phenylocta-2,6-dienoic acid (**4b**): Yellow oil, yield 99%. ¹H NMR (300 MHz, CDCl₃) δ: 10.42

(br, 1H), 7.44~7.13 (m, 5H), 6.01 (s, 0.4 H), 5.87 (s, 0.6 H), 5.10~5.02 (m, 1H), 3.08 (t, *J*=7.9 Hz, 0.6H), 2.47 (t, *J*=7.3 Hz, 1.2H), 2.12~2.01 (m, 2H), 1.71~1.63 (m, 3H), 1.54~1.48 (m, 3H); ESI-MS *m/z*: [M+H]⁺ 231. HR-ESI-MS calcd for C₁₅H₁₉O₂ [M+H]⁺ 231.1380, found 231.1384.

(*E*)-7-Methyl-3-(furan-2-yl)octa-2,6-dienoic acid (**5b**): Yellow oil, yield 88%. ¹H NMR (300 MHz, CDCl₃) δ: 7.69 (s, 1H), 7.28~7.25 (m, 1H), 6.70~6.46 (m, 1H), 5.75 (s, 1H), 5.21~5.10 (m, 1H), 2.93 (t, *J*=8.0 Hz, 0.6 H), 2.58 (t, *J*=7.6 Hz, 1.4H), 2.27~2.07 (m, 2H), 1.68 (s, 3H), 1.37 (s, 3H); ESI-MS *m/z*: [M+H]⁺ 221. HR-ESI-MS calcd for C₁₃H₁₇O₃ [M+H]⁺ 221.1172, found 221.1175.

3.2.2 General synthesis procedure of the olefin acid **6a**~**13a**

Under the protection of N₂, 5.4 g (40 mmol) of *o*-methyl benzoic acid, 150 mL of THF were added into a 500 mL Schlenk flask, cooled to -20 °C and kept for 20 min, then added 2 mol/L LDA (56 mL, 112 mmol) and stirred 4 h. After that, 7 mL (56 mmol) of 1-bromo-3-methylbut-2-ene was added at -20 °C, raised slowly to room temperature and reacted overnight. Then, the reaction mixture was treated as routine work to afford a white solid **6a**. In a similar way, compounds **7a**~**13a** was prepared.

2-(4-Methylpent-3-en-1-yl)benzoic acid (**6a**): White solid, yield 74%. m.p. 61~63 °C (lit.^[32] 61~63 °C); ¹H NMR (300 MHz, CDCl₃) δ: 8.02~8.05 (m, 1H), 7.44~7.50 (m, 1H), 7.26~7.31 (m, 2H), 5.18~5.24 (m, 1H), 3.06 (t, *J*=7.5 Hz, 2H), 2.28~2.35 (m, 2H), 1.69 (s, 3H), 1.54 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 173.42, 145.23, 132.76, 132.35, 131.55, 131.37, 128.32, 125.89, 123.72, 34.74, 30.30, 25.68, 17.44; HR-ESI-MS calcd for C₁₃H₁₇O₂ [M+H]⁺ 205.1223, found 205.1221.

5-Methyl-2-(4-methylpent-3-en-1-yl)benzoic acid (**7a**): White solid, yield 63%. m.p. 88~90 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.84 (s, 1H), 7.25~7.30 (m, 1H), 7.15~7.18 (m, 1H), 5.18~5.23 (m, 1H), 3.00 (t, *J*=8.1 Hz, 2H), 2.33 (s, 3H), 2.25~2.30 (m, 2H), 1.69 (s, 3H), 1.55 (s, 3H). HR-ESI-MS calcd for C₁₄H₁₉O₂ [M+H]⁺ 219.1380, found 219.1384.

5-Fluoro-2-(4-methylpent-3-en-1-yl)benzoic acid (**8a**): Yellow solid, yield 58%. m.p. 145~147 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.72 (dd, *J*=9.5, 2.7 Hz, 1H), 7.29~7.15 (m, 2H), 5.20~5.15 (m, 1H), 3.02 (t, *J*=7.7 Hz, 2H), 2.24~2.32 (m, 2H), 1.68 (s, 3H), 1.51 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.09, 160.57 (¹*J*_{CF}=243.5 Hz), 141.07 (⁴*J*_{CF}=3.4 Hz), 133.02 (³*J*_{CF}=7.2 Hz), 132.64, 129.53 (³*J*_{CF}=6.9 Hz), 123.38, 119.88 (²*J*_{CF}=20.6 Hz), 118.03 (²*J*_{CF}=23.1 Hz), 33.95, 30.25, 25.68, 17.43; HR-ESI-MS calcd for C₁₃H₁₆FO₂ [M+H]⁺ 223.1128, found 223.1123.

5-Chloro-2-(4-methylpent-3-en-1-yl)benzoic acid (**9a**): Little yellow solid, yield 65%. m.p. 132~134 °C; ¹H NMR (300 MHz, CDCl₃) δ: 8.01 (d, *J*=2.1 Hz, 1H), 7.43 (dd, *J*=8.4 Hz, 2.1 Hz, 1H), 7.21 (d, *J*=8.4 Hz, 1H), 5.20~5.15 (m, 1H), 3.05~2.99 (m, 2H), 2.32~2.25 (m, 2H), 1.75

(s, 3H), 1.52 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 172.29, 143.67, 132.81, 132.73, 131.70, 131.31, 130.32, 129.62, 123.26, 34.11, 30.08, 25.67, 17.45; HR-ESI-MS calcd for $\text{C}_{13}\text{H}_{16}\text{ClO}_2$ $[\text{M}+\text{H}]^+$ 239.0833, found 239.0827.

4-Methoxy-2-(4-methylpent-3-en-1-yl)benzoic acid (**10a**): White solid, yield 86%. m.p. 78~80 °C; ^1H NMR (300 MHz, CDCl_3) δ : 8.08~8.04 (m, 1H), 6.80~6.76 (m, 2H), 5.24~5.19 (m, 1H), 3.86 (s, 3H), 3.04 (t, $J=7.8$ Hz, 2H), 2.35~2.27 (m, 2H), 1.69 (s, 3H), 1.51 (s, 3H); ^{13}C NMR (300 MHz, CDCl_3) δ : 172.68, 163.01, 148.28, 134.24, 132.24, 123.82, 120.33, 116.62, 111.08, 55.31, 35.23, 30.09, 25.69, 17.50; HR-ESI-MS calcd for $\text{C}_{14}\text{H}_{19}\text{O}_3$ $[\text{M}+\text{H}]^+$ 235.1328, found 235.1322.

3-Methoxy-2-(4-methylpent-3-en-1-yl)benzoic acid (**11a**): White solid, yield 80%. m.p. 88~90 °C; ^1H NMR (300 MHz, CDCl_3) δ : 7.56~7.53 (m, 1H), 7.26~7.21 (m, 1H), 7.05~7.02 (m, 1H), 5.29~5.23 (m, 1H), 3.86 (s, 3H), 3.02 (t, $J=7.8$ Hz, 2H), 2.27~2.19 (m, 2H), 1.69 (s, 3H), 1.56 (s, 3H); ^{13}C NMR (300 MHz, CDCl_3) δ : 173.96, 158.03, 133.60, 131.98, 130.24, 126.33, 124.33, 122.85, 114.36, 55.79, 28.67, 26.74, 25.69, 17.29; HR-ESI-MS calcd for $\text{C}_{14}\text{H}_{19}\text{O}_3$ $[\text{M}+\text{H}]^+$ 235.1328, found 235.1323.

3-Chloro-2-(4-methylpent-3-en-1-yl)benzoic acid (**12a**): Little yellow solid, yield 55%. m.p. 46~48 °C; ^1H NMR (300 MHz, CDCl_3) δ : 7.89~7.86 (m, 1H), 7.58~7.55 (m, 1H), 7.25~7.20 (m, 1H), 5.27~5.22 (m, 1H), 3.20~3.15 (m, 2H), 2.34~2.26 (m, 2H), 1.70 (s, 3H), 1.56 (s, 3H); ^{13}C NMR (300 MHz, CDCl_3) δ : 173.11, 141.96, 136.07, 133.89, 132.77, 131.02, 129.76, 126.67, 123.37, 30.79, 28.25, 25.71, 17.36; HR-ESI-MS calcd for $\text{C}_{13}\text{H}_{16}\text{ClO}_2$ $[\text{M}+\text{H}]^+$ 239.0833, found 239.0827.

2,4-Dimethoxy-6-(4-methylpent-3-en-1-yl)benzoic acid (**13a**): Yellow solid, yield 82%. m.p. 173~174 °C; ^1H NMR (300 MHz, CDCl_3) δ : 6.42 (d, $J=2.4$ Hz, 1H), 6.28 (d, $J=2.4$ Hz, 1H), 5.22~5.16 (m, 1H), 3.91 (s, 3H), 3.84 (s, 3H), 2.87 (t, $J=7.8$ Hz, 2H), 2.34~2.26 (m, 2H), 1.68 (s, 3H), 1.57 (s, 3H); ^{13}C NMR (300 MHz, CDCl_3) δ : 171.43, 161.98, 158.94, 145.35, 132.39, 132.53, 113.85, 107.42, 96.50, 56.23, 55.33, 34.79, 29.91, 25.62, 17.52; HR-ESI-MS calcd for $\text{C}_{15}\text{H}_{21}\text{O}_4$ $[\text{M}+\text{H}]^+$ 265.1434, found 265.1432.

3.2.3 General procedure for compounds **1**~**13**

To a solution of *cis*-geranic acid **1b** (10.7 g, 63.9 mmol) in CH_2Cl_2 (300 mL) was added the peracetic acid solution (12.3 mL $\times$ 10) and anhydrous sodium carbonate (4.21 g $\times$ 10). The reaction mixture was stirred for 1.5 h at room temperature. After the reaction was finished as detected by thin-layer chromatography (TLC), the mixture was diluted with saturated $\text{Na}_2\text{S}_2\text{O}_3$ (50 mL) and extracted with CH_2Cl_2 (100 mL $\times$ 3). The combined organic phases were washed with brine (100 mL $\times$ 3), dried over anhydrous sodium sulfate, and concentrated in *vacuo* to give a white solid epoxide **1c** 10.5 g, yield 89%. m.p. 46~48 °C (Lit.^[30] 46~48 °C); ^1H NMR (300 MHz, CDCl_3) δ : 11.75 (br, 1H, OH), 5.74 (s, 1H, CH=), 2.89~2.66 (m, 3H, CHO+CH₂), 1.95 (s, 3H, CH₃), 1.81~1.66 (m, 2H, CH₂), 1.31 (s,

3H, CH₃), 1.28 (s, 3H, CH₃); ^{13}C NMR (75 MHz, CDCl_3) δ : 171.40, 162.32, 116.28, 63.84, 58.59, 30.24, 27.58, 25.51, 24.77, 18.56.

To a 1000 mL three-necked flask added 12.0 g (65.1 mmol) compound **1c**, 560 mL of CH_2Cl_2 and 0.33 g of (+)-camphor-sulfonic acid. The mixture was stirred for 4 h at room temperature. After the reaction was finished, the organic phase was washed with 5% Na_2CO_3 solution and brine, dried with anhydrous Na_2SO_4 , and concentrated in *vacuo*. Flash chromatography on silica gel (petroleum ether/EtOAc, $V:V=3:1$) afforded 3,7-dimethyl-7-hydroxy-2-octen-6-olide (**1**). In a similar way, compounds **2**~**13** were prepared.

3,7-Dimethyl-7-hydroxy-2-octen-6-olide (**1**): Colorless oil liquid, yield 65%. ^1H NMR (300 MHz, CDCl_3) δ : 5.86 (brq, $J=1.2$ Hz, 1H, CH=), 4.04 (dd, $J=9.0$, 2.2 Hz, 1H, CHO), 2.89 (br, 1H, OH), 2.53 (dt, $J=17.8$, 6.0 Hz, 1H, CH₂), 2.38~2.33 (m, 1H, CH₂), 2.19~2.09 (m, 1H, CH₂), 1.97 (s, 3H, CH₃), 1.96~1.87 (m, 1H, CH₂), 1.28 (s, 3H, CH₃), 1.26 (s, 3H, CH₃); ^{13}C NMR (75 MHz, CDCl_3) δ : 168.12, 154.92, 118.13, 84.34, 71.44, 33.52, 27.10, 26.06, 26.00, 24.79; HR-ESI-MS calcd for $\text{C}_{10}\text{H}_{17}\text{O}_3$ $[\text{M}+\text{H}]^+$ 185.1172, found 185.1171.

3-Trifluoromethyl-7-methyl-7-hydroxy-2-octen-6-olide (**2**): Light yellow oil, yield 90%. ^1H NMR (300 MHz, CDCl_3) δ : 6.50 (s, 1H), 4.07 (dd, $J=9.5$, 3.3 Hz, 1H), 2.93 (s, 1H), 2.58 (t, $J=6.9$ Hz, 2H), 2.26~2.03 (m, 2H), 1.30 (s, 3H), 1.26 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 165.87, 140.40 (q, $^2J_{\text{CF}}=30.9$ Hz), 123.32 (q, $^3J_{\text{CF}}=5.6$ Hz), 122.60 (q, $^1J_{\text{CF}}=274.5$ Hz), 84.23, 71.29, 26.45, 25.01, 24.83, 24.47; HR-ESI-MS calcd for $\text{C}_{10}\text{H}_{14}\text{F}_3\text{O}_3$ $[\text{M}+\text{H}]^+$ 239.0889, found 239.0890.

3-Methoxy-7-methyl-7-hydroxy-2-octen-6-olide (**3**): Colorless solid, yield 77%. m.p. 55~57 °C; ^1H NMR (300 MHz, CDCl_3) δ : 5.18 (s, 1H), 4.04 (dd, $J=9.7$, 3.2 Hz, 1H), 3.67 (s, 3H), 2.69~2.41 (m, 2H), 2.11~1.98 (m, 2H), 1.27 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 171.55, 169.47, 92.09, 84.41, 71.29, 55.62, 31.14, 25.55, 24.80, 24.76; HR-ESI-MS calcd for $\text{C}_{10}\text{H}_{17}\text{O}_4$ $[\text{M}+\text{H}]^+$ 201.1121, found 201.1121.

3-Phenyl-7-methyl-7-hydroxy-2-octen-6-olide (**4**): Light yellow oil, yield 62%. ^1H NMR (300 MHz, CDCl_3) δ : 7.48~7.38 (m, 5H), 6.24 (s, 1H), 4.17 (dd, $J=4.7$, 9.1 Hz, 1H), 2.86~2.82 (dd, $J=7.5$, 6.7 Hz, 2H), 2.46 (br, 1H), 2.23~2.14 (m, 2H), 1.30 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 169.04, 153.27, 139.93, 129.45, 128.75, 126.23, 118.77, 84.13, 71.62, 29.97, 27.79, 25.69, 24.61. HR-ESI-MS calcd for $\text{C}_{15}\text{H}_{19}\text{O}_3$ $[\text{M}+\text{H}]^+$ 247.1328, found 247.1328.

3-(Furan-2-yl)-7-methyl-7-hydroxy-2-octen-6-olide (**5**):^[32] Light yellow oil, yield 60%. ^1H NMR (300 MHz, CDCl_3) δ : 7.51 (d, $J=1.4$ Hz, 1H), 6.68 (d, $J=3.5$ Hz, 1H), 6.50~6.47 (m, 2H), 4.12 (dd, $J=9.9$, 4.2 Hz, 1H), 2.76 (t, $J=7.0$ Hz, 2H), 2.17~2.10 (m, 3H), 1.29 (s, 3H), 1.25 (s, 3H); HR-ESI-MS calcd for $\text{C}_{13}\text{H}_{17}\text{O}_4$ $[\text{M}+\text{H}]^+$ 237.1121, found 237.1123.

3-(2-Hydroxypropan-2-yl)-4,5-dihydrobenzo[*c*]oxepin-

1(3*H*)-one (**6**): Colorless solid, yield 86%. m.p. 45~47 °C (Lit.^[32] colorless oil). ¹H NMR (300 MHz, CDCl₃) δ: 7.71~7.74 (m, 1H), 7.46~7.52 (m, 1H), 7.35~7.40 (m, 1H), 7.21~7.28 (m, 1H), 3.86 (dd, *J*=4.8, 13.5 Hz, 1H), 2.89~2.96 (m, 1H), 2.77~2.84 (m, 1H), 2.25~2.36 (m, 1H), 1.92~2.01 (m, 1H), 1.25 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ: 171.20, 138.22, 132.74, 131.33, 130.14, 128.60, 127.29, 84.20, 71.55, 29.54, 28.47, 26.19, 24.30; HR-ESI-MS calcd for C₁₃H₁₇O₃ [M+H]⁺ 221.1172, found 221.1172.

3-(2-Hydroxypropan-2-yl)-8-methyl-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**7**): Yellow oil, yield 76%. ¹H NMR (300 MHz, CDCl₃) δ: 7.55 (s, 1H), 7.27~7.30 (m, 1H), 7.10~7.12 (m, 1H), 3.86 (dd, *J*=4.8, 12.1 Hz, 1H), 2.89~2.84 (m, 1H), 2.77~2.80 (m, 1H), 2.37 (br, 1H), 2.29 (s, 3H), 2.26~2.29 (m, 1H), 1.94 (m, 1H), 1.24 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ: 171.48, 137.20, 135.26, 133.46, 131.15, 130.64, 128.59, 84.30, 71.61, 29.10, 28.59, 26.30, 24.28, 20.77; HR-ESI-MS calcd for C₁₄H₁₉O₃ [M+H]⁺ 235.1328, found 235.1328.

3-(2-Hydroxypropan-2-yl)-8-fluoro-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**8**): Yellow oil, yield 75%. ¹H NMR (300 MHz, CDCl₃) δ: 7.49~7.41 (m, 1H), 7.22~7.14 (m, 2H), 3.87 (dd, *J*=12.1, 7.1 Hz, 1H), 2.95~2.77 (m, 2H), 2.35~2.23 (m, 1H), 2.00~1.87 (m, 1H), 1.26 (s, 6H); HR-ESI-MS calcd for C₁₃H₁₆FO₃ [M+H]⁺ 239.1078, found 239.1074.

3-(2-Hydroxypropan-2-yl)-8-chloro-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**9**): Light yellow solid, yield 70%. m.p. 89~91 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.72 (s, 1H), 7.44 (dd, *J*=8.7, 2.2 Hz, 1H), 7.18 (d, *J*=8.1 Hz, 1H), 3.87~3.82 (m, 1H), 2.96~2.76 (m, 2H), 2.37~2.26 (m, 2H), 1.98~1.86 (m, 1H), 1.25 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ: 169.75, 136.54, 133.19, 132.79, 132.58, 130.07, 129.95, 84.34, 71.44, 28.89, 28.19, 26.13, 24.31. HR-ESI-MS calcd for C₁₃H₁₆ClO₃ [M+H]⁺ 255.0782, found 255.0777.

3-(2-Hydroxypropan-2-yl)-7-methoxy-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**10**): White solid, yield 74%. m.p. 106~108 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.72 (d, *J*=8.6 Hz, 1H), 6.86 (dd, *J*=8.6, 2.5 Hz, 1H), 6.72 (s, 1H), 3.90~3.85 (m, 4H), 3.02~2.91 (m, 1H), 2.77~2.71 (m, 1H), 2.35~2.24 (m, 2H), 2.00~1.89 (m, 1H), 1.24 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ: 171.17, 163.03, 140.78, 132.59, 123.48, 114.40, 112.10, 84.21, 71.55, 55.36, 29.99, 28.39, 26.20, 24.23; HR-ESI-MS calcd for C₁₄H₁₉O₄ [M+H]⁺ 251.1277, found 251.1272.

3-(2-Hydroxypropan-2-yl)-6-methoxy-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**11**): White solid, yield 80%. m.p. 97~99 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.31~7.26 (m, 2H), 7.09~7.04 (m, 1H), 3.86~3.81 (m, 4H), 3.49~3.42 (m, 1H), 2.51~2.39 (m, 1H), 2.29~2.17 (m, 1H), 1.93~1.74 (m, 1H), 1.24 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ: 171.11, 155.97, 132.87, 127.83, 126.36, 121.60, 114.44, 84.46, 71.50, 55.80, 27.76, 26.15, 24.21, 20.25; HR-ESI-MS calcd for C₁₄H₁₉O₄ [M+H]⁺ 251.1277, found 251.1270.

3-(2-Hydroxypropan-2-yl)-6-chloro-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**12**): White solid, yield 78%. m.p. 55~57 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.62~7.55 (m, 2H), 7.33~7.26 (m, 1H), 3.81~3.76 (m, 1H), 3.52~3.45 (m, 1H), 2.75~2.64 (m, 1H), 2.36~2.24 (m, 1H), 1.99~1.86 (m, 1H), 1.25 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ: 169.94, 135.36, 133.56, 133.29, 128.45, 128.14, 84.22, 71.44, 27.39, 26.18, 25.32, 24.29; HR-ESI-MS calcd for C₁₃H₁₆ClO₃ [M+H]⁺ 255.0782, found 255.0777.

3-(2-Hydroxypropan-2-yl)-7,9-dimethoxy-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**13**): Yellow oil, yield 85%. ¹H NMR (300 MHz, CDCl₃) δ: 6.42 (s, 1H), 6.32 (s, 1H), 3.92~3.85 (m, 6H), 2.84~2.66 (m, 2H), 2.32 (br, 1H), 2.28~2.19 (m, 1H), 1.89~1.83 (m, 1H), 1.28 (s, 3H), 1.22 (s, 3H); HR-ESI-MS calcd for C₁₅H₂₁O₅ [M+H]⁺ 281.1389, found 281.1384.

3.2.4 Synthesis of 9-hydroxy-3-(2-hydroxypropan-2-yl)-7-methoxy-4,5-dihydrobenzo-[c]oxepin-1(3*H*)-one (**14**)

Under N₂ atmosphere, 0.28 g (1 mmol) of lactone **13** and 15 mL of CH₂Cl₂ were added into a 50 mL Schlenk flask, cooled to -78 °C, and dropped 0.1 mL of BBr₃ (2 mol·L⁻¹ CH₂Cl₂ solution). Then the reaction mixture was stirred, naturally raised to -10 °C, and reacted 48 h at -10 °C. After the reaction, the mixture solution was extracted with CH₂Cl₂ (10 mL×3), and the organic phase was combined. The organic phase was washed with brine, dried over anhydrous Na₂SO₄, and concentrated *in vacuo*. Flash chromatography on silica gel (petroleum ether/EtOAc, *V*:*V*=6:1) afforded lactone **14**. Light yellow oil, 0.20 g, yield 75%. ¹H NMR (300 MHz, CDCl₃) δ: 10.09 (s, 1H), 6.40~6.31 (m, 2H), 4.00~3.94 (m, 1H), 3.82 (s, 3H), 2.96~2.85 (m, 1H), 2.75~2.68 (m, 1H), 2.34~2.22 (m, 1H), 2.05~1.96 (m, 1H), 1.26 (s, 3H), 1.22 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 174.20, 164.88, 163.57, 142.70, 108.96, 106.51, 99.24, 86.20, 71.65, 55.38, 30.73, 28.55, 26.04, 24.03; HR-ESI-MS calcd for C₁₄H₁₉O₅ [M+H]⁺ 267.1227, found 267.1228.

3.3 Synthesis of *R*-**1** via Shi's asymmetric epoxidation

To a 100 mL three-necked round-bottom flask were added buffer (0.05 mol·L⁻¹ Na₂B₄O₇·10H₂O in 4×10⁻⁴ mol·L⁻¹ aqueous Na₂EDTA, 20 mL), acetonitrile (30 mL), *cis*-geranic acid **1b** (0.336 g, 2 mmol), tetrabutyl-ammonium hydrogen sulfate (0.03 g, 0.08 mmol), and catalyst (0.15 g, 0.6 mmol). The reaction mixture was cooled with an ice bath. A solution of Oxone (1.7 g, 2.76 mmol) in aqueous Na₂EDTA (4×10⁻⁴ mol·L⁻¹, 13 mL) and a solution of K₂CO₃ (1.6 g, 11.6 mmol) in water (13 mL) were added dropwise through two addition funnels over a period of 3 h. The reaction mixture of chiral 6,7-epoxy-geranic acid **15** was obtained. Then the mixture was acidified to pH 5 with 25% sulphuric acid and extracted with CH₂Cl₂. The CH₂Cl₂ phase was dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was dissolved in 25 mL of CH₂Cl₂, added (+)-CSA (10 mg) and stirred 3

h at room temperature. The solvent was evaporated in reduced pressure, and the residue was purified by flash silica gel chromatography and eluted with petroleum ether/EtOAc ($V:V=6:1$) to give colorless oil **R-1** 0.22 g, yield 60%. $[\alpha]_D^{25} -27$ (c 0.17, CH₃OH), 47% *ee*. The ¹H NMR, ¹³C NMR and MS data were consistent with the data of lactone **1** and that in references.^[1-2]

3.4 Synthesis of **R-1** and **S-1** via Sharpless asymmetric dihydroxylation

A mixture of *cis*-geranic acid (4.0 g, 22 mmol), methanol (15 mL), *N,N*-dicyclohexylcarbodiimide (4.5 g, 24 mmol), 4-*N*,*N*-dimethylaminopyridine (3.4 g, 27 mmol) and dry CH₂Cl₂ (140 mL) was stirred at room temperature under argon atmosphere for 5 h. A white solid was filtered off. The organic solution was washed with water, dried over anhydrous Na₂SO₄, removed the solid and concentrated under vacuum. Then the crude material was purified by column chromatography on silica gel (petroleum ether/EtOAc, $V:V=100:1$) to afford methyl (*Z*)-3,7-dimethyloct-2,6-dienoate (**16**). Light yellow oil, 3.7 g, yield 88%. ¹H NMR (300 MHz, CDCl₃) δ : 5.65 (t, $J=0.6$ Hz, 1H), 5.17~5.12 (m, 1H), 3.68 (s, 3H), 2.66~2.60 (t, $J=7.9$ Hz, 2H), 2.19~2.14 (m, 2H), 1.89 (s, 3H), 1.68 (s, 3H), 1.62 (s, 3H).

A 100 mL round-bottomed flask equipped with a magnetic stirring bar, was charged with AD-mix- β (6.6 g), water (35 mL) and *tert*-butyl alcohol (35 mL). The resulting mixture was stirred at room temperature till to produce two clear phases. Methanesulfonamide (0.31 g, 3.2 mmol) was added in one-portion and the reaction mixture was stirred for 1.5 h. The reaction mixture was cooled to 0 °C. Compound **6** (0.55 g, 3.0 mmol) was added at once, and the heterogeneous slurry was stirred vigorously at 0 °C for 40 h. The saturated Na₂S₂O₃ solution (60 mL) was added at 0 °C. The mixture was allowed to room temperature and stirred for 30 min. EtOAc (100 mL) and water (50 mL) were added to the reaction mixture. The organic layer was separated and the aqueous layer was re-extracted with EtOAc (20 mL $\times$ 3). The combined organic phases were dried over with anhydrous Na₂SO₄ and the solvent was removed to give the crude product. This product was purified by flash chromatography on silica gel with petroleum ether/EtOAc ($V:V=5:1$) as the eluent to give methyl (2*Z*,6*R*)-6,7-dihydroxy-3,7-dimethyloct-2-enoate (**17a**). Colorless oil, 0.52 g, yield 93%. $[\alpha]_D^{25} -44.0$ (c 0.22, CH₃OH). ¹H NMR (300 MHz, CDCl₃) δ : 5.77 (s, 1H), 4.10 (brs, 1H), 3.70 (s, 3H), 3.28 (d, $J=10.8$, 1H), 3.17~3.09 (m, 1H), 2.67 (brs, 1H), 2.37~2.30 (m, 1H), 1.90 (s, 3H), 1.73~1.53 (m, 2H), 1.18 (s, 3H), 1.16 (s, 3H). In similar way, methyl (2*Z*,6*S*)-6,7-dihydroxy-3,7-dimethyloct-2-enoate (**17b**) was prepared replacing AD-mix- β with AD-mix- α in 90% yield, $[\alpha]_D^{25} +42.5$ (c 0.14, CH₃OH). Its ¹H NMR was identical with that of **17a**.

To a solution of compound **17a** (303 mg, 1.5 mmol) in 50 mL of CH₃OH/H₂O/THF ($V:V:V=1:1:1$) was added LiOH $\cdot$ H₂O (100 mg, 2.4 mmol). After being stirred

at room temperature for 4 h, the reaction mixture was diluted with water and washed with ether. The aqueous layer was acidified with concentrated hydrochloric acid to pH 4 and extracted with ethyl acetate. The organic extracts were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated to give the pure hydroxyl acid in 89% yield (270 mg). To a stirred solution of hydroxyl acid (270 mg, 1.33 mmol) in tetrahydrofuran (THF, 26 mL) were added Et₃N (0.39 mL, 2.8 mmol) at 0 °C, and 2,4,6-trichlorobenzoyl chloride (0.6 mL, 2.84 mmol). After reacting 40 min, the mixture was diluted with THF (10 mL) and toluene (30 mL), and a solution of *N,N*-dimethylaminopyridine (DMAP) (0.95 g, 7.8 mmol) in 200 mL of toluene was added slowly at room temperature. The solution was heated to 100 °C for 12 h, and allowed to cool to room temperature. The solution was washed with saturated aqueous NaCl and extracted with EtOAc. The organic phases were dried over anhydrous Na₂SO₄, the solvent was removed *in vacuo*, and the residue was purified by chromatography on silica gel, eluted with petroleum ether/EtOAc ($V:V=6:1$) to give **R-1**. Light yellow oil, 195 mg, 60% yield, 92.5% *ee*. $[\alpha]_D^{20} -52.1$ (c 0.20, CH₃OH); $[\alpha]_D^{25} -56.0$ (c 0.10, CH₃OH) in the literature.^[2] Its ¹H NMR, ¹³C NMR data were identical with that in 3.2.1; HR-ESI-MS calcd for C₁₀H₁₇O₃ [M+H]⁺ 185.1172, found 185.1171. In similar way, compound **S-1** was prepared replacing **17a** with **17b** in 65% yield, 93.9% *ee*, $[\alpha]_D^{20} +52.7$ (c 0.24, CH₃OH). Its ¹H NMR, ¹³C NMR data were also identical with that in 3.2.1; HR-ESI-MS calcd for C₁₀H₁₇O₃ [M+H]⁺ 185.1172, found 185.1170.

3.5 Synthesis of **R-2** and **S-2**, **R-3** and **S-3** via Sharpless asymmetric dihydroxylation

Methyl 3-trifluoromethyl-7-methyloct-2-enoate (**2a**) was prepared via Wittig reaction according to the similar protocol in literature.^[18] A 100 mL round-bottomed flask equipped with a magnetic stirring bar, was charged with AD-mix- β (5.6 g), water (27 mL) and *tert*-butyl alcohol (27 mL). The mixture was stirred at room temperature to produce two phases, methanesulfonamide (0.28 g, 2.8 mmol) was added in one-portion and the reaction mixture was stirred for 1.5 h. Then the reaction mixture was cooled to 0 °C, compound **2a** (0.63 g, 2.7 mmol) was added at once, and the heterogeneous slurry was stirred vigorously at 0 °C for 40 h. The saturated Na₂S₂O₃ (60 mL) was added at 0 °C and the mixture was allowed to room temperature and stirred for 30 min. EtOAc (100 mL) and water (50 mL) were added to the reaction mixture. The organic layer was separated and the aqueous layer was extracted with EtOAc (20 mL $\times$ 3). The combined organic phase was dried over anhydrous Na₂SO₄, and concentrated to give the crude product. This crude product was purified by flash chromatography on silica gel and eluted with petroleum ether/EtOAc ($V:V=10:1$) to afford methyl (2*E*,6*R*)-6,7-dihydroxy-7-methyl-3-(trifluoromethyl)oct-2-enoate (**18a**). Colorless oil, 0.67 g, 93% yield. $[\alpha]_D^{25} -42.4$ (c 0.45, CH₃OH). ¹H NMR (300 MHz, CDCl₃) δ : 6.13 (s,

1H), 3.78 (s, 3H), 3.37 (dd, $J=10.5$, 2.1 Hz, 1H), 2.65~2.58 (m, 1H), 2.40~2.29 (m, 1H), 1.71~1.50 (m, 2H), 1.21 (s, 3H), 1.16 (s, 3H); HR-ESI-MS calcd for $C_{11}H_{18}F_3O_4$ $[M+H]^+$ 271.1151, found 271.1147. In similar way, methyl (2*E*,6*S*)-6,7-dihydroxy-7-methyl-3-(trifluoromethyl)oct-2-enoate (**18b**) was prepared replacing AD-mix- β with AD-mix- α in 90% yield, $[\alpha]_D^{25} + 42.5$ (c 0.14, CH₃OH). Its ¹H NMR was identical with that of **18a**; HR-ESI-MS calcd for $C_{11}H_{18}F_3O_4$ $[M+H]^+$ 271.1151, found 271.1148.

Methyl 3-methoxy-7-methyloct-2-enoate (**3a**) was prepared via four step reactions according to the similar protocol in literature.^[5] In a similar procedure as **18a** and **18b**, (*R/S*)-methyl 6,7-dihydroxy-3-methoxy-7-methyloct-2-enoates (**19a**, **19b**) were obtained as yellow oil. Compound **19a**: 79% yield, $[\alpha]_D^{25} - 13.5$ (c 0.11, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ : 5.07 (s, 1H), 3.95 (d, $J=4.3$ Hz, 1H), 3.68 (s, 3H), 3.64 (s, 3H), 3.34~3.30 (m, 1H), 3.08~2.98 (m, 2H), 2.73~2.64 (m, 1H), 1.81~1.73 (m, 1H), 1.62~1.53 (m, 1H), 1.17 (s, 3H), 1.14 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 176.11, 168.80, 90.74, 76.57, 72.35, 55.36, 50.88, 29.10, 28.53, 25.92, 23.35; HR-ESI-MS calcd for $C_{11}H_{21}O_5$ $[M+H]^+$ 233.1384, found 233.1383. Compound **19b**: 82% yield, $[\alpha]_D^{25} + 13.5$ (c 0.29, CH₃OH); ¹H NMR, ¹³C NMR data were identical with **19a**; HR-ESI-MS calcd for $C_{11}H_{21}O_5$ $[M+H]^+$ 233.1384, found 233.1380.

To a solution of compound **18a** (0.50 g, 1.75 mmol) in 54 mL of CH₃OH/H₂O/THF ($V:V:V=1:1:1$) was added LiOH·H₂O (150 mg, 3.6 mmol). After being stirred for 4 h at 0 °C, the reaction mixture was diluted with water and washed with ether. The aqueous layer was acidified with concentrated hydrochloric acid to pH=4 and extracted with ethyl acetate. The extracts were combined, dried with anhydrous Na₂SO₄, filtered, and concentrated to give the pure hydroxyl acid compound 380 mg, 79% yield. To a stirred solution of hydroxyl acid (112 mg, 0.44 mmol) in THF (26 mL) at 0 °C was added Et₃N (0.13 mL, 0.99 mmol) followed by 2,4,6-trichlorobenzoyl chloride (0.2 mL 1.28 mmol). After 40 min, the mixture was diluted with THF (5 mL) and toluene (10 mL), and was added slowly to a stirring solution of DMAP (0.32 g, 2.6 mmol) in toluene (70 mL) at room temperature. The solution was heated to 100 °C for 12 h, and allowed to cool to room temperature and diluted with EtOAc. The solution was washed with saturated aqueous NH₄Cl and extracted with EtOAc. The dried (over anhydrous Na₂SO₄) extract was concentrated *in vacuo*, and the residue was purified by chromatography on silica gel, eluted with petroleum ether/EtOAc ($V:V=6:1$) to give **R-2**. Yellow oil, 62 mg, yield 56%, 90.7% *ee*. $[\alpha]_D^{25} - 25.7$ (c 0.21, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ : 6.50 (s, 1H), 4.07 (dd, $J=9.5$, 3.3 Hz, 1H), 2.93 (s, 1H), 2.58 (t, $J=6.9$ Hz, 2H), 2.26~2.03 (m, 2H), 1.30 (s, 3H), 1.26 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 165.87, 140.40 (q, $^2J_{CF}=30.9$ Hz), 123.32 (q, $^3J_{CF}=5.6$ Hz), 122.60 (q, $^1J_{CF}=274.5$ Hz), 84.23, 71.29, 26.45, 25.01, 24.83, 24.47; HR-ESI-MS calcd for

$C_{10}H_{14}F_3O_3$ $[M+H]^+$ 239.0890, found 239.0890. In similar way, compound **S-2** was prepared replacing **18a** with **18b** in 55% yield, 95.4% *ee*, $[\alpha]_D^{25} + 25.9$ (c 0.30, CH₃OH). ¹H NMR, ¹³C NMR data were identical with that of **2a**; HR-ESI-MS calcd for $C_{10}H_{14}F_3O_3$ $[M+H]^+$ 239.0890, found 239.0889.

In similar way as **R-2** and **S-2**, compounds **R-3** and **S-3** were prepared using **19a** and **19b** as raw materials. Compound **R-3**: Yellow solid, 86% yield. m.p. 56~57 °C; $[\alpha]_D^{25} + 22.4$ (c 0.13, CH₃OH), 91.5% *ee*; ¹H NMR (300 MHz, CDCl₃) δ : 5.18 (s, 1H), 4.04 (dd, $J=9.7$, 3.2 Hz, 1H), 3.67 (s, 3H), 2.69~2.41 (m, 2H), 2.11~1.98 (m, 2H), 1.27 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ : 171.55, 169.47, 92.09, 84.41, 71.29, 55.62, 31.14, 25.55, 24.80, 24.76; HR-ESI-MS calcd for $C_{10}H_{17}O_4$ $[M+H]^+$ 201.1121, found 201.1124. Compound **S-3**: Yellow solid, yield 82%. m.p. 55~57 °C; $[\alpha]_D^{25} - 21.3$ (c 0.15, CH₃OH), 92.3% *ee*; ¹H NMR, ¹³C NMR data were identical with **3a**; HR-ESI-MS calcd for $C_{10}H_{17}O_4$ $[M+H]^+$ 201.1121, found 201.1120.

3.6 Synthesis of **R-6** and **S-6** via Sharpless asymmetric dihydroxylation

2-(4-Methylpent-3-en-1-yl)benzoic acid (**6a**) (2.0 g, 10 mmol) was dissolved in methanol (60 mL) in a 100 mL flask, 0.5 mL of conc. H₂SO₄ was added to the solution. After reaction 5 h at reflux temperature, the solvent was evaporated and the residue was purified by silica gel column chromatography (petroleum ether/EtOAc, $V:V=100:1$) to give methyl 2-(4-methylpent-3-en-1-yl)benzoate (**20**) (1.9 g, 87%). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ : 7.83~7.86 (m, 1H), 7.37~7.43 (m, 1H), 7.20~7.25 (m, 2H), 5.18 (t, $J=7.0$ Hz, 1H), 3.88 (s, 3H), 2.97 (t, $J=7.5$ Hz, 2H), 2.24~2.32 (m, 2H), 1.67 (s, 3H), 1.51 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 168.18, 143.93, 132.14, 131.68, 131.00, 130.45, 129.69, 125.68, 123.72, 51.77, 34.45, 30.21, 25.64, 17.52; HR-ESI-MS calcd for $C_{14}H_{19}O_2$ $[M+H]^+$ 219.1379, found 219.1379.

In the similar procedure as **17a**~**19a** and **17b**~**19b**, (*R*)- and (*S*)-methyl 2-(3,4-dihydroxy-4-methylpentyl)benzoate (**21a**, **21b**) were prepared utilizing compound **20** as raw material. Compound **21a**: Colorless oil. 77% yield, $[\alpha]_D^{25} - 32.5$ (c 0.22, CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ : 7.89~7.86 (m, 1H), 7.47~7.42 (m, 1H), 7.31~7.23 (m, 2H), 3.89 (s, 3H), 3.41 (br, 1H), 3.34 (dd, $J=10.6$, 1.4 Hz, 1H), 3.20~2.99 (m, 2H), 2.43 (br, 1H), 1.90~1.80 (m, 1H), 1.73~1.62 (m, 1H), 1.15 (s, 3H), 1.13 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 168.38, 143.98, 132.14, 131.00, 130.62, 129.16, 125.90, 77.32, 72.71, 52.07, 33.72, 30.80, 26.06, 23.32; HR-ESI-MS calcd for $C_{14}H_{21}O_4$ $[M+H]^+$ 253.1434, found 253.1431. Compound **21b**: Colorless oil, 70% yield, $[\alpha]_D^{25} + 32.5$ (c 0.24, CH₃OH); ¹H NMR and ¹³C NMR were identical with that of **21a**; HR-ESI-MS calcd for $C_{14}H_{21}O_4$ $[M+H]^+$ 253.1434, found 253.1430.

In similar protocol as **R-1** and **S-1**, **R-2** and **S-2**, **R-3** and **S-3**, compounds **R-6** and **S-6** were prepared using **21a** and

21b as raw materials. Compound **R-6**: Colorless crystal, 55% yield. m.p. 45 ~ 46 °C; $[\alpha]_D^{25} - 42.0$ (c 0.64, CH₃OH), 89.3% ee. ¹H NMR (300 MHz, CDCl₃) δ : 7.71 ~ 7.74 (m, 1H), 7.46 ~ 7.52 (m, 1H), 7.35 ~ 7.40 (m, 1H), 7.21 ~ 7.28 (m, 1H), 3.86 (dd, $J = 4.8, 13.5$ Hz, 1H), 2.89 ~ 2.96 (m, 1H), 2.77 ~ 2.84 (m, 1H), 2.25 ~ 2.36 (m, 1H), 1.92 ~ 2.01 (m, 1H), 1.25 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ : 171.20, 138.22, 132.74, 131.33, 130.14, 128.60, 127.29, 84.20, 71.55, 29.54, 28.47, 26.19, 24.30; HR-ESI-MS calcd for C₁₃H₁₇O₃ [M + H]⁺ 221.1172, found 221.1170. Compound **S-6**: Colorless crystal, 56% yield. m.p. 44 ~ 46 °C; $[\alpha]_D^{25} + 42.2$ (c 0.52, CH₃OH), 88.8% ee. ¹H NMR and ¹³C NMR were identical with that of **R-6**; HR-ESI-MS calcd for C₁₃H₁₇O₃ [M + H]⁺ 221.1172, found 221.1169.

3.7 Preparation of Mosher esters of **18a**, **19a** and **21a**

18a (12.5 mg, 0.046 mmol), DMAP (22 mg, 0.184 mmol) and dry CH₂Cl₂ (3 mL) were added into a 25 mL of Schlenk flask under N₂ protection, then (*R*)-configuration Mosher acid chloride (17 μ L, 0.092 mmol) was added quickly and stirred at room temperature for 3 h. Then removed the solvent and prepared by thin layer chromatography with petroleum ether/ethyl acetate (*V* : *V* = 3 : 1) as eluent to give methyl (2*E*,6*R*)-7-hydroxy-7-methyl-6-(*R*-(3,3,3-trifluoro-2-methoxy-2-phenylpropanoyl)oxy)-3-(trifluoromethyl)oct-2-enoate (**22a**) as colorless oil in 75% yield. ¹H NMR (300 MHz, CDCl₃) δ : 7.60 ~ 7.59 (m, 2H), 7.43 ~ 7.41 (m, 3H), 5.94 (s, 1H), 4.99 (dd, $J = 10.0, 2.0$ Hz, 1H), 3.77 (s, 3H), 3.56 (s, 3H), 2.16 (t, $J = 7.7$ Hz, 2H), 1.85 ~ 1.74 (m, 1H), 1.72 ~ 1.66 (m, 1H), 1.57 (brs, 1H), 1.21 (s, 3H), 1.16 (s, 3H). In a similar procedure, methyl (2*E*,6*R*)-7-hydroxy-7-methyl-6-(*S*-(3,3,3-trifluoro-2-methoxy-2-phenylpropanoyl)oxy)-3-(trifluoromethyl)oct-2-enoate (**22b**) of **18a** with (*S*)-configuration Mosher acid chloride was prepared as colorless oil in 80% yield. ¹H NMR (300 MHz, CDCl₃) δ : 7.59 ~ 7.56 (m, 2H), 7.45 ~ 7.40 (m, 3H), 6.01 (s, 1H), 4.99 (dd, $J = 9.6, 2.4$ Hz, 1H), 3.78 (s, 3H), 3.54 (s, 3H), 2.29 ~ 2.21 (m, 2H), 1.95 ~ 1.88 (m, 1H), 1.86 ~ 1.76 (m, 1H), 1.18 (s, 3H), 1.16 (s, 3H).

The Mosher esters **23a**, **23b**, **24a**, **24b** of **19a** and **21a** with (*R*)- and (*S*)-configuration Mosher acid chloride were synthesized in 70% ~ 86% yields in the same protocol as **22a** and **22b**.

Methyl (2*E*,6*R*)-7-hydroxy-3-methoxy-7-methyl-6-(*R*-(3,3,3-trifluoro-2-methoxy-2-phenylpropanoyl)oxy)oct-2-enoate (**23a**): Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ : 7.63 ~ 7.60 (m, 2H), 7.41 ~ 7.39 (m, 3H), 5.01 (dd, $J = 8.4, 3.4$ Hz, 1H), 5.00 (s, 1H), 3.65 (s, 3H), 3.60 (s, 3H), 3.56 (s, 3H), 2.73 ~ 2.67 (m, 2H), 1.85 ~ 1.80 (m, 3H), 1.23 (s, 3H), 1.18 (s, 3H); HR-ESI-MS calcd for C₂₁H₂₈F₃O₇ [M + H]⁺ 449.1782, found 449.1778.

Methyl (2*E*,6*R*)-7-hydroxy-3-methoxy-7-methyl-6-(*S*-(3,3,3-trifluoro-2-methoxy-2-phenylpropanoyl)oxy)oct-2-enoate (**23b**): Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ : 7.63 ~ 7.61 (m, 2H), 7.41 ~ 7.39 (m, 3H), 5.03 (s, 1H),

5.00 (dd, $J = 8.6, 3.2$ Hz), 3.66 (s, 3H), 3.62 (s, 6H), 2.82 (t, $J = 7.8$ Hz), 2.01 ~ 1.81 (m, 3H), 1.25 (s, 3H), 1.16 (s, 3H), 1.11 (s, 3H). HR-ESI-MS calcd for C₂₁H₂₈F₃O₇ [M + H]⁺ 449.1782, found 449.1783.

Methyl 3*R*-2-(4-hydroxy-4-methyl-3-(*R*-(3,3,3-trifluoro-2-methoxy-2-phenylpropanoyl)oxy)pentyl)benzoate (**24a**): Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ : 7.87 (dd, $J = 7.8, 1.3$ Hz, 1H), 7.68 ~ 7.65 (m, 2H), 7.42 ~ 7.37 (m, 4H), 7.28 ~ 7.22 (m, 1H), 7.07 (d, $J = 7.6$ Hz, 1H), 5.11 (dd, $J = 9.5, 2.6$ Hz, 1H), 3.85 (s, 3H), 3.61 (s, 3H), 2.98 ~ 2.90 (m, 1H), 2.79 ~ 2.73 (m, 1H), 2.01 ~ 2.00 (m, 1H), 1.80 ~ 1.74 (m, 1H), 1.22 (s, 3H), 1.16 (s, 3H). HR-ESI-MS calcd for C₂₄H₂₈F₃O₆ [M + H]⁺ 469.1832, found 469.1834.

Methyl 3*R*-2-(4-hydroxy-4-methyl-3-(*S*-(3,3,3-trifluoro-2-methoxy-2-phenylpropanoyl)oxy)pentyl)benzoate (**24b**): Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ : 7.90 (d, $J = 7.7$ Hz, 1H), 7.65 ~ 7.64 (m, 2H), 7.43 ~ 7.41 (m, 4H), 7.29 ~ 7.26 (m, 1H), 7.18 ~ 7.15 (m, 1H), 5.09 (dd, $J = 8.9, 2.8$ Hz, 1H), 3.87 (s, 3H), 3.62 (s, 3H), 2.99 ~ 2.93 (m, 2H), 2.05 ~ 2.01 (m, 1H), 1.98 ~ 1.85 (m, 1H), 1.17 (s, 3H), 1.13 (s, 3H). HR-ESI-MS calcd for C₂₄H₂₈F₃O₆ [M + H]⁺ 469.1382, found 469.1386.

3.8 X-ray diffraction analysis of compounds **10** and **S-6**

The crystals of **10** and **S-6** were obtained from their *n*-hexane/alcohol (*V* : *V* = 1 : 20) solutions. Their structure parameters were as following: compound **10**, crystal size: 0.40 mm × 0.35 mm × 0.07 mm, molecular formula C₁₄H₁₈O₄, molecular weight 250.28, triclinic crystal system, *P*-1 space group, *a* = 0.91440(19) nm, *b* = 1.2056(6) nm, *c* = 1.2831(3) nm, $\alpha = 73.80(3)^\circ$, $\beta = 78.301(19)^\circ$, $\gamma = 76.53(3)^\circ$, *V* = 1.3605(8) nm³, *Z* = 4, $\rho_{\text{calc}} = 1.272$ mg/mm³, $\mu = 0.093$ mm⁻¹, *F*(000) = 536, *S* = 1.044, *R*₁ = 0.0477, *wR*₂ = 0.0870 [*I* > 2 σ (*I*) i.e. *F*_o > 4 σ (*F*_o)], *R*₁ = 0.0701, *wR*₂ = 0.0985 [all data], completeness 0.998. Compound **S-6**, crystal size: 0.40 mm × 0.35 mm × 0.35 mm, molecular formula C₁₃H₁₆O₃, molecular weight 220.26, orthorhombic crystal system, *P*2₁2₁2₁ space group, *a* = 0.88933(2) nm, *b* = 0.91866(2) nm, *c* = 1.36381(3) nm, *V* = 1.11422(4) nm³, *Z* = 4, $\rho_{\text{calc}} = 1.313$ mg/mm³, $\mu = 0.751$ mm⁻¹, *F*(000) = 472, *S* = 1.111, *R*₁ = 0.0344, *wR*₂ = 0.0860 [*I* > 2 σ (*I*) i.e. *F*_o > 4 σ (*F*_o)], *R*₁ = 0.0344, *wR*₂ = 0.0861 [all data], flack parameters 0.13(19), completeness 0.995. The crystallographic data of **10** and **S-6** in this paper have been deposited with the Cambridge Crystallographic Data Centre having following accession number-CCDC 1472491 and 1470573. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

3.9 Bioassay of fungicidal activity

The fungicidal activities of compounds **1** ~ **14** against *R. Solani*, *A. Solani*, *F. graminearum*, *S. Sclerotiorum*, *B. cinerea* and *P. capsici* were evaluated using methods in the references^[48-50] by mycelium growth rate. Procedure for inhibition rate: The stock 2000 mg/L DMSO solution of tested compounds was prepared in advance. Then hot PDA

culture medium was added into a plate, added sample solution or blank DMSO to the plate and mix with PDA culture medium, made the final concentration as desired. When plate was made, put a 5 mm diameter fungus cake into the center of plate, incubated them at $(25 \pm 0.5)^\circ\text{C}$ for 24~48 h, checked the growth status and calculated the inhibition rate according to the reference. Three replicates were performed and the mean measurements were calculated from three replicates for each concentration. The EC_{50} values were determined from the inhibition rates of five different concentrations based on the statistics method for the compounds which had more than 70% inhibition rates in the preliminary evaluation. Carbendazim and chlorothalonil were used as the positive control in the mycelium growth rate test.

References

- [1] Schulz, S.; Hotling, S. *Nat. Prod. Rep.* **2015**, *32*, 1042.
- [2] Gabriela, L.; Gallardo, N. I. P.; Cabrera, G. M. *Phytochem. Lett.* **2008**, *1*, 155.
- [3] Yang, Y.; Jiang, J. Z.; Qimei, L. B.; Yan, X. J.; Zhao, J. X.; Yuan, H. Z.; Qin, Z. H.; Wang, M. A. *Molecules* **2010**, *15*, 7075.
- [4] Davies-Coleman, M. T.; Rivett, D. E. A. *Prog. Chem. Org. Nat. Prod.* **1989**, *55*, 1.
- [5] Collett, L. A.; Davies-Coleman, M. T.; Rivett, D. E. A. *Prog. Chem. Org. Nat. Prod.* **1998**, *75*, 182.
- [6] Pereda-Miranda, R.; Fragosó-Serrano, M.; Cerda-García-Rojas, C. M. *Tetrahedron* **2001**, *57*, 47.
- [7] Hoffmann, H. M. R.; Rabe, J. *Angew. Chem. Int. Ed.* **1985**, *24*, 94.
- [8] Marco, J. A.; Carda, M.; Murga, J.; Falomir, E. *Tetrahedron* **2007**, *63*, 2929.
- [9] Boucard, V.; Broustal, G.; Campagne, J. M. *Eur. J. Org. Chem.* **2007**, 225.
- [10] Gerard, R. *Tetrahedron* **1995**, *51*, 2777.
- [11] Helena, M. C.; Fernanda, I. B.; Myrian, K. S.; Luiz, S. L. J. *Quim. Nova* **2008**, *31*, 885.
- [12] Helena, M. C.; Fernanda, I. B.; Luiz, S. L. J. *Synthesis* **2007**, 3261.
- [13] Shiina, I. *Chem. Rev.* **2007**, *107*, 239.
- [14] Chang, H.-T.; Jayanth, T. T.; Wang, C.-C.; Cheng, C.-H. *J. Am. Chem. Soc.* **2007**, *129*, 12032.
- [15] Ebine, M.; Suga, Y.; Fuwa, H.; Sasaki, M. *Org. Biomol. Chem.* **2010**, *8*, 39.
- [16] Emily, B. P.; Tendai, G.; Sonbinh, T. N. *Org. Lett.* **2008**, *10*, 5613.
- [17] van As, B. A. C.; Chan, D.; Kivit, P. J. J.; Palmans, A. R. A.; Meijer, E. W. *Tetrahedron: Asymmetry* **2007**, *18*, 787.
- [18] Maxime, A. S.; Huub, K.; Anthony, L. S. *Acta Crystallogr.* **2008**, *E64*, o607.
- [19] Fukuzawa, A.; Masamune, T. *Tetrahedron Lett.* **1981**, *22*, 4081.
- [20] Cremer, D. J.; Pople, A. J. *Am. Chem. Soc.* **1975**, *97*, 1354.
- [21] Wandjia, J.; Wansi, J. D.; Fuendjie, V.; Dagne, E.; Mulholland, D. A.; Tillequin, F.; Fomum, Z. T.; Sondengam, B. L.; Nkeh, B. C.; Njamen, D. *Phytochemistry* **2000**, *54*, 811.
- [22] Nkeh, B. C.; Njamen, D.; Wandjia, J.; Fomum, Z. T.; Dongmo, A.; Nguelefack, T. B.; Wansi, J. D.; Kamanyi, A. *Pharm. Biol.* **2003**, *41*, 26.
- [23] Jhulki, S.; Seth, S.; Mondal, M.; Moorthy, J. N. *Tetrahedron* **2014**, *70*, 2286.
- [24] Kadnikov, D. V.; Larock, R. C. *Mendeleev Commun.* **2007**, *17*, 74.
- [25] Cheng, Y. A.; Chen, T.; Tan, C. K.; Heng, J. J.; Yeung, Y. Y. *J. Am. Chem. Soc.* **2012**, *134*, 16492.
- [26] Wang, L.; Ren, Z.-L.; Ding, M.-W. *J. Org. Chem.* **2015**, *80*, 641.
- [27] Arcadi, A.; Fabrizi, G.; Goggiamani, A.; Marinelli, F. *J. Org. Chem.* **2015**, *80*, 6986.
- [28] Matsuda, T.; Shigeno, M.; Murakami, M. *Org. Lett.* **2008**, *10*, 5219.
- [29] Zarrabi, S.; Mahmoodi, N. O.; Tabatabaieian, K.; Zanjanchi, M. A. *Chin. Chem. Lett.* **2009**, *20*, 1400.
- [30] Dong H. B.; Yang M. Y.; Jiang J. Z.; Wang M. A. *J. Asian Nat. Prod. Res.* **2013**, *15*, 880.
- [31] Dong H. B.; Yang M. Y.; Tang B.; Wang M. A. *Chin. J. Org. Chem.* **2014**, *34*, 2350 (in Chinese).
(董宏波, 杨明艳, 汤博, 王明安, 有机化学, **2014**, *34*, 2350.)
- [32] Zhao J.; Dong H. B.; Yang M. Y.; Du J.; Jiang J. Z.; Wang M. A. *J. Asian Nat. Prod. Res.* **2014**, *16*, 312.
- [33] Wang, W. W.; Zhao, Y.; Liu, X. L.; Geng, R.; Wang, M. A. *Chin. J. Org. Chem.*, **2019**, *39*, 1129 (in Chinese).
(王卫伟, 赵宇, 刘鑫磊, 耿瑞, 王明安, 有机化学, **2019**, *39*, 1129.)
- [34] Srivastava, A. K.; Song, H.; Park, S. B. *Synthesis* **2011**, 1708.
- [35] Gesinski, M. R.; Tadpetch, K.; Rychnovsky, S. D. *Org. Lett.* **2009**, *11*, 5342.
- [36] Matsuda, T.; Shigeno, M.; Murakami, M. *Org. Lett.* **2008**, *10*, 5219.
- [37] Chang, H. T.; Jeganmohan, M.; Cheng, C. H. *Chem. Commun.* **2005**, 4955.
- [38] Roux, M. C.; Paugam, R.; Rousseau, G. *J. Org. Chem.* **2001**, *66*, 4304.
- [39] Wong, O. A.; Shi, Y. *Chem. Rev.* **2008**, *108*, 3958.
- [40] Dong, H. B.; Yang, M. Y.; Liu, B.; Wang, M. A. *J. Asian Nat. Prod. Res.* **2014**, *16*, 847.
- [41] Kolb, H. C.; Van Nieuwenhze, M. S.; Sharpless, K. B. *Chem. Rev.* **1994**, *94*, 2483.
- [42] Dong, H. B.; Yang, M. Y.; Zhang, X. T.; Wang, M. A. *Tetrahedron: Asymmetry* **2014**, *25*, 610.
- [43] Jorge, G.-F.; Juan, M.; Miguel, C.; Alberto, M. J. *Org. Lett.* **2006**, *8*, 2695.
- [44] Narendar, T.; Sarkar, S.; Rajendar, K.; Tiwari, S. *Org. Lett.* **2011**, *13*, 6140.
- [45] Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989.
- [46] Yadav, J. S.; Rajendar, G.; Rao, R. S.; Pabbaraja, S. *J. Org. Chem.* **2013**, *78*, 8524.
- [47] Ikuko, O.; Takenori, K.; Yoel, K.; Hiroshi, K. *J. Am. Chem. Soc.* **1991**, *113*, 4092.
- [48] Tang, B.; Guan, A. Y.; Zhao, Y.; Jiang, J. Z.; Wang, M. A.; Zhou, L. G. *Chin. J. Chem.* **2017**, *35*, 1133.
- [49] Zhao, Y.; Tang, B.; Guan, A. Y.; Wang, W. W.; Zhang, Z. H.; Wang, M. A. *Synthesis* **2017**, *49*, 4663.
- [50] Geng, R.; Zhao, Y.; Li, Y. H.; Liu, X. L.; Wang, M. A. *Chin. J. Org. Chem.* **2019**, *39*, 3574 (in Chinese).
(耿瑞, 赵宇, 李益豪, 刘鑫磊, 王明安, 有机化学, **2019**, *39*, 3574.)

(Fan, Y.)