铜(I)催化邻炔基芳基异硫氰酸酯与(*E*)-2-(苄基氨基)乙酸酯串联双环化反应快速合成 5*H*-苯并咪唑[5,1-*b*][1,3]噻嗪

张亚辉 吴文锦 张可心 黎双双 郝文燕*

(江西师范大学化学与化工学院 功能有机小分子教育部重点实验室 江西省绿色化学重点实验室 南昌 330022)

摘要 以廉价的铜盐作为催化剂,快速高效地实现了邻炔基芳基异硫氰酸酯与(*E*)-2-(苄基氨基)乙酸酯的串联双环化反应,以中等至良好产率合成了一系列 5*H*-苯并[*d*]咪唑[5,1-*b*][1,3]噻嗪类化合物.该反应操作简单,利用一步[3+2]环加成继而分子内亲核加成串联机制,一锅法高效构建了两个新环.

关键词 铜催化;串联双环化;5*H*-苯并咪唑[5,1-*b*][1,3]噻嗪

A Rapid Synthesis of 5*H*-Benzo[*d*]imidazo[5,1-*b*][1,3]thiazines via Copper(I)-Catalyzed Cascade Bicyclization of *o*-Alkynylphenyl Isothiocyanates with Ethyl (*E*)-2-(Benzylideneamino)acetates

Zhang, Yahui Wu, Wenjin Zhang, Kexin Li, Shuangshuang Hao, Wenyan*

(Key Laboratory of Functional Small Organic Molecules of Ministry of Education, Jiangxi Key Laboratory of Green Chemistry, College of Chemistry & Chemical Engineering, Jiangxi Normal University, Nanchang 330022)

Abstract A rapid and efficient protocol for copper(I)-catalyzed tandem bicyclization of *o*-alkynylphenyl isothiocyanates with ethyl (*E*)-2-(benzylideneamino)acetates has been developed, affording a series of 5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazines in moderate to good yields. In this transformation, a [3+2] cycloaddition reaction mechanism was involved and two new rings were formed in one pot.

Keywords copper-catalysis; cascade bicyclization; 5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazine

1 Introduction

Imidazoles is an important class of five-membered heterocycles with significant biological activities.^[1] Among their derivatives, imidazobenzothiazines show a broad range of biological activities and attract considerable attention in medicinal and pharmaceutical sciences. Some of these compounds have been used as analgesic drugs with potential anti-inflammatory, antimicrobial and anti-helminthic activity,^[2] as selective inhibitors of COX-2 with affinity for the omega modulatory sites of GABAA receptors.^[3] Therefore, the synthesis of libraries of such compounds should be interesting. However, only a few practical methods have been developed for the synthesis of these imidazobenzothiazines.^[4] Sequential nucleophilic addition/salinization/cyclization reactions of 2-aminobenzhy-

drol with thiourea are used for the synthesis of 5*H*-benzo[*d*]imidazo[2,1-*b*][1,3]thiazines.^[5] Another approach to functionalized benzimidazobenzothiazines was getting a precursor via a S_N2 reaction and then adding NaH for cyclization.^[6] Ullmann coupling of 2-mercaptoimidazoles and bromobenzyl bromides provides an alternative method for synthesizing these frameworks.^[7] However, most of these mentioned reactions have more or less disadvantages such as narrow substrate scope, long reaction time and tedious multi-step manipulations.

Recently, transition-metal-catalyzed cascade bicyclization of *o*-alkynylphenyl isothiocyanates with various nucleophiles provides a new and powerful synthetic strategy to construct different heterocycles such as indoles,^[8] quinoline,^[9] thiazine^[10] and so on. For example, Saito group^[8c] developed an Rh-mediated Pauson-Khand (PK) reaction to

* Corresponding author. E-mail: wenyanhao@jxnu.edu.cn

Received December 2, 2020; revised January 18, 2021; published online January 28, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21762023, 22061021) and the Natural Science Foundation of Jiangxi Province (No. 2018BAB203006).

国家自然科学基金(Nos. 21762023, 22061021)和江西省自然科学基金(No. 2018BAB203006)资助项目.

construct a 3-substituted-2*H*-thieno[2,3-*b*]indol-2-ones by using *o*-alkynylphenyl isothiocyanates and CO as the substrates (Scheme 1, a). Li and co-workers^[8c,11] also developed two cases of cascade functionalization of isonitriles and diaryliodonium salts with *o*-alkynylphenyl isothiocyanates to construct thieno[2,3-*b*]indole and thiochromeno[2,3-*b*]indole heterocycles (Schemes 1, b and c). Wang group^[12] reported a Cu-catalyzed cascade bicyclization for the synthesis of 2-(1*H*-indol-1-yl)-4,5-dihydrothiazoles from *o*-alkynylphenyl isothiocyanates and propargylamine (Scheme 1, d). Recently, a Silver-catalyzed cascade bicyclization of *o*-alkynylphenyl isothiocyanates with sodium azide for the synthesis of 5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazines has been reported by our group (Scheme 1, e).^[13] Encouraged by these fascinating researches and our continuing interest in the transformation of *o*-alkynylphenyl isothiocyanates,^[14] we herein report an efficient copper-catalyzed cyclization of *o*-alkynylphenyl isothiocyanates with ethyl (*E*)-2-(benzylideneamino)acetates for the synthesis of 5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazines (Scheme 1, f).

2 Results and discussion

Similar to the process reported in previous literature,^[14] our initial study began with 1.0 equiv. of 1-isothiocyanato-2-(phenylethynyl)benzene (**1a**) and 2.0 equiv. of ethyl (*E*)-2-(benzylideneamino)acetates (**2a**) with 2.0 equiv. of Cs₂CO₃ in the presence of Cu(OAc)₂ (10 mol%) in 2 mL of anhydrous MeCN at 80 °C for 30 min. To our delight, desired product **3aa** was successfully isolated in 40% yield (Table 1, Entry 1). Then, other copper catalysts were screened, and CuCl was found to be the best choice (Table 1, Entries 2~7). A blank experiment indicated that the copper salt can promote the reaction, because desired product **3a** was obtained in only 30% yield in the absence of the copper catalyst (Table 1, Entry 8). The reaction also did not go well in the absence of a base (Table 1, Entry 9), which indicated that the combination of a copper catalyst and a base was helpful to improve the yield. Reducing the amount of copper salts to 5 mol%, the yield increased slightly (Table 1, Entry 10). The effect of base on the reaction was next investigated. The yield was improved to 81% if KOH was employed, whereas phosphorus salt failed to promote the reaction. Other bases such as Na₂CO₃, K₂CO₃, NaOH, *t*-BuOK and 1,5-diazabicyclo[5.4.0]undec-5-ene (DBU) were substantially less effective (Table 1, Entries 11~18). The solvent of the reaction was also screened. No improved yield was observed by using 1,2-dichloroethane (DCE), toluene, dioxane, tetrahydrofuran (THF), CH₃OH, *N,N*-dimethylformamide (DMF) and dimethyl sulfoxide (DMSO) as solvents (Table 1, Entries 19~25). Finally, the effect of temperature on the reaction was examined and it was found that 25 °C gave the best result (Table 1, Entries 26~30). Therefore, the optimized conditions include the use of a combination of CuCl (5 mol%) as the catalyst and KOH (2.0 equiv.) as the base in 2 mL of MeCN at 25 °C

Table 1 Initial studies for the tandem reaction of 1-isothiocyanato-2-(phenylethynyl)benzene (**1a**) with ethyl (*E*)-2-(benzylideneamino)acetate (**2a**)^a

Entry	Catalyst	Base	Solvent	Temp./°C	Yield ^b /%
1	Cu(OAc) ₂	Cs ₂ CO ₃	MeCN	80	40
2	Cu(OTf) ₂	Cs ₂ CO ₃	MeCN	80	36
3	CuBr ₂	Cs ₂ CO ₃	MeCN	80	42
4	CuO	Cs ₂ CO ₃	MeCN	80	47
5	CuI	Cs ₂ CO ₃	MeCN	80	42
6	CuBr	Cs ₂ CO ₃	MeCN	80	47
7	CuCl	Cs ₂ CO ₃	MeCN	80	52
8		Cs ₂ CO ₃	MeCN	80	30
9	CuCl		MeCN	80	32
10 ^c	CuCl	Cs ₂ CO ₃	MeCN	80	53
11 ^c	CuCl	K ₃ PO ₄	MeCN	80	Trace
12 ^c	CuCl	Na ₂ HPO ₄	MeCN	80	Trace
13 ^c	CuCl	K ₂ CO ₃	MeCN	80	53
14 ^c	CuCl	Na ₂ CO ₃	MeCN	80	56
15 ^c	CuCl	NaOH	MeCN	80	79
16 ^c	CuCl	KOH	MeCN	80	81
17 ^c	CuCl	<i>t</i> -BuOK	MeCN	80	65
18 ^c	CuCl	DBU	MeCN	80	70
19 ^c	CuCl	KOH	DCE	80	37
20 ^c	CuCl	KOH	Toluene	80	44
21 ^c	CuCl	KOH	1,4-Dioxane	80	33
22 ^c	CuCl	KOH	THF	80	40
23 ^c	CuCl	KOH	CH ₃ OH	80	36
24 ^c	CuCl	KOH	DMF	80	56
25 ^c	CuCl	KOH	DMSO	80	62
26 ^c	CuCl	KOH	MeCN	120	Trace
27 ^c	CuCl	KOH	MeCN	100	66
28 ^c	CuCl	KOH	MeCN	60	77
29 ^c	CuCl	KOH	MeCN	40	79
30 ^c	CuCl	KOH	MeCN	25	83

^a Reaction was performed with **1a** (0.2 mmol), **2a** (0.4 mmol), catalyst (0.02 mmol), base (0.4 mmol) in solvent (2 mL) for 30 min. ^b Isolated yield based on 1-isothiocyanato-2-(phenylethynyl)benzene (**1a**). ^c The amount of copper salt is 5 mol%.

(Table 1, Entry 30).

Under the optimal condition, a variety of *o*-alkynylphenyl isothiocyanates **1** with ethyl (*E*)-2-(benzylideneamino)acetate (**2a**) were investigated. As shown in Table 2, all reactions proceeded smoothly, leading to the desired 5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazines in moderate to good yields. For example, the substituents on the R² position of *o*-alkynylphenyl isothiocyanates showed obvious electronic effects on the reaction. When the R² group in the

Scheme 1 Synthesis of N,S-containing heterocycles through *o*-alkynylphenyl isothiocyanates

Table 2 Substrate scope of diverse *o*-alkynylphenyl isothiocyanates^{a,b}

^a Reaction conditions: *o*-alkynylphenyl isothiocyanates (**1**, 0.2 mmol), (*E*)-2-(benzylideneamino)acetate (**2a**, 0.4 mmol), KOH (0.4 mmol), and CuCl (5 mol%) in MeCN (2 mL) at r.t. ^b Isolated yield based on *o*-alkynylphenyl isothiocyanates **1**.

substrates **1** was an electron-deficient aryl, such as *p*-FC₆H₄, *p*-ClC₆H₄ and *p*-BrC₆H₄, the reactions with **2a** proceeded smoothly to afford the corresponding products **3ba**~**3da** in lower yields compared with the reactions of substrates **1** with an electron-rich aryl group such as *p*-MeC₆H₄ and *p*-MeOC₆H₄ at the R² position, which afforded the corresponding products **3ea** and **3fa** in 87% and 88% yields, respectively. No desired products were obtained when the R² group in the substrate *o*-alkynylphenyl isothiocyanates **1** was an alkyl group, such as *n*-butyl, *t*-butyl and *n*-hexyl.

This may be due to the fact that alkyl group cannot conjugate with an acetylenic bond and activate it. In addition, the reactions of *o*-alkynylphenyl isothiocyanates bearing various substituents such as fluoro, chloro, bromo, trifluoromethyl and methyl groups on the aryl rings, regardless of their substitution positions, gave the desired products **3ga**~**3oa** in moderate to good yields. Similarly, the substitution of electron-donating groups in the reaction was significantly better than that of electron-withdrawing groups (**3oa**/**3ja**). Later, the *o*-alkynylphenyl isothiocyanate was replaced to 2-isothiocyanato-3-(phenylethynyl)pyridine, the corresponding product **3pa** was also obtained in 70% yield. Products ethyl (Z)-7-bromo-5-(4-bromobenzylidene)-1-phenyl-5*H*-benzo[d]imidazo[5,1-*b*][1,3]thiazine-3-carboxylate (**3qa**)

and ethyl (Z)-5-(4-fluorobenzylidene)-1-phenyl-7-(trifluoromethyl)-5*H*-benzo[d]imidazo[5,1-*b*][1,3]thiazine-3-carboxylate (**3ra**) could also be conveniently obtained in 70% and 54% yields, respectively. It is noteworthy that only *Z*-isomers were formed which is determined by single-crystal X-ray analysis of **3aa** (Figure 1). This result may be due to kinetic effects according to Baldwin's rules.^[15]

To further assess the universality of this reaction, a variety of imines with *o*-alkynylphenyl isothiocyanates (**1a**) were examined. As shown in Table 3, methyl (*E*)-2-(benzylideneamino)acetates could get the corresponding product in 86% yield (**3ab**). Besides, this reaction tolerated lots of substituents on phenyl ring of (*E*)-2-(benzylideneamino)acetates containing electron-donating and electron-withdrawing groups in moderate yields (**3ac**~**3ai**). However, to our disappointment, the corresponding products could not be obtained when R³ was hydroxyl group and *n*-butyl group (**3aj**).

The structure of ethyl (Z)-5-benzylidene-1-phenyl-5*H*-benzo[d]imidazo[5,1-*b*][1,3]thiazine-3-carboxylate (**3aa**) was corroborated by X-ray diffraction analysis of the crystal structure of **3aa**, and its ORTEP diagram is displayed in Figure 1.

A plausible mechanism is proposed in Scheme 2. Firstly,

Table 3 Substrate scope of diverse (*E*)-2-(benzylideneamino)acetates^{a,b}

^a Reaction conditions: 1-isothiocyanato-2-(phenylethynyl)benzene (**1a**, 0.2 mmol), (*E*)-2-(benzylideneamino)acetates (**2**, 0.4 mmol), KOH (0.4 mmol), and CuCl (5 mol%) in MeCN (2 mL) at r.t. for 30 min. ^b Isolated yield based on *o*-alkynylphenyl isothiocyanate **1a**.

Figure 1 Single-crystal X-ray diffraction structure of **3aa**. Thermal ellipsoids are at a 30% probability level (CCDC number is 2047019)

in the presence of copper(I) chloride and the base, the C—H bond of ethyl (*E*)-2-(benzylideneamino)acetates (**2**) would be activated to afford the intermediate **A**. Then, intermediate **A** reacts with *o*-alkynylphenyl isothiocyanate through a concerted [3 + 2] cycloaddition process to generate the organo-copper intermediate **B** which undergoes subsequent isomerization to afford intermediate **C**. Next, protonation of the N—Cu bond of the intermediate **C** by ethyl (*E*)-2-(benzylideneamino)acetates **2** would produce the compound **D** with the regeneration of the copper intermediate **A** to end the catalytic cycle. Finally, the subsequent copper-catalyzed intra-molecular nucleophilic cyclization of compound **D** would occur to give product **3**.

3 Conclusions

In summary, a rapid method for the synthesis of 5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazines via the tandem bicyclization reaction of *o*-alkynylphenyl isothiocyanates with ethyl (*E*)-2-(benzylideneamino)acetates was developed. Different kinds of 5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazines could be synthesized in moderate to good yields. In this reaction, a [3 + 2] cycloaddition was proposed to be in-

volved and two new rings were formed in a one-pot reaction. This present cascade bicyclization strategy represents an effective way to construct small molecular N,S-heterocycles with great value.

4 Experimental section

4.1 Instruments and reagents

¹H NMR and ¹³C NMR spectra were recorded in 400 MHz apparatus and the frequencies for ¹H NMR and ¹³C NMR test are 400 MHz and 100 MHz, respectively. The mass analyzer type used for the HRMS measurements is micro TOF. Yields refer to chromatographically and spectroscopically pure compounds, unless otherwise indicated. All reagents and metal catalysts were obtained from commercial sources without further purification, and commercially available solvents were purified before use. All reactions were performed in reaction tubes. All new compounds were fully characterized. Silica gel plate GF254 was used for thin layer chromatography (TLC) and silica gel H or 300~400 mesh was used for flash column chromatography. Thin layer chromatography plates were visualized by exposure to ultraviolet light.

4.2 General procedure for the synthesis of product **3**

A mixture of *o*-alkynylphenyl isothiocyanates **1** (0.20 mmol), ethyl (*E*)-2-(benzylideneamino)acetates **2** (0.4 mmol) and KOH (0.40 mmol) was added into a tube. Subsequently MeCN (2 mL) was added. Then, the sealed tube was heated at room temperature for 30 min. After completion of reaction as indicated by TLC, the mixture was concentrated and directly purified by flash column chromatography (EtOAc/petroleum ether, *V* : *V* = 1 : 2) to give the desired product **3**.

Ethyl (*Z*)-5-benzylidene-1-phenyl-5*H*-benzo[*d*]imidazo[5,1-*b*][1,3]thiazine-3-carboxylate (**3aa**): Yellow solid, 70.4 mg, 83% yield. m.p. 92.5~94.5 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.62~7.56 (m, 4H), 7.45 (t, *J* = 7.2 Hz, 2H), 7.36~7.42 (m, 5H), 7.31~7.33 (m, 2H), 7.13 (t, *J* =

Scheme 2 Plausible mechanism

7.8 Hz, 1H), 6.98 (d, $J=8.0$ Hz, 1H), 4.41 (q, $J=7.2$ Hz, 2H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 147.3, 134.8, 133.6, 132.8, 132.0, 130.99, 129.8, 129.6, 129.2, 128.9, 128.7, 128.7, 128.5, 127.8, 127.4, 124.2, 122.1, 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{21}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 425.1318, found 425.1323.

Ethyl (Z)-5-(4-fluorobenzylidene)-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ba**): Yellow solid, 56.6 mg, 64% yield. m.p. 101.1~102.8 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.62~7.55 (m, 5H), 7.46~7.38 (m, 4H), 7.33 (t, $J=7.2$ Hz, 1H), 7.16 (d, $J=8.4$ Hz, 3H), 7.00 (d, $J=8.0$ Hz, 1H), 4.43 (q, $J=7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.57, 162.54 (d, $^1J_{\text{CF}}=248$ Hz), 147.3, 133.2, 132.8, 131.6 (d, $^3J_{\text{CF}}=9$ Hz), 131.05, 131.02, 130.9, 130.8, 129.6, 129.5, 128.9, 128.7, 128.7, 127.9, 127.5 (d, $^2J_{\text{CF}}=26$ Hz), 124.1, 122.1, 115.6 (d, $^2J_{\text{CF}}=22$ Hz), 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{FN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 443.1224, found 443.1234.

Ethyl (Z)-5-(4-chlorobenzylidene)-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ca**): Yellow solid, 65.0 mg, 71% yield. m.p. 115.2~117.2 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.58~7.62 (m, 3H), 7.52 (d, $J=8.4$ Hz, 2H), 7.41~7.45 (m, 5H), 7.34 (t, $J=7.6$ Hz, 1H), 7.25 (s, 1H), 7.16 (t, $J=7.6$ Hz, 1H), 7.00 (d, $J=8.4$ Hz, 1H), 4.43 (q, $J=7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.5, 147.4, 134.4, 133.3, 133.0, 132.8, 131.0, 130.9, 130.5, 129.7, 129.4, 129.0, 128.9, 128.8, 128.8, 127.9, 127.7, 127.4, 125.1, 122.1, 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{ClN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 459.0929, found 459.0930.

Ethyl (Z)-5-(4-bromobenzylidene)-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3da**): Yellow solid, 69.3 mg, 69% yield. m.p. 121.6~123.1 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.50~7.55 (m, 4H), 7.38~7.31 (m, 5H), 7.26 (t, $J=7.6$ Hz, 1H), 7.17 (d, $J=16.4$ Hz, 2H), 7.08 (d, $J=7.2$ Hz, 1H), 6.92 (d, $J=8.4$ Hz, 1H), 4.35 (q, $J=7.2$ Hz, 2H), 1.34 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.5, 147.4, 133.7, 132.9, 132.8, 131.7, 131.2, 130.9, 130.6, 129.7, 129.4, 129.00, 128.9, 128.8, 127.9, 127.7, 127.5, 125.2, 122.7, 122.1, 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{BrN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 503.0423, found 503.0428.

Ethyl (Z)-5-(4-methylbenzylidene)-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ea**): Yellow solid, 76.2 mg, 87% yield. m.p. 108.4~109.4 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.58~7.62 (m, 3H), 7.48 (d, $J=8.0$ Hz, 2H), 7.38~7.43 (m, 3H), 7.32 (t, $J=7.2$ Hz, 1H), 7.27 (d, $J=6.0$ Hz, 2H), 7.20~7.08 (m, 2H), 6.98 (d, $J=8.4$ Hz, 1H), 4.42 (q, $J=7.2$ Hz, 2H), 2.41 (s, 3H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 147.2, 138.7, 132.1, 132.0, 131.0, 129.9, 129.7, 129.6, 129.3, 128.9, 128.7, 128.5, 128.5, 127.8, 127.7, 127.3, 123.1, 122.0, 60.8, 21.4, 14.5; HRMS calcd for $\text{C}_{27}\text{H}_{23}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 439.1475, found 439.1485.

Ethyl (Z)-5-(4-methoxybenzylidene)-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3fa**): Yellow solid, 79.9 mg, 88% yield. m.p. 115.5~117.5 °C;

^1H NMR (400 MHz, CDCl_3) δ : 7.55~7.61 (m, 4H), 7.36~7.45 (m, 4H), 7.32 (d, $J=7.2$ Hz, 1H), 7.25 (d, $J=8.4$ Hz, 1H), 7.06~7.18 (m, 2H), 6.97~7.00 (m, 2H), 4.43 (q, $J=7.2$ Hz, 2H), 3.87 (s, 3H), 1.42 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 159.8, 147.2, 133.9, 132.8, 131.7, 131.3, 131.1, 130.1, 129.5, 129.2, 128.9, 128.7, 128.3, 127.7, 127.6, 127.3, 122.0, 121.8, 114.0, 60.8, 55.4, 14.5; HRMS calcd for $\text{C}_{27}\text{H}_{23}\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$ 455.1424, found 455.1434.

Ethyl (Z)-5-benzylidene-7-fluoro-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ga**): Yellow solid, 54.8 mg, 62% yield. m.p. 100.5~101.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.58 (d, $J=7.2$ Hz, 3H), 7.37~7.49 (m, 6H), 7.28~7.35 (m, 3H), 6.96~6.99 (m, 1H), 6.84~6.89 (m, 1H), 4.42 (q, $J=7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.5, 161.1 (d, $^1J_{\text{CF}}=248$ Hz), 147.1, 134.4, 133.0, 132.9, 131.8, 130.8, 129.8, 129.7, 129.2, 128.9, 128.6, 128.4, 128.3, 128.0, 123.8 (d, $^3J_{\text{CF}}=9$ Hz), 123.4, 115.7 (d, $^2J_{\text{CF}}=23$ Hz), 114.3 (d, $^2J_{\text{CF}}=24$ Hz), 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{FN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 443.1224, found 443.1237.

Ethyl (Z)-5-benzylidene-7-chloro-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ha**): Yellow solid, 68.7 mg, 75% yield. m.p. 116.0~116.9 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.59 (dd, $J=13.6, 4.8$ Hz, 3H), 7.48~7.36 (m, 8H), 7.30 (s, 1H), 7.10 (dd, $J=8.4, 2.0$ Hz, 1H), 6.92 (d, $J=8.4$ Hz, 1H), 4.41 (q, $J=7.2$ Hz, 2H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 147.2, 134.4, 133.2, 133.0, 132.9, 131.8, 131.4, 131.2, 130.7, 129.8, 129.6, 128.9, 128.8, 128.6, 128.4, 128.0, 127.5, 123.2, 123.0, 61.0, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{ClN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 459.0929, found 459.0939.

Ethyl (Z)-5-benzylidene-7-bromo-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ia**): Yellow solid, 74.3 mg, 74% yield. m.p. 120.8~121.8 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.69 (d, $J=2.0$ Hz, 1H), 7.50 (d, $J=7.2$ Hz, 4H), 7.42~7.24 (m, 6H), 7.09~7.19 (m, 2H), 6.78 (d, $J=8.4$ Hz, 1H), 4.34 (q, $J=7.2$ Hz, 2H), 1.33 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 147.2, 134.4, 133.1, 131.9, 131.8, 131.5, 131.4, 130.7, 130.4, 129.8, 128.9, 128.8, 128.6, 128.4, 128.3, 128.0, 123.4, 122.9, 120.8, 61.0, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{BrN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 503.0423, found 503.0427.

Ethyl (Z)-5-benzylidene-1-phenyl-7-(trifluoromethyl)-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ja**): Yellow solid, 53.1 mg, 54% yield. m.p. 106.5~107.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.89 (s, 1H), 7.59 (d, $J=6.8$ Hz, 3H), 7.38~7.51 (m, 8H), 7.29~7.33 (m, 1H), 7.11 (d, $J=6.8$ Hz, 1H), 4.43 (q, $J=7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 147.6, 135.4, 134.3, 133.7, 131.8, 130.5, 130.2, 130.0, 129.9, 129.0, 128.8, 128.8, 128.6, 128.5, 128.4, 128.2, 125.3 (q, $J_{\text{CF}_3}=4$ Hz), 125.1 (q, $J_{\text{CF}_3}=4$ Hz), 122.9, 122.5, 61.0, 14.5; HRMS calcd for $\text{C}_{27}\text{H}_{20}\text{F}_3\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 493.1192, found 493.1193.

Ethyl (Z)-5-benzylidene-7-methyl-1-phenyl-5H-benzo-[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ka**): Yel-

low solid, 78.0 mg, 89% yield. m.p. 106.6~108.6 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.57~7.60 (m, 3H), 7.35~7.49 (m, 7H), 7.31 (s, 2H), 6.95 (d, $J=7.6$ Hz, 1H), 6.85~6.90 (m, 1H), 4.42 (q, $J=7.2$ Hz, 2H), 2.39 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 147.1, 137.4, 134.9, 133.3, 131.6, 131.1, 130.6, 129.8, 129.5, 129.4, 129.0, 128.9, 128.8, 128.7, 128.5, 128.0, 127.7, 124.4, 121.9, 60.8, 21.0, 14.56; HRMS calcd for $\text{C}_{27}\text{H}_{23}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 439.1475, found 439.1482.

Ethyl (Z)-5-benzylidene-8-fluoro-1-phenyl-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3la**): Yellow solid, 53.0 mg, 60% yield. m.p. 102.6~103.9 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.63~7.53 (m, 5H), 7.55~7.61 (m, 5H), 7.38 (t, $J=7.2$ Hz, 1H), 7.25 (s, 1H), 7.05 (td, $J=8.6, 2.5$ Hz, 1H), 6.69 (dd, $J=10.0, 2.4$ Hz, 1H), 4.42 (q, $J=7.2$ Hz, 2H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.5, 162.0 (d, $^1J_{\text{CF}}=248$ Hz), 147.4, 134.6, 133.9 (d, $^3J_{\text{CF}}=11$ Hz), 133.5, 132.1, 130.4, 130.0, 129.7, 129.5 (d, $^3J_{\text{CF}}=9$ Hz), 129.0, 128.9, 128.6, 128.6, 127.9, 125.7, 123.4, 114.6 (d, $^2J_{\text{CF}}=21$ Hz), 109.5 (d, $^2J_{\text{CF}}=27$ Hz), 61.0, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{FN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 443.1224, found 443.1234.

Ethyl (Z)-5-benzylidene-8-chloro-1-phenyl-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ma**): Yellow solid, 66.0 mg, 72% yield. m.p. 115.2~117.2 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.59 (dd, $J=13.6, 4.8$ Hz, 4H), 7.49~7.37 (m, 7H), 7.31 (s, 1H), 7.11 (dd, $J=8.4, 2.0$ Hz, 1H), 6.92 (d, $J=8.4$ Hz, 1H), 4.42 (q, $J=7.2$ Hz, 2H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 147.2, 134.4, 133.2, 133.0, 132.9, 131.8, 131.4, 131.2, 130.7, 129.8, 129.6, 128.9, 128.8, 128.6, 128.3, 127.5, 123.2, 123.0, 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{ClN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 459.0929, found 459.0939.

Ethyl (Z)-5-benzylidene-8-bromo-1-phenyl-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3na**): Yellow solid, 71.3 mg, 71% yield. m.p. 123.5~124.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (d, $J=2.0$ Hz, 1H), 7.58 (d, $J=7.2$ Hz, 4H), 7.49~7.37 (m, 7H), 7.31 (s, 1H), 6.85 (d, $J=8.8$ Hz, 1H), 4.41 (q, $J=7.2$ Hz, 2H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 147.2, 134.4, 133.1, 131.9, 131.8, 131.5, 131.4, 130.7, 130.4, 129.8, 128.9, 128.8, 128.8, 128.6, 128.4, 128.0, 123.4, 122.9, 120.7, 61.0, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{BrN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 503.0423, found 503.0437.

Ethyl (Z)-5-benzylidene-8-methyl-1-phenyl-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3oa**): Yellow solid, 75.3 mg, 86% yield. m.p. 107.3~109.6 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.52~7.49 (m, 4H), 7.40~7.27 (m, 7H), 7.22 (s, 1H), 6.86 (dd, $J=8.4, 1.2$ Hz, 1H), 6.79 (d, $J=8.4$ Hz, 1H), 4.34 (q, $J=7.2$ Hz, 2H), 2.30 (s, 3H), 1.32 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 147.1, 137.3, 134.9, 133.3, 131.6, 131.1, 130.6, 129.8, 129.5, 129.4, 129.4, 128.9, 128.7, 128.5, 128.5, 128.0, 127.7, 124.4, 121.9, 60.8, 21.0, 14.5; HRMS calcd for $\text{C}_{27}\text{H}_{23}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 439.1475, found 439.1487.

Ethyl (Z)-5-(4-chlorobenzylidene)-9-phenyl-5H-imidazo[5,1-b]pyrido[2,3-d][1,3]thiazine-7-carboxylate

(**3pa**): Yellow solid, 64.3 mg, 70% yield. m.p. 109.4~111.4 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.17 (d, $J=2.8$ Hz, 1H), 7.96 (d, $J=8.0$ Hz, 1H), 7.65~7.50 (m, 4H), 7.44 (d, $J=8.4$ Hz, 2H), 7.29~7.41 (m, 4H), 7.19 (d, $J=5.6$ Hz, 1H), 4.44 (td, $J=13.8, 6.8$ Hz, 2H), 1.42 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.7, 148.2, 145.1, 135.6, 134.7, 133.0, 132.3, 131.0, 130.9, 130.1, 129.2, 129.0, 128.9, 128.8, 128.7, 127.89, 127.82, 127.7, 124.1, 123.2, 123.1, 61.0, 14.5; HRMS calcd for $\text{C}_{25}\text{H}_{19}\text{ClN}_3\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 460.0881, found 460.0892.

Ethyl (Z)-7-bromo-5-(4-bromobenzylidene)-1-phenyl-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3qa**): Yellow solid, 81.1 mg, 70% yield. m.p. 175.5~177.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (d, $J=2.0$ Hz, 1H), 7.57 (d, $J=6.0$ Hz, 1H), 7.52 (d, $J=8.4$ Hz, 2H), 7.39~7.48 (m, 5H), 7.24~7.34 (m, 3H), 6.86 (d, $J=8.8$ Hz, 1H), 4.43 (dt, $J=14.3, 7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 147.3, 134.8, 132.9, 132.6, 131.9, 131.7, 131.6, 131.1, 131.0, 130.6, 130.3, 129.9, 128.9, 128.8, 128.4, 127.4, 123.7, 123.5, 120.8, 61.0, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{19}\text{Br}_2\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 580.9529, found 580.9534.

Ethyl (Z)-5-(4-fluorobenzylidene)-1-phenyl-7-(trifluoromethyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ra**): Yellow solid, 53.1 mg, 54% yield. m.p. 130.5~132.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.87 (s, 1H), 7.58 (d, $J=7.6$ Hz, 3H), 7.49~7.38 (m, 5H), 7.32 (s, 1H), 7.18 (t, $J=8.4$ Hz, 2H), 7.11 (d, $J=8.8$ Hz, 1H), 4.43 (q, $J=7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.8 (d, $^1J_{\text{CF}}=250$ Hz), 162.4, 147.7, 135.4, 133.2, 132.4, 131.8 (d, $^3J_{\text{CF}}=9$ Hz), 130.5, 130.5, 130.1, 130.0, 129.3, 129.0, 128.9, 128.2, 125.4 (q, $J_{\text{CF}_3}=4$ Hz), 125.0 (q, $J_{\text{CF}_3}=4$ Hz), 122.8, 122.5, 115.7 (d, $^2J_{\text{CF}}=22$ Hz), 61.1, 14.5; HRMS calcd for $\text{C}_{27}\text{H}_{19}\text{CF}_4\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 511.1098, found 511.1107.

Methyl (Z)-5-benzylidene-1-phenyl-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ab**): Yellow solid, 70.5 mg, 86% yield. m.p. 87.1~87.9 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.49~7.55 (m, 3H), 7.38 (t, $J=7.2$ Hz, 2H), 7.35~7.23 (m, 7H), 7.18 (s, 1H), 7.06 (t, $J=7.8$ Hz, 1H), 6.91 (d, $J=8.0$ Hz, 1H), 3.84 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.0, 147.3, 134.8, 133.8, 132.8, 132.2, 131.7, 130.9, 129.8, 129.7, 128.8, 128.8, 128.7, 128.6, 128.4, 128.2, 127.9, 127.5, 124.0, 122.0, 51.9; HRMS calcd for $\text{C}_{25}\text{H}_{19}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 411.1162, found 411.1171.

Ethyl (Z)-5-benzylidene-1-(4-fluorophenyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ac**): Yellow solid, 57.5 mg, 65% yield. m.p. 100.5~102.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.62~7.55 (m, 5H), 7.46~7.37 (m, 4H), 7.33 (d, $J=7.6$ Hz, 1H), 7.15 (t, $J=8.4$ Hz, 3H), 6.99 (d, $J=7.6$ Hz, 1H), 4.43 (q, $J=7.2$ Hz, 2H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 162.5 (d, $^1J_{\text{CF}}=248$ Hz), 147.4, 133.2, 132.8, 131.6 (d, $^3J_{\text{CF}}=8$ Hz), 131.1, 130.9, 130.8, 129.6, 129.5, 128.9, 128.7, 127.8, 127.7, 127.4, 124.1, 122.1, 115.72, 115.51 (d, $^2J_{\text{CF}}=21$ Hz), 60.9, 14.5; HRMS calcd for $\text{C}_{26}\text{H}_{20}\text{FN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$

443.1224, found 443.1229.

Ethyl (Z)-5-benzylidene-1-(4-chlorophenyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ad**): Yellow solid, 67.0 mg, 73% yield. m.p. 118.1~119.6 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.60 (t, *J*=8.8 Hz, 3H), 7.52 (d, *J*=8.4 Hz, 2H), 7.44~7.38 (m, 5H), 7.33 (t, *J*=7.6 Hz, 1H), 7.26 (s, 1H), 7.16 (t, *J*=7.6 Hz, 1H), 7.00 (d, *J*=8.0 Hz, 1H), 4.43 (q, *J*=7.2 Hz, 2H), 1.41 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.6, 147.4, 134.4, 133.3, 133.0, 132.8, 131.0, 130.9, 130.6, 129.7, 129.4, 128.9, 128.8, 128.8, 127.9, 127.7, 127.4, 125.1, 122.1, 61.0, 14.5; HRMS calcd for C₂₆H₂₀ClN₂O₂S [M+H]⁺ 459.0929, found 459.0931.

Ethyl (Z)-5-benzylidene-1-(2-chlorophenyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ae**): Yellow solid, 68.7 mg, 75% yield. m.p. 116.5~117.8 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.60 (t, *J*=8.8 Hz, 3H), 7.52 (d, *J*=8.4 Hz, 2H), 7.44~7.39 (m, 5H), 7.34 (t, *J*=7.6 Hz, 1H), 7.25 (s, 1H), 7.16 (t, *J*=8.0 Hz, 1H), 7.00 (d, *J*=8.4 Hz, 1H), 4.43 (q, *J*=7.2 Hz, 2H), 1.41 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.6, 147.4, 134.4, 133.3, 133.0, 132.8, 131.3, 131.0, 130.9, 130.6, 129.7, 128.9, 128.8, 128.8, 128.8, 127.9, 127.7, 127.4, 125.1, 122.1, 61.0, 14.5; HRMS calcd for C₂₆H₂₀ClN₂O₂S [M+H]⁺ 459.0929, found 459.0937.

Ethyl (Z)-5-benzylidene-1-(4-bromophenyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3af**): Yellow solid, 71.3 mg, 71% yield. m.p. 127.5~128.9 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.52 (dd, *J*=14.0, 4.8 Hz, 5H), 7.38~7.30 (m, 5H), 7.25 (t, *J*=7.6 Hz, 1H), 7.14 (s, 1H), 7.08 (t, *J*=8.4 Hz, 1H), 6.92 (d, *J*=8.0 Hz, 1H), 4.35 (q, *J*=7.2 Hz, 2H), 1.33 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.6, 147.4, 133.7, 132.9, 132.8, 131.7, 131.2, 130.9, 130.6, 129.7, 129.4, 128.9, 128.8, 127.9, 127.7, 127.5, 125.2, 122.7, 122.1, 60.9, 14.6; HRMS calcd for C₂₆H₂₀BrN₂O₂S [M+H]⁺ 503.0423, found 503.0434.

Ethyl (Z)-5-benzylidene-1-(2-bromophenyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ag**): Yellow solid, 72.3 mg, 72% yield. m.p. 127.0~129.0 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.60 (dd, *J*=14.0, 4.8 Hz, 4H), 7.46~7.36 (m, 5H), 7.33 (t, *J*=7.2 Hz, 2H), 7.22 (s, 1H), 7.16 (t, *J*=7.2 Hz, 1H), 7.00 (d, *J*=8.0 Hz, 1H), 4.43 (q, *J*=7.2 Hz, 2H), 1.41 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.6, 147.4, 133.7, 132.9, 133.0, 132.8, 131.7, 131.2, 130.9, 130.6, 129.7, 128.9, 128.8, 128.4, 128.2, 128.1, 127.7, 127.5, 125.2, 122.7, 122.1, 61.0, 14.6; HRMS calcd for C₂₆H₂₀BrN₂O₂S [M+H]⁺ 503.0423, found 503.0424.

Ethyl (Z)-5-benzylidene-1-(*p*-tolyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ah**): Yellow solid, 71.0 mg, 81% yield. m.p. 107.9~108.8 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.53 (t, *J*=7.6 Hz, 3H), 7.41 (d, *J*=8.0 Hz, 2H), 7.37~7.30 (m, 3H), 7.24 (t, *J*=7.6 Hz, 1H), 7.19 (d, *J*=6.4 Hz, 2H), 7.00~7.13 (m, 2H), 6.91 (d, *J*=8.4 Hz, 1H), 4.35 (q, *J*=7.2 Hz, 2H), 2.33 (s, 3H), 1.34 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.6, 147.2, 138.7, 133.9, 132.8, 132.0, 131.0, 129.8, 129.7, 129.6,

129.2, 129.0, 128.9, 128.7, 128.5, 127.8, 127.7, 127.4, 123.1, 122., 60.9, 21.4, 14.5; HRMS calcd for C₂₇H₂₃N₂O₂S⁺ [M+H]⁺ 439.1475, found 439.1481.

Ethyl(Z)-5-(4-fluorobenzylidene)-1-(3-(trifluoromethyl)-phenyl)-5H-benzo[d]imidazo[5,1-b][1,3]thiazine-3-carboxylate (**3ai**): Yellow solid, 54.1 mg, 53% yield. m.p. 138.2~140.2 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.87 (s, 1H), 7.59 (t, *J*=7.6 Hz, 3H), 7.49~7.38 (m, 5H), 7.32 (s, 1H), 7.18 (t, *J*=8.4 Hz, 2H), 7.11 (d, *J*=8.4 Hz, 1H), 4.43 (q, *J*=7.2 Hz, 2H), 1.41 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.8 (d, ¹*J*_{CF}=248.0 Hz), 162.4, 147.7, 135.4, 133.8, 133.3, 132.5, 131.8 (d, ³*J*_{CF}=8.0 Hz), 130.5, 130.5, 130.5, 129.0, 128.8, 128.2, 125.4 (q, *J*_{CF3}=4.0 Hz), 125.0 (q, *J*_{CF3}=4.0 Hz), 122.8, 122.5, 115.7 (d, ²*J*_{CF}=22.0 Hz), 61.1, 14.5; HRMS calcd for C₂₇H₁₉F₄N₃O₂S [M+H]⁺ 511.1098, found 511.1099.

Supporting Information ¹H NMR and ¹³C NMR spectra for compounds **3aa**~**3ra** and **3ab**~**3ai**. The Supporting Information is available free of charge via the Internet at <http://siocjournal.cn>.

References

- [1] Huang, X.; Cong, X.; Mi, P.; Bi, X. *Chem. Commun.* **2017**, 53, 3858.
- [2] (a) Palagiano, F.; Arenare, L.; De Caprariis, P.; Grandolini, G.; Ambrogio, V.; Perioli, L.; Filippelli, W.; Falcone, G.; Rossi, F. *Farmacologia* **1996**, 51, 483.
(b) Shridhar, D. R.; Jogibhukta, M.; Krishnan, V. S. H. *Ind. J. Chem.* **1982**, 21B, 130.
(c) Raddatz, S.; Bartel, S.; Guarnieri, W.; Rosentreter, U.; Ruppelt, M.; Wild, H.; Endermann, R.; Kroll, H.-P.; Henninger, K. *WO 1999/040094*, **1999**.
- [3] De Peretti, D.; George, P.; Sevrin, M.; Roy, J.; Demande, Fr. *FR 2759700*, **1998**.
- [4] (a) Fryer, R. I.; Lauer, R. F.; Trybulski, E. J.; Vitone, S.; Walser, A.; Zenchoff, G. J. *Chem.* **1983**, 20, 1605.
(b) Rao, K. V.; Sundaramurthy, V. *Sulfur Lett.* **1988**, 8, 137.
(c) Rao, K. V.; Sundaramurthy, V. *J. Org. Chem.* **1992**, 57, 2737.
(d) Petrova, K.; Petrov, O.; Antonova, A.; Kalcheva, V. *Heterocycl. Commun.* **2003**, 9, 593.
(e) Tu, X.; Li, T.; Zhou, L.; Zhang, Y.; Zeng, Q. *Fresenius Environ. Bull.* **2013**, 22, 3857.
(f) Omar, M. A.; Conrad, J.; Beifuss, U. *Tetrahedron* **2014**, 70, 5682.
(g) Tan, F.; Xiao, W. J.; Zeng, G. P. *Chin. J. Org. Chem.* **2017**, 37, 824 (in Chinese).
(谭芬, 肖文精, 曾国平, 有机化学, **2017**, 37, 824.)
(h) Sun, Y. W.; Zhou, L. Y.; Wang, L. Z. *Chin. J. Org. Chem.* **2019**, 39, 3516 (in Chinese).
(孙悦玮, 周来运, 王兰芝, 有机化学, **2019**, 39, 3516.)
(i) Yang, W. C.; Zhang, M. M.; Chen, W.; Yang, X. H.; Feng, J. G. *Chin. J. Org. Chem.* **2020**, 40, 4060 (in Chinese).
(杨文超, 张明明, 陈旺, 杨小虎, 冯建国, 有机化学, **2020**, 40, 4060.)
(j) Sun, X. H.; Sun, C. C.; Feng, L. J.; Kang, C. M. *Chin. J. Org. Chem.* **2020**, 40, 2626 (in Chinese).
(孙晓华, 孙传策, 冯立军, 康从民, 有机化学, **2020**, 40, 2626.)
- [5] Gauthier, J.; Duceppe, J. S. *J. Heterocycl. Chem.* **1984**, 21, 1081.
- [6] Dolbier, W. R.; Burkholder, J. C.; Abboud, K. A.; Loehle, D. J. *J. Org. Chem.* **1994**, 59, 7688.
- [7] Wang, R.; W.; Bao, W. *Tetrahedron Lett.* **2012**, 53, 442.
- [8] (a) Benati, L.; Calestani, G.; Leardini, R.; Minozzi, M.; Nanni, D.;

- Spagnolo, P.; Strazzari, S.; Zanardi, G. *J. Org. Chem.* **2003**, *68*, 3454.
- (b) Minozzi, M.; Nanni, D.; Zanardi, G.; Calestani, G. *ARKIVOC* **2006**, *6*, 6.
- (c) Saito, T.; Nihei, H.; Otani, T.; Suyama, T.; Furukawa, N.; Saito, M. *Chem. Commun.* **2008**, 172.
- (d) Xie, J. L.; Guo, Z. L.; Huang, Y. Q.; Qu, Y.; Song, H. J.; Song, H. B.; Liu, Y. X.; Wang, Q. M. *Adv. Synth. Catal.* **2019**, *361*, 490.
- (e) Wen, L.-R.; Shen, Q.-Y.; Guo, W.-S.; Li, M. *Org. Chem. Front.* **2016**, *3*, 870.
- [7] (a) Otani, T.; Kunimatsu, S.; Nihei, H.; Abe, Y.; Saito, T. *Org. Lett.* **2007**, *9*, 5513.
- (b) Otani, T.; Kunimatsu, S.; Takahashi, T.; Nihei, H.; Saito, T. *Tetrahedron Lett.* **2009**, *50*, 3853.
- [8] (a) Kaname, M.; Sashida, H. *Tetrahedron Lett.* **2012**, *53*, 748.
- (b) Sashida, H.; Kaname, M.; Minoura, M. *Tetrahedron* **2013**, *69*, 6478.
- [9] Liu, R.-J.; Wang, P.-F.; Yuan, W.-K.; Wen, L.-R.; Li, M. *Adv. Synth. Catal.* **2017**, *359*, 1373.
- [10] Xie, J. L.; Guo, Z. L.; Huang, Y. Q.; Qu, Y.; Song, H. J.; Song, H. B.; Liu, Y. X.; Wang, Q. M. *Adv. Synth. Catal.* **2019**, *361*, 490.
- [11] Miao, J. K.; Zhang, Y. H.; Sang, X. Y.; Hao, W. Y. *Org. Biomol. Chem.* **2019**, *17*, 2336.
- [12] (a) Hao, W. Y.; Zeng, J. B.; Cai, M. Z. *Chem. Commun.* **2014**, *50*, 11686.
- (b) Hao, W. Y.; Jiang, Y. Y.; Cai, M. Z. *J. Org. Chem.* **2014**, *79*, 3634.
- (c) Hao, W. Y.; Huang, J.; Jie, S. S.; Cai, M. Z. *Eur. J. Org. Chem.* **2015**, 6655.
- (d) Hao, W. Y.; Sang, X. Y.; Jiang, J.; Cai, M. Z. *Tetrahedron Lett.* **2016**, *57*, 1511.
- (e) Hao, W. Y.; Sang, X. Y.; Jiang, J.; Cai, M. Z. *Tetrahedron Lett.* **2016**, *57*, 4207.
- (f) Miao, J. K.; Sang, X. Y.; Wang, Y.; Deng, S. F.; Hao, W. Y. *Org. Biomol. Chem.* **2019**, *17*, 6994.
- (g) Zhang, Y. H.; Liu, Y.; Miao, J. K.; Hao, W. Y. *Chin. J. Org. Chem.* **2020**, *40*, 2426 (in Chinese).
- (张亚辉, 刘洋, 缪建康, 郝文燕, 有机化学, **2020**, *40*, 2426.)
- (h) Liu, Y.; Yao, S. J.; Wang, C. L.; Zhang, Y. H.; Hao, W. Y. *RSC Adv.* **2020**, *10*, 32211.
- [13] (a) Uchiyama, M.; Ozawa, H.; Takuma, K.; Matsumoto, Y.; Yonehara, M.; Hiroya, K.; Sakamoto, T. *Org. Lett.* **2006**, *8*, 5517.
- (b) Park, J. H.; Bhilare, S. V.; Youn, S. W. *Org. Lett.* **2011**, *13*, 2228.
- (c) Gilmore, K.; Alabugin, I. V. *Chem. Rev.* **2011**, *111*, 6513.

(Lu, Y.)