“一锅”三组分合成新型 1,5-苯并二氮杂萘类化合物与 抗牛病毒性腹泻病毒(BVDV)活性

韩超^{*,a} 聂磊^a 韩晓^b 张岩^a 孙克磊^a
石磊^a 崔广华^a 孟伟^{*,a}

(^a 华北理工大学化学工程学院 河北唐山 063210)

(^b 邯郸学院化学化工与材料学院 河北邯郸 056005)

摘要 以特氨酸、邻苯二胺和芳香醛为原料, 乙醇为溶剂, 对甲苯磺酸为催化剂, 多组分一锅法构建了一系列含有吡咯烷酮的 1,5-苯并二氮杂萘类化合物。该方法反应条件温和, 操作简单, 产物收率高。探索了抗牛病毒性腹泻病毒(BVDV)病毒活性, 结果显示合成的苯二氮杂萘类衍生物具有显著的抗 BVDV 活性, EC_{50} 值好, 且无明显的细胞毒性, 为抗 BVDV 制剂的研究提供了有意义的参考物。

关键词 “一锅”三组分反应; 1,5-苯并二氮杂萘类化合物; 对甲苯磺酸; 吡咯烷酮; 抗 BVDV

One-Pot Three-Component Synthesis of Novel 1,5-Benzodiazepine Derivatives and Their *anti*-BVDV (Bovine Viral Diarrhea Virus) Activity

Han, Chao^{*,a} Nie, Lei^a Han, Xiao^b Zhang, Yan^a Sun, Kelei^a Shi, Lei^a
Cui, Guanghua^a Meng, Wei^{*,a}

(College of Chemical Engineering, North China University of Science and Technology, Tangshan, Hebei 063210)

(College of Chemical Engineering & Material, Handan University, Handan, Hebei 056005)

Abstract The “one-pot” three-component reaction is established to synthesize 1,5-benzodiazepine derivatives containing pyrrolidinone. These reactions were carried out readily with *p*-toluenesulfonic acid (*p*-TSA) as the catalyst in good yields. Reaction conditions were mild, and environmental friendly. The benzodiazepine derivatives displayed prominent anti-BVDV (bovine viral diarrhea virus) activity with excellent EC_{50} values and no significant cytotoxicity. Our research could afford significant compounds for the research of anti-BVDV agents.

Keywords “one-pot” three-component reaction; 1,5-benzodiazepine derivatives; *p*-TSA; pyrrolidinone; anti-BVDV

1 Introduction

Viruses result in clinically serious diseases in humans and animals.^[1] Among various diseases, the global scope of hepatitis C virus (HCV) infection is a major concern for human health.^[2] The disease can lead to liver fibrosis, cirrhosis, hepatocellular carcinoma,^[3] or even death if treatment is not provided. At present, no vaccine is serviceable to prevent HCV infection. Bovine viral diarrhea virus (BVDV),^[4] which is similar to HCV, is the archetype of the Pestivirus genus of the Flaviviridae family. As a valua-

ble virus model, BVDV is used to identify and characterize antiviral agents for the treatment of HCV.^[5]

Among the large varieties of heterocycles, 1,5-benzodiazepine compounds display good biological activity and occupy a very important position,^[6-7] which gain extensive interest in recent years. This class of compounds is extensively consumed psychoactive drugs worldwide as platelet aggregation inhibitors, anticonvulsant agents, antitumor compounds and anxiolytic compounds.^[8-10] The benzodiazepine skeleton is a core structure in a lot of drug molecules, for instance, clozapine tablets,^[11] olanzapine,^[12]

* Corresponding authors. E-mail: hanchao@ncst.edu.cn; mengwei@ncst.edu.cn

Received April 12, 2020; revised June 17, 2020; published online September 29, 2020.

Project supported by the Natural Science Foundation of Hebei Province (No. B2017209119) and the Education Department of Hebei Province (No. Z2017013).

河北省自然科学基金(No. B2017209119)、河北省教育厅基金(No. Z2017013)资助项目。

clabazam^[13] and triflubazam (Figure 1).^[14]

Figure 1 Some drug molecules with a benzodiazepine structure

Furthermore, Raboisson's group^[15] discovered a novel class of 1,5-benzodiazepines (1,5-BZDs) derivative (NS-5B) (Figure 2). The identification of potent and selective NS5B inhibitors got a lot of attention. Intense drug discovery efforts were initiated and have been subsequently validated in the clinic. They are described as excellent samples of HCV inhibitors.^[16]

Figure 2 Previously reported 1,5-BZD HCV NS5B inhibitors

The development of effective chemical reactions is an important area for the synthesis of heterocyclic scaffolds.^[17-18] Over the years, multicomponent reactions (MCRs) play an important role as powerful tools for the creation of molecular complexity and diversity.^[19] MCRs are efficient, diversity, environmentally friendly and atom economic.^[20-21] Furthermore, organic and medical chemists can construct new compounds by MCRs, which contain two or more heterocyclic scaffolds fused to enhanced biological activity.^[22-23]

On the other hand, tetramic acids (2,4-pyrrolidinedione) display important biological activities^[24] and possess insecticidal, antibacterial properties, antineoplastic and antiviral.^[25-27] Therefore, tetramic acids are chosen as a significant substrate, and an efficient and novel green three-component reaction to synthesize new compounds which contain both benzodiazepine and pyrrole motifs is reported.^[28] The reactions were carried out by reacting tetramic acids, 1,2-phenylenediamine, and aromatic aldehyde in EtOH with *p*-TSA as the catalyst. The *anti*-BVDV activities of these synthesized compounds have been evaluated and discussed in detail.

2 Results and discussion

The three-component reaction of benzaldehyde, (*S*)-5-benzyltetramic acid and 1,2-phenylenediamine was taken as a model reaction. To justify the significance of catalyst in this process, the reaction was first carried out without the catalyst. Regrettably, the desired product was obtained in only 10% yield. (Table 1, Entry 1). Several Lewis and Bronsted acid were employed and the results were shown in Table 1. In the presence of HCO₂H, AlCl₃, FeCl₃ and BF₃·Et₂O, the reaction gave isolated yield of product after 3 h (Table 1, Entries 8 ~ 11). In addition, AcOH, CF₃CO₂H, CF₃SO₃H, HCl, H₂SO₄ and CH₃SO₃H could also catalyze this transformation, but no improvement was clearly observed (Table 1, Entries 3 ~ 7). To our delight, the benzodiazepine compound **4a** was furnished in 81% yield when the reaction was accomplished with *p*-TSA as catalyst (Table 1, Entry 2).

Table 1 Optimization of catalyst for the synthesis of **4a**^a

Entry	Catalyst ^b	T/°C	Time/h	Yield ^c /%
1	No	40	6	10
2	<i>p</i> -TSA	40	3	81
3	AcOH	40	3	73
4	CF ₃ CO ₂ H	40	3	69
5	HCl	40	3	60
6	H ₂ SO ₄	40	3	53
7	CH ₃ SO ₃ H	40	3	65
8	HCO ₂ H	40	3	55
9	AlCl ₃	40	3	41
10	FeCl ₃	40	3	32
11	BF ₃ ·Et ₂ O	40	3	30

^a The reaction was carried out with **1a** (1 mmol), **2** (1 mmol) and **3a** (1 mmol), in 2 mL of ethanol. ^b 20 mol% catalyst. ^c Isolated yields.

Next, by using *p*-TSA as the catalyst, the effect of different solvents and catalysts loading was screened. The results were shown in Table 2. The raw materials dissolved well in organic solvents, so we got a higher yield than that in water (Table 2, Entries 3 ~ 11). Further studies showed that EtOH was the best choice of the solvent for this transformation. On the other hand, the benzodiazepine products were insoluble in EtOH and the raw materials were soluble, which could simplify the reaction operation.

The reactions were also carried out using different amounts of the catalyst. The results showed that 15 mol% catalyst was the optimum choice to complete the reaction in ethanol. By decreasing the amount of catalyst to 10 mol% relative to substrate, the yield decreased (Table 2, Entry 14).

Table 2 Optimization of solvent and amount for the synthesis of **4a**^a

Entry	Solvent	<i>p</i> -TSA/mol%	Time/h	Yield ^b /%
1	No	20	5	5
2	H ₂ O	20	3	20
3	EtOH	20	3	81
4	Toluene	20	3	50
5	MeOH	20	3	71
6	<i>i</i> -PrOH	20	3	64
7	CH ₃ CN	20	3	52
8	THF	20	3	55
9	CHCl ₃	20	3	61
10	DMF	20	3	46
11	Dioxane	20	3	58
12	EtOH	30	3	82
13	EtOH	15	3	81
14	EtOH	10	3	72

^a The reaction was carried out with **1a** (1 mmol), **2** (1 mmol) and **3a** (1 mmol), at 40 °C in 2 mL of ethanol. ^b Isolated yields.

The effects of temperature and reaction time were also investigated. Variation of the reaction temperature proved that 60 °C was the optimal result for this reaction (Table 3, Entries 1~4). Prolonging the reaction time did not improve the yield (Table 3, Entries 5~6).

Table 3 Optimization of time and temperature for the synthesis of **4a**^a

Entry	<i>T</i> /°C	Time/h	Yield ^b /%
1	25	3	72
2	40	3	81
3	60	3	88
4	Reflux	3	88
5	60	5	88
6	60	7	88

^a The reaction was carried out with **1a** (1 mmol), **2** (1 mmol) and **3a** (1 mmol), *p*-TSA (15 mol%) in 2 mL of ethanol. ^b Isolated yields.

Overall, the reactions were conducted at 60 °C for 3 h in EtOH in the presence of 15 mol% *p*-TSA as catalyst. With the optimal multicomponent reaction conditions in hand, a series of benzodiazepine compounds were synthesized in high yields with good diastereoselectivities by using tetramic acids, 1,2-phenylenediamine, and aromatic aldehyde as starting materials (Table 4).

All the benzodiazepine derivatives **4** were characterized by ¹H NMR, and ¹³C NMR spectroscopies. To further illustrate the structure of these products, one of fused heterocyclic structure (**4a**) was also characterized by the single crystal x-ray diffraction. It confirmed that the compound was the *S,S*-diastereoisomer as shown in Figure 3.

Taking the synthesis of the benzodiazepine derivatives **4a** as an example, a proposed mechanism is summarized in Scheme 1. First, benzene-1,2-diamine condenses with **3a** to give enaminone **6**, which reacts with benzaldehyde to form intermediate **7**. Undergoing an intramolecular Michael addition reaction and isomerization, the intermediate **7** is converted into the final product **4a** (Scheme 1).

Table 4 synthesis of product **4**

Entry	Compd.	R ¹	R ²	R ³	Yield ^a /%	Dr ^b
1	4a	Bn	H	H	88	18 : 1
2	4b	Bn	H	3-OH	82	16 : 1
3	4c	Bn	H	2-OMe	86	15 : 1
4	4d	Bn	H	4-OH	87	16 : 1
5	4e	Bn	H	3,4-(OMe) ₂	85	>20 : 1
6	4f	Bn	H	4-Br	86	9 : 1
7	4g	Bn	H	4-NMe ₂	83	>20 : 1
8	4h	Bn	H	4-F	86	8 : 1
9	4i	Bn	H	2-NO ₂	86	>20 : 1
10	4j	H	Bn	H	87	— ^c
11	4k	H	Bn	2-OMe	87	— ^c
12	4l	H	Bn	4-Cl	81	— ^c
13	4m	H	Bn	3-OH	85	— ^c
14	4n	H	Bn	2-NO ₂	81	— ^c
15	4o	H	Bn	4-NMe ₂	82	— ^c
16	4p	H	Bn	4-Br	80	— ^c
17	4q	H	Bn	4-NO ₂	80	— ^c

^a Isolated yield of the major diastereoisomer. ^b Determined by ¹H NMR analysis of the reaction system. ^c Not determined.

Figure 3 X-ray crystal structure of **4a**

Prodrugs **4a**~**4q** were tested antiviral activity against BVDV in MDBK cells and cytotoxicity by methyl thiazolyl tetrazolium (MTT) method. Ribavirin was chosen as the positive control. Excitingly, compounds **4c**, **4e**, **4g** and **4o** showed excellent antiviral activities, and the results were shown in Table 5.

Preliminary structure-activity relationship (SAR) indicated that substituents of the compounds displayed different effect about antiviral activities. Compounds **4i**, **4n** and

Scheme 1 Plausible mechanism

Table 5 Anti-BVDV activity of benzodiazepine derivatives 4a~4o

Compd.	R ¹	R ²	R ³	EC ₅₀ ^a / (μmol·L ⁻¹)	CC ₅₀ ^b / (μmol·L ⁻¹)
4a	Bn	H	H	1.25	>20
4b	Bn	H	3-OH	>1.30	— ^c
4c	Bn	H	2-OMe	0.20	>20
4d	Bn	H	4-OH	>1.30	— ^c
4e	Bn	H	3,4-(OMe) ₂	0.25	No ^d
4f	Bn	H	4-Br	>1.30	No
4g	Bn	H	4-NMe ₂	0.12	No
4h	Bn	H	4-F	>1.30	No
4i	Bn	H	2-NO ₂	>1.30	>20
4j	H	Bn	H	1.50	>20
4k	H	Bn	2-OMe	0.28	No
4l	H	Bn	4-Cl	>1.3	>20
4m	H	Bn	3-OH	>1.3	— ^c
4n	H	Bn	2-NO ₂	>1.30	>20
4o	H	Bn	4-NMe ₂	0.25	No
4p	H	Bn	4-Br	>1.3	>20
4q	H	Bn	4-NO ₂	>1.30	>20
Ribavirin				1.30	5.08

^a The 50% cytotoxic concentration. ^b The 50% effective concentration. ^c The cytotoxicity was not tested. ^d No cytotoxicity was observed in 25 μmol·L⁻¹.

4q produced a marked decrease in the bioactivity when the benzene ring contained electrondrawing group such as nitro. The antiviral activities of halo-substituted compounds 4f, 4h, 4l and 4p were markedly reduced. While hydroxyl were introduced to different position of compounds 4b, 4d and 4m, their antiviral activities were decreased. It was delightful when R³ was the electron donating group OMe and NMe₂, compounds 4c, 4e, 4g, 4k and 4o showed better antiviral activities. In the benzodiazepine compounds, 4g (R³=NMe₂, EC₅₀=0.12 μmol·L⁻¹) had the best activity, which was 10 fold more active than that of positive control of ribavirin (EC₅₀=1.3 μmol·L⁻¹).

3 Conclusions

In short, we have found an efficient and green reaction for the preparation of benzodiazepine derivatives with *p*-TSA as the catalyst in ethanol. It was the first time that the multicomponent reaction was used to prepare 1,5-benzodiazepine derivatives containing pyrrolidinone. The synthesis method displays attractive advantage, for example one-pot conditions, easy purification, as well as reduced waste production. At the same time, compounds exhibited good antiviral activities against BVDV with no significant cytotoxicity. Compound 4g (R³ = 4-NMe₂) showed the best antiviral activity (EC₅₀=0.12 μmol/L).

4 Experimental section

4.1 Materials

Reagents and solvents were purchased from commercial sources and were used without further purification. Reactions were monitored by thin-layer chromatography (TLC) on 0.25 mm silicagel plates (GF254). Melting points were determined on a Beijing Keyi XT5 apparatus and are uncorrected. NMR spectra were obtained on a Bruker AVANCE III systems using DMSO-*d*₆ as solvent, TMS as internal standard substance, at 400 MHz for ¹H NMR, 100 MHz for ¹³C NMR. IR spectra were recorded as KBr pellets on a Thermo Nicolet (IR 200) spectrometer. High-resolution mass spectra were recorded on a Waters Micromass Q-ToF Micro instrument operated in the ESI mode.

4.2 Chemistry

(*S*)-5-Benzyltetramic acid (1a, 1 mmol), 1,2-phenylenediamine (2, 1 mmol), *p*-TSA (15% mmol) were mixed and stirred for 30 min in EtOH (2 mL). Then, benzaldehyde 3a (1 mmol) was added to the reaction mixture and stirred for 3 h at 60 °C. After completion, the reaction mixture was cooled to room temperature, filtered and washed by ethanol to give 4a. Compounds 4b~4q were synthesized as compound 4a.

Crystallographic data of compound 4a have been deposited with the Cambridge Crystallographic Data Centre with CCDC number of 1992698, which are free of charge via e-mail: deposit@ccdc.cam.ac.uk or <http://www.ccdc.cam.ac.uk>.

(3*S*)-3-Benzyl-10-[(*S*)-phenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (4a): White powder, m.p. 224~225 °C; [α]_D²⁵ -0.2 (*c* 0.333, MeOH); ¹H NMR (400 MHz, DMSO-*d*₆) δ : 8.95 (s, 1H, NH), 7.37 (s, 1H, CONH), 7.26~7.16 (m, 5H, ArH), 7.08~7.05 (m, 1H, ArH), 6.95~6.87 (m, 3H, ArH), 6.71~6.67 (m, 1H, ArH), 6.56~6.52 (m, 1H, ArH), 6.43~6.40 (m, 3H, ArH), 5.73 (d, *J*=4.4 Hz, 1H, NH), 4.90 (s, 1H, CH), 4.45 (s, 1H, CONHCH), 3.39~3.34 (m, 1H, CH₂), 3.12~3.08 (m, 1H, CH₂); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 172.6, 154.0, 143.8, 137.1, 135.7, 133.5, 130.0, 127.7, 127.4, 127.0, 126.3, 125.7, 122.8, 121.6, 120.7, 119.2, 103.9, 57.2, 55.4, 36.3; IR (KBr) ν : 3272, 3200, 1650, 1562, 1493, 1390,

1329, 698 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{22}\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 368.1757, found 368.1755.

(3*S*)-3-Benzyl-10-[(*S*)-3-hydroxyphenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4b**): White powder, m.p. 258~259 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ -6.0 (*c* 0.100, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.05 (s, 1H, OH), 8.97 (s, 1H, NH), 7.25~7.18 (m, 3H, ArH), 7.19~7.18 (m, 3H, ArH), 7.06~7.04 (m, 1H, ArH), 6.71~6.67 (m, 1H, ArH), 6.61~6.56 (m, 3H, ArH), 6.47~6.44 (m, 1H, ArH), 6.39~6.37 (m, 1H, ArH), 5.69 (s, 1H, ArH), 5.51~5.49 (m, 1H, NH), 4.80 (s, 1H, CH), 4.40~4.38 (m, 1H, CONHCH), 3.19~3.18 (m, 2H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.7, 156.6, 154.1, 145.6, 137.3, 136.3, 133.1, 129.7, 128.3, 127.8, 126.2, 122.7, 121.6, 120.4, 119.2, 117.5, 114.72, 113.0, 104.1, 57.2, 55.6, 37.3; IR (KBr) ν : 3413, 3361, 3182, 1651, 1570, 1347, 1252, 1038, 760 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{22}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 384.1707, found 384.1707.

(3*S*)-3-Benzyl-10-[(*S*)-2-methoxyphenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4c**): White powder, m.p. 220~221 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ +2.9 (*c* 0.333, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.01 (s, 1H, NH), 7.43 (s, 1H, CONH), 7.27~7.19 (m, 5H, ArH), 7.07 (d, $J=8.0$ Hz, 1H, ArH), 6.93 (t, $J=7.6$ Hz, 1H, ArH), 6.81 (d, $J=8.0$ Hz, 1H, ArH), 6.69 (t, $J=7.6$ Hz, 1H, ArH), 6.51 (t, $J=7.6$ Hz, 1H, ArH), 6.28 (d, $J=8.0$ Hz, 1H, ArH), 6.18 (t, $J=7.6$ Hz, 1H, ArH), 5.13 (s, 1H, ArH), 5.12~5.11 (m, 1H, NH), 5.03~5.02 (m, 1H, CH), 4.48 (s, 1H, CONHCH), 3.85 (s, 3H, CH_3), 3.45~3.40 (m, 1H, CH_2), 3.10~3.06 (m, 1H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.5, 156.8, 154.9, 137.0, 135.7, 133.3, 130.7, 130.1, 127.8, 127.4, 126.7, 126.3, 122.6, 121.7, 121.0, 119.6, 119.1, 109.6, 103.2, 55.5, 55.3, 53.2, 36.2; IR (KBr) ν : 3330, 3273, 1636, 1562, 1505, 1344, 1201, 701 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{24}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 398.1863, found 398.1868.

(3*S*)-3-Benzyl-10-[(*S*)-4-hydroxyphenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4d**): White powder, m.p. 265~266 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ -3.7 (*c* 0.333, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.99 (s, 1H, OH), 8.93 (s, 1H, NH), 7.32 (s, 1H, NH), 7.23~7.19 (m, 5H, ArH), 7.04~7.02 (m, 1H, ArH), 6.67 (t, $J=7.2$ Hz, 1H, ArH), 6.53 (t, $J=7.2$ Hz, 1H, ArH), 6.41~6.39 (m, 1H, ArH), 6.27~6.25 (m, 2H, ArH), 6.20~6.16 (m, 2H, ArH), 5.62 (d, $J=4.0$ Hz, 1H, NH), 4.83 (s, 1H, CH), 4.45 (s, 1H, CONHCH), 3.37~3.33 (m, 1H, CH_2), 3.07~3.04 (m, 1H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.8, 155.2, 153.9, 137.5, 135.9, 134.4, 133.5, 130.0, 128.1, 127.7, 126.2, 122.8, 121.5, 120.5, 119.1, 114.2, 104.5, 56.7, 55.4, 36.4; IR (KBr) ν : 3422, 3305, 3280, 1649, 1542, 1383, 1084, 707 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{22}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 384.1707, found 384.1702.

(3*S*)-3-Benzyl-10-[(*S*)-3,4-dimethoxyphenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4e**): Yellow powder, m.p. 269~270 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ +2.0 (*c* 0.333, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.91 (s, 1H, NH), 7.31 (s, 1H, NH), 7.23~7.19 (m, 5H, ArH), 7.03

(d, $J=7.6$ Hz, 1H, ArH), 6.93~6.91 (m, 1H, ArH), 6.68 (t, $J=7.6$ Hz, 1H, ArH), 6.56 (t, $J=7.2$ Hz, 1H, ArH), 6.47 (d, $J=7.2$ Hz, 1H, ArH), 6.22 (d, $J=8.0$ Hz, 1H, ArH), 5.75 (d, $J=4.0$ Hz, 1H, ArH), 5.22 (d, $J=8.4$ Hz, 1H, NH), 4.82 (d, $J=4.0$ Hz, 1H, CH), 4.42~4.39 (m, 1H, CONHCH), 3.63 (s, 3H), 3.60 (s, 3H, CH_3), 3.29~3.25 (m, 1H, CH_2), 3.14~3.11 (m, 1H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.8, 154.1, 148.0, 146.8, 135.9, 135.2, 133.4, 129.9, 129.8, 127.8, 126.3, 122.8, 121.7, 120.7, 119.2, 118.4, 111.7, 110.3, 104.2, 56.8, 55.4, 55.3, 55.1, 36.7; IR (KBr) ν : 3349, 3256, 1660, 1558, 1477, 1384, 1010, 699 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{26}\text{N}_3\text{O}_3$ $[\text{M} + \text{H}]^+$ 428.1969, found 428.1972.

(3*S*)-3-Benzyl-10-[(*S*)-4-bromophenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4f**): White powder, m.p. 252~253 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ +2.9 (*c* 0.333, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.99 (s, 1H, NH), 7.40 (s, 1H, NH), 7.24~7.20 (m, 5H, ArH), 7.05~7.03 (m, 3H, ArH), 6.72~6.41 (m, 3H, ArH), 6.30~6.29 (m, 2H, ArH), 5.74 (s, 1H, NH), 4.84 (s, 1H, CH), 4.46 (s, 1H, CONHCH), 3.40~3.36 (m, 1H, CH_2), 3.08~3.05 (m, 1H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.4, 154.4, 142.50, 135.48, 133.71, 132.09, 130.22, 130.01, 129.26, 127.78, 126.37, 123.03, 121.98, 121.72, 119.44, 119.21, 103.10, 56.82, 55.42, 36.09; IR (KBr) ν : 3282, 3205, 1651, 1548, 1495, 1371, 1224, 695 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{BrN}_3\text{O}$ $[\text{M} + \text{H}]^+$ 446.0863, found 446.0860.

(3*S*)-3-Benzyl-10-[(*S*)-4-(dimethylamino)phenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4g**): White powder, m.p. 230~231 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ -0.3 (*c* 0.333, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.87 (s, 1H, NH), 7.31 (s, 1H, NH), 7.23~7.20 (m, 5H, ArH), 7.04~7.02 (m, 1H, ArH), 6.67~6.44 (m, 3H, ArH), 6.27~6.25 (m, 4H, ArH), 5.62 (s, 1H, NH), 4.80 (s, 1H, CH), 4.40 (s, 1H, CONHCH), 3.33~3.29 (m, 1H, CH_2), 3.12~3.09 (m, 1H, CH_2), 2.74 (s, 6H, CH_3); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.7, 153.7, 148.5, 137.6, 136.0, 133.4, 132.0, 129.9, 127.8, 127.7, 126.2, 122.8, 121.5, 120.3, 119.1, 111.7, 104.8, 56.5, 55.4, 40.2, 36.7; IR (KBr) ν : 3230, 3070, 1630, 1570, 1465, 1387, 1265, 1123, 692 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{27}\text{N}_4\text{O}$ $[\text{M} + \text{H}]^+$ 411.2179, found 411.2175.

(3*S*)-3-Benzyl-10-[(*S*)-4-fluorophenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4h**): Faint Yellow powder, m.p. 218~219 $^{\circ}\text{C}$; $[\alpha]_{\text{D}}^{25}$ +0.2 (*c* 0.333, MeOH); ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.03 (s, 1H, NH), 7.45 (s, 1H, NH), 7.24~7.18 (m, 5H, ArH), 7.10~7.08 (m, 1H, ArH), 6.74~6.65 (m, 3H, ArH), 6.58~6.55 (m, 1H, ArH), 6.42~6.35 (m, 3H, ArH), 5.88 (s, 1H, NH), 4.92 (s, 1H, CH), 4.47 (s, 1H, CONHCH), 3.41~3.38 (m, 1H, CH_2), 3.09~3.06 (m, 1H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.5, 159.2, 154.1, 139.8, 135.7, 133.7, 130.0, 128.9, 128.8, 127.8, 126.4, 122.9, 121.8, 119.3, 114.1, 113.9, 103.8, 56.6, 55.4, 36.3; IR (KBr) ν : 3321, 3221, 16536, 1532, 1358, 1200, 747, 657 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{FN}_3\text{O}$ $[\text{M} + \text{H}]^+$ 386.1663,

found 386.1669.

(3*S*)-3-Benzyl-10-[(*S*)-2-nitrophenyl]-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4i**): Orange powder, m.p. 276~277 °C; $[\alpha]_D^{25} = -0.5$ (c 0.333, MeOH); ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.25 (s, 1H, NH), 7.84~7.82 (m, 1H, ArH), 7.53 (s, 1H, NH), 7.23~7.19 (m, 7H, ArH), 6.93~6.80 (m, 2H, ArH), 6.58~6.56 (m, 1H, ArH), 6.37~6.35 (m, 1H, ArH), 5.53~5.51 (m, 1H, ArH), 5.38 (s, 1H, NH), 5.08 (s, 1H, CH), 4.54 (s, 1H, CONHCH), 3.47~3.43 (m, 1H, CH₂), 3.08~3.05 (m, 1H, CH₂); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 172.0, 154.7, 148.6, 137.8, 135.6, 135.5, 133.8, 132.6, 130.0, 127.8, 127.7, 127.5, 126.5, 124.3, 123.4, 122.2, 121.8, 119.4, 102.9, 55.5, 53.5, 36.1; IR (KBr) ν : 3313, 3253, 1670, 1565, 1410, 1246, 1132, 780 cm⁻¹. HRMS (ESI) calcd for C₂₄H₂₁N₄O₃ [M+H]⁺ 413.1608, found 413.1612.

2-Benzyl-10-phenyl-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4j**): White powder, m.p. 216~217 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.24 (s, 1H, NH), 7.37~7.34 (m, 2H, ArH), 7.29~7.25 (m, 1H, ArH), 7.22~7.12 (m, 6H, ArH), 7.10~7.06 (m, 1H, ArH), 6.80~6.78 (m, 1H, ArH), 6.67~6.63 (m, 1H, ArH), 6.58~6.55 (m, 2H, ArH), 5.97 (s, 1H, NH), 5.12 (s, 1H, CH), 4.55~4.41 (m, 2H, NCH₂), 3.99~3.87 (m, 2H, CH₂); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 171.2, 151.1, 138.3, 133.0, 129.5, 129.2, 128.6, 127.9, 127.9, 127.4, 127.3, 127.1, 126.6, 123.1, 121.8, 119.0, 102.6, 57.8, 48.7, 44.8; IR (KBr) ν : 3280, 3205, 1665, 1610, 1560, 1489, 1335, 1026, 735, 689 cm⁻¹. HRMS (ESI) calcd for C₂₄H₂₂N₃O [M+H]⁺ 368.1757, found 368.1759.

2-Benzyl-10-(2-methoxyphenyl)-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4k**): White powder, m.p. 224~226 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.26 (s, 1H, NH), 7.38~7.34 (m, 2H, ArH), 7.29~7.26 (m, 1H, ArH), 7.23~7.21 (m, 2H, ArH), 7.12~7.08 (m, 1H, ArH), 6.98~6.96 (m, 1H, ArH), 6.79~6.77 (m, 1H, ArH), 6.66~6.58 (m, 3H, ArH), 6.55~6.51 (m, 1H, ArH), 6.41~6.40 (m, 1H, ArH), 5.41 (d, *J*=4.0 Hz, 1H, NH), 5.25 (d, *J*=4.0 Hz, 1H, CH), 4.54~4.44 (m, 2H, NCH₂), 3.97 (q, *J*=16.8 Hz, 2H, CH₂), 3.93 (s, 3H, CH₃); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 171.2, 157.1, 152.3, 138.4, 136.7, 132.8, 131.2, 128.6, 128.0, 127.4, 127.1, 126.5, 122.7, 121.7, 120.9, 119.5, 118.6, 110.4, 102.1, 55.5, 53.1, 48.7, 44.8; IR (KBr) ν : 3295, 3186, 1647, 1600, 1560, 1520, 1367, 1136, 743, 692 cm⁻¹. HRMS (ESI) calcd for C₂₅H₂₄N₃O₂ [M+H]⁺ 398.1863, found 398.1870.

2-Benzyl-10-(4-chlorophenyl)-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4l**): White powder, m.p. 226~227 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.27 (s, 1H, NH), 7.37~7.34 (m, 2H, ArH), 7.29~7.20 (m, 5H, ArH), 7.15~7.13 (m, 2H, ArH), 6.81~6.79 (m, 1H, ArH), 6.70~6.66 (m, 1H, ArH), 6.63~6.59 (m, 1H, ArH), 6.55~6.53 (m, 1H, ArH), 5.97 (s, 1H, NH), 5.11 (s, 1H, NH), 4.55~4.41 (m, 2H, NCH₂), 3.99~3.88 (m, 2H, CH₂); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 171.1, 151.2, 143.6, 138.4, 136.6, 132.7, 130.8, 128.9, 128.6,

127.8, 127.4, 127.1, 122.6, 121.9, 120.7, 118.9, 102.6, 56.9, 48.7, 44.8; IR (KBr) ν : 3323, 3134, 1672, 1571, 1390, 1133, 820, 697 cm⁻¹. HRMS (ESI) calcd for C₂₄H₂₁ClN₃O [M+H]⁺ 402.1368, found 402.1370.

2-Benzyl-10-(3-hydroxyphenyl)-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4m**): White powder, m.p. 224~226 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.22 (s, 1H, OH), 9.16 (s, 1H, NH), 7.37~7.33 (m, 2H, ArH), 7.29~7.21 (m, 3H, ArH), 6.97~6.94 (m, 1H, ArH), 6.80~6.78 (m, 1H, ArH), 6.66~6.48 (m, 6H, ArH), 5.89~5.88 (m, 1H, NH), 5.03~5.02 (m, 1H, CH), 4.55~4.42 (m, 2H, NCH₂), 3.97~3.87 (m, 2H, CH₂); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 171.3, 156.8, 150.8, 146.1, 138.4, 137.0, 132.4, 128.7, 128.6, 127.5, 127.1, 122.6, 121.6, 120.3, 118.7, 118.1, 114.3, 113.1, 103.2, 57.4, 48.6, 44.8; IR (KBr) ν : 3357, 3248, 16676, 1499, 1365, 1190, 1053, 723 cm⁻¹. HRMS (ESI) calcd for C₂₄H₂₂N₃O₂ [M+H]⁺ 384.1707, found 384.1705.

2-Benzyl-10-(2-nitrophenyl)-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4n**): Yellow powder, m.p. 246~247 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.51 (s, 1H, NH), 7.97~7.96 (m, 1H, ArH), 7.39~7.18 (m, 7H, ArH), 6.93~6.91 (m, 2H, ArH), 6.79~6.75 (m, 1H, ArH), 6.64~6.60 (m, 1H, ArH), 6.55~6.53 (m, 1H, ArH), 5.68~5.67 (m, 1H, NH), 5.42~5.41 (m, 1H, CH), 4.52~4.36 (m, 2H, NCH₂), 4.04~3.94 (m, 2H, CH₂); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 170.7, 151.9, 149.1, 138.3, 138.2, 135.3, 133.0, 132.9, 128.6, 128.0, 128.0, 127.5, 127.1, 124.6, 123.4, 122.2, 121.5, 119.1, 101.8, 53.3, 48.8, 44.8; IR (KBr) ν : 3350, 3286, 1646, 1566, 1405, 1242, 1012, 697 cm⁻¹. HRMS (ESI) calcd for C₂₄H₂₁N₄O₃ [M+H]⁺ 413.1608, found 413.1610.

2-Benzyl-10-(4-(dimethylamino)phenyl)-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4o**): White powder, m.p. 234~235 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.15 (s, 1H, NH), 7.35~7.20 (m, 5H, ArH), 6.97~6.95 (m, 2H, ArH), 6.78~6.76 (m, 1H, ArH), 6.64~6.51 (m, 5H, ArH), 5.82 (s, 1H, NH), 5.03 (s, 1H, CH), 4.53~4.43 (m, 2H, NCH₂), 3.95~3.85 (m, 2H, CH₂), 2.78 (s, 6H, CH₃); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 171.4, 150.6, 148.9, 138.5, 137.3, 132.6, 132.5, 128.6, 127.8, 127.4, 127.0, 122.7, 121.5, 120.2, 118.6, 111.9, 103.9, 56.8, 48.6, 44.8, 40.2; IR (KBr) ν : 3352, 3247, 1670, 1595, 1365, 1062, 775, 680 cm⁻¹. HRMS (ESI) calcd for C₂₆H₂₇N₄O [M+H]⁺ 411.2179, found 411.2180.

2-Benzyl-10-(4-bromophenyl)-3,4,9,10-tetrahydrobenzo[*b*]pyrrolo[3,4-*e*][1,4]diazepin-1(2*H*)-one (**4p**): Yellow powder, m.p. 253~254 °C; ^1H NMR (400 MHz, DMSO-*d*₆) δ : 9.27 (s, 1H, NH), 7.38~7.34 (m, 4H, ArH), 7.27~7.20 (m, 3H, ArH), 7.09~7.07 (m, 2H, ArH), 6.81~6.79 (m, 1H, ArH), 6.70~6.66 (m, 1H, ArH), 6.61~6.59 (m, 1H, ArH), 6.55~6.53 (m, 1H, ArH), 5.97~5.96 (m, 1H, NH), 5.10~5.09 (m, 1H, CH), 4.54~4.41 (m, 2H, NCH₂), 3.99~3.88 (m, 2H, CH₂); ^{13}C NMR (100 MHz, DMSO-*d*₆) δ : 171.1, 151.2, 144.0, 138.4, 136.6, 132.7, 130.7, 129.3, 128.6, 127.4, 127.1, 122.6, 121.9, 120.7, 119.4, 118.9, 102.5, 56.9, 48.7, 44.8; IR (KBr) ν : 3283,

3132, 1646, 1564, 1408, 1269, 1098, 700 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{BrN}_3\text{O}$ $[\text{M}+\text{H}]^+$ 446.0863, found 446.0866.

2-Benzyl-10-(4-nitrophenyl)-3,4,9,10-tetrahydrobenzo-[b]pyrrolo[3,4-e][1,4]diazepin-1(2H)-one (**4q**): White powder, m.p. 241~241 $^{\circ}\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.40 (s, 1H, NH), 8.08~8.06 (m, 2H, ArH), 7.40~7.34 (m, 4H, ArH), 7.29~7.21 (m, 3H, ArH), 6.86~6.84 (m, 1H, ArH), 6.73~6.69 (m, 1H, ArH), 6.63~6.55 (m, 2H, ArH), 6.15~6.14 (m, 1H, NH), 5.24~5.23 (m, 1H, CH), 4.56~4.42 (m, 2H, NCH_2), 4.03~3.92 (m, 2H, CH_2); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 171.0, 152.4, 151.4, 146.0, 138.3, 136.2, 132.6, 128.6, 128.2, 127.5, 127.1, 123.2, 122.6, 122.1, 120.9, 119.1, 101.8, 57.1, 48.8, 44.8; IR (KBr) ν : 3413, 3252, 1678, 1553, 1400, 1326, 1058, 813, 726 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{N}_4\text{O}_3$ $[\text{M}+\text{H}]^+$ 413.1608, found 413.1610.

4.3 Cytotoxicity assay

In RPMI 1640-FBS medium, MDBK cells, with a density of 7×10^3 per well in 96-well cell culture plates, were seeded. After incubation for 24 h under the condition of 37 $^{\circ}\text{C}$, 5% CO_2 , the supernatant was replaced by the maintenance medium. Adding 20 μL of MTT (0.5 mg/mL) after 48 h, the reaction was ended by removing the supernatant. Then the product was dissolved by 200 μL of DMSO in each well. By using Power WaveX Microplate Scanning Spectrophotometer (Bio-tek Instruments, Inc), optical density (OD) was tested at 570 nm. Finally, the inhibitory percentage of each compound to the cell proliferation was calculated.

Supporting Information ^1H NMR and ^{13}C NMR spectra of compounds **4a**~**4q**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Han, C.; Guo, Y.-C.; Wang, D.-D.; Dai, X.-J.; Wu, F.-J.; Liu, H.-F.; Dai, G.-F.; Tao, J.-C. *Chin. Chem. Lett.* **2015**, 26, 534.
- [2] Freedman, H.; Logan, M. R.; Law, J. K. M.; Houghton M. *ACS Infect. Dis.* **2016**, 2, 749.
- [3] Wetzel, D.; Barbican, A.; Jenzelewski, V.; Schembecker, G.; Merz, J.; Michael Piontek, M. *J. Biotechnol.* **2019**, 306, 203.
- [4] Darweesh, M. F.; Rajput, M. K. S.; Braun, L. J.; Rohila, J. S.; Chase, C. C. L. *Microb. Pathog.* **2018**, 121, 341.
- [5] (a) Buckwold, V. E.; Wei, J.; Wenzel-Mathers, M.; Russell, J. *Antimicrob. Agents Chemother.* **2003**, 47, 2293.
- (b) Yanagida, K.; Baba, C.; Baba, M. *Antiviral Res.* **2004**, 64, 195.
- [6] Wang, L.-Z.; Hua, Z.-X.; Niu, W.-G. *Chin. J. Org. Chem.* **2010**, 30, 1664 (in Chinese).
(王兰芝, 花中霞, 牛文刚, 有机化学, **2010**, 30, 1664.)
- [7] Yin, L.-Y.; Wang, L.-Z.; Li, X.-Q.; An, Y.-S. *Chin. J. Org. Chem.* **2016**, 36, 711 (in Chinese).
(尹刘燕, 王兰芝, 李晓庆, 安迎双, 有机化学, **2016**, 36, 711.)
- [8] Thurston, D. E.; Bose, D. S.; Thompson, A. S.; Howard, P. W.; Leoni, A.; Croker, S. J.; Jenkins, T. C.; Neidle, S.; Hartley, J. A.; Hurley, L. H. *J. Org. Chem.* **1996**, 61, 8141.
- [9] Jaafar, Z.; Chniti, S.; Sassi, A. B.; Dziri, H.; Marque, S.; Lecouvey, M.; Gharbi, R.; Msaddek, M. *J. Mol. Struct.* **2019**, 1195, 689.
- [10] (a) Drummer, O. H. *Forensic Sci. Rev.* **2002**, 14, 1.
(b) Hadjipavlou-Litina, D.; Garg, R.; Hansch, C. *Chem. Rev.* **2004**, 104, 3751.
- [11] Grossi, G.; Braccio, M. D.; Roma, G.; Ballabeni, V.; Tognolini, M.; Calcina, F.; Barocelli, E. *Eur. J. Med. Chem.* **2002**, 37, 933.
- [12] Adegoke, O. A.; Thomas, O. E.; Makanjuola, D. M.; Adewole, O. *J. Taibah Univ. Sci.* **2014**, 8, 248.
- [13] Kruse, H. *Drug Dev. Res.* **1982**, 2, 145.
- [14] Nicholson, A. N.; Stone, B. M.; Clarke, C. H. *J. Clin. Pharmacol.* **1977**, 4, 567.
- [15] Vendeville, S.; Vandyck, K.; Broeck, W. V.; Boutton, C. W.; Bondt, H. D.; Quirynen, L.; Amssoms, K.; Bonfanti, J. F.; Last, S.; Rombaux, K.; Tahri, A.; Hu, L. L.; Delouvroy, F.; Vermeiren, K.; Vandercruyssen, G.; Helm, L. V.; Cleiren, E.; Mostmans, W.; Lory, P.; Pille, G.; Emelen, K. V.; Fanning, G.; Pauwels, F.; Lin, T.; Simmen, K.; Raboisson, P. *Bioorg. Med. Chem. Lett.* **2009**, 19, 2492.
- [16] Nyanguile, O.; Pauwels, F.; Van den Broeck, W.; Boutton, C. W.; Quirynen, L.; Ivens, T.; Van der Helm, L.; Vandercruyssen, G.; Mostmans, W.; Delouvroy, F.; Dehertogh, P.; Cummings, M. D.; Bonfanti, J.-F.; Simmen, K. A.; Raboisson, P. *Antimicrob. Agents Chemother.* **2008**, 52, 4420.
- [17] Lu, L.-Q.; Chen, J.-R.; Xiao, W.-J. *Acc. Chem. Res.* **2012**, 45, 1278.
- [18] Li, X.-M.; Jia, X.-S.; Yin, L. *Chin. J. Org. Chem.* **2017**, 37, 2237 (in Chinese).
(李修明, 贾学顺, 殷亮, 有机化学, **2017**, 37, 2237.)
- [19] Liu, B.-Y.; Xu, X.-J.; Huang, L.-L.; Feng, H.-D. *Chin. J. Org. Chem.* **2020**, 40, 1 (in Chinese).
(刘博瑜, 徐仙君, 黄立梁, 冯煌迪, 有机化学, **2020**, 40, 1.)
- [20] Zhang, M.; Liu, Y.-H.; Shang, Z.-R.; Hu, H.-C.; Zhang, Z.-H. *Catal. Commun.* **2017**, 88, 39.
- [21] Deng, J.; Mo, L.-P.; Zhao, F.-Y.; Hou, L.-L.; Yang, L.; Zhang, Z.-H. *Green Chem.* **2011**, 13, 2576.
- [22] Toure, B. B.; Hall, D. G. *Chem. Rev.* **2009**, 109, 4439.
- [23] Ramon, D. J.; Yus, M. *Angew. Chem., Int. Ed.* **2005**, 44, 1602.
- [24] Royles, B. J. L. *Chem. Rev.* **1995**, 95, 1981.
- [25] Schobert, R.; Schlenk, A. *Bioorg. Med. Chem.* **2008**, 16, 4203.
- [26] Jeong, Y. C.; Moloney, M. G. *Beilstein J. Org. Chem.* **2013**, 9, 1899.
- [27] (a) Ishida, T.; Kobayashi, R.; Yamada, T. *Org. Lett.* **2014**, 16, 2430.
(b) Huang, P. Q.; W.; Ye, J. L. *Chin. J. Chem.* **2015**, 33, 655.
- [28] Sorokina, I. K.; Alekseeva, L. M.; Pashin, V. A.; Asnina, V. V.; Yuzhakov, S. D.; Parimbetova, R. B.; Granik, V. G. *Pharm. Chem. J.* **1991**, 25, 768.

(Fan, Y.)