铜卟咯催化烯丙基 $\text{sp}^3\text{-C-H}$ 键与羧酸的酯化反应

田婉群 李梦媛 杨霜 章浩 刘海洋* 肖新颜*

(华南理工大学化学与化工学院 广东省功能分子工程重点实验室 广州 510641)

摘要 研究了 5,10,15-三(五氟苯基)卟咯铜配合物(CuTPFC)催化的烯丙基 sp^3 杂化 C—H 键与羧酸的氧化交叉脱氢偶联反应(CDC). 该反应可以平稳地生成烯丙基酯, 且具有广泛的底物范围和良好的官能团耐受性. 克级实验中, 在催化剂的用量仅为 0.01 mol% 的情况下, 转换量(TON)可以达到 5100, 表明目前的催化体系在烯丙基酯合成方面具有潜在的应用前景.

关键词 铜; 卟咯; 交叉脱氢偶联; 烯丙基化合物; 催化

Copper Corrole as an Efficient Catalyst for Esterification of Allylic $\text{sp}^3\text{-C-H}$ Bonds with Carboxylic AcidsTian, Wanqun Li, Mengyuan Yang, Shuang Zhang, Hao
Liu, Haiyang* Xiao, Xinyan*

(Key Laboratory of Functional Molecular Engineering of Guangdong Province, School of Chemistry and Chemical Engineering, South China University of Technology, Guangzhou 510641)

Abstract Copper 5,10,15-tris(pentafluorophenyl)corrole (CuTPFC) catalyzed oxidative cross dehydrogenative coupling (CDC) of allylic $\text{sp}^3\text{-C-H}$ bonds and acids had been investigated in a homogeneous system, which provided allylic esters smoothly. This reaction proceeded with broad substrate scope and good functional group tolerance. In a gram scale test, the turnover number (TON) reached 5100 at only 0.01 mol% catalyst loading, indicating the current catalytic system has potential uses in the allylic esters.

Keywords copper; corrole; cross dehydrogenative coupling; allylic compounds; catalysis

1 Introduction

Allylic esters are considered important structural motifs in lots of natural products and bioactive molecules (Figure 1).^[1] The first synthetic reaction of allylic esters, called Kharasch-Sosnovsky reaction, was reported in 1958,^[2] which is achieved via the oxidation of allylic C—H by *tert*-butyl peroxy benzoate in the presence of copper or cobalt salt catalysts (Scheme 1a). An alternative is to use carboxylic acids as the donors of acyloxy free radicals to yield the corresponding allylic esters directly.^[3] Cross-dehydrogenative coupling (CDC) reaction, as a direct C—H functionalization method, avoids the pre-functionalization of starting materials and has become a fundamental, high atom-economic, and highly valuable tool in organic synthesis.^[4] In the past few decades, great efforts have been made to develop new bond-formation strategies for differ-

ent hybrid C—H substrates by using CDC reactions.^[5] For the esterification of allylic $\text{sp}^3\text{-C-H}$ bonds, although tetra-n-butylammonium iodide^[6] and transition metal catalysts such as cobalt,^[7] iron,^[8] copper,^[9] and copper-aluminum^[10] have been developed, high catalyst loading, poor regioselectivity, and stereoselectivity are the main drawbacks of these developed catalysts, especially when the reaction is carried out in a large scale.

Corrole is a trivalent anionic ligand in which there are three pyrrolic N—H protons.^[11] Great advances in the application of metal corrole complex have been made in catalytic oxidation of olefins,^[12] oxidation of sulfides,^[13] hydrogen-atom-transfer reaction,^[14] CO_2 fixation with epoxides,^[15] Ar-Me conversion to Ar-aldehyde,^[16] and epoxidation reaction.^[17] And recently, we had successfully used 5,10,15-tris(pentafluorophenyl)corrole (TPFC) cobalt (Scheme 1d) in the hydration of terminal alkynes (Scheme

* Corresponding authors. E-mail: chhyliu@scut.edu.cn; cexyxiao@scut.edu.cn

Received January 15, 2021; revised March 9, 2021; published online March 25, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21671068, 21878115).

国家自然科学基金(Nos. 21671068, 21878115)资助项目.

Figure 1 Examples of the natural product or precursor with a motif of allylic ester

Scheme 1 Overview of the relevant work

1b).^[18] Despite many metal corrole-catalyzed reactions have been documented, the utilization of metal corrole catalysts in CDC reactions is scarcely explored. In this article, we wish to report the copper TPFC (CuTPFC) complex catalyzed the CDC esterification reaction between allylic C(sp³)—H and carboxylic acids (Scheme 1c). It turned out that CuTPFC complex exhibited good performance with 0.1 mol% catalyst loading with broad substrate scope under optimized conditions. Gram scale test showed that 5100 TON might be achieved, demonstrating its potential practical use. Furthermore, we had also checked the regioselectivity of allylic esterification by using asymmetric allylic olefins.

2 Results and discussion

The optimization of reaction conditions was begun by using benzoic acid **1a** and cyclohexene **2a** as model substrates. When **1a** and **2a** were in the reaction tube with 1 mol% CuTPFC (vs. benzoic acid) as the catalyst and di-*tert*-butyl peroxide (DTBP) as the oxidant in toluene at 120 °C for 12 h, cyclohex-2-en-1-yl benzoate (**3a**) was observed to be the main product with the yield of 47% (Table 1, Entry 1). Then the screening of solvents was

started. The employment of 1,2-dichloroethane (DCE) and ethyl acetate (EA) increased the yield of **3a** to 95% (Table 1, Entry 7) and 93% (Table 1, Entry 6), respectively, whereas using other solvents, including CH₃CN, CHCl₃, MeOH and tetrahydrofuran (THF), did not bring any improvement to the yield of **3a** (Table 1, Entries 2~5). When the catalyst loading was decreased to 0.5 mol%, the yield of **3a** in DCE reached 94% (Table 1, Entry 8). To our surprise, when we continued to decrease the concentration to 0.1 mol%, the yield of **3a** in DCE instead rose to 97% (Table 1, Entry 9), which confirmed DCE as the solvent. These preliminary results motivated us to further decrease the catalyst loading to 0.01 mol%, and **3a** was still obtained a 75% yield with 7500 TON in 12 h (Table 1, Entry 10). Using *tert*-butyl hydroperoxide (TBHP) instead of DTBP, the yield slightly dropped to 87% (Table 1, Entry 11). When TBHP was replaced with other oxidants such as 3-chloroperbenzoic acid (*m*-CPBA), H₂O₂, and 2,3-dicyano-5,6-dichlorobenzoquinone (DDQ), almost no products were detected (Table 1, Entries 12~14). Using MnTPFC, CoTPFC*PPh₃ and Cu(OAc)₂ instead of CuTPFC to catalyze the reaction, the yield of **3a** was only 9%, 15% and 23% (Table 1, Entries 15~16, 18), respectively. When a fluorine-free complex of copper 5,10,15-triphenyl-corrole (CuTPC) was used (Table

1, Entry 17), the yield of **3a** reached 91%, a bit lower than that of CuTPFC, possibly because the electron-withdrawing effect of fluorine makes CuTPFC complex more stable. Using cuprous salt CuOAc instead of CuTPC (Table 1, Entry 19) as the catalyst, the yield of **3a** could reach 84%. And when the reaction was carried out in the absence of catalyst, a 10% yield was obtained (Table 1, Entry 20), which indicated that catalysts had made an important role in the condition.

After the optimized condition was established (Table 1, Entry 9), our attention was then focused on the scope and limitation of this CuTPFC-catalyzed CDC allylic esterification reaction. As shown in Table 2, a series of electron-donating and electron-deficient aryl carboxylic acids were used to react with cyclohexene **2a** and produced good yields of the products of allylic esterification. The yields of **3b**~**3d** decreased slightly as the position of methoxy in phenyl ring was *para*-, *meta*-, and *ortho*-position in turn, and when both *ortho*-positions of benzoic acid were substituted, the yield of **3e** dropped slightly to a lower level. It might prove that the methoxy group in *ortho* had a hindrance to the formation of the product. Moderate to good yields were obtained when the substituents such as methyl, phenyl, *tert*-butyl, hydroxyl, fluoro, chloro, bromo, iodo, cyano groups were located at *para*-position (**3f**~**3n**). Surprisingly, 4-hydroxybenzoic acid and hippuric acid exhibited very high selectivity, merely affording the mono-esterifi-

cation products **3i** and **3v**, respectively. However, in the presence of nitro group or pyridyl group in the substrate, the esterification reaction could not proceed normally (**3o**~**3p**). Non-aromatic carboxylic acids, *e.g.* *trans*-cinnamic acid and 3-phenylpropanoic acid, other aryl carboxylic acids, *e.g.* 1-naphthoic acid, 2-furoic acid and 2-thiophene carboxylic acid, and benzoic anhydride afforded target products in good yield (**3q**~**3u**, **3w**).

After a broad carboxylic acid substrate scope has been established, the possibility of various alkenes having α -H was next examined (Table 3). The reaction of benzoic acid and cyclopentene was achieved to the corresponding esterification product (**4a**) in a 59% yield. Similarly, the esterification product of expanded cyclic olefin substrate cyclooctene was obtained in 69% yield (**4b**). Interestingly, internal and terminal olefins exhibited different regioselectivity of esterification (**4ca**~**4fb**). Reaction of symmetric internal alkene 2,3-dimethylbut-2-ene afforded 2,3-dimethylbut-3-en-2-yl benzoate **4cb** and 2,3-dimethylbut-2-en-1-yl benzoate **4ca** at a ratio of 11.6 to 1, and asymmetric internal alkene 2-methylbut-2-ene provided the products in 63% yield with moderate selectivity (**4da** : **4db** : **4dc** : **4dd** = 1 : 6.6 : 1.5 : 3.2). The terminal alkene substrates hex-1-ene and oct-1-ene was studied and gave products respectively in 50% and 77% yield (**4ea** : **4eb** = 7.3 : 1, **4fa** : **4fb** = 7.4 : 1). With increasing C—H bond strength, esterification products **4h** and **4i** were obtained respectively

Table 1 Optimization of the reaction conditions^a

Entry	Catalyst (mol%)	Solvent	Oxidant	Yield ^b /%	TON
1	CuTPFC (1.0)	Toluene	DTBP	47	47
2	CuTPFC (1.0)	CHCl ₃	DTBP	36	36
3	CuTPFC (1.0)	MeOH	DTBP	20	20
4	CuTPFC (1.0)	CH ₃ CN	DTBP	Trace	—
5	CuTPFC (1.0)	THF	DTBP	N.D. ^c	—
6	CuTPFC (1.0)	EA	DTBP	93	93
7	CuTPFC (1.0)	DCE	DTBP	95	95
8	CuTPFC (0.5)	DCE	DTBP	94	188
9	CuTPFC (0.1)	DCE	DTBP	97 (95 ^d)	970
10	CuTPFC (0.01)	DCE	DTBP	75	7500
11	CuTPFC (0.1)	DCE	TBHP	87	870
12	CuTPFC (0.1)	DCE	<i>m</i> -CPBA	15	150
13	CuTPFC (0.1)	DCE	H ₂ O ₂	17	170
14	CuTPFC (0.1)	DCE	DDQ	N.D. ^c	—
15	MnTPFC (0.1)	DCE	DTBP	9	90
16	CoTPFC (0.1) ^e	DCE	DTBP	15	150
17	CuTPC (0.1)	DCE	DTBP	91	910
18	Cu(OAc) ₂ (0.1)	DCE	DTBP	23	230
19	CuOAc (0.1)	DCE	DTBP	84	840
20	—	DCE	DTBP	10	—

^a Reaction condition: benzoic acid (0.3 mmol), cyclohexene (3 mmol), catalyst (0.1~1.0 mol%), oxidant (0.6 mmol), and solvent (1 mL) at 120 °C for 12 h. ^b Determined by ¹H NMR (internal standard: PhTMS, 10 μ L) and based on carboxylic acid. ^c Not detected. ^d Isolated yield based on carboxylic acid. ^e CoTPFC*PPh₃.

Table 2 CuTPFC-catalyzed CDC reaction of acids with cyclohexene^a

^a Reaction condition: benzoic acid (0.3 mmol), cyclohexene (3 mmol), catalyst CuTPFC (0.1 mol%), oxidant DTBP (0.6 mmol) and DCE (1 mL) at 120 °C for 12 h. N.D. = not detected.

Table 3 CuTPFC-catalyzed CDC reaction of acids with cyclohexene^a

^a Reaction condition: benzoic acid (0.3 mmol), alkene (3 mmol), catalyst CuTPFC (0.1 mol%), oxidant DTBP (0.6 mmol) and DCE (1 mL) at 120 °C for 12 h. N.D. = not detected.

in moderate yields of 41% and 21%, and **4g** were not detected because of a large number of byproducts. When norbornene with only one hydrogen at α -C of double bond was used to react with benzoic acid, esterification products were not detected, no matter α -esterification or β -esterification. Besides, a gram-level reaction experiment was done for benzoic acid and cyclohexene CDC reaction of which TON reached 5100 with 0.01 mol% catalyst loading for 48 h (Scheme 2).

Scheme 2 Gram-level reaction and radical trapping experiment

To further explore the mechanism of allylic esterification, a series of experiments was conducted. When the reaction was carried out in the presence of radical scavenger 2,2,6,6-tetramethylpyridine *N*-oxide (TEMPO), no desired product was observed and compound **6a** trapped by TEMPO was found (Scheme 2). It indicated that the oxidative coupling reaction proceeded through the formation of allylic radical intermediates. The TEMPO was detected by electron paramagnetic resonance (EPR) in the 5-hour reaction mixture of benzoic acid, cyclohexene, CuTPFC, and DCE as shown in lines a, b, and c in Figure 2 (only line c with DTBP). The reaction in line a was stirring at 25 °C and lines b, c was at 120 °C. In the spectra, the free radicals of line b were slightly weaker than those of line a, and almost no EPR signal of line c was observed. The disappearance of radicals indicated that the reactants produced free radicals at high temperatures and confirmed the free radical reaction.

When benzoyl peroxide (BPO) was used to react with cyclohexene, the reactions generated two products at the same time, allylic esterification products **3a** and **5a** (Table 4). When the reactions proceeded without (Table 4, Entry 1) and with (Table 4, Entry 3) 0.1 mol% CuTPFC, **5a** and **3a** were produced at the ratio of 1.4 : 1 and 1 : 4.2, respectively. And when DTBP was added to the two systems as the additional oxidant, the yield ratio of **5a** to **3a** didn't change without catalyst (Table 4, Entry 2) and changed to 1 : 2.7 in the presence of catalyst (Table 4, Entry 4). This indicated that CuTPFC might improve the selectivity of allylic esterification of this reaction and the additional

Figure 2 EPR spectra of (a) Benzoic acid (0.3 mmol), cyclohexene (3 mmol), CuTPFC (0.1 mol%), TEMPO (0.3 mmol) and DCE (1 mL) in room temperature; (b) Benzoic acid (0.3 mmol), cyclohexene (3 mmol), CuTPFC (0.1 mol%), TEMPO (0.3 mmol) and DCE (1 mL) in 120 °C after 5 h; (c) the same condition as b, except adding DTBP (0.6 mmol)

DTBP might improve the yield slightly (Table 4, Entry 3~5).

Table 4 Reactions of benzoyl peroxide and cyclohexene^a

Entry	Catalyst	DTBP (equiv.)	Yield/%		Sum/%
			5a	3a	
1	—	—	26	19	45
2	—	2	28	19	47
3	CuTPFC	—	10	42	52
4	CuTPFC	2	17	46	63
5	CuTPFC	4	16	49	65

^a Reaction condition: benzoyl peroxide (0.15 mmol), cyclohexene (3 mmol), catalyst CuTPFC (0.1 mol%), and DCE (1 mL) at 120 °C for 12 h. Yield determined by ¹H NMR (internal standard: PhTMS, 10 μ L) and based on benzoyl peroxide.

As far as I have investigated the reaction mechanism,^[2a,19] there are three pathways assumed to explain the mechanism of the Kharasch-Sosnovsky reaction: Beckwith-Zavitsas mechanism, Slough mechanism, and synergistic coordination mechanism.^[20] As free radical scavengers succeeded to block the esterification reaction and cyclohexene allylic radical was trapped by TEMPO successfully, synergistic coordination mechanism should not be the main pathway to produce esterification products. A suggested reaction mechanism is shown in Scheme 3. CuTPFC catalyzed the splitting of DTBP to generate ^tBuO-CuTPFC **A** and *t*-butoxy free radical. Then **A** reacted with a carboxylic acid to produce RCOO-CuTPFC **B**, which would further turn into RCOO-allyl-CuTPFC **C** through the Beckwith-Zavitsas or Slough pathway. Finally, through reduc-

tive elimination, **C** would release the target product and complete the catalytic circle. It is necessary to point out that capturing these transient copper corrole intermediate species is of a great challenge and further studies on the detailed reaction mechanism will be our future work.

Scheme 3 Proposed mechanism for esterification

3 Conclusions

In summary, an efficient oxidative coupling of allylic $\text{sp}^3\text{-C-H}$ bonds and carboxylic acids has been achieved successfully by using copper corrole complex as catalyst and DTBP as oxidant. It was observed that CuTPFC was effective for this CDC reaction, while the other tested metal (cobalt and manganese) complexes of TPFC were not effective. Interestingly, in the gram scale test (0.01 mol% CuTPFC catalyst loading at 120 °C, with DTBP as the oxidant), the TON of this CDC reaction reached up to 5100. The reaction can be applied to a wide range of substrates with outstanding functional group tolerance including aryl and aliphatic carboxylic acids, symmetric and asymmetric alkenes. Besides, the regioselectivity of the asymmetric alkene esterification was also explored. CuTPFC-catalyzed oxidative esterification of allylic $\text{sp}^3\text{-C-H}$ bond by acid progressed through a radical mechanism. Further studies on the detailed reaction mechanism and the application of chiral metal corrole complexes in the reaction are currently underway in our laboratory.

4 Experimental section

4.1 Materials and methods

All reagents were of commercial grade and analytical purity. Extracts were dried over anhydrous sodium sulfate and solvents were removed in rotary evaporators under vacuum. The resulting residues were purified using silica gel columns. UV-Visible absorption spectra were measured using a Hitachi 3900H spectrometer in a quartz cell of 1 cm optical path length at room temperature. NMR spectra were

recorded in CDCl_3 with trimethylchlorosilane as the internal standard for ^1H NMR (400 MHz) and ^{13}C NMR (101 MHz). The solvent peak was used as a reference value, for ^1H NMR: CDCl_3 δ 7.26, for ^{13}C NMR: CDCl_3 δ 77.16. HRMS spectra were recorded using an Agilent 1290/Bruker maXis impact mass spectrometer with an ESI source. EPR spectra were recorded using a Bruker Elexsys-II E500 CW-EPR. IR spectra were recorded using a Nicolet IS50-Nicolet Continuum infrared spectrometer.

4.2 Synthetic procedure

Benzoic acids **1** (0.3 mmol), alkene **2** (3 mmol), DTBP (0.6 mmol), CuTPFC (0.1 mol%) and DCE (1 mL) were added to a Schlenk tube and stirred at 120 °C for 12 h. After the reaction mixture was cooled to room temperature and monitored by thin layer chromatography (TLC), which employed glass 0.25 mm GF 254 silica gel plates, it was concentrated to remove the solvent in a rotary evaporator. The crude products were purified by silica gel (300~400 mesh size) column chromatography (hexane/ethyl acetate ($V:V=50:1$)).

All free base corroles were prepared according to the reported procedures.^[21] The metallocorrole 5,10,15-tris(pentafluorophenyl)corrole manganese (III) (MnTPFC), 5,10,15-tris(pentafluorophenyl)corrole copper(III) (CuTPFC), and 5,10,15-tris(pentafluorophenyl)corrole (III) cobalt (CoTPFC*PPh₃) were prepared by using methods described in literatures.^[21-22]

5,10,15-Tris(pentafluorophenyl)corrole copper:^[22a] ^1H NMR (CDCl_3 , 500 MHz) δ : 7.98 (d, $J=4.0$ Hz, 2H), 7.38 (d, $J=4.0$ Hz, 2H), 7.21 (d, $J=4.6$ Hz, 2H), 7.04~7.00 (m, 2H); ^{19}F NMR (CDCl_3 , 471 MHz) δ : -136.65~-136.79 (m), -137.59 (d, $J=24.0$ Hz), -151.95 (q, $J=18.5$ Hz), -160.42~-160.65 (m).

Cyclohex-2-en-1-yl benzoate (**3a**):^[23] 57.6 mg, yield 95%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.14~8.03 (m, 2H), 7.55 (t, $J=7.4$ Hz, 1H), 7.44 (t, $J=7.7$ Hz, 2H), 6.07~5.99 (m, 1H), 5.86 (ddd, $J=8.0, 3.3, 1.7$ Hz, 1H), 5.59~5.50 (m, 1H), 2.15 (ddt, $J=15.2, 7.3, 3.7$ Hz, 1H), 2.07 (ddt, $J=7.5, 5.4, 2.1$ Hz, 1H), 2.05~1.95 (m, 1H), 1.95~1.81 (m, 2H), 1.77~1.67 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.27, 132.88, 132.80, 130.87, 129.65, 128.32, 125.81, 68.67, 28.49, 25.02, 19.02.

Cyclohex-2-en-1-yl 2-methoxybenzoate (**3b**):^[23] 63.7 mg, yield 92%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.77 (dd, $J=7.8, 1.7$ Hz, 1H), 7.47~7.39 (m, 1H), 6.95 (dt, $J=7.3, 3.0$ Hz, 2H), 6.01~5.94 (m, 1H), 5.87~5.79 (m, 1H), 5.52~5.46 (m, 1H), 3.88 (s, 3H), 2.16~1.96 (m, 2H), 1.96~1.90 (m, 1H), 1.90~1.76 (m, 2H), 1.68 (dddd, $J=13.4, 10.2, 7.2, 4.1$ Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.87, 159.24, 133.33, 132.78, 131.54, 125.90, 120.78, 120.13, 112.12, 68.43, 56.06, 28.47, 25.03, 18.98.

Cyclohex-2-en-1-yl 3-methoxybenzoate (**3c**):^[24] 65.6 mg, yield 94%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.64 (d, $J=7.6$ Hz, 1H), 7.56 (s, 1H), 7.32 (t, $J=7.9$ Hz, 1H), 7.07 (dd, $J=8.2, 2.4$ Hz, 1H), 6.02~5.95 (m, 1H), 5.82 (d, $J=9.9$ Hz, 1H), 5.53~5.45 (m, 1H), 3.84 (s, 3H),

2.18~2.04 (m, 2H), 1.97 (ddd, $J=17.8, 8.7, 3.8$ Hz, 2H), 1.91~1.78 (m, 2H), 1.70 (ddt, $J=13.4, 7.9, 4.4$ Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.17, 159.60, 132.92, 132.21, 129.37, 125.79, 122.08, 119.23, 114.25, 68.83, 55.51, 28.48, 25.03, 19.04.

Cyclohex-2-en-1-yl 4-methoxybenzoate (**3d**):^[23] 69 mg, yield 99%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.00 (d, $J=8.9$ Hz, 2H), 6.89 (d, $J=8.8$ Hz, 2H), 6.02~5.94 (m, 1H), 5.85~5.78 (m, 1H), 5.47 (d, $J=1.8$ Hz, 1H), 3.84 (s, 3H), 2.17~2.05 (m, 1H), 2.01~1.91 (m, 1H), 1.83 (dddd, $J=19.6, 16.5, 9.0, 3.0$ Hz, 3H), 1.74~1.64 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.99, 163.24, 132.61, 131.59, 125.98, 123.24, 113.50, 68.24, 55.42, 28.48, 24.97, 19.01.

Cyclohex-2-en-1-yl 2,6-dimethoxybenzoate (**3e**):^[9b] 66.9 mg, yield 85%. Yellow solid, m.p. 69~70 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.22 (d, $J=8.4$ Hz, 1H), 6.52 (d, $J=8.4$ Hz, 2H), 6.00~5.92 (m, 1H), 5.85~5.77 (m, 1H), 5.58~5.50 (m, 1H), 3.78 (s, 6H), 2.09~1.88 (m, 4H), 1.78 (dq, $J=9.4, 3.6$ Hz, 1H), 1.70~1.58 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.32, 157.39, 132.73, 130.89, 125.85, 113.65, 104.08, 69.06, 56.07, 56.04, 28.30, 24.95, 18.93.

Cyclohex-2-en-1-yl 4-methylbenzoate (**3f**):^[23] 62.3 mg, yield 96%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.94 (d, $J=8.1$ Hz, 2H), 7.22 (d, $J=8.0$ Hz, 2H), 6.03~5.96 (m, 1H), 5.83 (d, $J=7.8$ Hz, 1H), 5.53~5.46 (m, 1H), 2.39 (s, 3H), 2.19~2.08 (m, 1H), 2.05 (s, 1H), 1.97 (ddd, $J=17.9, 8.8, 3.8$ Hz, 2H), 1.85 (dddd, $J=18.8, 15.9, 9.0, 3.2$ Hz, 3H), 1.75~1.64 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.37, 143.42, 132.76, 129.70, 129.05, 128.14, 125.97, 68.43, 28.53, 25.04, 21.72, 19.07.

Cyclohex-2-en-1-yl [1,1'-biphenyl]-4-carboxylate (**3g**):^[7] 81.1 mg, yield 97%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.14 (d, $J=8.4$ Hz, 2H), 7.65 (dd, $J=13.4, 7.9$ Hz, 4H), 7.47 (t, $J=7.4$ Hz, 2H), 7.40 (t, $J=7.3$ Hz, 1H), 6.06~6.00 (m, 1H), 5.91~5.85 (m, 1H), 5.59~5.53 (m, 1H), 2.22~2.12 (m, 1H), 2.12~2.06 (m, 1H), 2.01 (ddd, $J=13.4, 8.5, 3.7$ Hz, 1H), 1.97~1.82 (m, 2H), 1.74 (ddd, $J=10.5, 8.5, 5.4$ Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.19, 145.57, 140.17, 132.93, 130.21, 129.64, 129.00, 128.17, 127.36, 127.05, 125.88, 68.71, 28.55, 25.07, 19.08.

Cyclohex-2-en-1-yl 4-(*tert*-butyl)benzoate (**3h**):^[23] 46.6 mg, yield 60%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.99 (d, $J=8.5$ Hz, 2H), 7.45 (d, $J=8.5$ Hz, 2H), 6.03~5.96 (m, 1H), 5.87~5.79 (m, 1H), 5.51 (d, $J=1.9$ Hz, 1H), 2.19~2.03 (m, 2H), 1.97 (ddd, $J=18.0, 8.2, 3.6$ Hz, 1H), 1.86 (dddd, $J=16.7, 13.7, 9.0, 4.8$ Hz, 2H), 1.75~1.64 (m, 1H), 1.34 (s, 9H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.35, 156.48, 132.77, 129.59, 128.14, 126.02, 125.34, 68.42, 35.15, 31.24, 28.57, 25.08, 19.09.

Cyclohex-2-en-1-yl 4-hydroxybenzoate (**3i**): 41.6 mg, yield 64%. Yellow solid, m.p. 109~110 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.95 (d, $J=8.8$ Hz, 2H), 7.22 (s, 1H), 6.89 (d, $J=8.8$ Hz, 2H), 6.00 (dt, $J=10.0, 3.3$ Hz, 1H), 5.82 (ddt, $J=10.1, 4.1, 2.2$ Hz, 1H), 5.48 (tdd, $J=5.7, 3.6, 1.9$ Hz, 1H), 2.12 (dtdd, $J=13.5, 5.7, 3.8, 1.9$ Hz, 1H), 2.02 (dddt, $J=19.4, 6.3, 4.6, 2.2$ Hz, 1H), 1.98~1.91 (m, 1H),

1.85 (dddd, $J=22.2, 10.9, 5.9, 3.3$ Hz, 2H), 1.69 (dddd, $J=13.5, 8.0, 5.6, 3.2$ Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.96, 166.91, 160.70, 160.65, 133.00, 131.99, 125.63, 122.53, 115.31, 68.77, 28.44, 24.95, 18.92. IR (KBr) ν : 3381, 1690, 1595, 1278 cm^{-1} . HRMS calcd for $\text{C}_{13}\text{H}_{14}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 241.0943, found 241.0827.

Cyclohex-2-en-1-yl 4-fluorobenzoate (**3j**):^[23] 61.6 mg, yield 93%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.10~8.00 (m, 2H), 7.08 (t, $J=8.6$ Hz, 2H), 6.03~5.95 (m, 1H), 5.85~5.78 (m, 1H), 5.52~5.45 (m, 1H), 2.19~2.08 (m, 1H), 2.04 (dd, $J=6.3, 2.5$ Hz, 1H), 1.96 (ddd, $J=17.9, 8.2, 4.1$ Hz, 1H), 1.91~1.77 (m, 2H), 1.75~1.63 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.93 (C-1), 165.24, 164.41 (C-1), 132.96, 132.15 (C-3), 132.06 (C-3), 127.05 (C-4), 127.02 (C-4), 125.60, 115.47 (C-2), 115.26 (C-2), 68.75, 28.40, 24.93, 18.92; ^{19}F NMR (CDCl_3 , 471 MHz) δ : -106.14 (p, $J=7.0, 6.6$ Hz).

Cyclohex-2-en-1-yl 4-chlorobenzoate (**3k**):^[23] 70.1 mg, yield 99%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.97 (d, $J=8.5$ Hz, 2H), 7.38 (d, $J=8.5$ Hz, 2H), 6.04~5.96 (m, 1H), 5.85~5.78 (m, 1H), 5.48 (d, $J=2.1$ Hz, 1H), 2.18~2.08 (m, 1H), 2.04 (dd, $J=4.6, 2.5$ Hz, 1H), 1.96 (ddd, $J=18.0, 8.9, 4.0$ Hz, 1H), 1.91~1.76 (m, 2H), 1.74~1.64 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.43, 139.26, 133.15, 131.09, 129.35, 128.69, 125.61, 69.04, 28.47, 25.03, 19.00.

Cyclohex-2-en-1-yl 4-bromobenzoate (**3l**):^[23] 63.9 mg, yield 76%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.90 (d, $J=8.5$ Hz, 2H), 7.55 (d, $J=8.5$ Hz, 2H), 6.04~5.96 (m, 1H), 5.81 (dd, $J=9.4, 2.3$ Hz, 1H), 5.52~5.45 (m, 1H), 2.19~2.04 (m, 2H), 1.96 (dq, $J=13.5, 4.9, 4.4$ Hz, 1H), 1.84 (dddd, $J=25.8, 16.0, 9.0, 3.0$ Hz, 2H), 1.70 (ddd, $J=10.4, 8.5, 5.3$ Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.58, 133.19, 131.70, 131.24, 129.80, 127.93, 125.58, 69.08, 28.47, 25.03, 19.01.

Cyclohex-2-en-1-yl 4-iodobenzoate (**3m**):^[9a] 91.9 mg, yield 93%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.80~7.67 (m, 4H), 5.99 (dd, $J=9.6, 2.7$ Hz, 1H), 5.85~5.76 (m, 1H), 5.52~5.44 (m, 1H), 2.19~2.07 (m, 1H), 2.07~2.01 (m, 1H), 2.01~1.91 (m, 1H), 1.90~1.75 (m, 2H), 1.69 (dtd, $J=10.2, 5.1, 2.3$ Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.74, 137.69, 133.14, 131.15, 130.35, 125.58, 100.61, 69.05, 69.00, 28.45, 25.02, 19.00.

Cyclohex-2-en-1-yl 4-cyanobenzoate (**3n**):^[6] 57.5 mg, yield 84%. Yellow solid, m.p. 68~69 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 8.13 (d, $J=8.4$ Hz, 2H), 7.72 (d, $J=8.4$ Hz, 2H), 6.03 (dt, $J=8.9, 3.3$ Hz, 1H), 5.85~5.77 (m, 1H), 5.54~5.47 (m, 1H), 2.19~2.04 (m, 2H), 1.98 (ddd, $J=18.0, 8.3, 3.8$ Hz, 1H), 1.92~1.76 (m, 2H), 1.75~1.65 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 164.65, 134.74, 133.64, 132.24, 130.21, 125.15, 118.15, 116.29, 69.74, 28.39, 25.00, 18.90.

Cyclohex-2-en-1-yl 1-naphthoate (**3q**):^[6] 39.5 mg, yield

52%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.93 (d, $J=8.5$ Hz, 1H), 8.19 (dd, $J=7.3$, 1.2 Hz, 1H), 8.01 (d, $J=8.2$ Hz, 1H), 7.88 (d, $J=8.1$ Hz, 1H), 7.62 (ddd, $J=8.5$, 6.8, 1.4 Hz, 1H), 7.57~7.45 (m, 2H), 6.09~6.02 (m, 1H), 5.94 (ddt, $J=10.0$, 3.8, 2.0 Hz, 1H), 5.68~5.61 (m, 1H), 2.23~2.13 (m, 1H), 2.10 (ddd, $J=7.2$, 5.6, 3.4 Hz, 1H), 2.07~1.94 (m, 2H), 1.88 (ddt, $J=11.2$, 5.6, 2.5 Hz, 1H), 1.75 (ddt, $J=11.2$, 5.6, 2.7 Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 167.40, 133.94, 133.19, 133.08, 131.44, 130.16, 128.60, 127.93, 127.71, 126.21, 125.95, 125.83, 124.59, 68.86, 28.58, 25.09, 19.11.

Cyclohex-2-en-1-yl thiophene-2-carboxylate (**3r**):^[6] 53.2 mg, yield 85%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.79 (dd, $J=3.7$, 1.2 Hz, 1H), 7.52 (dd, $J=5.0$, 1.2 Hz, 1H), 7.08 (dd, $J=4.9$, 3.8 Hz, 1H), 6.04~5.94 (m, 1H), 5.81 (ddd, $J=7.9$, 3.4, 1.7 Hz, 1H), 5.51~5.42 (m, 1H), 2.18~2.07 (m, 1H), 2.08~1.97 (m, 1H), 1.97~1.76 (m, 3H), 1.68 (dddt, $J=13.7$, 8.3, 5.6, 3.0 Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 162.00, 134.56, 133.25, 133.06, 132.21, 127.70, 125.61, 69.04, 28.45, 24.98, 18.96.

Cyclohex-2-en-1-yl furan-2-carboxylate (**3s**):^[9b] 47.2 mg, yield 82%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.59~7.53 (m, 1H), 7.18~7.12 (m, 1H), 6.48 (dd, $J=3.5$, 1.7 Hz, 1H), 6.03~5.96 (m, 1H), 5.82~5.75 (m, 1H), 5.53~5.43 (m, 1H), 2.17~2.06 (m, 1H), 2.06~1.97 (m, 1H), 1.96~1.84 (m, 2H), 1.84~1.74 (m, 1H), 1.72~1.61 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 158.54, 146.19, 145.15, 133.23, 125.46, 117.73, 111.79, 68.88, 28.42, 24.95, 18.96.

Cyclohex-2-en-1-yl cinnamate (**3t**):^[6] 55.8 mg, yield 82%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.69 (d, $J=16.0$ Hz, 1H), 7.52 (dd, $J=6.6$, 2.8 Hz, 2H), 7.41~7.34 (m, 3H), 6.45 (d, $J=16.0$ Hz, 1H), 6.03~5.96 (m, 1H), 5.84~5.75 (m, 1H), 5.46~5.37 (m, 1H), 2.18~2.07 (m, 1H), 2.02 (s, 1H), 1.95 (ddt, $J=13.5$, 9.0, 3.7 Hz, 1H), 1.86~1.75 (m, 2H), 1.68 (ddt, $J=13.2$, 7.8, 3.0 Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.72, 144.61, 134.63, 132.87, 130.27, 128.96, 128.14, 125.90, 118.75, 68.27, 28.51, 25.02, 19.04.

Cyclohex-2-en-1-yl 3-phenylpropanoate (**3u**):^[6] 65.5 mg, yield 95%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 7.31~7.24 (m, 2H), 7.20 (d, $J=7.4$ Hz, 3H), 5.93 (ddd, $J=10.1$, 3.8, 1.3 Hz, 1H), 5.67 (ddt, $J=10.1$, 4.1, 2.2 Hz, 1H), 5.27 (ddt, $J=5.3$, 3.7, 1.6 Hz, 1H), 2.95 (t, $J=7.8$ Hz, 2H), 2.62 (dd, $J=8.5$, 7.1 Hz, 2H), 2.12~1.92 (m, 2H), 1.89~1.78 (m, 1H), 1.75~1.55 (m, 3H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 172.61, 140.60, 132.69, 128.48, 128.36, 126.24, 125.73, 68.16, 36.25, 31.12, 28.32, 24.90, 18.88.

Cyclohex-2-en-1-yl benzoylglycinate (**3v**):^[6] 28.8 mg, yield 37%. Yellow solid, m.p. 69~70 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.87~7.79 (m, 2H), 7.53 (t, $J=7.3$ Hz, 1H), 7.46 (t, $J=7.5$ Hz, 2H), 6.73 (s, 1H), 6.05~5.98 (m, 1H), 5.79~5.71 (m, 1H), 5.39 (d, $J=3.3$ Hz, 1H), 4.25 (dd, $J=5.0$, 1.6 Hz, 2H), 2.19~1.97 (m, 2H), 1.91 (ddd, $J=13.5$, 9.3, 5.4 Hz, 1H), 1.86~1.73 (m, 2H), 1.73~1.63 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 169.92, 133.69, 131.89, 128.74, 127.20, 125.01, 69.83, 42.24, 28.31, 24.94,

18.78.

cyclopent-2-en-1-yl benzoate (**4a**):^[10] 33.3 mg, yield 59%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.08~7.97 (m, 2H), 7.58~7.50 (m, 1H), 7.42 (dd, $J=8.4$, 7.0 Hz, 2H), 6.16 (dt, $J=4.5$, 2.2 Hz, 1H), 5.99~5.90 (m, 2H), 2.66~2.54 (m, 1H), 2.46~2.32 (m, 2H), 2.02~1.92 (m, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.71, 137.81, 132.87, 130.85, 129.70, 129.52, 128.39, 31.31, 30.04.

cyclooct-2-en-1-yl benzoate (**4b**):^[10] 47.8 mg, yield 69%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.08~8.02 (m, 2H), 7.57~7.51 (m, 1H), 7.43 (dd, $J=8.4$, 6.9 Hz, 2H), 5.91 (dddd, $J=8.4$, 6.6, 4.9, 2.5 Hz, 1H), 5.72 (dddd, $J=10.6$, 8.7, 7.0, 1.5 Hz, 1H), 5.62 (ddd, $J=10.8$, 6.9, 1.2 Hz, 1H), 2.41~2.28 (m, 1H), 2.16 (ddt, $J=13.9$, 7.6, 4.1 Hz, 1H), 2.06 (qt, $J=8.3$, 2.9 Hz, 1H), 1.79~1.52 (m, 5H), 1.44 (tdd, $J=12.9$, 6.5, 3.6 Hz, 2H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.07, 132.85, 130.88, 130.79, 129.96, 129.67, 128.37, 73.09, 35.25, 28.94, 26.52, 26.01, 23.52.

2,3-dimethylbut-2-en-1-yl benzoate (**4ca**):^[7] + 2,3-dimethylbut-3-en-2-yl benzoate (**4cb**): 49.3 mg, yield 81%. **4cb** : **4ca** = 11.6 : 1, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.11 (dd, $J=8.3$, 1.4 Hz, 2H), 8.09~8.05 (m, 23.2H), 7.62~7.55 (m, 12.6H), 7.51~7.44 (m, 25.2H), 5.07 (s, 11.6H), 4.97 (p, $J=1.4$ Hz, 11.6H), 4.91 (s, 2H), 1.87 (d, $J=1.5$ Hz, 3H), 1.85 (t, $J=1.0$ Hz, 34.8H), 1.84 (s, 3H), 1.79 (s, 3H), 1.74 (s, 69.6H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.35, 148.44, 132.85, 132.70, 131.59, 129.68, 129.58, 128.40, 128.36, 110.29, 83.36, 26.66, 21.03, 18.86, 16.83; IR (KBr) ν : 1719, 1283, 1109, 712 cm^{-1} ; HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{O}_2\text{Na}$ [$\text{M} + \text{Na}$] $^+$ 227.1150, found 227.1038.

2-Methylbut-2-en-1-yl benzoate (**4da**):^[25] + 3-methylbut-3-en-2-yl benzoate (**4db**):^[26] + 3-methylbut-2-en-1-yl benzoate (**4dc**):^[27] + 2-methylbut-3-en-2-yl benzoate (**4dd**):^[28] 62%, 35.4 mg. **4da** : **4db** : **4dc** : **4dd** = 1 : 6.6 : 1.5 : 3.2, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.07 (d, $J=7.0$ Hz, 14H), 8.06~7.99 (m, 13H), 7.58~7.50 (m, 13H), 7.47~7.38 (m, 27H), 6.21 (dd, $J=17.5$, 10.9 Hz, 3.2H), 5.68~5.61 (m, 1H), 5.54 (q, $J=6.5$ Hz, 6.6H), 5.51~5.45 (m, 1.5H), 5.28 (d, $J=17.5$ Hz, 3.2H), 5.14 (dd, $J=10.9$, 0.6 Hz, 3.2H), 5.06 (s, 6.6H), 4.91 (s, 6.6H), 4.83 (d, $J=7.2$ Hz, 3H), 4.72 (s, 2H), 1.83 (s, 19.8H), 1.79 (s, 4.5H), 1.78 (s, 4.5H), 1.74 (s, 3H), 1.69 (d, $J=3.2$ Hz, 3H), 1.67 (s, 19.2H), 1.47 (s, 9.9H), 1.45 (s, 9.9H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.87, 144.73, 142.72, 139.22, 132.95, 132.89, 132.68, 130.80, 129.70, 129.59, 128.45, 128.41, 128.34, 112.94, 111.78, 74.11, 70.80, 62.00, 26.72, 25.90, 19.35, 18.50, 18.23.

Hex-1-en-3-yl benzoate (**4ea**):^[29] + hex-2-en-1-yl benzoate (**4eb**):^[30] 30.4 mg, yield 50%. **4ea** : **4eb** = 7.3 : 1, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.09~8.03 (m, 16.6H), 7.55 (s, 8.3H), 7.44 (d, $J=15.2$ Hz, 16.6H), 5.90 (ddd, $J=17.0$, 10.5, 6.2 Hz, 7.3H), 5.83 (d, $J=6.6$ Hz, 1H), 5.72~5.65 (m, 1H), 5.51 (q, $J=6.1$ Hz, 7.3H), 5.34 (s, 7.3H), 5.20 (d, $J=10.5$ Hz, 7.3H), 4.77 (d, $J=6.4$ Hz, 2H), 2.07 (d, $J=7.1$ Hz, 2H), 1.83~1.71 (m, 14.6H), 1.45 (ddd, $J=15.6$, 11.8, 7.2 Hz, 19.6H), 0.96 (t, $J=7.4$ Hz, 21.9H);

^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.01, 136.78, 132.96, 130.77, 129.73, 129.71, 128.48, 128.45, 128.40, 116.63, 36.60, 22.20, 18.52, 14.01.

Oct-1-en-3-yl benzoate (**4fa**)^[31]+oct-2-en-1-yl benzoate (**4fb**): 77%, 53.8 mg. **4fa** : **4fb** = 7.4 : 1, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.10~8.03 (m, 16.8H), 7.54 (d, J =7.4 Hz, 8.4H), 7.44 (t, J =7.6 Hz, 16.8H), 5.88 (dd, J =10.6, 6.4 Hz, 7.4H), 5.84 (d, J =6.6 Hz, 1H), 5.72~5.64 (m, 1H), 5.50 (d, J =7.0 Hz, 7.4H), 5.32 (d, J =17.2 Hz, 7.4H), 5.20 (d, J =10.5 Hz, 7.4H), 4.77 (d, J =5.7 Hz, 2.2H), 2.08 (dd, J =14.9, 7.9 Hz, 5H), 1.85~1.66 (m, 22.2H), 1.46~1.37 (m, 22.2H), 0.88 (t, J =6.9 Hz, 43H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.97, 136.78, 132.93, 130.77, 129.69, 128.43, 123.89, 116.65, 75.48, 65.86, 34.42, 31.71, 24.87, 22.78, 22.63, 14.22, 14.09; IR (KBr) ν : 2957, 2930, 2859, 1721, 1271, 1111, 712 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{20}\text{O}_2\text{Na}$ [$\text{M}+\text{Na}$]⁺ 255.3230, found 255.1356.

1-Phenylethyl benzoate (**4h**)^[6] 27.6 mg, yield 41%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.12 (dt, J =7.1, 1.4 Hz, 2H), 7.64~7.56 (m, 1H), 7.51~7.44 (m, 6H), 7.40 (ddt, J =7.9, 5.8, 1.6 Hz, 5H), 7.34 (td, J =7.4, 7.0, 1.5 Hz, 3H), 6.18 (q, J =6.6 Hz, 1H), 1.71 (d, J =6.6 Hz, 3H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 165.84, 141.83, 132.93, 130.58, 129.67, 128.58, 128.36, 127.91, 126.08, 72.94, 22.44.

2,3-Dihydro-1*H*-inden-1-yl benzoate (**4i**)^[10] 15 mg, yield 21%, yellow oil. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.08~8.02 (m, 2H), 7.57~7.52 (m, 1H), 7.51 (d, J =7.5 Hz, 1H), 7.42 (dd, J =8.4, 7.1 Hz, 2H), 7.34~7.31 (m, 2H), 7.28~7.24 (m, 1H), 6.47 (dd, J =7.1, 4.1 Hz, 1H), 3.20 (ddd, J =15.0, 8.6, 5.9 Hz, 1H), 2.96 (ddd, J =16.0, 8.5, 5.3 Hz, 1H), 2.65 (dddd, J =14.3, 8.6, 7.0, 5.9 Hz, 1H), 2.25 (dddd, J =13.9, 8.9, 5.2, 4.0 Hz, 1H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 166.73, 144.55, 141.26, 133.02, 130.64, 129.82, 129.09, 128.43, 126.87, 125.81, 124.97, 32.63, 30.43.

1-(Cyclohex-2-en-1-yloxy)-2,2,6,6-tetramethylpiperidine (**6a**): HRMS calcd for $\text{C}_{15}\text{H}_{28}\text{NO}$ [$\text{M}+\text{H}$]⁺ 238.2093, found 238.2167.

Supporting Information Supplemental result and spectra for corroles, metallocorroles, compounds **3** and **4** mentioned in the article. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Saito, T.; Fuwa, H.; Sasaki, M. *Org. Lett.* **2009**, *11*, 5274.
(b) Kolodziej, H.; Burmeister, A.; Trun, W.; Radtke, O. A.; Kiderlen, A. F.; Ito, H.; Hatano, T.; Yoshida, T.; Foo, L. Y. *Bioorg. Med. Chem.* **2005**, *13*, 6470.
(c) Takeuchi, Y.; Shi, Q. W.; Sugiyama, T.; Oritani, T. *Biosci. Biotechnol. Biochem.* **2002**, *66*, 537.
(d) Kijjoa, A.; Bessa, J.; Pinto, M. M.; Anatachoke, C.; Silva, A. M.; Eaton, G.; Herz, W. *Phytochemistry* **2002**, *59*, 543.
(e) Covell, D. J.; Vermeulen, N. A.; Labenz, N. A.; White, M. C. *Angew. Chem., Int. Ed.* **2006**, *45*, 8217.
(f) Huong, D. T.; Kamperdick, C.; Sung, T. V. *J. Nat. Prod.* **2004**, *67*, 445.
- [2] (a) Kharasch, M. S.; Sosnovsky, G.; Yang, N. C. *J. Am. Chem. Soc.* **1959**, *81*, 5819.
(b) Kharasch, M. S.; Sosnovsky, G. *J. Am. Chem. Soc.* **1958**, *80*, 756.
- [3] (a) Shi, E. B.; Shao, Y.; Chen, S. L.; Hu, H. Y.; Liu, Z. J.; Zhang, J.; Wan, X. B. *Org. Lett.* **2012**, *14*, 3384.
(b) Aakermark, B.; Larsson, E. M.; Oslob, J. D. *J. Org. Chem.* **1994**, *59*, 5729.
- [4] (a) Ashenhurst, J. A. *Chem. Soc. Rev.* **2010**, *39*, 540.
(b) Scheuermann, C. J. *Chem.-Asian J.* **2010**, *5*, 436.
(c) Lv, L.; Li, Z. *Top. Curr. Chem.* **2016**, *374*, 38.
(d) Li, C. J. *Acc. Chem. Res.* **2009**, *42*, 335.
(e) Girard, S. A.; Knauber, T.; Li, C. J. *Angew. Chem., Int. Ed.* **2014**, *53*, 74.
(f) Faisca Phillips, A. M.; Pombeiro, A. J. L. *ChemCatChem* **2018**, *10*, 3354.
- [5] (a) Young, A. J.; White, M. C. *J. Am. Chem. Soc.* **2008**, *130*, 14090.
(b) Su, X.; Surry, D. S.; Spandl, R. J.; Spring, D. R. *Org. Lett.* **2008**, *10*, 2593.
(c) Meng, Z.; Sun, S.; Yuan, H.; Lou, H.; Liu, L. *Angew. Chem., Int. Ed.* **2014**, *53*, 543.
(d) Liu, Z. Q.; Zhao, L.; Shang, X.; Cui, Z. *Org. Lett.* **2012**, *14*, 3218.
(e) Lin, S.; Song, C. X.; Cai, G. X.; Wang, W. H.; Shi, Z. J. *J. Am. Chem. Soc.* **2008**, *130*, 12901.
(f) Li, Z.; Bohle, D. S.; Li, C. J. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 8928.
(g) Zhao, S.; Yuan, J.; Li, Y. C.; Shi, B. F. *Chem. Commun.* **2015**, *51*, 12823.
(h) Zhao, J.; Fang, H.; Zhou, W.; Han, J.; Pan, Y. *J. Org. Chem.* **2014**, *79*, 3847.
(i) Tsang, A. S.; Kapat, A.; Schoenebeck, F. *J. Am. Chem. Soc.* **2016**, *138*, 518.
(j) Sun, S.; Li, C.; Floreancig, P. E.; Lou, H.; Liu, L. *Org. Lett.* **2015**, *17*, 1684.
(k) Liang, Y. F.; Jiao, N. *Angew. Chem., Int. Ed.* **2014**, *53*, 548.
(l) Pan, X.; Liu, X.; Sun, S.; Meng, Z.; Liu, L. *Chin. J. Chem.* **2018**, *36*, 1187.
(m) Lu, F.; Yang, Z.; Wang, T.; Wang, T.; Zhang, Y.; Yuan, Y.; Lei, A. *Chin. J. Chem.* **2019**, *37*, 547.
(n) Xiao, Y. X.; Liu, Z. Q. *Acta Chim. Sinica* **2019**, *77*, 874 (in Chinese).
(肖莹霞, 柳忠全, 化学学报, **2019**, *77*, 874.)
- [6] Shi, E.; Shao, Y.; Chen, S.; Hu, H.; Liu, Z.; Zhang, J.; Wan, X. *Org. Lett.* **2012**, *14*, 3384.
- [7] Lu, B.; Zhu, F.; Wang, D.; Sun, H.; Shen, Q. *Tetrahedron Lett.* **2017**, *58*, 2490.
- [8] Ren, T.-L.; Xu, B.-H.; Mahmood, S.; Sun, M.-X.; Zhang, S.-J. *Tetrahedron* **2017**, *73*, 2943.
- [9] (a) Wang, C. Y.; Song, R. J.; Wei, W. T.; Fan, J. H.; Li, J. H. *Chem. Commun.* **2015**, *51*, 2361.
(b) Tran, B. L.; Driess, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2014**, *136*, 17292.
(c) Zhou, J.; Jin, C.; Li, X.; Su, W. *RSC Adv.* **2015**, *5*, 7232.
- [10] Garcia-Cabeza, A. L.; Marin-Barrios, R.; Moreno-Dorado, F. J.; Ortega, M. J.; Massanet, G. M.; Guerra, F. M. *Org. Lett.* **2014**, *16*, 1598.
- [11] (a) Ghosh, A. *Chem. Rev.* **2017**, *117*, 3798.
(b) Barata, J. F. B.; Neves, M. G. P. M. S.; Faustino, M. A. F.; Tomé, A. C.; Cavaleiro, J. A. S. *Chem. Rev.* **2016**, *117*, 3192.
- [12] (a) Ali, A.; Akram, W.; Liu, H. Y. *Molecules* **2018**, *24*, 78.
(b) Huang, L.-T.; Ali, A.; Wang, H.-H.; Cheng, F.; Liu, H.-Y. *J. Mol. Catal. A: Chem.* **2017**, *426*, 213.
- [13] (a) Chen, T. H.; Kwong, K. W.; Lee, N. F.; Ranburger, D.; Zhang, R. *Inorg. Chim. Acta* **2016**, *451*, 65.
(b) Ranburger, D.; Willis, B.; Kash, B.; Jeddi, H.; Alcantar, C.; Zhang, R. *Inorg. Chim. Acta* **2019**, *487*, 41.
- [14] Sacramento, J. J. D.; Goldberg, D. P. *Acc. Chem. Res.* **2018**, *51*, 2641.
- [15] (a) Li, P.; Cao, Z. *Organometallics* **2018**, *37*, 406.

- (b) Tiffner, M.; Gonglach, S.; Haas, M.; Schofberger, W.; Waser, M. *Chem. Asian J.* **2017**, *12*, 1048.
- [16] Zhan, X.; Kolanu, S.; Fite, S.; Chen, Q. C.; Lee, W.; Churchill, D. G.; Gross, Z. *Photochem. Photobiol. Sci.* **2020**, *19*, 996.
- [17] Nayak, M.; Nayak, P.; Sahu, K.; Kar, S. *J. Org. Chem.* **2020**, *85*, 11654.
- [18] Lai, J.-W.; Liu, Z.-Y.; Chen, X.-Y.; Zhang, H.; Liu, H.-Y. *Tetrahedron Lett.* **2020**, *61*.
- [19] Schaffer, R.; Isbell, H. S. *J. Am. Chem. Soc.* **1958**, *80*, 756.
- [20] Mayoral, J. A.; Rodriguez-Rodriguez, S.; Salvatella, L. *Chem.-Eur. J.* **2008**, *14*, 9274.
- [21] Koszarna, B.; Gryko, D. T. *J. Org. Chem.* **2006**, *71*, 3707.
- [22] (a) Luobeznova, I.; Simkhovich, L.; Goldberg, I.; Gross, Z. *Eur. J. Inorg. Chem.* **2004**, *2004*, 1724.
(b) Chen, Y.; Fan, Q.-H.; Hossain, M. S.; Zhan, S.-Z.; Liu, H.-Y.; Si, L.-P. *Eur. J. Inorg. Chem.* **2020**, *2020*, 491.
- [23] Li, J.; Jia, X.; Deng, H.; Li, C.; Jin, T.; Li, C. *Synlett* **2018**, *29*, 840.
- [24] Liu, G.; Jin, C.; Wu, G. *Synlett* **2014**, *25*, 2908.
- [25] Wright, B. J.; Hartung, J.; Peng, F.; Van de Water, R.; Liu, H.; Tan, Q. H.; Chou, T. C.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2008**, *130*, 16786.
- [26] Werkhoven, T. M.; Siebum, A. H. G.; Lugtenburg, J. *Eur. J. Org. Chem.* **2000**, *2000*, 2113.
- [27] Tanaka, S.; Nakashima, T.; Satou, N.; Oono, H.; Kon, Y.; Tamura, M.; Sato, K. *Tetrahedron Lett.* **2019**, *60*, 2009.
- [28] Sedighi, M.; Calimsiz, S.; Lipton, M. A. *J. Org. Chem.* **2006**, *71*, 9517.
- [29] Mamone, P.; Grunberg, M. F.; Fromm, A.; Khan, B. A.; Goossen, L. *J. Org. Lett.* **2012**, *14*, 3716.
- [30] Lim, S.; Ji, M.; Wang, X.; Lee, C.; Jang, H.-Y. *Eur. J. Org. Chem.* **2015**, *2015*, 591.
- [31] Whittaker, A. M.; Lalic, G. *Org. Lett.* **2013**, *15*, 1112.

(Cheng, F.)