一种大环类双核非离子型磁共振对比剂的合成及弛豫性能

周进^{†,a,b} 孙宏顺^{†,*,a,b} 李玉龙^b 蒋蕪^b
郭成^{*,a} 沈临江^c

(a 南京工业大学化学与分子工程学院 南京 211816)

(b 南京科技职业学院 江苏省磁共振靶向显像剂工程实验室 南京 210048)

(c 南京工业大学数理科学学院 南京 211816)

摘要 磁共振成像技术被广泛应用于诊断医学和软组织成像,而磁共振对比剂有助于提高成像对比度。报道了一种基于钆-DOTA-酰肼结构(DOTA = 1,4,7,10-四氮杂环十二烷-1,4,7,10-四乙酸)的新型双核非离子型磁共振对比剂(Gd-DOTA)₂-SBDC的设计、合成及弛豫性能。在0.5 T磁场下,测得其纵向弛豫率为每分子10.6 L·mmol⁻¹·s⁻¹或5.3 L·mmol⁻¹·Gd⁻¹·s⁻¹,高于目前临床使用的单核大环对比剂钆-DOTA。体外磁共振成像研究显示该磁共振对比剂具有提高诊断灵敏度和准确度的特点。同时,该对比剂分子中含有能与髓鞘特异性结合的二苯乙烯结构,具有成为髓鞘靶向磁共振对比剂的应用潜力。另外,探讨了该对比剂的两条合成路线(A和B),总收率分别为70%和75%。综合考虑适用性,合成路线B更优。

关键词 磁共振成像; 对比剂; 合成; 弛豫率; DOTA-酰肼; 钆-DOTA; 髓鞘

Synthesis and Relaxivity of One Macrocyclic Binuclear Nonionic Magnetic Resonance Contrast Agent

Zhou, Jin^{†,a,b} Sun, Hongshun^{†,*,a,b} Li, Yulong^b Jiang, Hong^b
Guo, Cheng^{*,a} Shen, Linjiang^c

(a College of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816)

(b Targeted MRI Contrast Agents Laboratory of Jiangsu Province, Nanjing Polytechnic Institute, Nanjing 210048)

(c School of Physical and Mathematical Sciences, Nanjing Tech University, Nanjing 211816)

Abstract Magnetic resonance imaging (MRI) is widely employed in diagnostic medicine and soft tissue imaging. Contrast agents (CAs) can improve the specificity of enhanced MRI images. Herein, the design, synthesis and relaxivity of a novel binuclear nonionic Gd-based DOTA-hydrazide (DOTA = 1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraacetic acid) derived MRI contrast agents, (Gd-DOTA)₂-SBDC, were reported. The improved longitudinal relaxivity was determined as 10.6 L·mmol⁻¹·s⁻¹ per molecule or 5.3 L·mmol⁻¹·Gd⁻¹·s⁻¹ at 0.5 T, which is higher than that of the mononuclear clinical macrocyclic agent Gd-DOTA. *In vitro* MRI studies confirmed its suitability for use as a MRI contrast agent with improved diagnostic sensitivity and accuracy. Furthermore, the stilbene moiety endows it with myelin-targeting capabilities. In addition, two synthetic routes (A and B) for this contrast agent were compared, providing overall yields of 70% and 75%, respectively, and route B was deemed superior.

Keywords magnetic resonance imaging; contrast agents; synthesis; relaxivity; DOTA-hydrazide; Gd-DOTA; myelin

* Corresponding authors. E-mail: njutshs@126.com; guocheng@njtech.edu.cn

Received February 5, 2021; revised March 12, 2021; published online April 9, 2021.

Project supported by the Natural Science Foundation of Jiangsu Province (No. BK20181486), the Natural Science Foundation of the Jiangsu Higher Education Institutions (No. 17KJB320001), the Overseas Training Program for Excellent Young Teachers and Principals of Jiangsu Province and the Qing Lan Project of Jiangsu Province.

江苏省自然科学基金(No. BK20181486)、江苏省高校自然科学基金(No. 17KJB320001)、江苏省高校优秀中青年骨干教师和校长境外研修计划和江苏省青蓝工程资助项目。

† 共同第一作者(These authors contributed equally to this work).

1 Introduction

Magnetic resonance imaging (MRI) has become one of the most commonly implemented diagnostic techniques as it is nonionizing, noninvasive, and deep penetrating, and it provides high spatial resolution for whole-body diagnosis.^[1-2] To improve specificity and acquire high-intensity MR signals, contrast agents (CAs) are widely used to facilitate the acquisition and interpretation of MRI images. Clinical-used CAs are generally coordination complexes in which a Gd^{III} ion is encapsulated within octadentate chelators derived from polyaminocarboxylate anions, such as DTPA (diethylenetriaminepentaacetic acid) and DOTA (1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraacetic acid). According to a recent report, Gd-based CAs (GBCAs) are employed in approximately 40% of all MRI exams and 60% of neuro-MRI exams.^[3] However, the release of Gd^{3+} ions from GBCAs causes nephrogenic systemic fibrosis (NSF) in patients with impaired renal clearance and gadolinium accumulation in the brains.^[4-7] In recent years, considerable efforts have been directed toward the development of tissue specific MRI CAs with higher proton relaxivities, as this allows for the use of lower doses.

According to the Solomon-Bloembergen-Morgan (SBM) equation, the relaxivity of GBCAs is primarily determined by residence time (τ_{M}), the number of first-shell (inner) water molecules (q), rotational correlation time (τ_{R}), and electron spin relaxation time ($T_{1\text{e}}$). For small molecular

weight GBCAs, enhancing the relaxivity can be achieved by either increasing q or prolonging τ_{R} .^[8-10] As an increase in q is often accompanied by a decrease in thermodynamic and kinetic stability, prolonging τ_{R} is generally practiced. One of the most efficient approaches for achieving this is by constructing multiple-nuclear agents by fusing several small-molecule agents.^[11-13]

Myelin is the axon-protecting sheath surrounding the nerves of vertebrates, and it facilitates impulse conduction. Damage to myelin or demyelination is associated with various neurodegenerative diseases such as multiple sclerosis (MS).^[14] Over the past decades, MRI has played an important role in the diagnosis and disease management of MS. However, due to the lack of myelin-specific MRI CAs, demyelination and inflammation can not be differentiated by conventional MRI. To address this problem, efforts have been devoted to improving the MR specificity of myelin imaging. Luca *et al.*^[15-16] designed and synthesized two types of myelin-specific MRIs, namely CAs Gd-DODAS (Chart 1, **A**) and MIC (Chart 1, **B**), based on a stilbene binding moiety and demonstrated their efficacy in binding specifically to myelin *in vitro* and *in vivo*.

Over the past decade, our group has been extensively exploring magnetic resonance imaging and near infrared imaging. We have designed and synthesized two nonionic binuclear MRI CAs, $(\text{Gd-DOTA})_2\text{-DYMB}$ (Chart 1, **C**) and $(\text{Gd-DOTA})_2\text{-DYMBP}$ (Chart 1, **D**), using diyl-

Chart 1 Structures of some compounds concerned

denemethylbenzene and diylidenemethylbiphenyl, respectively, as the linkers. The longitudinal relaxivities of the two CAs were 11.4 and 11.7 L·mmol⁻¹·s⁻¹ per molecule at 0.5 T, respectively.^[17] In addition, we have reported a series of Gd-based DOTA-hydrazide (DOTAH) derived MRI contrast agents (Chart 1, E), some of which exhibited marginally higher r_1 relaxivities and ideal contrast enhancements in T_1 -weighted MR images compared with the clinical contrast agent Gd-DOTA and could be used as liver-targeting MRICAs. These types of binuclear structures can ameliorate the occurrence of NSF resulting from the use of linear agents.^[18] For myelin imaging, we designed and synthesized a series of Nile Red Analogs (Chart 1, F), bearing structural features resembling stilbene. *In vitro*, *ex vivo* and *in vivo* experiments indicated that they were capable of crossing the blood-brain barrier (BBB) and binding specially to myelin tracts, and were therefore suitable for use as near-infrared fluorescence (NIRF) probes for myelin-targeted imaging.^[19]

Herein, we report a novel dimeric nonionic MRI CAs, (Gd-DOTAH)₂-SBDC (SBDC: 4,4'-stilbenedicarboxylic, Chart 1, G), in which the two DOTAH units that function as Gd³⁺ chelators are affixed to a stilbenedicarbonyl moiety, to investigate its potential application as a myelin-targeting MRI CA. Relaxometric studies and T_1 -weighted MR imaging were performed for this complex.

2 Results and discussion

2.1 Design and synthesis

To improve the specificity of enhanced MR images for MS diagnosis, we aimed to develop a myelin-targeting MRI CA with a high relaxivity. The structure-activity relationship (SAR) studies of a library of compounds suggested that the stilbene framework is crucial for binding to myelin.^[20-24] On the other hand, we decided to construct binuclear CAs to attain a high relaxivity. Accordingly, 4,4'-stilbenedicarboxylic acid was chosen as the starting material for the construction of binuclear MRI CAs that could potentially be utilized for myelin targeting. DOTAH was selected as the Gd³⁺ chelator, because this macrocyclic ligand exhibits superior thermodynamic stability and slower dissociation kinetics, which are beneficial for preventing transmetallation *in vivo*, compared to linear ligands, such as DTPA. In addition, Gd³⁺ binding neutralizes the charges of three acetic groups of DOTAH, resulting in nonionic CAs, beneficial for lowering osmotic pressure. Moreover, complex (Gd-DOTAH)₂-SBDC is more stable than (Gd-DOTAH)₂-DYMB and (Gd-DOTAH)₂-DYMBP due to the susceptibility of the C=N bonds of (Gd-DOTAH)₂-DYMB and (Gd-DOTAH)₂-DYMBP toward hydrolysis under acidic conditions.^[27]

The synthetic procedure for complex (Gd-DOTAH)₂-SBDC is outlined in Scheme 1. Two different routes (A and

Reagents and conditions: (i) SOCl₂, DMF; (ii) CH₂Cl₂, K₂CO₃; (iii) TFA, r.t., 24 h; (iv) GdCl₃, 50 °C, 24 h; (v) TFA, r.t., 24 h; (vi) GdCl₃, 50 °C, 24 h; (vii) DMF, K₂CO₃

Scheme 1 Synthetic routes (A and B) for complex 8

B) for its synthesis are considered. In route A, the first step entailed the formation of an acid chloride from 4,4'-stibenedicarboxylic acid and SOCl_2 , affording 4,4'-stibenedicarbonyl chloride (**2**), which was used directly in the next step without purification. *t*-Bu-DOTA (**3**) was synthesized from cyclen according to our previously published protocol.^[25] Intermediate $(t\text{-Bu-DOTA})_2\text{-SBDC}$ (**6**) was obtained from the coupling of **2** and **3** in 90% yield. Then, **6** was deprotected using trifluoroacetic acid (TFA) to deliver $(\text{DOTA})_2\text{-SBDC}$ (**7**). The target complex $(\text{Gd-DOTA})_2\text{-SBDC}$ (**8**) was obtained by combining **7** and GdCl_3 in aqueous media at 50 °C. The overall yield of route A was 70%. In route B, the DOTA gadolinium complex Gd-DOTA (**5**) was first synthesized, which could then be used for the synthesis of various gadolinium-based contrast agents in one step. Gd-DOTA (**5**) was obtained by combining GdCl_3 and DOTA (**4**), which was generated from the TFA-catalyzed ester hydrolysis of *t*-Bu-DOTA (**3**). Finally, the Gd complex precursor **5** was reacted with 4,4'-stibenedicarbonyl chloride (**2**) to afford the target product $(\text{Gd-DOTA})_2\text{-SBDC}$ (**8**). Complex **8** was purified on a C-18 SPE column. The overall yield of route B was 75%. Thus, route B was higher yielding than route A. Moreover, the use of Gd-DOTA (**5**) as an intermediate is more convenient for the facile preparation of various gadolinium-based contrast agents. Therefore, synthetic route B was deemed more suitable for the preparation of Gd-DOTA-based MRI contrast agents in this study.

2.2 In vitro MRI

To evaluate the level of contrast enhancement of the

developed agent, longitudinal relaxivity r_1 and transverse relaxivity r_2 were calculated from the linear fitting of $1/T_1$ and $1/T_2$ as a function of complex concentrations (Figure 1, A, B)^[28-29]. Commercially available Gd-DOTA, self-prepared $(\text{Gd-DOTA})_2\text{-DYMB}$, and $(\text{Gd-DOTA})_2\text{-DYMBP}$ were used for comparison. The longitudinal relaxivity (r_1) of $(\text{Gd-DOTA})_2\text{-SBDC}$ ($M_r=1377$) was $10.6 \text{ L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$ per molecule or $5.3 \text{ L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$ per Gd at 0.5 T (Table 1), which is higher than that of the current clinical mononuclear macrocyclic CA, Gd-DOTA ($3.6 \text{ L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$, $M_r=563$). The improvement in relaxivity was principally attributed to the increase in molecular weight and the curbed rotational correlation time (τ_R) resulting from increased molecular rigidity. However, the relaxivity was lower than that of our previously reported $(\text{Gd-DOTA})_2\text{-DYMB}$ ($11.4 \text{ L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$, $M_r=1244$) and $(\text{Gd-DOTA})_2\text{-DYMBP}$ ($11.7 \text{ L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$, $M_r=1319$), despite its molecular weight being the highest among them (Table 1). Presumably the presence of the vinyl group in the molecule allows for a higher degree of molecular rotation, which affects its relaxation performance. The r_2/r_1 ratio was calculated as 1.16 (Table 1); the low r_2/r_1 value indicates that the complex can be used as a T_1 -weighted MRI CA. As shown in Figure 1 C and D, **8** presents excellent positive T_1 contrast enhancement akin to $(\text{Gd-DOTA})_2\text{-DYMB}$ and $(\text{Gd-DOTA})_2\text{-DYMBP}$, and the brightness of the MR images increased with the increasing concentrations, showing a clear dose-dependent color change, because the relaxation of the proton increases with increasing concentration. Moreover, as a potential myelin-targeting binuclear

Figure 1 MR relaxivities and T_1 -weighted MR images at 0.5 T
 r_1 and r_2 relaxivity curves of **8** (A, B); T_1 -weighted MR images at varying concentrations in the range of 0.25 to 2.0 mmol/L (C, D).

Table 1 The r_1 and r_2 values and the ratios of r_2/r_1 of the synthesized complex at 0.5 T^a

Entry	Gd-complexes	$r_2/(\text{L} \cdot \text{mmol}^{-1} \cdot \text{s}^{-1})$	$r_1/(\text{L} \cdot \text{mmol}^{-1} \cdot \text{s}^{-1})$	r_2/r_1	Ref.
1	(Gd-DOTA) ₂ -SBDC	12.3	10.6	1.16	This work
2	Gd-DOTA	4.3	3.6	1.19	[26]
3	(Gd-DOTA) ₂ -DYMB	13.2	11.4	1.16	[17]
4	(Gd-DOTA) ₂ -DYMBP	14.0	11.7	1.20	[17]

^a Gd-DOTA, (Gd-DOTA)₂-DYMB and (Gd-DOTA)₂-DYMBP were as compares.

MR probe, (Gd-DOTA)₂-SBDC exhibited a higher relaxivity compared to that of reported MIC and Gd-DODAS ($5.8 \text{ L} \cdot \text{mmol}^{-1} \cdot \text{s}^{-1}$ at 0.5 T for both^[15-16]). Hence, the developed binuclear macrocyclic contrast agent can be used at lower dosages and consequently provide reduced clinical toxicity.

3 Conclusions

In summary, a nonionic binuclear MRI CAs, (Gd-DOTA)₂-SBDC was designed and synthesized via two different synthetic routes (A and B). Route B was found to be more suitable for the preparation of Gd-DOTA-based MRI contrast agents as it provided a higher yield and a more useful intermediate. Meanwhile, stilbenedicarbonyl was selected as the linker for (Gd-DOTA)₂-SBDC, because the stilbene motif is crucial for myelin binding. Thus, the developed complex has considerable potential for use as a myelin-targeting MRI CA. In addition, the feasibility of utilizing it as a T_1 -weighted MRI CA was explored. The prepared agent provided a good positive imaging effect and a low r_2/r_1 ratio, confirming its suitability for use as a T_1 -weighted MRI CA. The longitudinal relaxivity was found to be $10.6 \text{ L} \cdot \text{mmol}^{-1} \cdot \text{s}^{-1}$ per molecule or $5.3 \text{ L} \cdot \text{mmol}^{-1} \cdot \text{Gd}^{-1} \cdot \text{s}^{-1}$ at 0.5 T, which is higher than that of the FDA approved mononuclear macrocyclic agent Gd-DOTA and the reported myelin-targeting probes MIC and Gd-DODAS. Therefore, the developed contrast agent can be used at lower dosages, allowing for reduced clinical toxicity. Moreover, complex (Gd-DOTA)₂-SBDC is more stable than our previously reported (Gd-DOTA)₂-DYMB and (Gd-DOTA)₂-DYMBP, as the C=N bonds of the two latter complexes are susceptible to hydrolysis under acidic conditions. Animal experiments related to myelin imaging are on the way.

4 Experimental section

4.1 General procedures

All chemicals and reagents were used as received without further purification. Unless specially mentioned, all the reactions were performed open to air. Reactions were monitored by thin-layer chromatography (TLC) and visualized by a dual short/long wavelength UV lamp. Flash column chromatography was performed using 200~400 mesh silica gel (Fisher). NMR spectra were recorded on a Bruker AVANCE AV-300 NMR spectrometer at room temperature. Spectra were referenced to internal standards (TMS in CDCl₃ and TMSP in D₂O). Multiplicity and coupling constants (J) were calculated automatically on Mes-

tReNova 10.0, a NMR processing software from Mestrelab Research. MS (ESI) was determined by using a micro-TOF-Q II HRMS/MS instrument (Bruker). Element analysis was determined by using a Perkin-Elmer 240c elemental analysis instrument. *In vitro* MRI testing and T_1 relaxation time measurements were tested at a 0.5 T NMR120-Analyst NMR Analyzing and Imaging system (Niuman Corporation, Shanghai, China). Melting points were determined in open capillaries and were uncorrected.

4.2 Synthesis

4.2.1 Synthesis of DOTA (4)

A solution of **3** (1.17 g, 2 mmol) in TFA (15 mL) was stirred at room temperature for 24 h. TFA was removed under reduced pressure and the residue was dissolved in ethyl acetate (5 mL). Freezing for 12 h, filtered to give compound **4**^[27] as a white solid (quantitative). m.p. 182.6~183.8 °C; ¹H NMR (300 MHz, D₂O) δ : 4.03~2.93 (m, 24H); ¹³C NMR (126 MHz, D₂O) δ : 174.5, 170.5, 56.3, 53.1, 52.5, 51.6, 51.0, 48.7, 48.2. HRMS calcd for C₁₆H₃₀N₆O₇ [M+H]⁺ 419.2249, found 419.2251.

4.2.2 Synthesis of Gd-DOTA (5)

To a solution of **4** (0.83 g, 2 mmol) in water (20 mL) was added a solution of GdCl₃ in water (0.05 mol/L, 40 mL, 2 mmol) while maintaining the pH at 6.5 by addition of 1 mol/L NaOH. The solution was stirred at 50 °C for 24 h. The solution was loaded on a Hypersep C-18 cartridge, and the stationary phase was washed extensively with water. The product was recovered after gradient elution with MeOH/H₂O ($V:V=1:9$ to $3:2$). The fractions containing the product were evaporated under reduced pressure to give **5**^[27] (1.01 g, yield 88%) as a white solid. m.p. > 300 °C; ESI-MS m/z : 596.1 [M+Na]⁺. A correct Gd isotope pattern was observed. Anal. calcd for C₁₆H₂₇GdN₆O₇: C 33.56, H 4.75, N 14.68; found C 33.52, H 4.76, N 14.60.

4.2.3 Synthesis of (*t*-Bu-DOTA)₂-SBDC (6)

To a 250 mL round-bottom flask fitted with a magnetic stir bar were added 4,4'-stilbene dicarboxylic acid (**1**, 1.34 g, 5.0 mmol), freshly distilled sulfoxide chloride (85 mL) and *N,N*-dimethylformamide (DMF) (0.5 mL). The mixture was stirred at 75 °C for 4 h. After completion of the reaction, the unreacted SOCl₂ was recovered at 110 °C, the residual SOCl₂ was evaporated *in vacuo* and taken out by dichloromethane. The resulting residue was recrystallized with ethyl acetate, filtered, and washed with cold ethyl acetate. The product 4,4'-stilbenedicarbonyl chloride (**2**) was got as green solid of 1.40 g, yield 92%. It was used without further purification. Then, to a suspension of compounds **2** (0.31 g,

1 mmol) and *t*-Bu-DOTA (3, 1.17 g, 2 mmol) in CH₂Cl₂ (30 mL) was added K₂CO₃ (0.27 g, 2 mmol) as acid-binding agent. The mixture was stirred at room temperature for 1 h, then refluxed for 2 h (TLC, 10% CH₃OH-CH₂Cl₂). After cooled, the solid was filtered, the filtrate was evaporated under reduced pressure and got beige solid. The crude product was purified over a silica flash column using 5% CH₃OH in CH₂Cl₂ as eluent, which resulted in the pure light yellow solid of **6** (1.26 g, yield, 90%). m.p. 172.4 ~ 174.5 °C; ¹H NMR (300 MHz, CDCl₃) δ: 7.88 (d, *J*=7.9 Hz, 4H), 7.43 (d, *J*=7.8 Hz, 4H), 7.06 (s, 2H), 4.48 (s, 4H), 3.41 (s, 4H), 3.16~2.02 (m, 44H), 1.55~1.02 (m, 54H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.2, 171.4, 165.4, 140.0, 131.40, 129.4, 128.3, 126.4, 81.8, 81.6, 55.7, 55.6, 55.1, 28.0, 27.8; ESI-MS *m/z*: 1427.7 [M+Na]⁺. Anal. calcd for C₇₂H₁₁₆N₁₂O₁₆: C 61.52, H 8.32, N 11.96; found C 61.46, H 8.34, N 11.99.

4.2.4 Synthesis of (DOTA)₂-SBDC (7)

A solution of **6** (0.70 g, 0.5 mmol) in TFA (10 mL) was stirred at room temperature for 24 h. TFA was removed under reduced pressure and the residue was dissolved in ethyl acetate (5 mL). Freezing for 12 h, filtered to give compound **7** as a light yellow solid (quantitative). m.p. 159.6~162.4 °C; ¹H NMR (300 MHz, D₂O) δ: 7.61 (d, *J*=5.8 Hz, 4H), 7.33 (d, *J*=6.4 Hz, 4H), 6.86 (s, 2H), 4.00~2.93 (m, 48H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 171.0, 165.6, 165.1, 143.5, 132.4, 132.2, 130.6, 129.6, 120.9, 117.1, 57.3, 55.9, 53.1, 51.8; ESI-MS *m/z*: 1145.4 [M+2K-H]⁺. Anal. calcd for C₄₈H₆₈N₁₂O₁₆: C 53.92, H 6.41, N 15.72; found C 53.97, H 6.40, N 15.69.

4.2.5 Synthesis of (Gd-DOTA)₂-SBDC (8)

Method A: A solution of **7** (0.27 g, 0.25 mmol) in water (20 mL) was mixed with a solution of GdCl₃ in water (0.05 mol/L, 10 mL, 0.5 mmol) while maintaining the pH at 6.5 by addition of 1 mol/L NaOH. The solution was stirred at 50 °C for 24 h. The pH was raised to 8.0. The solution was filtered through a 0.2 μm filter, and the pH was lowered to 7.0 by adding 1 mol/L HCl. The solution was loaded on a Hypersep C-18 cartridge, and the stationary phase was washed extensively with water. The product was recovered after gradient elution with MeOH/H₂O (*V*: *V*=1:9 to 3:2). The fractions containing the product were evaporated under reduced pressure to give **8** with a light yellow solid, (0.29 g, yield 85%).

Method B: To a suspension of compounds of **2** (0.08 g, 0.25 mmol) and Gd-DOTA (**5**, 0.29 g, 0.5 mmol) in DMF (10 mL) was added K₂CO₃ (0.07 g, 0.5 mmol) as acid-binding agent. The mixture was stirred at room temperature for 4 h. After completion of the reaction, the solid was filtered, the filtrate was evaporated under reduced pressure to obtain beige solid. The crude product was redissolved in water and loaded on a Hypersep C-18 cartridge, and the stationary phase was washed extensively with water. The product was recovered after gradient elution with MeOH/H₂O from 1:9 to 3:2. The fractions containing the product were evaporated under reduced pressure to give **8**

with a light yellow solid (0.32 g, yield 92%). m.p. >300 °C; ESI-MS *m/z*: 1442.3 [M+CH₃CN+Na]⁺. A correct Gd isotope pattern was observed. Anal. calcd for C₄₈H₆₂Gd₂N₁₂O₁₆: C 41.85, H 4.54, N 12.20; found C 41.80, H 4.55, N 12.23.

4.3 In vitro MRI

To obtain *T*₁ relaxation times of the aqueous solution of the new agents at different concentrations, following measurement parameters were used: repetition time (TR)=6000 ms, number of data=25, and number of averages (NA)=2. While the *T*₂ relaxation times were determined with the following parameters: TR=6000 ms, echo time (TE)=1 ms, echo count=6000 and NA=2. Both *r*₁ and *r*₂ relaxivities were calculated by the linear fitting of 1/*T*₁ or 1/*T*₂ as a function of metal concentration. For *T*₁-weighted MR images, the instrument parameters were set as follows; TR=100 ms, TE=18.2 ms, imaging matrix=192×256, slice thickness=5 mm, field of view (FOW)=100 mm×100 mm and NA=2.

Supporting Information ¹H NMR and ¹³C NMR spectra, mass spectra of the synthesized compounds. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Wei, R.; Gong, X.; Lin, H.; Zhang, K.; Li, A.; Liu, K.; Shan, H.; Chen, X.; Gao, J. *Nano Lett.* **2019**, *19*, 5394.
- [2] Mi, P.; Kokuryo, D.; Cabral, H.; Wu, H.; Terada, Y.; Saga, T.; Aoki, I.; Nishiyama, N.; Kataoka, K. *Nat. Nanotechnol.* **2016**, *11*, 724.
- [3] Wahsner, J.; Gale, E. M.; Rodríguez-Rodríguez, A.; Caravan, P. *Chem. Rev.* **2019**, *119*, 957.
- [4] Idée, J.-M.; Port, M.; Robic, C.; Medina, C.; Sabatou, M.; Corot, C. *J. Magn. Reson. Imaging.* **2009**, *30*, 1249.
- [5] Di Gregorio, E.; Gianolio, E.; Stefania, R.; Barutello, G.; Digilio, G.; Aime, S. *Anal. Chem.* **2013**, *85*, 5627.
- [6] Marckmann, P.; Skov, L.; Rossen, K.; Dupont, A.; Damholt, M.-B.; Heaf, J.-G.; Thomsen, H.-S. *J. Am. Soc. Nephrol.* **2006**, *17*, 2359.
- [7] Jost, G.; Lenhard, D.-C.; Sieber, M.-A.; Lohrke, J.; Frenzel, T.; Pietsch, H. *Invest Radiol.* **2016**, *51*, 83.
- [8] Lauffer, R.-B. *Chem. Rev.* **1987**, *87*, 901.
- [9] Caravan, P. *Acc. Chem. Res.* **2009**, *42*, 851.
- [10] Hermann, P.; Kotek, J.; Kubicek, V.; Lukes, I. *Dalton Trans.* **2008**, *23*, 3027.
- [11] Luo, J.; Li, W.-S.; Xu, P.; Zhang, L.-Y.; Chen, Z.-N. *Inorg. Chem.* **2012**, *51*, 9508.
- [12] Zhao, G.; Lu, C.; Li, H.; Xiao, Y.; Zhang, W.; Fang, X.; Wang, P.; Fang, X.; Xu, J.; Yang, W. *Inorg. Chim. Acta* **2013**, *406*, 146.
- [13] Zhao, G.; Li, H.; Lu, C.; Xiao, Y.; Fang, X.; Wang, P.; Fang, X.; Zhao, K.; Li, X.; Yin, S.; Xu, J.; Yang, W. *RSC Adv.* **2012**, *2*, 6404.
- [14] Tiwari, A.-D.; Zhu, J.; You, J.; Eck, B.; Zhu, J.; Wang, X.; Wang, X.; Wang, B.; Silver, J.; Wilson, D.; Wu, C.; Wang, Y. *J. Med. Chem.* **2019**, *62*, 4902.
- [15] Frullano, L.; Zhu, J.; Miller, R.-H.; Wang, Y. *J. Med. Chem.* **2013**, *56*, 1629.
- [16] Frullano, L.; Zhu, J.; Wang, C.; Wu, C.; Miller, R.-H.; Wang, Y. *J. Med. Chem.* **2012**, *55*, 94.
- [17] Sun, H.-S.; Li, Y.-L.; Jiang, H.; Guo, C.; Shen, L.-J. *Chin. J. Org. Chem.* **2018**, *38*, 1779 (in Chinese). (孙宏顺, 李玉龙, 蒋燕, 郭成, 沈临江, 有机化学, **2018**, *38*, 1779.)
- [18] Sun, H.-S.; Zhou, J.; Li, Y.-L.; Jiang, H.; Zhang, Y.; Wang, J.; Guo,

- C.; Shen, L.-J. *Chin. J. Org. Chem.* **2019**, 39, 778 (in Chinese).
(孙宏顺, 周进, 李玉龙, 蒋蕪, 张艳, 王建强, 郭成, 沈临江, 有机化学, **2019**, 39, 778.)
- [19] Sun, H.-S.; Li, Y.-L.; Jiang, H.; Xu, N.; Lu, X.; Chen, Y. *CN 1077595333*, **2017** [*Chem. Abstr.* **2018**, 168, 326713].
- [20] Wang, C.; Wu, C.; Popescu, D.-C.; Zhu, J.; Macklin, W.-B.; Miller, R.-H.; Wang, Y. *J. Neurosci.* **2011**, 31, 2382.
- [21] Wang, C.; Wu, C.; Zhu, J.; Miller, R.-H.; Wang, Y. *J. Med. Chem.* **2011**, 54, 2331.
- [22] Frullano, L.; Wang, C.; Miller, R.-H.; Wang, Y. *J. Am. Chem. Soc.* **2011**, 133, 1611.
- [23] Tiwari, A.-D.; Wu, C.; Zhu, J.; Zhang, S.; Zhu, J.; Wang, W.-R.; Zhang, J.; Tatsuoka, C.; Matthews, P.-M.; Miller, R.-H.; Wang, Y. *J. Med. Chem.* **2016**, 59, 3705.
- [24] Wu, C.; Eck, B.; Zhang, S.; Zhu, J.; Tiwari, A.-D.; Zhang, Y.; Zhu, Y.; Zhang, J.; Wang, B.; Wang, X.; Wang, X.; You, J.; Wang, J.; Guan, Y.; Liu, X.; Yu, X.; Trapp, B.-D.; Miller, R.; Silver, J.; Wilson, D.; Wang, Y. *J. Med. Chem.* **2017**, 60, 987.
- [25] Zhou, J.; Yang, H.-D.; Sun, H.-S.; Li, Y.-L.; Jiang, H. *Chin. J. Synth. Chem.* **2017**, 25, 159 (in Chinese).
(周进, 杨海东, 孙宏顺, 李玉龙, 蒋蕪, 合成化学, **2017**, 25, 159.)
- [26] Aime, S.; Caravan, P. *J. Magn. Reson. Imaging.* **2009**, 30, 1259.
- [27] Frullano, L.; Tejerina, B.; Meade, T. J. *Inorg. Chem.* **2006**, 45, 8489.
- [28] Saini, N.; Varshney, R.; Tiwari, A. K.; Kaul, A.; Allard, M.; Ishar, M. P. S.; Mishra, A. K. *Dalton Trans.* **2013**, 42, 4994.
- [29] Gao, L.; Yu, J.; Liu, Y.; Zhou, J.; Sun, L.; Wang, J.; Zhu, J.; Peng, H.; Lu, W.; Yu, L.; Yan, Z.; Wang, Y. *Theranostics* **2018**, 8, 92.

(Fan, Y.)