铜催化合成 2-亚氨基-1,2-二氢喹啉化合物

祝改革^a 赖晓辉^a 王思雨^a 林彩霞^{*,a} 袁耀锋^{*,a}

(福州大学化学学院 福州 350108)

摘要 报道了室温条件下磺酰叠氮、2-氨基苯甲醛、末端炔烃通过一价铜催化合成 2-亚氨基-1,2-二氢喹啉化合物的方法。该反应条件温和, 具有较好的底物普适性, 当更改底物炔烃的类型时, 可以选择控制不同产物的合成, 包括二氢喹啉和 *N*-磺酰脲类化合物。

关键词 2-亚氨基-1,2-二氢喹啉; 三组分反应; 磺酰叠氮; 炔烃

Synthesis of 2-Imino-1,2-dihydroquinolines via Copper Catalysis

Zhu, Gaige^a Lai, Xiaohui^a Wang, Siyu^a Lin, Caixia^{*,a} Yuan, Yaofeng^{*,a}

(College of Chemistry, Fuzhou University, Fuzhou 350108)

Abstract A mild three-component method for the synthesis of 2-imino-1,2-dihydroquinoline compounds has been successfully developed. In the presence of CuCl, various terminal alkynes, sulfonyl azides and 2-aminobenzaldehydes were coupled. When changing the type of substrate alkyne, controlled synthesis of dihydroquinoline and sulfonyl amidine compounds were achieved.

Keywords 2-imino-1,2-dihydroquinoline; three-component synthesis; sulfonyl azides; alkynes

1 Introduction

1,2-Dihydroquinolines (1,2-DHQs), a class of well-known nitrogen-containing heterocyclic compounds, are widely used as important pharmacophores in many drug molecules.^[1] These compounds have widely range of biological properties, such as antibacterial,^[2] antimalarial,^[3] esophageal squamous cell cancer,^[4] anti-tumor agents,^[5] effective plant growth stimulant,^[6] and antioxidants.^[7] At the same time, 1,2-DHQs have widely used in the preparation of various functional materials, which can also serve as effective antioxidants,^[8] sensors and dyes.^[9] Up to now, significant progress has been made in the synthesis of 1,2-quinolines.^[10] Early in 1880, Skraup^[11] reported the classical reaction to construct quinoline scaffolds. Wang and co-workers^[12] reported a conjugate addition-aldol-dehydration reaction of α,β -unsaturated aldehydes with 2-*N*-protected amino benzaldehydes to afford synthetically useful 1,2-dihydroquinolines in high enantioselectivities. In 2020, Lambert *et al.*^[13] reported the synthesis of 1,2-dihydroquinolines by the hydrazine-catalyzed ring-closing carbonyl-olefin metathesis of *N*-prenylated 2-amino-benzaldehydes. Recently, Hwu *et al.*^[14] disclosed domino re-

actions of arynes reacting with Schiff bases, alkynes to generate 1,2-dihydroquinolines. However, these methods often are plagued by using excessive strong acid, alkali, expensive metal catalysts or tedious procedures.

Much work^[15] showed that the copper-triazole intermediate is formed in the cycloaddition reaction of terminal alkynes and sulfonyl azides catalyzed by copper(I), which is so fragile and easy to produce *N*-ketimine. In this case, ketimine is an intermediate similar to the accumulated diene, which has high activity in the reaction and can be captured by other nucleophiles, such as water,^[16] alcohol^[17] and amine.^[18] It can undergo cycloaddition reaction with many olefin structures to prepare many new compounds.

In 2008, Wang and co-workers^[19] reported the effective synthesis of 2-imino-1,2-dihydroquinoline and 2-imino-thiochromone through copper-catalyzed domino reaction (Scheme 1a), starting from 2-acetyl aniline. However, no report on the use of 2-aminobenzaldehyde was reported. Herein, we established that terminal alkynes, sulfonyl azides and 2-aminobenzaldehyde can be used to generate imino-1,2-dihydroquinoline via a three-component coupling reaction catalyzed by monovalent copper, which is one of the powerful and practical strategy for the construc-

* Corresponding authors. E-mail: yaofeng_yuan@fzu.edu.cn; lincaixia_fzu@fzu.edu.cn

Received February 21, 2021; revised March 23, 2021; published online April 9, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21772023, 22071025).

国家自然科学基金(Nos. 21772023, 22071025)资助项目。

tion of 1,2-dihydroquinoline (Scheme 1b).

a) Wang's work:

b) This work:

Scheme 1 Three-component synthesis of 2-imino-1,2-dihydroquinolines

2 Results and discussion

Inspired by Wang's study, we initially attempted the reaction of *p*-toluenesulfonyl azide, phenylacetylene and 2-aminobenzaldehyde in the presence of Et₃N (1.2 equiv.) in CH₂Cl₂ under air at room temperature using a series of Cu catalysts, as presented in Table 1. To our delight, we isolated the desired product in 85% yield after 12 h (Entry 3). The other bases, such as K₂CO₃, Cs₂CO₃, pyridine, *N,N*-diisopropylethylamine (DIEA) and 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU), were proved to be less effective than triethylamine (TEA) (Entries 5~9). Later on, different solvents were investigated, and CH₂Cl₂ was yet the best one (Entries 10~14). Next, controlling experiments demonstrated that the target product was not obtained in the absence of either catalyst CuCl or base (Entries 15~16). Increasing the amount of TEA had no effect on the results, but reducing the amount of TEA would lead to lower yield (Entries 17~18). With the reducing of the catalyst dose, the reaction activity got lower (Entries 19~20).

With the optimal conditions in hand, the scope and limitation of this protocol were further evaluated (Table 2). The scope of 2-aminobenzaldehydes was investigated firstly. Functional groups at different positions on the phenyl, such as methyl (**4a**, **4d**, **4f**), fluoro (**4b**), chloro (**4c**), methoxy (**4e**), and 3-amino-2-naphthaldehyde (**4g**), were well tolerated in the reaction. Moreover, various sulfonyl azides involved in the reaction were studied. Both alkyl- and aryl-substituted sulfonyl azides could furnish to the corresponding products (**4h**~**4t**) in good yields. It was worth noting that the electronic properties and the position of the substituents on the phenyl group did not have much

Table 1 Reaction condition optimization^a

Entry ^a	Catalyst	Base	Solvent	Yield ^b /%
1	CuI	TEA	DCM	73
2	CuBr	TEA	DCM	78
3	CuCl	TEA	DCM	85
4	CuCl ₂ ·2H ₂ O	TEA	DCM	N.P. ^c
5	CuCl	K ₂ CO ₃	DCM	27
6	CuCl	Cs ₂ CO ₃	DCM	23
7	CuCl	Pyridine	DCM	19
8	CuCl	DIEA	DCM	34
9	CuCl	DBU	DCM	trace
10	CuCl	TEA	CH ₃ CN	73
11	CuCl	TEA	DMF	49
12	CuCl	TEA	THF	33
13	CuCl	TEA	1,4-Dioxane	47
14	CuCl	TEA	DCE	66
15	CuCl	—	DCM	N.P. ^c
16	—	TEA	DCM	N.P. ^c
17	CuCl	TEA (2 equiv.)	DCM	85
18	CuCl	TEA (0.5 equiv.)	DCM	45
19	CuCl (5 mol%)	TEA	DCM	55
20	CuCl (1 mol%)	TEA	DCM	23

^a Reaction conditions: **1a** (0.6 mmol), **2a** (0.5 mmol), **3a** (0.6 mmol), [Cu] (10 mol%), solvent (2 mL), r.t., 12 h, under air atmosphere. ^b Isolated yield. ^c N.P. = No product.

effect on the reaction. Strong electron-withdrawing groups (such as nitro, trifluoromethyl) also showed good reactivity.

In addition, naphthalenesulfonyl azide (**2q**) could also give the desired product **4q** in 85% yield. Alkyl sulfonyl azides (**2r**~**2t**) could also be tolerated. Unfortunately, employing benzylsulfonyl azide did not provide the desired product under standard reaction conditions. A single crystal of compound **4o** (CCDC: 2057810, Figure 1) was also obtained. From the crystal structure it can be clearly confirmed that the product is a dihydroquinoline structure.

The substrate scope of terminal alkynes was also studied (Table 3). A series of aromatic alkynes with electron donating groups and electron withdrawing groups on the aromatic ring, gave moderate to good yields (**4u**~**4ab**). Then, the influence of the position of the substituents on the arene ring was explored, when the substituent was in the *meta*-position of aryl group (**4ab**), the yield of the target product is higher than it was in the *para*-position (**4x**). *o*-Substituent aromatic alkynes were not applicable the reaction. Alternatively, ferrocene acetylene (**3y**) was also a suitable substrate, providing the target product **4y** in a yield of 47%.

Table 2 Substrate scope of amino-aldehydes and azides

Finally, the aliphatic alkynes, such as cyclopropylacetylene (**5a**), 1-hexyne (**5b**), 3,3-dimethyl-1-butyne (**5c**) and 3-methylbutynol (**5d**) and so on, were evaluated. To our surprise, the reaction produced new structure *N*-sulfonyl amidine in good yields rather than 1,2-dihydroquinoline under the optimal conditions (Table 4). This might due to the poor reactivity of their intermediate *N*-sulfonyl amidine.

On the basis of the above results and the previous reports,^[16,18,20] a plausible mechanism for this tandem reaction to construct imino-1,2-dihydroquinoline or *N*-sulfonyl amidine is illustrated in Scheme 2. Under the catalysis of alkali and copper reagents, sulfonyl azide (**2**) reacts with alkyne (**3**) forming intermediate **C** that transforms into intermediate **D** through rearrangement, and intermediate ketenimine

G is obtained from **D** by the release of N₂ through two ways. Then, 2-aminobenzaldehyde, as a nucleophile, reacts with ketenimine **G** leading to intermediate **H**. When R² is a phenyl, the intermediate **H** undergoes intramolecular 1,5-*H* migration and cyclization dehydration leading to product **4**. When R² is an alkyl, the intermediate **H** undergoes intramolecular 1,3-*H* migration leading to product **5**.

3 Conclusions

In conclusion, a novel copper-catalyzed three-component coupling has been developed. A variety of terminal alkynes, sulfonyl azides and 2-aminobenzaldehydes were compatible under the catalytic conditions. This reaction provided

Figure 1 X-ray crystal structure of compound 4o

Table 3 Scope of aromatic alkynes

rapid access to imino-1,2-dihydroquinoline derivatives in high yields. More importantly, the reaction can be controlled to prepare different products including dihydroquinoline and *N*-sulfonyl amidine highly selectively. Moreover, the protocol features a simple experimental procedure, mild reaction conditions and broad functional group compatibility.

4 Experimental section

4.1 General experimental information

All experiments were carried out under air atmosphere. Substituted 2-aminobenzaldehydes were prepared.^[21] The other chemicals were purchased and used without further

purification, unless otherwise noted. Melting points were measured on XT-4 digital micro melting point apparatus and are uncorrected. Proton nuclear magnetic resonance (¹H NMR) spectra were recorded on a Bruker AV-400 MHz or a JEOL JNM-ECZ500R/S1. NMR samples were dissolved in CDCl₃ and chemical shifts are reported in ppm referenced to residual nondeuterated solvent. Carbon nuclear magnetic resonance (¹³C NMR)/fluorine nuclear magnetic resonance (¹⁹F NMR) spectra were recorded on the same Bruker instrument (101 (126) MHz/376 (471) MHz). HRMS were obtained on a Exactive Plus LC-MS (ESI) mass spectrometers.

Table 4 Scope of aliphatic alkynes

Scheme 2 Proposed mechanistic routes

4.2 General procedure for the synthesis of imino-1,2-dihydroquinoline or *N*-sulfonyl amidine

To a 25 mL round-bottom flask were added sulfonyl azide (0.5 mmol), alkyne (0.6 mmol) and 2-aminobenzaldehyde (0.6 mmol), CuCl (10 mol%), TEA (1.2 equiv.) and

DCM (2 mL). Then the mixture was stirred at the room temperature for 12 h in air. The resulting mixture was filtered with celite, the solvent in the acquired solution was removed under reduced pressure. The residue was subjected to flash silica gel column chromatography to

provide pure products with the elution of mixed petroleum ether/ethyl acetate ($V:V=5:1$).

4.3 Spectral data of the products

N-(6-Fluoro-3-phenylquinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4a**): White solid, 151 mg, 78% yield. m.p. 180~181 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.23 (s, 1H), 7.83 (d, $J=8.0$ Hz, 2H), 7.81 (s, 1H), 7.51 (d, $J=5.6$ Hz, 2H), 7.41 (d, $J=7.3$ Hz, 2H), 7.35 (d, $J=7.2$ Hz, 3H), 7.29 (d, $J=8.9$ Hz, 1H), 7.23 (d, $J=7.9$ Hz, 2H), 2.44 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 152.46, 142.42, 140.45, 139.54, 135.78, 134.64, 133.64, 132.98, 132.93, 129.55, 129.29, 128.31, 127.97, 127.69, 126.02, 121.39, 116.29, 21.51, 21.07. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 389.1318; found 389.1318.

N-(6-Fluoro-3-phenylquinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4b**): White solid, 163 mg, 90% yield. m.p. 168~169 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.29 (s, 1H), 7.89~7.76 (m, 3H), 7.54~7.48 (m, 2H), 7.42~7.35 (m, 4H), 7.31 (dd, $J=8.3, 2.6$ Hz, 2H), 7.27~7.22 (m, 2H), 2.39 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 160.15, 158.19, 152.46, 142.6, 140.16, 138.66, 135.31, 134.50, 132.18, 129.54, 129.37, 128.70, 128.08, 126.10, 119.83 (d, $J=24.9$ Hz), 118.28, 112.90 (d, $J=22.4$ Hz), 21.58; ^{19}F NMR (471 MHz, Chloroform-*d*) δ : -116.05. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{17}\text{FN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 393.1068; found 393.1067.

N-(6-Chloro-3-phenylquinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4c**): White solid, 162 mg, 80% yield. m.p. 186~187 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.25 (s, 1H), 7.81 (d, $J=7.9$ Hz, 2H), 7.77 (s, 1H), 7.60 (s, 1H), 7.54~7.48 (m, 3H), 7.41~7.30 (m, 4H), 7.24 (d, $J=7.9$ Hz, 2H), 2.38 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 152.53, 142.79, 140.04, 138.26, 135.27, 134.55, 134.07, 131.56, 130.03, 129.54, 129.41, 128.74, 128.11, 127.26, 126.15, 122.26, 117.93, 21.60. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{ClN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 409.0772; found 409.0772.

N-(6-Fluoro-3-phenylquinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4d**): White solid, 144 mg, 74% yield. m.p. 205~206 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.24 (s, 1H), 7.83 (d, $J=7.9$ Hz, 3H), 7.51 (d, $J=7.7$ Hz, 3H), 7.35 (d, $J=6.9$ Hz, 3H), 7.25~7.19 (m, 3H), 7.17 (d, $J=8.1$ Hz, 1H), 2.50 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 152.74, 142.59, 142.46, 140.48, 139.69, 135.86, 135.76, 132.04, 129.58, 129.34, 128.27, 128.00, 126.45, 126.08, 119.37, 116.36, 77.42, 77.16, 76.91, 22.00, 21.56. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 389.1318; found 389.1318.

N-(7-Methoxy-3-phenylquinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4e**): White solid, 153 mg, 76% yield. m.p. 197~198 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.29 (s, 1H), 7.83 (d, $J=7.9$ Hz, 2H), 7.81 (s, 1H), 7.51 (dd, $J=8.6, 4.4$ Hz, 3H), 7.38~7.30 (m, 3H), 7.23 (d, $J=7.9$ Hz, 2H), 6.93 (d, $J=8.8$ Hz, 1H), 6.82 (s, 1H), 3.90 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 162.52, 152.83, 142.43, 140.54, 139.79, 137.45, 135.93, 129.93, 129.54, 129.33, 128.09, 127.98, 126.03, 115.78, 115.12,

98.44, 55.98, 21.55. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$: 405.1267; found 405.1267.

4-Methyl-*N*-(8-methyl-3-phenylquinolin-2(1*H*)-ylidene)-benzenesulfonamide (**4f**): White solid, 103 mg, 53% yield. m.p. 156~157 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.52 (s, 1H), 7.92~7.83 (m, 3H), 7.55 (dd, $J=7.7, 1.9$ Hz, 2H), 7.49 (d, $J=7.9$ Hz, 1H), 7.44 (d, $J=7.4$ Hz, 1H), 7.37 (d, $J=6.9$ Hz, 3H), 7.24 (d, $J=8.3$ Hz, 3H), 2.61 (s, 3H), 2.38 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 152.48, 142.28, 140.17, 139.95, 135.39, 134.28, 132.37, 132.03, 129.29, 129.10, 128.12, 127.78, 125.97, 125.79, 124.16, 123.88, 120.97, 21.29, 16.17. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 389.1318; found 389.1318.

4-Methyl-*N*-(3-phenylbenzo[*g*]quinolin-2(1*H*)-ylidene)-benzenesulfonamide (**4g**): Yellow solid, 132 mg, 62% yield. m.p. 219~220 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.06 (s, 1H), 8.14 (s, 1H), 7.97 (s, 1H), 7.92 (dd, $J=16.1, 8.3$ Hz, 2H), 7.88~7.84 (m, 2H), 7.78 (s, 1H), 7.58 (ddd, $J=8.3, 6.8, 1.3$ Hz, 1H), 7.56~7.53 (m, 2H), 7.49 (ddd, $J=8.1, 6.8, 1.2$ Hz, 1H), 7.39 (d, $J=2.1$ Hz, 1H), 7.38 (d, $J=1.9$ Hz, 2H), 7.26 (d, $J=8.1$ Hz, 2H), 2.39 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 153.18, 142.82, 140.09, 139.34, 135.81, 134.51, 133.61, 132.74, 130.51, 129.72, 129.46, 128.54, 128.50, 128.31, 128.13, 128.08, 127.36, 126.29, 125.86, 121.06, 112.52, 21.64. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{21}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 425.1318; found 425.1318.

4-Methyl-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4h**): White solid, 159 mg, 85% yield. m.p. 147~149 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.26 (s, 1H), 7.88 (s, 1H), 7.83 (d, $J=6.8$ Hz, 2H), 7.61 (dd, $J=18.2, 7.8$ Hz, 2H), 7.55~7.48 (m, 2H), 7.45~7.32 (m, 5H), 7.25 (d, $J=8.8$ Hz, 2H), 2.38 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.6, 142.6, 140.3, 139.7, 135.6, 133.0, 131.3, 129.5, 129.3, 128.4, 128.2, 128.0, 126.1, 124.8, 121.3, 116.4, 21.5. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{19}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 375.1162; found 375.1157.

N-(3-Phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4i**): White solid, 144 mg, 80% yield. m.p. 183~185 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.24 (s, 1H), 7.94 (d, $J=6.8$ Hz, 2H), 7.85 (s, 1H), 7.63~7.53 (m, 2H), 7.51~7.47 (m, 2H), 7.46~7.39 (m, 3H), 7.38~7.29 (m, 5H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.8, 143.1, 139.8, 135.6, 135.4, 133.0, 132.0, 131.4, 129.5, 128.7, 128.4, 128.2, 128.0, 126.0, 124.8, 121.4, 116.5. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 361.1005; found 361.1001.

4-Methoxy-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4j**): White solid, 148 mg, 76% yield. m.p. 214~216 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.24 (s, 1H), 7.95~7.84 (m, 3H), 7.61 (dd, $J=18.4, 8.0$ Hz, 2H), 7.55~7.49 (m, 2H), 7.46~7.32 (m, 5H), 6.92 (d, $J=8.8$ Hz, 2H), 3.83 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 162.4, 139.4, 135.7, 131.2, 129.5, 128.4, 128.1, 124.6, 113.8, 55.5. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{19}\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$: 391.1111; found 391.1105.

4-Bromo-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4k**): White solid, 175 mg, 80% yield. m.p. 200~202 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.19 (s, 1H),

7.89 (s, 1H), 7.77 (d, $J=8.0$ Hz, 2H), 7.65~7.56 (m, 2H), 7.53 (d, $J=8.8$ Hz, 2H), 7.51~7.46 (m, 2H), 7.43~7.31 (m, 5H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.7, 142.2, 140.1, 135.5, 135.4, 133.0, 131.9, 131.5, 128.5, 128.2, 128.1, 127.7, 126.6, 125.0, 121.4, 116.6. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{16}\text{BrN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 439.0110; found 439.0109.

4-Nitro-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4l**): Yellow solid, 111 mg, 55% yield. m.p. 155~157 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.22 (s, 1H), 8.41~8.19 (m, 2H), 8.17~7.93 (m, 3H), 7.78~7.60 (m, 2H), 7.59~7.31 (m, 7H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.9, 149.5, 148.7, 140.5, 135.3, 131.8, 129.4, 128.7, 128.3, 128.1, 127.3, 125.3, 124.0, 121.7, 116.7. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{16}\text{N}_3\text{O}_4\text{S}$ $[\text{M}+\text{H}]^+$: 406.0856; found 406.0852.

2-Methyl-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4m**): Slight yellow solid, 144 mg, 77% yield. m.p. 171~173 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.27 (s, 1H), 8.06 (d, $J=7.6$ Hz, 1H), 7.82 (s, 1H), 7.56 (dd, $J=18.8, 7.6$ Hz, 2H), 7.51~7.46 (m, 2H), 7.38~7.30 (m, 6H), 7.21 (t, $J=8.2$ Hz, 2H), 2.56 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.6, 141.0, 139.6, 137.6, 135.6, 132.2, 132.0, 131.3, 129.6, 128.4, 128.2, 128.0, 127.2, 125.6, 124.7, 121.3, 116.5, 20.4. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{19}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 375.1162; found 375.1153.

2-Fluoro-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4n**): White solid, 162 mg, 86% yield. m.p. 169~171 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.47 (s, 1H), 7.95 (t, $J=7.2$ Hz, 1H), 7.89 (s, 1H), 7.64 (d, $J=8.0$ Hz, 1H), 7.57 (t, $J=7.8$ Hz, 1H), 7.53~7.47 (m, 2H), 7.45~7.33 (m, 3H), 7.31~7.24 (m, 3H), 7.15 (t, $J=7.6$ Hz, 1H), 7.10~7.03 (m, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 159.4 (d, $J=254.8$ Hz), 153.2, 140.3, 135.5, 135.3, 134.1, 132.8, 131.5, 130.8 (d, $J=14.1$ Hz), 129.5, 129.0, 128.3 (d, $J=10.2$ Hz), 128.0, 125.0, 124.1 (d, $J=3.6$ Hz), 121.5, 116.8, 116.6. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{16}\text{FN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 379.0911; found 379.0908.

N-(3-Phenylquinolin-2(1*H*)-ylidene)-2-(trifluoromethyl)benzenesulfonamide (**4o**): White solid, 182 mg, 85% yield. m.p. 190~192 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.30 (s, 1H), 8.31 (d, $J=5.6$ Hz, 1H), 7.91 (s, 1H), 7.74 (d, $J=7.2$ Hz, 1H), 7.65 (d, $J=8.0$ Hz, 1H), 7.62~7.52 (m, 3H), 7.51~7.45 (m, 2H), 7.41 (d, $J=8.0$ Hz, 1H), 7.36 (t, $J=7.6$ Hz, 1H), 7.33~7.26 (m, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.9, 141.4, 140.1, 135.5, 132.9, 132.1, 132.0, 131.5, 130.51, 129.4, 128.4, 128.2, 128.0, 127.9 (q, $J=6.3$ Hz), 127.1, 127.0 (q, $J=33.2$ Hz), 125.0, 122.9 (q, $J=275.1$ Hz), 121.5, 116.6. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{16}\text{F}_3\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 429.0879; found 429.0878.

3-Chloro-*N*-(3-phenylquinolin-2(1*H*)-ylidene)benzenesulfonamide (**4p**): White solid, 157 mg, 82% yield. m.p. 162~164 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.18 (s, 1H), 7.96~7.88 (m, 2H), 7.80 (d, $J=7.6$ Hz, 1H), 7.67~7.56 (m, 2H), 7.51~7.46 (m, 2H), 7.45~7.32 (m, 7H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.8, 144.8, 140.1, 135.4, 134.7, 133.0, 132.0, 131.6, 130.1, 129.5, 128.5, 128.3, 128.1, 126.3, 125.1, 124.1, 121.5, 116.6. HRMS (ESI) calcd

for $\text{C}_{21}\text{H}_{16}\text{ClN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 395.0616; found 395.0614.

N-(3-Phenylquinolin-2(1*H*)-ylidene)naphthalene-2-sulfonamide (**4q**): White solid, 174 mg, 85% yield. m.p. 190~192 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.22 (s, 1H), 8.49 (s, 1H), 7.93 (dd, $J=8.8, 0.8$ Hz, 1H), 7.89~7.80 (m, 4H), 7.59~7.47 (m, 6H), 7.39 (d, $J=8.0$ Hz, 1H), 7.36~7.27 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.5, 139.9, 139.8, 135.8, 135.6, 134.5, 132.9, 132.1, 131.4, 129.5, 129.2, 129.0, 128.4, 128.3, 128.2, 128.1, 127.8, 127.2, 126.6, 124.8, 122.3, 121.5, 116.8. HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{19}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 411.1162; found 411.1158.

N-(3-Phenylquinolin-2(1*H*)-ylidene)methanesulfonamide (**4r**): White solid, 94 mg, 63% yield. m.p. 182~184 °C; ^1H NMR (400 MHz, CDCl_3) δ : 11.98 (s, 1H), 7.89 (s, 1H), 7.68~7.53 (m, 4H), 7.47~7.38 (m, 3H), 7.34 (t, $J=6.4$ Hz, 2H), 3.05 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.9, 139.7, 135.7, 135.5, 132.8, 131.3, 129.5, 128.5, 128.1, 124.7, 121.2, 116.4, 42.9. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 299.0849; found 299.0843.

N-(3-Phenylquinolin-2(1*H*)-ylidene)propane-2-sulfonamide (**4s**): White solid, 168 mg, 86% yield. m.p. 145~146 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (s, 1H), 7.86 (s, 1H), 7.62 (d, $J=7.8$ Hz, 1H), 7.59 (d, $J=6.4$ Hz, 2H), 7.57~7.52 (m, 1H), 7.37 (dd, $J=10.3, 7.1$ Hz, 3H), 7.33~7.27 (m, 2H), 3.19 (p, $J=6.8$ Hz, 1H), 1.36 (d, $J=6.8$ Hz, 6H); ^{13}C NMR (126 MHz, Chloroform-*d*) δ : 153.81, 139.29, 135.80, 135.58, 132.99, 131.21, 129.50, 128.33, 128.14, 127.90, 124.47, 121.08, 116.27, 54.69, 16.50. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{19}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 327.1162; found 327.1158.

N-(3-Phenylquinolin-2(1*H*)-ylidene)cyclopropanesulfonamide (**4t**): White solid, 133 mg, 82% yield. m.p. 161~162 °C; ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (s, 1H), 7.87 (s, 1H), 7.62 (d, $J=7.5$ Hz, 3H), 7.55 (t, $J=7.5$ Hz, 1H), 7.39 (d, $J=7.7$ Hz, 3H), 7.31 (t, $J=8.5$ Hz, 2H), 2.60 (s, 1H), 1.17 (s, 2H), 0.93 (d, $J=8.0$ Hz, 2H); ^{13}C NMR (126 MHz, Chloroform-*d*) δ : 152.98, 139.52, 135.74, 135.52, 132.87, 131.26, 129.49, 128.38, 128.16, 127.98, 124.55, 121.16, 116.30, 32.34, 5.29. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 325.1005; found 325.1002.

4-Methyl-*N*-(3-(*p*-tolyl)quinolin-2(1*H*)-ylidene)benzenesulfonamide (**4u**): White solid, 157 mg, 81% yield. m.p. 220~222 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.26 (s, 1H), 7.90~7.79 (m, 3H), 7.59 (dd, $J=19.4, 7.8$ Hz, 2H), 7.46~7.30 (m, 4H), 7.24 (d, $J=8.4$ Hz, 2H), 7.17 (d, $J=7.2$ Hz, 2H), 2.38 (s, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.8, 142.6, 140.2, 139.2, 138.3, 135.4, 133.0, 132.7, 131.1, 129.4, 129.3, 128.8, 128.1, 126.1, 124.7, 121.5, 116.4, 21.5, 21.3. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 389.1318; found 389.1314.

N-(3-(4-Methoxyphenyl)quinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4v**): White solid, 84 mg, 42% yield. m.p. 181~183 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.24 (s, 1H), 7.90~7.80 (m, 3H), 7.64~7.54 (m, 2H), 7.48 (d, $J=8.4$ Hz, 2H), 7.39 (d, $J=7.6$ Hz, 1H), 7.33 (t, $J=7.4$ Hz, 1H), 7.24 (d, $J=8.0$ Hz, 2H), 6.89 (d, $J=8.4$ Hz, 2H), 3.82 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (101 MHz,

CDCl_3) δ : 159.8, 142.6, 138.8, 131.0, 130.8, 129.3, 128.0, 127.9, 126.1, 124.7, 113.5, 55.3, 21.5. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{N}_2\text{O}_3\text{S}$ $[\text{M} + \text{H}]^+$: 405.1267; found 405.1262.

N-(3-(4-Bromophenyl)quinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4w**): White solid, 160 mg, 71% yield. m.p. 233~235 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.25 (s, 1H), 7.86 (s, 1H), 7.82 (d, $J=8.0$ Hz, 2H), 7.66~7.58 (m, 2H), 7.49 (d, $J=8.4$ Hz, 2H), 7.43~7.33 (m, 4H), 7.27~7.23 (m, 2H), 2.39 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 142.7, 140.0, 139.5, 134.5, 131.6, 131.2, 131.1, 130.2, 129.4, 128.7, 128.2, 126.1, 124.8, 122.7, 121.2, 116.6, 21.5. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{BrN}_2\text{O}_2\text{S}$ $[\text{M} + \text{H}]^+$: 453.0267; found 453.0264.

N-(3-(4-Ferrocenylphenyl)naphthalen-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4x**): White solid, 129 mg, 66% yield. m.p. 217~219 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.24 (s, 1H), 7.85 (s, 1H), 7.82 (d, $J=7.6$ Hz, 2H), 7.66~7.56 (m, 2H), 7.52~7.45 (m, 2H), 7.42~7.31 (m, 2H), 7.24 (d, $J=7.6$ Hz, 2H), 7.03 (t, $J=8.0$ Hz, 2H), 2.37 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 162.8 (d, $J=249.1$ Hz), 152.5, 142.7, 140.1, 139.6, 135.5, 132.0, 131.6 (d, $J=3.4$ Hz), 131.5, 131.3 (d, $J=8.3$ Hz), 129.4, 128.2, 126.0, 124.8, 121.2, 116.5, 115.0 (d, $J=21.6$ Hz), 21.5. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{FeN}_2\text{O}_2\text{S}$ $[\text{M}]^+$: 392.0989; found 392.0971.

N-(3-(4-Fluorophenyl)naphthalen-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4y**): Orange solid, 113 mg, 47% yield. 197 °C decompose; ^1H NMR (400 MHz, CDCl_3) δ : 12.34 (s, 1H), 8.04~7.92 (m, 3H), 7.58 (dd, $J=19.0$, 7.8 Hz, 2H), 7.38 (d, $J=8.0$ Hz, 1H), 7.35~7.29 (m, 3H), 4.96 (s, 2H), 4.37 (s, 2H), 4.00 (s, 5H), 2.39 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.8, 142.8, 140.4, 135.4, 134.5, 131.4, 130.3, 129.5, 127.3, 126.1, 124.6, 121.4, 116.4, 80.0, 69.8, 69.7, 68.8, 21.6. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{23}\text{FeN}_2\text{O}_2\text{S}$ $[\text{M} + \text{H}]^+$: 483.0824; found 483.0818.

4-Methyl-*N*-(3-(thiophen-3-yl)quinolin-2(1*H*)-ylidene)-benzenesulfonamide (**4z**): Slight yellow solid, 152 mg, 80% yield. m.p. 187~189 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.30 (s, 1H), 8.05 (s, 1H), 7.97 (s, 1H), 7.88 (d, $J=7.6$ Hz, 2H), 7.63 (d, $J=8.0$ Hz, 1H), 7.57 (t, $J=7.8$ Hz, 1H), 7.43 (d, $J=4.8$ Hz, 1H), 7.38~7.31 (m, 2H), 7.30~7.22 (m, 3H), 2.38 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.1, 142.7, 140.2, 137.8, 135.1, 131.2, 129.4, 128.1, 127.9, 127.1, 126.2, 126.1, 124.8, 121.2, 116.4, 21.5. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{17}\text{N}_2\text{O}_2\text{S}_2$ $[\text{M} + \text{H}]^+$: 381.0726; found 381.0722.

4-Methyl-*N*-(3-(4-nitrophenyl)quinolin-2(1*H*)-ylidene)-benzenesulfonamide (**4aa**): Yellow solid, 180 mg, 86% yield. m.p. >250 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.30 (s, 1H), 8.22 (d, $J=8.8$ Hz, 2H), 7.96 (s, 1H), 7.81 (d, $J=8.0$ Hz, 2H), 7.75~7.64 (m, 4H), 7.49~7.36 (m, 2H), 7.28 (d, $J=8.4$ Hz, 2H), 2.41 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.9, 147.5, 143.0, 142.2, 140.6, 139.7, 136.0, 132.3, 130.8, 130.5, 129.5, 128.6, 126.0, 125.1, 123.2, 120.9, 116.7, 21.6. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{N}_3\text{O}_4\text{S}$ $[\text{M} + \text{H}]^+$: 420.1013; found 420.1010.

N-(3-(3-Fluorophenyl)quinolin-2(1*H*)-ylidene)-4-methylbenzenesulfonamide (**4ab**): White solid, 149 mg, 76%

yield. m.p. 184~186 °C; ^1H NMR (400 MHz, CDCl_3) δ : 12.25 (s, 1H), 7.88 (s, 1H), 7.84 (d, $J=7.6$ Hz, 2H), 7.61 (dd, $J=18.0$, 8.0 Hz, 2H), 7.46~7.30 (m, 3H), 7.27~7.13 (m, 4H), 7.03 (d, $J=8.0$ Hz, 1H), 2.38 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 163.3 (d, $J=245.8$ Hz), 152.2, 142.7, 140.0, 137.6 (d, $J=8.4$ Hz), 135.6, 131.7, 129.5 (d, $J=8.0$ Hz), 129.4, 128.3, 126.0, 125.2, 124.9, 121.1, 116.8, 116.6, 115.3 (d, $J=21.1$ Hz), 21.5. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{FN}_2\text{O}_2\text{S}$ $[\text{M} + \text{H}]^+$: 393.1068; found 393.1063.

2-Cyclopropyl-*N*-(2-formylphenyl)-*N'*-tosylacetimidamide (**5a**): Colorless oil, 128 mg, 72% yield. ^1H NMR (400 MHz, CDCl_3) δ : 11.61 (s, 1H), 9.91 (s, 1H), 8.68 (d, $J=8.4$ Hz, 1H), 7.87 (d, $J=7.6$ Hz, 2H), 7.71 (d, $J=7.6$ Hz, 1H), 7.52 (t, $J=8.0$ Hz, 1H), 7.34~7.24 (m, 3H), 3.10 (d, $J=7.2$ Hz, 2H), 2.42 (s, 3H), 1.27~1.16 (m, 1H), 0.84 (d, $J=7.6$ Hz, 2H), 0.40 (d, $J=4.8$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.4, 166.5, 142.7, 140.2, 139.8, 136.4, 135.9, 129.4, 126.4, 124.2, 123.1, 121.5, 39.9, 21.6, 6.7, 5.6. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{21}\text{N}_2\text{O}_3\text{S}$ $[\text{M} + \text{H}]^+$: 357.1267; found 357.1264.

N-(2-Formylphenyl)-*N'*-tosylhexanimidamide (**5b**): Colorless oil, 140 mg, 73% yield. ^1H NMR (400 MHz, CDCl_3) δ : 11.28 (s, 1H), 9.91 (s, 1H), 8.63 (d, $J=8.4$ Hz, 1H), 7.88 (d, $J=7.6$ Hz, 2H), 7.70 (d, $J=7.2$ Hz, 1H), 7.53 (t, $J=8.0$ Hz, 1H), 7.34~7.23 (m, 3H), 3.05 (t, $J=7.8$ Hz, 2H), 2.42 (s, 3H), 1.87 (q, $J=7.4$ Hz, 2H), 1.49~1.36 (m, 4H), 0.92 (t, $J=6.6$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.8, 167.0, 142.7, 140.2, 140.1, 136.4, 136.1, 129.4, 126.5, 124.0, 122.6, 121.3, 35.8, 31.4, 27.0, 22.2, 21.6, 13.9. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{25}\text{N}_2\text{O}_3\text{S}$ $[\text{M} + \text{H}]^+$: 373.1580; found 373.1594.

N-(2-Formylphenyl)-3,3-dimethyl-*N'*-tosylbutanimidamide (**5c**): Colorless oil, 84 mg, 45% yield. ^1H NMR (400 MHz, CDCl_3) δ : 11.08 (s, 1H), 9.90 (s, 1H), 8.66 (d, $J=8.4$ Hz, 1H), 7.87 (d, $J=8.0$ Hz, 2H), 7.68 (d, $J=6.8$ Hz, 1H), 7.52 (t, $J=8.0$ Hz, 1H), 7.32~7.24 (m, 3H), 3.07 (s, 2H), 2.41 (s, 3H), 1.21 (s, 9H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.6, 164.2, 142.5, 140.6, 140.0, 136.4, 136.1, 129.3, 126.5, 123.8, 122.5, 121.2, 47.4, 32.4, 30.5, 21.5. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{25}\text{N}_2\text{O}_3\text{S}$ $[\text{M} + \text{H}]^+$: 373.1580; found 373.1990.

N-(2-Formylphenyl)-3-hydroxy-3-methyl-*N'*-tosylbutanimidamide (**5d**): Colorless oil, 140 mg, 75% yield. ^1H NMR (400 MHz, CDCl_3) δ : 11.35 (s, 1H), 9.90 (s, 1H), 8.57 (d, $J=7.2$ Hz, 1H), 7.87 (d, $J=8.0$ Hz, 2H), 7.68 (d, $J=7.2$ Hz, 1H), 7.50 (d, $J=7.8$ Hz, 1H), 7.31 (d, $J=8.0$ Hz, 2H), 7.28~7.23 (m, 1H), 3.29 (s, 3H), 2.43 (s, 2H), 1.47 (s, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.1, 163.6, 142.8, 140.0, 139.4, 136.2, 135.8, 129.4, 126.4, 124.2, 123.2, 121.8, 70.3, 47.8, 30.3, 21.6. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{23}\text{N}_2\text{O}_4\text{S}$ $[\text{M} + \text{H}]^+$: 375.1373; found 375.1164.

N-(2-Formylphenyl)-3-phenyl-*N'*-tosylpropanimidamide (**5e**): White solid, 114 mg, 56% yield. m.p. 133~134 °C; ^1H NMR (500 MHz, CDCl_3) δ : 11.20 (s, 1H), 9.84 (s, 1H), 8.60 (s, 1H), 7.91 (d, $J=8.3$ Hz, 2H), 7.66 (dd, $J=7.6$, 1.7 Hz, 1H), 7.55~7.50 (m, 1H), 7.36~7.30 (m, 4H), 7.29 (d, $J=7.3$ Hz, 2H), 7.28~7.22 (m, 2H), 7.20 (t, $J=7.1$ Hz,

1H), 3.38~3.29 (m, 2H), 3.23 (dd, $J=9.8, 6.1$ Hz, 2H), 2.43 (s, 3H); ^{13}C NMR (126 MHz, Chloroform- d) δ : 195.60, 165.66, 142.83, 140.23, 140.01, 139.43, 136.30, 136.09, 129.48, 128.73, 128.64, 126.66, 126.57, 124.12, 122.63, 121.30, 38.31, 33.74, 21.60. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$: 407.1424; Found: 407.1429.

N-(2-Formylphenyl)-3-methyl-*N'*-tosylbut-2-enimidamide (**5f**): White solid, 79 mg, 44% yield. m.p. 157~158 °C; ^1H NMR (500 MHz, CDCl_3) δ : 11.14 (s, 1H), 9.89 (s, 1H), 8.95~8.75 (m, 1H), 7.82 (d, $J=7.8$ Hz, 2H), 7.70 (d, $J=9.4$ Hz, 1H), 7.61 (t, $J=8.0$ Hz, 1H), 7.34~7.22 (m, 4H), 6.35 (s, 1H), 2.41 (s, 3H), 1.88 (s, 3H), 1.66 (s, 3H); ^{13}C NMR (126 MHz, Chloroform- d) δ : 195.58, 162.05, 145.83, 142.64, 139.98, 139.86, 136.44, 136.34, 129.20, 127.09, 124.25, 122.80, 121.73, 119.72, 26.05, 21.61, 20.70. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{21}\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$: 355.1267; found 355.1270.

N-(2-Formylphenyl)-2-(1-hydroxycyclohexyl)-*N'*-tosylacetimidamide (**5g**): White solid, 119 mg, 58% yield. m.p. 126~127 °C; ^1H NMR (500 MHz, CDCl_3) δ : 11.30 (s, 1H), 9.87 (s, 1H), 8.52 (d, $J=8.5$ Hz, 1H), 7.84 (d, $J=8.0$ Hz, 2H), 7.65 (d, $J=6.1$ Hz, 1H), 7.50~7.44 (m, 1H), 7.28 (d, $J=8.0$ Hz, 2H), 7.22 (t, $J=7.5$ Hz, 1H), 3.29 (s, 1H), 3.22 (s, 2H), 2.40 (s, 3H), 1.82~1.72 (m, 2H), 1.64 (d, $J=9.1$ Hz, 4H), 1.56 (s, 3H), 1.34~1.23 (m, 1H); ^{13}C NMR (126 MHz, Chloroform- d) δ : 195.13, 163.65, 142.84, 140.10, 139.42, 136.20, 135.80, 129.64, 129.43, 126.47, 124.19, 123.30, 121.86, 77.41, 77.16, 76.91, 71.29, 47.69, 38.08, 25.37, 21.92, 21.56. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{27}\text{N}_2\text{O}_4\text{S}$ $[\text{M}+\text{H}]^+$: 415.1686; found 415.1690.

5-Chloro-*N*-(2-formylphenyl)-*N'*-tosylpentanimidamide (**5h**): Colorless oil, 89 mg, 46% yield. ^1H NMR (500 MHz, CDCl_3) δ : 11.30 (s, 1H), 9.91 (s, 1H), 8.60 (d, $J=8.5$ Hz, 1H), 7.87 (d, $J=8.3$ Hz, 2H), 7.70 (dd, $J=7.6, 1.7$ Hz, 1H), 7.53 (ddd, $J=8.8, 7.4, 1.7$ Hz, 1H), 7.30 (d, $J=7.7$ Hz, 2H), 7.26 (td, $J=7.5, 1.0$ Hz, 2H), 3.60 (t, $J=6.5$ Hz, 2H), 3.10 (t, $J=7.7$ Hz, 2H), 2.42 (s, 3H), 2.11~2.01 (m, 2H), 1.99~1.92 (m, 2H); ^{13}C NMR (126 MHz, Chloroform- d) δ : 195.96, 166.11, 142.92, 140.17, 136.45, 136.25, 129.52, 126.58, 124.19, 122.72, 121.41, 44.26, 35.14, 32.01, 24.79, 21.64. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{22}\text{ClN}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$: 393.1034; found 393.1029.

2-(Cyclohex-1-en-1-yl)-*N*-(2-formylphenyl)-*N'*-tosylacetimidamide (**5i**): White solid, 136 mg, 69% yield. m.p. 101~102 °C; ^1H NMR (500 MHz, CDCl_3) δ : 11.17 (s, 1H), 9.86 (s, 1H), 8.71 (d, $J=8.5$ Hz, 1H), 7.87 (d, $J=8.2$ Hz, 2H), 7.66 (d, $J=7.6$ Hz, 1H), 7.50 (t, $J=7.1$ Hz, 1H), 7.29 (d, $J=8.0$ Hz, 2H), 7.24 (t, $J=7.4$ Hz, 1H), 5.86 (s, 1H), 3.76 (s, 2H), 2.41 (s, 3H), 2.20 (s, 2H), 1.95 (s, 2H), 1.66~1.60 (m, 4H); ^{13}C NMR (126 MHz, Chloroform- d) δ : 194.92, 165.81, 142.70, 140.31, 139.77, 136.26, 135.82, 131.32, 130.01, 129.40, 126.59, 124.07, 123.21, 121.62, 43.75, 28.31, 25.61, 22.48, 21.69, 21.58. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{25}\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$: 397.1580; found 397.1579.

Supporting Information Crystal data of **4o** and NMR spectra of **4a**~**5i** are available free of charge via the In-

ternet at <http://sioc-journal.cn/>.

References

- (a) Katritzky, A. R.; Rachwal, S.; Rachwal, B. *Tetrahedron* **1996**, 52, 15031.
(b) Kariba, R. M.; Houghton, P. J.; Yenesew, A. *J. Nat. Prod.* **2002**, 65, 566.
(c) Khadtare, N.; Stephani, R.; Korlipara, V. *Bioorg. Med. Chem. Lett.* **2017**, 27, 2281.
(d) Rabbani, S. M. I. M.; Vahabpour, R.; Hajimahdi, Z.; Zarghi, A. *Iran. J. Pharm. Res.* **2019**, 18, 1790.
(e) Zhang, H. S.; Harmon, M.; Radoshitzky, S. R.; Soloveva, V.; Kane, C. D.; Duplantier, A. J.; Ogungbe, I. V. *ACS Med. Chem. Lett.* **2020**, 11, 2139.
(f) Yang, Y.; Guo, J.; Liu, Z. Z. *Chin. J. Org. Chem.* **2019**, 39, 1913 (in Chinese).
(杨扬, 郭举, 刘站柱, 有机化学, **2019**, 39, 1913.)
(g) Gul, N. S.; Khan, T. M.; Liu, Y. C.; Choudhary, M. I.; Chen, Z. F.; Hong, L. *CCS Chem.* **2020**, 2, 1626.
- (a) Laurentiz, R. S.; Gomes, W. P.; Pissurno, A. P. R.; Santos, F. A.; Santos, V. C. O.; Martins, C. H. G. *Med. Chem. Res.* **2018**, 27, 1074.
(b) Ushiyama, F.; Amada, H.; Takeuchi, T.; Tanaka-Yamamoto, N.; Kanazawa, H.; Nakano, K.; Mima, M.; Masuko, A.; Takata, I.; Hitaka, K.; Iwamoto, K.; Sugiyama, H.; Ohtake, N. *ACS Omega* **2020**, 5, 10145.
(c) Chen, W.; Zuo, H. L.; Li, Y. X.; Liu, J.; Zhou, X. L. *Chin. J. Org. Chem.* **2019**, 39, 2317 (in Chinese).
(陈伟, 左怀龙, 李玉新, 刘江, 周先礼, 有机化学, **2019**, 39, 2317.)
(d) Yang, R.; Ma, Y. N.; Huang, T.; Xie, W.; Zhang, X.; Huang, G. S.; Liu, X. D. *Chin. J. Org. Chem.* **2018**, 38, 2143 (in Chinese).
(杨锐, 马艳妮, 黄婷, 解伟, 张霞, 黄国双, 刘小东, 有机化学, **2018**, 38, 2143.)
- Theerlalanon, C.; Arisawa, M.; Nishida, A.; Nakagawa, M. *Tetrahedron* **2004**, 60, 3017.
- (a) Shahin, M. I.; Roy, J.; Hanafi, M.; Wang, D. Y.; Luesakul, U.; Chai, Y. F.; Muangsin, N.; Lasheen, D. S.; Abou El Ella, D. A.; Abouzid, K. A.; Neamati, N. *Eur. J. Med. Chem.* **2018**, 155, 516.
(b) Tran, S. T. P.; Hipolito, C. J.; Suzuki, H.; Xie, R.; Kim Tuyen, H. D.; Dijke, P. T.; Terasaka, N.; Goto, Y.; Suga, H.; Kato, M. *Biochem. Biophys. Res. Commun.* **2019**, 516, 445.
- (a) Wu, B. W.; Cui, X. X.; Zhu, T.; Wang, S. H.; Lu, C. F.; Wang, J. J.; Dang, H. X.; Zhang, S. Y.; Ding, L. N.; Jin, C. Y. *Chin. J. Org. Chem.* **2020**, 40, 978 (in Chinese).
(吴博文, 崔鑫鑫, 朱挺, 王胜辉, 陆超凡, 王金杰, 党贺祥, 张赛扬, 丁丽娜, 金成允, 有机化学, **2020**, 40, 978.)
(b) Gu, Y. Y.; Lv, X. Q.; Ma, X. D.; Zhang, H. J.; Ji, Y. Y.; Ding, W. J.; Sheng, L. *Chin. J. Org. Chem.* **2020**, 40, 95 (in Chinese).
(顾依钰, 吕晓庆, 马晓东, 张浩健, 嵇媛媛, 丁婉婧, 沈立, 有机化学, **2020**, 40, 95.)
(c) Hu, Y.; Li, Z. Y.; Ding, Y. J.; Li, Z. Y.; Liu, Z. Y.; Shen, Y. M. *Chin. J. Org. Chem.* **2019**, 39, 3230 (in Chinese).
(胡园, 李震宇, 丁艳娇, 李志颖, 刘志勇, 沈月毛, 有机化学, **2019**, 39, 3230.)
- Vostrikova, T. V.; Kalaev, V. N.; Medvedeva, S. M.; Novichikhina, N. P.; Shikhaliev, K. S. *Period. Tche. Quim.* **2020**, 17, 327.
- Brazhnikova, D. A.; Popova, T. N.; Krylskii, E. D.; Shulgin, K. K.; Matasova, L. V.; Shikhaliev, H. S.; Popov, S. S. *Bioorg. Chem.* **2020**, 14, 70.
- Oncel, S.; Kurtoglu, B.; Karaagac, B. *J. Elastom. Plast.* **2019**, 51, 440.
- (a) Matvienko, I. V.; Bayramov, V. M.; Parygina, N. A.; Kurochkin, V. E.; Alekseev, Y. I. *Russ. J. Bioorg. Chem.* **2020**, 46, 349.
(b) Vijay, K.; Nandi, C.; Samant, S. D. *RSC Adv.* **2016**, 6, 49724.
(c) Nizamov, S.; Sednev, M. V.; Bossi, M. L.; Hebisch, E.; Fraundorf, H.; Lehnart, S. E.; Belov, V. N.; Hell, S. W. *Chem.-Eur. J.*

- 2016, 22, 11631.
- [10] (a) Praja-pati, S. M.; Patel, K. D.; Vekariya, R. H.; Panchal, S. N.; Patel, H. D. *Curr. Org. Synth.* **2019**, 16, 671.
(b) Prajapati, S. M.; Patel, K. D.; Vekariya, R. H.; Panchal, S. N.; Patel, H. D. *RSC Adv.* **2014**, 4, 24463.
(c) Wang, D. S.; Chen, Q. A.; Lu, S. M.; Zhou, Y. G. *Chem. Rev.* **2011**, 112, 2557.
(d) Sharma, R.; Kour, P.; Kumar, A. *J. Chem. Sci.* **2018**, 130, 73.
- [11] Skraup, Z. H. *Ber. Dtsch. Chem. Ges.* **1880**, 13, 2086.
- [12] Li, H.; Wang, J.; Xie, H. X.; Zu, L. S.; Jiang, W.; Duesler, E. N.; Wang, W. *Org. Lett.* **2007**, 9, 965.
- [13] Zhang, Y. F.; Sim, J. H.; MacMillan, S. N.; Lambert, T. H. *Org. Lett.* **2020**, 22, 6026.
- [14] Hwu, J. R.; Panja, A.; Gupta, N. K.; Hu, Y. C.; Hwu, J. R. *Eur. J. Org. Chem.* **2021**, 683.
- [15] (a) Jiang, Y.; Wang, Q.; Sun, R.; Tang, X. Y.; Shi, M. *Org. Chem. Front.* **2016**, 3, 744.
(b) Massaro, N. P.; Chatterji, A.; Sharma, I. *Org. Chem.* **2019**, 84, 13676.
- [16] (a) Cho, S. H.; Yoo, E. J.; Bae, L.; Chang, S. *J. Am. Chem. Soc.* **2005**, 127, 16046.
(b) Cho, S. H.; Chang, S. B. *Angew. Chem. Int. Ed.* **2007**, 46, 1897.
(c) Cassidy, M. P.; Raushel, J.; Fokin, V. V. *Angew. Chem. Int. Ed.* **2006**, 45, 3154.
- [17] Yoo, E. J.; Bae, I.; Cho, S. H.; Han, H.; Chang, S. *Org. Lett.* **2006**, 8, 1347.
- [18] Bae, I.; Han, H.; Chang, S. *J. Am. Chem. Soc.* **2005**, 127, 2038.
- [19] Cui, S. L.; Wang, J.; Wang, Y. G. *Tetrahedron* **2008**, 64, 487.
- [20] (a) Yang, W. G.; Yu, Z.; Zhou, Z. T.; Li, L.; Cui, L.; Luo, H. *RSC Adv.* **2021**, 11, 8701.
(b) Wu, J. Y.; Liao, W. J.; Lin, X. Y.; Liang, C. F. *Org. Biomol. Chem.* **2020**, 18, 8881.
(c) Ding, Z. C.; An, X. M.; Zeng, J. H.; Tang, Z. N.; Zhan, Z. P. *Adv. Synth. Catal.* **2017**, 359, 3319.
(d) Liu, T. L.; Li, Q. H.; Wei, L.; Xiong, Y.; Wang, C. J.; *Adv. Synth. Catal.* **2017**, 359, 1854.
- [21] Sanchez-Diez, E.; Vesga, D. L.; Reyes, E.; Uria, U.; Vicario, J. L. *Org. Lett.* **2016**, 47, 1270.

(Li, L.; Fan, Y.)