钯催化 2-炔基芳基叠氮与丙烯酸衍生物反应合成 1*H*-吲哚-3-丙烯酸酯

李 萍 周志强 杨 帆 徐 瑶 张小祥*

(南京林业大学化学工程学院 江苏省林业资源高效加工利用协同创新中心 南京 210037)

摘要 报道了一种钯催化 2-炔基芳基叠氮化合物与丙烯酸衍生物合成吲哚-3-丙烯酸酯的新方法。该方法实现了在温和的反应条件下合成多种 1*H*-吲哚-3-丙烯酸酯衍生物。

关键词 钯催化; α -亚胺钯卡宾; 吲哚-3-丙烯酸酯; 丙烯酸

Palladium-Catalyzed Synthesis of 1*H*-Indol-3-yl Acrylates from 2-Alkynyl Arylazides and Acrylic Acids

Li, Ping Zhou, Zhiqiang Yang, Fan Xu, Yao Zhang, Xiaoxiang*

(Jiansu Co-Innovation Center of Efficient Processing and Utilization of Forest Resources, College of Chemical Engineering, Nanjing Forestry University, Nanjing 210037)

Abstract A novel method to prepare 1*H*-indol-3-yl acrylates from 2-alkynyl arylazides with acrylic acids catalyzed by Pd has been described. A variety of 1*H*-indol-3-yl acrylate derivatives were prepared under mild reaction conditions.

Keywords Pd-catalysis; α -imino palladium carbene; indole-3-acrylate; acrylic acid

1 Introduction

1*H*-Indol-3-yl acetate derivatives^[1] are widely distributed in nature, which usually present biological activity and medicine potential. At the same time, they are also a very important class of organic intermediates. Furthermore, indole-3-acrylates^[2] with an unsaturated bond as a special class of 1*H*-indol-3-yl acetates have promising application value in the field of organic synthesis and bioorganic chemistry.^[3] However, little information has been done with respect to the synthesis of such skeleton. In 2016, O'Donohue and co-workers^[4] reported the synthesis of *N*-acetyl-indol-3-yl acrylates from *N*-acetyl-indoxyl-3-ols and acrylic acids via a traditional condensation method, which could be used as chromogenic probes for efficient screening and evaluation of feruloyl esterase-like activities (Scheme 1, a). In 2018, one preparation example of [2,3'-biindol]-3-yl cinnamate was developed by Liu's group^[5] in the work of palladium-catalyzed directly oxidative esterification reaction of indoles (Scheme 1, b). In 2018, Du's group^[6] disclosed a *N*-heterocyclic carbene-catalyzed formal conjugate hydroacylation for the synthesis of 1*H*-indol-3-yl esters from indolin-3-ones and aldehydes. In their work, only one

example about the formation of indole-3-acrylate from the cinnamaldehyde was shown (Scheme 1, c). Therefore, novel methods for the preparation of indole-3-acrylates are still needed.

Recently, numerous studies on the formation of *N*-containing heterocycles from transition-metal catalyzed 2-alkynyl arylazides via α -imino metal carbene have been developed by us^[7] and many other groups.^[8] For example, our group previously reported an efficient gold-catalyzed tandem reaction of 2-alkynyl arylazides with carboxylic acids to synthesize 1*H*-indol-3-yl esters.^[9] Then, an efficient method for the synthesis of 1*H*-indole-3-sulfonates via Pd-catalyzed tandem reactions of 2-alkynyl arylazides with sulfonic acids has been developed by our group.^[10] As our ongoing interest on transformation of α -imino metal carbenes, we envisaged that the α -imino palladium carbene generated *in situ* from 2-alkynyl arylazides could be trapped by various acrylic acids to afford the indole-3-acrylates (Scheme 1, d).

2 Results and discussion

Initially, 4-(2-azido-3,5-dichlorophenyl)-2-methylbut-3-

* Corresponding author. E-mail: zhangxiaoxiang@njfu.edu.cn

Received January 27, 2021; revised April 10, 2021; published online May 7, 2021.

Project supported by the National Natural Science Foundation of China (No. 21302096), the Natural Science Foundation of Jiangsu Province (No. BK20171449) and the National First-class Disciplines (PNFD).

国家自然科学基金(No. 21302096)、江苏省自然科学基金(No. BK20171449)以及国家一流学科资助项目。

Scheme 1 Synthetic strategies to prepare indole-3-acrylates
 yn-2-ol (**1a**) and *trans*-crotonic acid (**2a**) were chosen as model substrates to establish the tandem reaction conditions (Table 1). Reaction of **1a** with **2a** in 1,2-dichloroethane

Table 1 Optimization of the reaction conditions^a

Entry	Catalyst	Solvent	Time/h	Yield ^b /%
1	Pd(OAc) ₂	DCE	2	91
2	PdCl ₂	DCE	6	59
3	PdCl ₂ (PPh ₃) ₂	DCE	6	20
4	Pd(NO ₃) ₂ ·2H ₂ O	DCE	1	70 ^c
5	Pd(PPh ₃) ₄	DCE	6	36
6	Pd(OAc) ₂	DCE	3	91 ^d
7	Pd(OAc) ₂	DCE	6	70 ^e
8	Pd(OAc) ₂	DCE	18	72 ^c
9	Pd(OAc) ₂	DCE	2	82 ^f
10	Pd(OAc) ₂	PhCl	3	83
11	Pd(OAc) ₂	MeCN	3	75
12	—	DCE	24	NR ^g

^a Reaction conditions: **1a** (0.1 mmol), **2a** (10 equiv.), catalyst (10 mol%), dry DCE (0.4 mL) and 60 °C; ^b Isolated yield; ^c Reaction was conducted at room temperature; ^d 5 mol% catalyst was employed; ^e 5.0 equiv. of **2a** was used; ^f Reaction was conducted at 80 °C; ^g No catalyst was used. No reaction based on ¹H NMR analysis of crude mixture and **1a** was recovered.

(DCE) with 10 mol% Pd(OAc)₂ at 60 °C for 2 h gave the desired product 5,7-dichloro-2-(2-hydroxypropan-2-yl)-1H-indol-3-yl but-2-enoate (**3a**) in 91% yield (Entry 1).

Later, other palladium catalysts such as PdCl₂, PdCl₂(PPh₃)₂, Pd(NO₃)₂·2H₂O and Pd(PPh₃)₄ were examined, which afforded the desired product **3a** in yields of 20%~70% (Entries 2~5). Repeating the reaction with the catalyst loading of 5 mol% gave the corresponding product **3a** in 91% yield (Entry 6). However, the product yield was decreased to 70% when 5 equiv. of but-2-enoic acid (**2a**) was employed (Entry 7). Upon changing the reaction temperature to room temperature or 80 °C, a low product yield of 72% or 82% was obtained, respectively (Entries 8, 9). When the reaction solvent was changed from DCE to PhCl or MeCN, the desired product **3a** was also obtained in 83% and 75% yields, respectively (Entries 10, 11). Finally, the product **3a** was not detected after 24 h without the presence of catalyst and only the starting material **1a** was recovered (Entry 12).

To define the generality of the Pd-catalyzed tandem reactions for the synthesis of 1H-indole-3-acrylates **3**, we next turned our attention to the reactions of a series of 2-alkynyl arylazides **1**. As shown in Table 2, various substituent groups on the benzene ring (R¹) were first examined. Under the optimal reaction conditions, it was found that the substituted groups such as halogen, CN and CF₃ were all tolerable for the reaction, which afforded the corresponding 1H-indol-3-yl acrylates products in moderate to excellent yields (**3a**~**3d**). It was noted that the reaction carried out with **1d** as starting material underwent a dehydration process to give product **3d**. The substrate **1e** with electron-donating group (R¹=Me) was also employed for the reaction. However the desired product **3e** could not be identified by the analysis of analytical thin layer chromatography (TLC) and crude ¹H NMR after stirring for 24 h under optimal conditions and most of the starting material **1e** was left. Then different substituted groups on the alkyne carbon (R²) were screened. Reaction of substrates with long straight alkyl chain (*n*-Bu and *n*-C₅H₁₁) on the alkyne carbon needed longer reaction time and afforded the desired products in moderate yields (**3f**~**3g**). However, when R²=cyclopropyl, **3h** could be obtained within 0.5 h in 53% yield under the standard reaction conditions. The reaction with steric hindered group (R²=*t*-Bu) was also examined, which resulted in the formation of **3i** in 79% yield. It was found that the desired product **3j** was still obtained in 71% yield when the substituted group on the alkyne carbon contained alkoxy substitute group. The effect of an aromatic substituted group on the alkyne carbon was also checked. When R²=phenyl or substituted phenyl, the desired products (**3k**~**3m**) could be afforded in moderate yields.

In addition, substrates scope of the acrylic acids was next examined under the optimal reaction conditions. The experiment results are shown in Table 3. Reactions with

Table 2 Substrates scope of the 2-alkynyl arylazides **1**^{a,b}

^a Reaction conditions: **1a** (0.1 mmol), **2a** (10 equiv.), $\text{Pd}(\text{OAc})_2$ (5 mol%), dry DCE (0.4 mL) and 60 °C. ^b Isolated yield. ^c Starting material **1e** was left.

Table 3 Substrates scope of the acrylic acids **2**^{a,b}

^a Reaction conditions: **1a** (0.1 mmol), **2** (10 equiv.), $\text{Pd}(\text{OAc})_2$ (5 mol%), dry DCE (0.4 mL) and 60 °C; ^b Isolated yield.

different acrylic acids such as 3,3-dimethylacrylic acid (**2b**), *trans*-2-hexenoic acid (**2c**), *trans*-2-methyl-2-pentenoic acid (**2d**), sorbic acid (**2e**) and *trans*-cinnamic acid (**2f**) could proceed well and provide the corresponding products **3n**~**3s** in moderate to excellent yields.

According to our previous work^[7,9-11] and above experiment results, a possible mechanism of Pd-catalyzed synthesized 1*H*-indol-3-yl acrylates from 2-alkynyl arylazides and acrylic acids was tentatively proposed and outlined in

Scheme 2. Initially, the triple bond of **1a** was activated by palladium catalyst to form the activated complex **A**, which was followed by intramolecular amination of N to the activated triple bond to give vinyl palladium **B**. After releasing N_2 from species **B**, α -imino palladium carbene **C**^[8-12] was generated, which was then attacked by the nucleophilic *trans*-crotonic acid (**2a**) to afford the complex **D**. Final demetalation and protonation of **D** led to the formation of product **3a**.

Scheme 2 Proposed mechanism

3 Conclusions

In conclusion, a new method for the synthesis of 1*H*-indole-3-acrylate derivatives via palladium-catalyzed tandem reactions of 2-alkynyl arylazides and acrylic acids was developed. The reaction was conducted under mild conditions and had good tolerance of various functional groups. Further applications of this strategy are currently underway and will be reported in the future.

4 Experimental section

4.1 Instruments and reagents

^1H NMR and ^{13}C NMR spectra were measured on 400 or 600 MHz spectrometers. Chemical shifts were recorded with tetramethylsilane (TMS) as the internal reference standard. High resolution mass spectra were obtained on a LTQ Orbitrap LC/HRMS mass spectrometer. The starting materials **1** were prepared by reported methods.^[8b] Acrylic acid derivatives **2a**~**2f** were purchased from commercial sources and checked by ^1H NMR. Flash chromatography was performed using silica gel and a gradient solvent system (ethyl acetate/petrol ether as eluent). Unless specified, all reagents were purchased from commercial sources and used as received. Solvents were purified following standard literature procedures. TLC was performed using pre-coated silica gel plate. Visualization was achieved by UV light (254 nm).

4.2 General experimental procedure for the synthesis of 1*H*-indole-3-acrylates **3**

A 10 mL test tube was charged with **1** (0.1 mmol, 1 equiv.), **2** (1 mmol, 10 equiv.) and $\text{Pd}(\text{OAc})_2$ (5 mol%). Sequentially DCE (0.4 mL) was added by a syringe. The test tube was submerged in an oil bath that was already heated to 60 °C. When the reaction was completed checked by TLC, the reaction mixture was cooled to room temperature and then purified directly by silica gel flash chromatography to afford the desired product **3**.

The reaction mixture was neutralized with NaHCO_3 (aq.) and extracted with EtOAc for some products that can not be separated directly (**3b**~**3d**, **3j**~**3l**). The organic layer was

dried with MgSO_4 and then filtered off the MgSO_4 . The filtrate was concentrated under reduced pressure and purified by column chromatography to afford the product **3**.

5,7-Dichloro-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3a**): Isolated yield 91%, yellow solid. m.p. 96.6~98.2 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 1.63 (s, 6H), 2.01 (dd, $J=7.2, 1.6$ Hz, 3 H), 2.38 (br, 1H), 6.12 (dq, $J=14, 1.6$ Hz, 1H), 7.16 (d, $J=1.6$ Hz, 1H), 7.20 (d, $J=1.2$ Hz, 1H), 7.27 (dq, $J=15.6, 6.8$ Hz, 1H), 8.54 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.4, 29.7, 69.6, 115.9, 117.4, 121.2, 121.8, 123.6, 123.9, 125.6, 127.5, 135.5, 148.3, 164.7; IR (KBr) ν : 3479, 3432, 3294, 3083, 2984, 2922, 2853, 1725, 1659, 1557, 1478, 1458, 1263, 1206, 1117, 1098, 968 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{15}\text{H}_{16}\text{Cl}_2\text{NO}_3$ $[\text{M}+\text{H}]^+$ 328.0507, found 328.0508.

5-Cyano-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3b**): Isolated yield 61%, red solid. m.p. 170.6~173.2 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 1.64 (s, 6 H), 2.03 (dd, $J=6.8, 1.6$ Hz, 3H), 6.12~6.16 (dq, $J=14.0, 1.6$ Hz, 1H), 7.29~7.37 (m, 3H), 7.66 (s, 1H), 8.77 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.4, 29.7, 69.6, 103.1, 112.3, 120.4, 121.1, 122.1, 123.4, 123.8, 125.0, 133.2, 135.4, 148.4, 164.7; IR (KBr) ν : 3463, 3403, 3055, 2976, 2923, 2851, 2215, 1722, 1659, 1477, 1441, 1315, 1267, 1193, 1167, 983, 969, 955, 813 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{16}\text{H}_{17}\text{N}_2\text{O}_3$ $[\text{M}+\text{H}]^+$ 285.1239, found 285.1239.

2-(2-Hydroxypropan-2-yl)-5-(trifluoromethyl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3c**): Isolated yield 54%, brown solid. m.p. 124.0~125.6 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 1.64 (s, 6H), 2.03 (dd, $J=6.8, 1.6$ Hz, 3H), 6.16 (dq, $J=15.6, 1.6$ Hz, 1H), 7.27~7.38 (m, 3H), 7.60 (s, 1H), 8.61 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.4, 29.7, 69.6, 111.7, 115.3 (q, $J=4.3$ Hz), 119.0 (q, $J=3.3$ Hz), 121.3, 122.3, 122.6, 123.8, 124.1 (q, $J=236.7$ Hz), 133.0, 134.8, 148.2, 165.0; IR (KBr) ν : 3400, 3264, 3038, 2969, 2925, 2854, 1707, 1640, 1595, 1463, 1334, 1280, 1105, 1048, 965, 736 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{16}\text{H}_{17}\text{F}_3\text{NO}_3$ $[\text{M}+\text{H}]^+$ 328.1161, found 328.1157.

5-Chloro-2-(prop-1-en-2-yl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3d**): Isolated yield 52%, green solid. m.p. 94.4~

98.1 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 2.04 (d, $J=6.8$ Hz, 3H), 2.15 (s, 3H), 5.15 (s, 1H), 5.35 (s, 1H), 6.16 (d, $J=15.2$ Hz, 1H), 7.11~7.18 (m, 2H), 7.27~7.34 (m, 2H), 7.97 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.4, 20.8, 112.3, 113.5, 117.3, 121.5, 123.2, 123.5, 125.9, 126.8, 128.2, 131.4, 133.7, 147.8, 164.6; IR (KBr) ν : 3388, 3051, 2959, 2925, 2853, 1723, 1653, 1630, 1459, 1309, 1284, 1161, 1117, 1098, 1053, 968, 799 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{15}\text{H}_{15}\text{ClNO}_2$ $[\text{M}+\text{H}]^+$ 276.0791, found 276.0788.

2-Butyl-1*H*-indol-3-yl (*E*)-but-2-enoate (**3f**): Isolated yield 61%, brown solid. m.p. 92.4~96.1 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 0.92 (t, $J=7.2$ Hz, 3H), 1.33~1.42 (m, 2H), 1.59~1.67 (m, 2H), 2.00 (dd, $J=6.8$, 1.6 Hz, 3H), 2.65 (t, $J=7.6$ Hz, 2H), 6.16 (dq, $J=15.6$, 1.6 Hz, 1H), 7.05~7.14 (m, 2H), 7.20~7.29 (m, 2H), 7.33 (d, $J=8.0$ Hz, 1H), 7.72 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 13.8, 18.3, 22.3, 24.7, 30.8, 110.9, 116.9, 119.8, 121.6, 121.7, 121.8, 126.3, 128.3, 132.9, 146.7, 164.9; IR (KBr) ν : 3363, 3061, 2955, 2924, 2853, 1723, 1655, 1459, 1317, 1290, 1173, 1102, 983, 737 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{16}\text{H}_{20}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 258.1494, found 258.1495.

2-Pentyl-1*H*-indol-3-yl (*E*)-but-2-enoate (**3g**): Isolated yield 59%, brown liquid. ^1H NMR (CDCl_3 , 400 MHz) δ : 0.88 (t, $J=5.8$ Hz, 3H), 1.31~1.35 (m, 4H), 1.61~1.69 (m, 2H), 2.00 (dd, $J=6.8$, 1.6 Hz, 3H), 2.64 (t, $J=7.6$ Hz, 2H), 6.16 (dq, $J=15.2$, 1.6 Hz, 1H), 7.05~7.14 (m, 2H), 7.20~7.29 (m, 2H), 7.34 (d, $J=7.6$ Hz, 1H), 7.72 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 13.9, 18.3, 22.4, 24.9, 28.4, 31.4, 110.9, 116.9, 119.8, 121.6, 121.7, 121.8, 126.3, 128.3, 132.9, 146.7, 164.9; IR (neat) ν : 3364, 3055, 2955, 2926, 2857, 1720, 1460, 1258, 1154, 970, 741 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{17}\text{H}_{22}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 272.1651, found 272.1653.

2-Cyclopropyl-1*H*-indol-3-yl (*E*)-but-2-enoate (**3h**): Isolated yield 53%, yellow liquid. ^1H NMR (CDCl_3 , 400 MHz) δ : 0.78 (q, $J=4.4$ Hz, 2H), 0.93 (q, $J=4$ Hz, 2H), 1.90~1.93 (m, 1H), 2.00 (dd, $J=6.8$, 1.6 Hz, 3H), 6.17 (dq, $J=15.6$, 1.6 Hz, 1H), 7.04~7.13 (m, 2H), 7.20~7.28 (m, 2H), 7.31 (d, $J=8.8$ Hz, 1H), 7.65 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 5.9, 6.5, 18.2, 110.8, 116.8, 119.9, 121.8, 121.9, 126.1, 127.1, 128.6, 132.4, 146.7, 164.8; IR (neat) ν : 3359, 3013, 2959, 2924, 2853, 1736, 1637, 1607, 1527, 1449, 1292, 1220, 1162, 1107, 967, 773 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{15}\text{H}_{16}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 242.1181, found 242.1184.

2-(*tert*-Butyl)-5,7-dichloro-1*H*-indol-3-yl (*E*)-but-2-enoate (**3i**): Isolated yield 79%, yellow solid. m.p. 96.6~98.2 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 1.42 (s, 9H), 2.01 (dd, $J=6.8$, 1.6 Hz, 3H), 6.13 (dq, $J=15.6$, 1.6 Hz, 1H), 7.13 (d, $J=2$ Hz, 1H), 7.17 (d, $J=1.2$ Hz, 1H), 7.21~7.31 (m, 1H), 7.87 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.3, 29.3, 32.2, 115.7, 116.9, 121.2, 121.5, 124.2, 125.3, 125.6, 127.5, 137.4, 147.7, 164.6; IR (KBr) ν : 3370, 3093, 2969, 2923, 2862, 1730, 1655, 1480, 1311, 1224, 1161, 1100, 986, 976, 875 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{16}\text{H}_{18}\text{Cl}_2\text{NO}_2$ $[\text{M}+\text{H}]^+$ 326.0715, found 326.0719.

2-(2-(Benzyloxy)ethyl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3j**): Isolated yield 71%, green liquid. ^1H NMR (CDCl_3 , 400

MHz) δ : 1.99 (dd, $J=6.8$, 1.6 Hz, 3H), 2.95 (t, $J=5.6$ Hz, 2H), 3.79 (t, $J=5.6$ Hz, 2H), 4.58 (s, 2H), 6.14 (dq, $J=15.6$, 1.6 Hz, 1H), 7.05~7.15 (m, 2H), 7.21~7.26 (m, 2H), 7.33~7.41 (m, 6H), 8.45 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.2, 25.1, 69.4, 73.4, 111.0, 117.0, 119.7, 121.1, 121.7, 121.8, 126.0, 126.6, 127.8, 127.9, 128.6, 133.0, 138.0, 146.8, 164.8; IR (neat) ν : 3372, 3087, 3031, 2922, 2858, 1737, 1654, 1494, 1459, 1310, 1237, 1158, 1099, 968, 741 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{21}\text{H}_{22}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 336.1600, found 336.1603.

2-Phenyl-1*H*-indol-3-yl (*E*)-but-2-enoate (**3k**): Isolated yield 45%, red liquid. ^1H NMR (CDCl_3 , 400 MHz) δ : 2.04 (d, $J=6.8$ Hz, 3H), 6.22 (d, $J=15.6$ Hz, 1H), 7.12~7.16 (m, 1H), 7.19~7.24 (m, 1H), 7.28~7.35 (m, 3H), 7.42~7.46 (m, 3H), 7.65 (d, $J=7.6$ Hz, 2H), 8.13 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.4, 111.4, 118.1, 120.5, 121.8, 122.5, 123.1, 126.2, 127.1, 127.8, 129.0, 130.8, 133.7, 147.4, 164.7; IR (neat) ν : 3368, 3060, 2952, 2924, 2853, 1727, 1655, 1605, 1491, 1233, 1079, 1008, 968, 740 cm^{-1} ; HRMS (ESI) calcd. For $\text{C}_{18}\text{H}_{16}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 278.1181, found 278.1183.

2-(4-Chlorophenyl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3l**): Isolated yield 51%, red liquid. ^1H NMR (CDCl_3 , 400 MHz) δ : 2.02 (dd, $J=6.8$, 1.6 Hz, 3H), 6.20 (dq, $J=15.6$, 1.6 Hz, 1H), 7.10~7.13 (m, 1H), 7.16~7.20 (m, 1H), 7.26~7.40 (m, 5H), 7.51 (d, $J=8.4$ Hz, 2H), 8.10 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 18.4, 111.4, 118.0, 120.5, 121.5, 122.3, 123.4, 125.0, 127.2, 127.3, 129.2, 133.5, 133.8, 147.7, 164.6; IR (neat) ν : 3366, 3055, 2924, 2853, 1718, 1655, 1488, 1251, 1156, 1012, 968, 883, 739 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{18}\text{H}_{15}\text{ClNO}_2$ $[\text{M}+\text{H}]^+$ 312.0791, found 312.0794.

2-(*p*-Tolyl)-1*H*-indol-3-yl (*E*)-but-2-enoate (**3m**): Isolated yield 40%, red liquid. ^1H NMR (CDCl_3 , 600 MHz) δ : 2.00 (dd, $J=6.8$, 1.6 Hz, 3H), 2.38 (s, 3H), 6.19 (dq, $J=15.6$, 1.6 Hz, 1H), 7.09~7.13 (m, 1H), 7.16~7.20 (m, 1H), 7.24 (d, $J=7.8$ Hz, 2H), 7.28~7.33 (m, 2H), 7.32 (d, $J=8.4$ Hz, 1H), 7.39 (d, $J=7.8$ Hz, 2H), 8.04 (br, 1H); ^{13}C NMR (CDCl_3 , 150 MHz) δ : 18.4, 21.3, 111.4, 117.9, 120.4, 121.8, 122.5, 122.9, 126.1, 126.3, 127.9, 129.7, 133.5, 137.7, 147.4, 164.8; IR (neat) ν : 3377, 3035, 2952, 2924, 2854, 1758, 1724, 1608, 1508, 1457, 1233, 1158, 1099, 1062, 969, 819, 740 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{19}\text{H}_{18}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 292.1338, found 292.1341.

5,7-Dichloro-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl 3-methylbut-2-enoate (**3n**): Isolated yield 92%, brown solid. m.p. 98.1~101.1 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 1.63 (s, 6H), 2.03 (s, 3H), 2.25 (s, 3H), 5.99 (s, 1H), 7.15 (d, $J=2.0$ Hz, 1H), 7.21 (d, $J=1.2$ Hz, 1H), 8.55 (br, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 20.7, 27.7, 29.7, 69.6, 114.2, 116.0, 117.4, 121.7, 123.7, 124.2, 125.5, 127.5, 135.5, 161.5, 164.7; IR (KBr) ν : 3520, 3464, 3326, 3080, 2975, 2924, 2853, 1722, 1649, 1556, 1463, 1296, 1240, 1190, 1139, 1061, 985, 949, 850 cm^{-1} ; HRMS (ESI) calcd. for $\text{C}_{16}\text{H}_{17}\text{Cl}_2\text{NO}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 364.0483, found 364.0485.

5,7-Dichloro-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl (*E*)-hex-2-enoate (**3o**): Isolated yield 84%, red liquid. ^1H

NMR (CDCl₃, 400 MHz) δ : 1.00 (t, $J=7.6$ Hz, 3H), 1.55~1.61 (m, 2H), 1.63 (s, 6H), 2.28~2.34 (m, 2H), 6.07~6.11 (m, 1H), 7.14~7.15 (m, 1H), 7.19~7.20 (m, 1H), 7.22~7.29 (m, 1H), 8.60 (br, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 13.7, 21.2, 29.7, 34.6, 69.6, 115.9, 117.4, 119.7, 121.8, 123.6, 123.9, 125.6, 127.5, 135.5, 153.0, 164.9; IR (neat) ν : 3457, 3327, 3080, 2960, 2928, 2871, 1728, 1699, 1649, 1465, 1288, 1154, 1076, 982, 853 cm⁻¹; HRMS (ESI) calcd. for C₁₇H₁₉Cl₂NO₃Na [M + Na]⁺ 378.0640, found 378.0641.

5,7-Dichloro-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl (*E*)-2-methylpent-2-enoate (**3p**): Isolated yield 83%, black solid. m.p. 66.7~68.4 °C; ¹H NMR (CDCl₃, 400 MHz) δ : 1.14 (t, $J=7.6$ Hz, 3H), 1.63 (s, 6H), 1.99 (s, 3H), 2.28~2.35 (m, 2H), 7.05~7.09 (m, 1H), 7.16 (d, $J=1.6$ Hz, 1H), 7.20 (d, $J=1.6$ Hz, 1H), 8.55 (br, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 12.5, 12.9, 22.4, 29.7, 69.6, 116.0, 117.4, 121.7, 123.9, 124.0, 125.6, 126.1, 127.5, 135.4, 147.3, 166.4; IR (KBr) ν : 3516, 3317, 3077, 2970, 2927, 2850, 1713, 1645, 1519, 1474, 1293, 1206, 1132, 1092, 984 cm⁻¹; HRMS (ESI) calcd. for C₁₇H₁₉Cl₂NO₃Na [M + Na]⁺ 378.0640, found 378.0642.

5,7-Dichloro-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl (*2E,4E*)-hexa-2,4-dienoate (**3q**): Isolated yield 68%, yellow solid. m.p. 113.2~114.6 °C; ¹H NMR (CDCl₃, 400 MHz) δ : 1.61 (s, 6H), 1.92 (d, $J=5.2$ Hz, 3H), 2.53 (br, 1H), 6.02 (d, $J=15.2$ Hz, 1H), 6.26~6.34 (m, 2H), 7.15 (d, $J=2.0$ Hz, 1H), 7.21 (d, $J=1.2$ Hz, 1H), 7.46~7.52 (m, 1H), 8.56 (br, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 18.8, 29.7, 69.6, 116.0, 116.8, 117.4, 121.7, 123.7, 124.0, 125.6, 127.4, 129.7, 135.6, 141.7, 148.0, 165.7; IR (KBr) ν : 3461, 3301, 3077, 3026, 2979, 2922, 2853, 1718, 1639, 1616, 1556, 1476, 1378, 1329, 1135, 1000, 984, 868 cm⁻¹; HRMS (ESI) calcd. for C₁₇H₁₇Cl₂NO₃Na [M + Na]⁺ 376.0483, found 376.0485.

5,7-Dichloro-2-(2-hydroxypropan-2-yl)-1*H*-indol-3-yl cinnamate (**3r**): Isolated yield 93%, brown solid. m.p. 116.9~119.6 °C; ¹H NMR (CDCl₃, 400 MHz) δ : 1.67 (s, 6H), 6.70 (d, $J=16.0$ Hz, 1H), 7.18 (d, $J=2.0$ Hz, 1H), 7.26 (d, $J=2.8$ Hz, 1H), 7.44~7.46 (m, 3H), 7.61~7.64 (m, 2H), 7.95 (d, $J=16.0$ Hz, 1H), 8.60 (br, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 29.7, 69.6, 115.9, 116.1, 114.1, 117.5, 121.8, 123.7, 123.9, 125.7, 127.5, 128.5, 129.1, 131.1, 133.9, 135.6, 139.3, 147.7, 165.3; IR (KBr) ν : 3507, 3419, 3071, 2961, 2922, 2853, 1762, 1634, 1553, 1467, 1287, 1145, 1077, 982, 759 cm⁻¹; HRMS (ESI) calcd. for C₂₀H₁₇Cl₂NO₃Na [M + Na]⁺ 412.0483, found 412.0481.

5,7-Dichloro-2-(1-hydroxy-1-phenylethyl)-1*H*-indol-3-yl cinnamate (**3s**): Isolated yield 65%, brown liquid. ¹H NMR (CDCl₃, 400 MHz) δ : 2.01 (s, 3H), 6.45 (d, $J=16$ Hz, 1H), 7.20 (d, $J=2.0$ Hz, 1H), 7.25~7.28 (m, 2H), 7.31~7.35 (m, 2H), 7.43~7.45 (m, 3H), 7.48 (d, $J=7.6$ Hz, 2H), 7.53~7.56 (m, 2H), 7.70 (d, $J=16.0$ Hz, 1H), 8.48 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 29.3, 72.8, 115.9, 116.2, 117.5, 122.3, 123.7, 125.1, 125.4, 125.9, 127.8, 127.9, 128.4, 128.5, 129.1, 130.9, 133.9, 134.4, 144.7, 147.4, 165.4; IR (KBr) ν : 3444, 3061, 2926, 2855, 1728, 1633,

1557, 1494, 1449, 1284, 1170, 1135, 982, 765, 701 cm⁻¹; HRMS (ESI) calcd. for C₂₅H₁₉Cl₂NO₃Na [M + Na]⁺ 474.0640, found 474.0633.

Supporting Information ¹H NMR and ¹³C NMR spectra of **3a**~**3d** and **3f**~**3s**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Bandini, M.; Eichholzer, A. *Angew. Chem., Int. Ed.* **2009**, *48*, 9608.
(b) Joucla, L.; Djakovitch, L. *Adv. Synth. Catal.* **2009**, *351*, 673.
(c) Hong, J.; Xun, L.; Hong, W.; Huashan, Z.; Jieke, C. *J. Wuhan Univ. (Nat. Sci. Ed.)* **1998**, *44*, 175.
(d) Arnold, R. D.; Nutter, W. M.; Stepp, W. L. *J. Org. Chem.* **1959**, *24*, 117.
(e) Liu, K.; Wen, P.; Liu, J.; Huang, G. *Synthesis* **2010**, 3623.
(f) Pohanka, M.; Drtinova, L. *Talanta* **2013**, *106*, 281.
(g) Li, X.; Yu, L.-M.; Wang, B.-J.; Xia, S.-W.; Zhao, H.-Z. *Acta Chim. Sinica* **2008**, *66*, 2507 (in Chinese).
(李霞, 于良民, 王宝娟, 夏树伟, 赵海洲, 化学学报, **2008**, *66*, 2507.)
- [2] (a) Sassano, M. F.; Schlesinger, A. P.; Jarstfer, M. B. *Open Med. Chem. J.* **2012**, *6*, 20.
(b) Masterova, N. S.; Ryabova, S. Y.; Alekseeva, L. M.; Evstratova, M. I.; Kiselev, S. S.; Granik, V. G. *Russ. Chem. Bull.* **2010**, *59*, 637.
(c) Dorjsuren, D.; Burnette, A.; Gray, G. N.; Chen, X.; Zhu, W.; Roberts, P. E.; Currens, M. J.; Shoemaker, R. H.; Ricciardi, R. P.; Sei, S. *Antiviral Res.* **2006**, *69*, 9.
(d) Perlmutter, P.; Puniani, E.; Westman, G. *Tetrahedron Lett.* **1996**, *37*, 1715.
(e) Krajnc, P.; Štefanec, D.; Brown, J. F.; Cameron, N. R. *J. Polym. Sci., Part A: Polym. Chem.* **2005**, *43*, 296.
(f) Štefanec, D.; Krajnc, P. *Polym. Inter.* **2007**, *56*, 1313.
(g) Vasconcellos, M. L.; Silva, T. M.; Camara, C. A. *Pest. Manage. Sci.* **2006**, *62*, 288.
(h) Littke, A. F.; Fu, G. C. *J. Org. Chem.* **1999**, *64*, 10.
- [3] (a) Pohanka, M.; Drtinova, L. *Talanta* **2013**, *106*, 281.
(b) Nirogia, R. V. S.; Kambhampati, R.; Daulatabad, A. V.; Gudla, P.; Shaikh, M.; Achanta, P. K.; Shinde, A. K.; Dubey, P. K. *J. Enzyme Inhib. Med. Chem.* **2011**, *26*, 341.
- [4] Gherbovet, O.; Fauré, R.; Ferreira, F.; Durand, J.; Ragon, M.; Hostyn, G.; Record, E.; Bozonnet, S.; O'Donohue, M. J. *J. Mol. Catal. B: Enzym.* **2016**, *126*, 24.
- [5] Song, J.; Cui, J.; Liang, H.-B.; Liu, Q.; Dong, Y.-H.; Liu, H. *Asian J. Org. Chem.* **2018**, *7*, 341.
- [6] Zhu, J.-D.; Fang, S.-S.; Sun, K.-W.; Fang, C.; Lu, T.; Du, D. *J. Org. Chem.* **2018**, *83*, 10430.
- [7] (a) Li, P.; Zhu, B.; Xu, Y.; Zhou, Z.; Hu, G.; Yang, F.; Xu, S.; Zhang, X. *Org. Chem. Front.* **2020**, *7*, 3480.
(b) Li, P.; Yong, W.; Sheng, R.; Rao, W.; Zhu, X.; Zhang, X. *Adv. Synth. Catal.* **2019**, *361*, 201.
- [8] For recent selected examples of α -imino metal carbenes from 2-alkynyl arylazides: (a) Lu, B.; Luo, Y.-D.; Liu, L.-Z.; Ye, L.-W.; Wang, Y.-Z.; Zhang, L.-M. *Angew. Chem., Int. Ed.* **2011**, *50*, 8358.
(b) Wetzel, A.; Gagosz, F. *Angew. Chem., Int. Ed.* **2011**, *50*, 7354.
(c) Hou, Z.; Oishi, S.; Suzuki, Y.; Kure, T.; Nakanishi, I.; Hirasawa, A.; Tsujimoto, G.; Ohno, H.; Fujii, N. *Org. Biomol. Chem.* **2013**, *11*, 3288.

- (d) Tokimizu, Y.; Oishi, S.; Fujii, N.; Ohno, H. *Org. Lett.* **2014**, *16*, 3138.
- (e) Zhou, Q.; Zhang, Z.; Zhou, Y.; Li, S.; Zhang, Y.; Wang, J. *J. Org. Chem.* **2017**, *82*, 48.
- (f) Li, T.; Chen, B.-L.; Zhu, L.-L.; Chen, Z. *Tetrahedron Lett.* **2020**, *61*, 151851.
- (g) Li, N.; Lian, X.-L.; Li, Y.-H.; Wang, T.-Y.; Han, Z.-Y.; Zhang, L.; Gong, L.-Z. *Org. Lett.* **2016**, *18*, 4178.
- (h) Gronnier, C.; Boissonnat, G.; Gagosz, F. *Org. Lett.* **2013**, *15*, 4234.
- (i) Shen, W.-B.; Sun, Q.; Li, L.; Liu, X.; Zhou, B.; Yan, J.-Z.; Lu, X.; Ye, L.-W. *Nat. Commun.* **2017**, *8*, 1748.
- For recent selected examples of metal carbenes: (j) Li, M.-L.; Chen, M.-Q.; Xu, B.; Zhu, S.-F.; Zhou, Q.-L. *Acta Chim. Sinica* **2018**, *76*, 883 (in Chinese).
- (李茂霖, 陈梦青, 徐彬, 朱守非, 周其林, 化学学报, **2018**, *76*, 883.)
- (k) Qiu, D.; Qiu, M.-L.; Ma, R.; Zhang, Y.; Wang, J.-B. *Acta Chim. Sinica* **2016**, *74*, 472 (in Chinese).
- (邱颀, 邱孟龙, 马戎, 张艳, 王剑波, 化学学报, **2016**, *74*, 472.)
- [9] Zhang, X.-X.; Sun, X.-P.; Zhang, H.-F.; Cui, X.-L.; Ma, M.-T. *Chin. J. Org. Chem.* **2015**, *35*, 1469 (in Chinese).
- (张小祥, 孙小萍, 张海飞, 崔杏丽, 马猛涛, 有机化学, **2015**, *35*, 1469.)
- [10] Zhang, X.-X.; Li, P.; Lyu, C.; Yong, W.-X.; Li, J.; Zhu, X.-B.; Rao, W.-D. *Org. Biomol. Chem.* **2017**, *15*, 6080.
- [11] (a) Zhang, X.-X.; Sun, X.-P.; Fan, H.; Li, P.; Lyu, C.; Rao, W.-D. *Eur. J. Org. Chem.* **2016**, *25*, 426.
- (b) Zhang, X.-X.; Sun, X.-P.; Fan, H.; Lyu, C.; Li, P.; Zhang, H.-F.; Rao, W.-D. *RSC Adv.* **2016**, *6*, 56319.
- [12] For recent selected examples on the generation of palladium carbenes: (a) Ding, H.; Bai, S.; Lu, P.; Wang, Y. *Org. Lett.* **2017**, *19*, 4604.
- (b) Shao, Z.; Zhang, H. *Chem. Soc. Rev.* **2012**, *41*, 560.
- (c) Xiao, Q.; Zhang, Y.; Wang, J. *Acc. Chem. Res.* **2013**, *46*, 236.
- (d) Xia, Y.; Zhang, Y.; Wang, J. *ACS Catal.* **2013**, *3*, 2586.
- (e) Liu, Z.; Tan, H.; Fu, T.; Xia, Y.; Qiu, D.; Zhang, Y.; Wang, J. *J. Am. Chem. Soc.* **2015**, *137*, 12800.

(Zhao, C.)