膦氧导向钯催化邻碳硼烷 Heck 类型的官能团化反应

连凌翔 尹静怡 林彩霞* 叶克印 袁耀锋*

(福州大学化学学院 福州 350108)

摘要 邻-碳硼烷的选择性官能团化是碳硼烷化学中的一个热点。由于邻-碳硼烷特殊的三维芳香性, 其在聚集诱导发光材料、非线性发光材料以及其他功能材料领域有着广泛的应用。在邻碳硼烷上引入 π -共轭体系能进一步扩大其在发光材料领域的应用。报道了膦氧导向钯催化邻碳硼烷 Heck 类型的单烯基化反应。

关键词 钯; Heck 反应; 碳硼烷; B—H 活化

Phosphine Oxide-Directed Palladium-Catalyzed Heck-Type Functionalization of *o*-Carboranes

Lian, Lingxiang Yin, Jingyi Lin, Caixia* Ye, Keyin Yuan, Yaofeng*

(College of Chemistry, Fuzhou University, Fuzhou 350108)

Abstract The selective functionalization of *o*-carboranes is a popular topic in carborane chemistry. *o*-Carborane has been widely used in aggregation-induced luminescent materials, nonlinear luminescent materials and other function materials due to its special 3D-aromaticity. The π conjugated systems-tethered *o*-carboranes could further broaden its applications in the field of luminescent materials. Herein, we have developed a palladium-catalyzed Heck-type mono-alkenylation of *o*-carboranes featuring phosphine oxide as the directing group.

Keywords palladium; Heck reaction; carborane; B—H activation

1 Introduction

The rich properties of carborane lead to a series of applications in many fields.^[1] It can be used in coordination chemistry^[2] and the construction of functional molecules^[1a,3] because of rigid structure, 3d-aromaticity and unique electronic property. It also can be introduced into pharmaceutical molecules^[4] due to its non-toxic and high boron content. Generally speaking, the π conjugated systems-tethered *o*-carboranes^[5] have many unexpected optical properties.^[6] Recently, Yan and her co-worker^[6a] reported many studies about luminescent materials containing carborane. Thereinto, alkenylation of *o*-carborane,^[7] like Heck-type reaction, is one of the most effective ways to give the π conjugated systems-tethered *o*-carboranes.

Heck reaction, *i.e.*, palladium-catalyzed alkenylation of aryl halides and alkenes, is arguably one of the most widely used catalytic reactions to form diverse C—C bonds.^[8] Recent investigations also show that the unactivated aromatics are also promising candidates involved in this type

transformation via an interesting C—H activation step.^[9] Analogously, the σ -aromatic carboranes^[10] are anticipated to undergo similar Heck-type functionalization with olefins via a B—H activation step.^[11] Indeed, Cao and co-workers^[11a] have demonstrated that the *o*-carboranes could be employed in the palladium-catalyzed alkenylation reaction, in which the most electron-rich B(8)/B(9) sites were mono-alkenylated almost non-selectively (Schemes 1A and 1B).^[11a] The installment of a carboxylic acid directing group provided opportunities to outcompete the innate reactivity of the *o*-carboranes.^[12] Both the palladium^[10b-10c] and rhodium^[10d] catalysts have been shown to be capable to catalyze the carboxylic acid directed Heck-type functionalization toward mono-alkenylation and bis-alkenylation at B(4) and B(4,5), respectively (Scheme 1C).

We have recently unveiled a palladium-catalyzed B(3)-H Suzuki-type arylation of *o*-carboranes^[13] featuring the easily prepare, air and moisture stable phosphine oxide^[14] as the competent directing group. Herein, we showed that the phosphine oxide-tethered *o*-carboranes could be further

* Corresponding authors. E-mail: yaofeng_yuan@fzu.edu.cn; lincaixia_fzu@fzu.edu.cn

Received March 6, 2021; revised April 16, 2021; published online May 7, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21772023, 21901041).

国家自然科学基金(Nos. 21772023, 21901041)资助项目.

(A) Numbering system of *o*-carborane chemistry (B) Undirected Heck-type functionalization of *o*-carborane (previous work) (C) Carboxylic acid directed Heck-type functionalization of *o*-carborane (previous work) (D) Phosphine oxide-directed directed Heck-type functionalization of *o*-carborane (this work) **Scheme 1** Transition-metal-catalyzed B—H bond functionalization of *o*-carboranes

applied in the Heck-type mono-functionalization of *o*-carboranes with styrenes in a highly site selective manner (Scheme 1D).

2 Results and discussion

Initial trial on the phosphine oxide-directed palladium-catalyzed Heck-type reaction between phosphine oxide-tethered *o*-carborane (**1a**) and styrene (**2a**) [reaction conditions: 10 mol% Pd(OAc)₂ as catalyst, 2 equiv. of AgOAc as oxidant in toluene at 80 °C for 24 h] only led to a marginal yield (Entry 1, Table 1). Extensively reaction condition optimization revealed that the trifluoroacetate (TFA) was crucial to the desired reactivity since improved yields were observed (Entries 2–3). Interestingly, the implementation of both Pd(TFA)₂ as catalyst and AgTFA as oxidant afforded the desired Heck-type product in 32% yield (Entry 4). Further improved results could be obtained by the employment of 3 equiv. of styrene and 20 mol% palladium catalyst (Entries 5–6). Accordingly, the desired alkenyl functionalized *o*-carborane (**3**) was isolated in 46% yield. Unfortunately, the efforts to further improve reaction yield were not fruitful (see Supporting Information for details). The X-ray diffraction analysis of compound **3** (CCDC 2057524) unambiguously confirmed that this phosphine oxide-directed Heck-type reaction took place highly selectively at B(4) vertex in the *o*-carborane scaffold (Figure 1).

The scope and limitation of this phosphine oxide-di-

Table 1 Reaction condition optimization^a

Entry	Cat.	Ag salt	x	Yield ^b /%
1	Pd(OAc) ₂	AgOAc	2	<5
2	Pd(TFA) ₂	AgOAc	2	13
3	Pd(OAc) ₂	AgTFA	2	21
4	Pd(TFA) ₂	AgTFA	2	32
5 ^c	Pd(TFA) ₂	AgTFA	3	46
6 ^{c,d}	Pd(TFA) ₂	AgTFA	3	54 (46) ^e

^a 0.05 mmol of **1a**, 0.1 mmol of **2a**, 10 mol% Pd(TFA)₂ in 1 mL of toluene under an argon atmosphere. ^b Yield determined by ³¹P NMR of the crude reaction mixture. ^c 0.15 mmol of **2a**. ^d 20 mol% Pd(TFA)₂. ^e Isolated yield.

Figure 1 X-ray diffraction analysis of **3**

rected palladium-catalyzed Heck-type reaction were further evaluated (Table 2). Styrene and its derivatives bearing electron-donating group (4-Me, 3-Me) or electron-withdrawing group (4-F, 4-Cl) were generally well tolerated under current reaction conditions (**4**–**7**, 34%–48%). The strong electron-withdrawing groups, like CF₃, could dramatically reduce the yield (see Supporting Information for details). However, there was no conversion with reaction between **1a** and 4-vinylpyridine or styrenes which bore a strong electron-withdrawing groups (such as NO₂). 2-Naphthalene (**8**) and 4-phenylstyrene (**9**) could also be efficiently incorporated into the *o*-carborane scaffold. In addition, *p*-tolyl substituted phosphine oxide-tethered *o*-carboranes were also amenable to this Heck-type reaction with a variety of substituted styrenes (**10**–**13**, 27%–42%). The reaction between alkyl groups substituted diphenylphosphinoyl-*o*-carborane (**1f**) and 4-vinyl-1,1'-bi-phenyl could lead to three products with similar polarity. Unfortunately, they could not be isolated.

Table 2 Substrate scope of the palladium-catalyzed B(4)-alkenylation of *o*-carboranes^a

^a 0.05 mmol 1, 3.0 equiv. 2, 20 mol% Pd(TFA)₂, 3.0 equiv. AgTFA, 1.0 mL toluene for 24 h under an argon atmosphere.

Consistent with our previous observation in the Suzuki coupling reaction, this Heck-type reaction was also very sensitive to the substitution *ortho* (**14**) to the phenyl group. Both the hydron (**15**) and *m*-tolyl (**16**) substituted phosphine oxide-tethered *o*-carboranes were sluggish under current reaction conditions. Activated alkenes, such as ethyl acrylate (**17**), were not sufficiently reactive in this transformation (Scheme 2).

Scheme 2 Unsuccessful substrates

A plausible mechanism was then proposed based on our experimental observation and literature precedence (Scheme 3). The use of phosphine oxide as the directing group was previously found to prefer the B(3)-selectivity in the Suzuki coupling reaction. The B(4)-site selectivity **II** in this study might be due to the cooperative effects between palladium catalyst and silver salt though much efforts still needed to elucidate the reaction mechanism.^[13c] Migratory insertion of the Pd—B bond into styrene then produced the seven-membered palladacycle **III**. Finally, β -H elimination delivered the intermediate **IV**, which could be further oxidized by the superstoichiometric amount of AgTFA to give the alkenylated *o*-carborane close the catalytic cycle.

3 Conclusions

In summary, we have developed a phosphine oxide-directed highly regioselective palladium-catalyzed Heck-type B(4)-alkenylation of *o*-carboranes with styrenes under mild reaction conditions. Further extension of the substrate scope and synthetic application of this protocol are ongoing in our lab.

4 Experimental section

4.1 General experimental information

All reactions were performed under an atmosphere of

argon using standard Schlenk techniques, unless otherwise indicated. All commercial reagents were used without further purification, unless otherwise noted. Reactions were

Scheme 3 Proposed mechanism

monitored by thin layer chromatography (TLC) analysis. TLC plates were viewed under UV light and stained with potassium permanganate. Yields refer to products isolated after purification by column chromatography, unless otherwise stated. Proton nuclear magnetic resonance (^1H NMR) spectra were recorded on a Bruker AV-400 MHz or a JEOL JNM-500 MHz. NMR samples were dissolved in CDCl_3 (unless specified otherwise) and chemical shifts are referenced to residual nondeuterated solvent. Carbon nuclear magnetic resonance (^{13}C NMR)/boron nuclear magnetic resonance (^{11}B NMR)/phosphorus nuclear magnetic resonance (^{31}P NMR)/fluorine nuclear magnetic resonance (^{19}F NMR) spectra were recorded on the same Bruker instrument (101/128/162/376 MHz) or the same JEOL instrument (126/160/1202/471 MHz). HRMS were obtained on a Waters Micromass GCT Premier (EI) and Exactive Plus LC-MS (ESI) mass spectrometers.

4.2 Procedure for the synthesis of material 1f

To a 50 mL dried Schlenk tube with a stir bar were sequentially added 2-hexyl-*o*-carborane^[15] (2 mmol), tetrahydrofuran (THF, 5 mL), *n*-BuLi (1.3 mol·L⁻¹, 2.2 mmol) at -78°C under argon. After 2 h, Ph_2PCl (2.2 mmol) was added and the reaction mixture was kept stirring for 2 h. The reaction was quenched by adding water, to which an excess amount of 30% H_2O_2 was then added (6 mmol) at 0°C . The mixture was stirred at room temperature for 4~6 h. The mixture was washed by saturated aqueous Na_2SO_3 , extracted with ethyl acetate 3 times. The organic layer was then dried by using Na_2SO_4 and filtered. After evaporation of the solvent, the residue was purified by column chromatography on 200~300 mesh silica gel with

petroleum ether/dichloromethane ($V:V=5:1\sim 3:1$) as eluent and afforded 1-diphenylphosphinoyl-2-hexyl-*o*-carborane (**1f**),^[15] white solid, yield 86%. m.p. $167\sim 168^\circ\text{C}$; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.15~8.04 (m, 4H), 7.65~7.58 (m, 2H), 7.58~7.51 (m, 4H), 2.57~2.47 (m, 2H), 1.42~1.31 (m, 2H), 1.29~1.21 (m, 2H), 1.21~1.13 (m, 4H), 0.86 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ : 133.1 (d, $^3J_{\text{P,C}}=2.9$ Hz), 132.4 (d, $^2J_{\text{P,C}}=8.9$ Hz), 130.1, 129.0, 128.6 (d, $^1J_{\text{P,C}}=12.8$ Hz), 85.0, 75.8 (d, $^1J_{\text{P,C}}=55.9$ Hz), 36.0, 31.2, 29.8, 28.7, 22.4, 14.0; ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 0.34 (d, $J=149.5$ Hz, 1B), -3.62 (d, $J=146.6$ Hz, 1B), -6.34~-16.74 (m, 8B); ^{31}P NMR (162 MHz, Chloroform-*d*) δ : 21.9.

4.3 General procedure for the synthesis of 1-aryl-2-diphenylphosphinoyl-(*E*)-7-alkenyl-*o*-carborane

To a 10 mL dried Schlenk tube with a stir bar were sequentially added **4** (0.05 mmol), styrene (0.15 mmol), $\text{Pd}(\text{TFA})_2$ (20 mol%), toluene (1 mL) and AgTFA (0.15 mmol) under argon. After the reaction mixture was stirred at 90°C for 24 h, the reaction mixture was cooled to room temperature and filtered through a short silica gel column using dichloromethane as eluent. After evaporation of the solvent, the residue was purified by preparative TLC with petroleum ether/1,4-dioxane ($V:V=20:1$) as eluent to give the crude product which was purified further by recrystallization with petroleum ether.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-styryl-*o*-carborane (**3**): White solid, 36.04 mg, yield 46%. m.p. $218\sim 220^\circ\text{C}$; ^1H NMR (500 MHz, Chloroform-*d*) δ : 7.73~7.68 (m, 2H), 7.60~7.54 (m, 2H), 7.44~7.38 (m, 1H), 7.37~7.32 (m, 3H), 7.35~7.24 (m, 1H), 7.30~7.27 (m, 2H), 7.27~7.23 (m, 2H), 7.23 (d, $J=3.6$ Hz, 1H), 7.21 (d, $J=3.2$ Hz, 1H), 7.19 (d, $J=3.7$ Hz, 1H), 7.18~7.17 (m, 1H), 7.17~7.14 (m, 1H), 7.05 (t, $J=7.7$ Hz, 2H), 6.76 (d, $J=18.0$ Hz, 1H), 5.62 (d, $J=18.0$ Hz, 1H); ^{13}C NMR (126 MHz, Chloroform-*d*) δ : 144.5, 137.7, 132.9 (d, $^2J_{\text{P,C}}=9.0$ Hz), 132.6 (d, $^2J_{\text{P,C}}=9.3$ Hz), 132.43 (d, $^3J_{\text{P,C}}=3.0$ Hz), 132.35 (d, $^3J_{\text{P,C}}=3.0$ Hz), 131.9, 130.3, 130.1, 129.9, 129.0, 128.5, 128.3, 128.0, 127.89 (d, $^1J_{\text{P,C}}=12.6$ Hz), 127.88 (d, $^1J_{\text{P,C}}=12.7$ Hz), 127.8, 126.6, 86.3, 77.6 (d, $^1J_{\text{P,C}}=53.7$ Hz); ^{11}B NMR (160 MHz, Chloroform-*d*) δ : 0.8 (d, $J=144.5$ Hz, 2B), +2.2 (d, $J=139.5$ Hz, 2B), -6.8~+18.3 (m, 6B); ^{31}P NMR (202 MHz, Chloroform-*d*) δ : 18.7. HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{32}\text{B}_{10}\text{OP}$ [$\text{M}+\text{H}$]⁺: 523.3188; found 523.3188.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-(*p*-fluorostyryl)-*o*-carborane (**4**): White solid, 29.99 mg, yield 37%. m.p. $262\sim 263^\circ\text{C}$; ^1H NMR (500 MHz, Chloroform-*d*) δ : 7.85~7.68 (m, 2H), 7.70~7.51 (m, 2H), 7.44~7.38 (m, 1H), 7.38~7.31 (m, 3H), 7.30~7.26 (m, 1H), 7.27~7.16 (m, 4H), 7.15~7.09 (m, 2H), 7.09~7.03 (m, 2H), 7.03~6.94 (m, 2H), 6.69 (d, $J=17.8$ Hz, 1H), 5.53 (d, $J=17.9$ Hz, 1H); ^{13}C NMR (126 MHz, Chloroform-*d*) δ : 162.5 (d, $^1J_{\text{F,C}}=247.4$ Hz), 143.0, 134.0 (d, $^4J_{\text{F,C}}=3.1$ Hz), 132.9 (d, $^2J_{\text{P,C}}=9.0$ Hz), 132.5 (d, $^2J_{\text{P,C}}=9.2$ Hz), 132.4 (d, $^3J_{\text{P,C}}=$

2.9 Hz), 132.3 (d, $^3J_{\text{P,C}}=3.0$ Hz), 131.9, 130.3, 130.0, 129.8, 128.9 (d, $^3J_{\text{F,C}}=12.2$ Hz), 128.2, 128.1, 127.90 (d, $^1J_{\text{P,C}}=12.3$ Hz), 127.89 (d, $^1J_{\text{P,C}}=12.7$ Hz), 127.8, 115.1 (d, $^2J_{\text{F,C}}=21.5$ Hz), 86.3, 77.7; ^{11}B NMR (160 MHz, Chloroform-*d*) δ : 0.8 (d, $J=180.6$ Hz, 2B), -2.5 (d, $J=97.5$ Hz, 2B), $-6.7\sim-29.7$ (m, 6B); ^{31}P NMR (202 MHz, Chloroform-*d*) δ : 18.3; ^{19}F NMR (471 MHz, Chloroform-*d*) δ : -113.7 . HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{31}\text{B}_{10}\text{OPF} [\text{M}+\text{H}]^+$: 541.3094; found 541.3099.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-(*p*-chlorostyryl)-*o*-carborane (**5**): White solid, 28.37 mg, yield 34%. m.p. $222\sim 224$ °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.81~7.72 (m, 2H), 7.63~7.52 (m, 2H), 7.47~7.42 (m, 1H), 7.42~7.34 (m, 3H), 7.32~7.20 (m, 7H), 7.15~7.03 (m, 4H), 6.71 (d, $J=17.9$ Hz, 1H), 5.63 (d, $J=17.9$ Hz, 1H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ : 142.8, 136.2, 133.5, 132.9 (d, $^2J_{\text{P,C}}=8.9$ Hz), 132.5 (d, $^2J_{\text{P,C}}=8.9$ Hz), 132.4 (d, $^3J_{\text{P,C}}=3.0$ Hz), 132.3 (d, $^3J_{\text{P,C}}=3.2$ Hz), 131.9, 130.3, 130.0, 129.8, 129.0, 128.8, 128.4, 127.92 (d, $^1J_{\text{P,C}}=11.7$ Hz), 127.89 (d, $^1J_{\text{P,C}}=11.4$ Hz), 127.84, 127.80, 86.3, 77.7; ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 1.8 (d, $J=157.0$ Hz, 2B), -1.1 (d, $J=125.7$ Hz, 2B), $-5.5\sim-14.6$ (m, 6B); ^{31}P NMR (162 MHz, Chloroform-*d*) δ : 17.5. HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{31}\text{B}_{10}\text{ClOP} [\text{M}+\text{H}]^+$: 557.2794; found 557.2799.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-(*p*-methylstyryl)-*o*-carborane (**6**): White solid, 38.61 mg, yield 48%. m.p. $180\sim 181$ °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.74~7.65 (m, 2H), 7.64~7.56 (m, 2H), 7.47~7.40 (m, 1H), 7.41~7.32 (m, 3H), 7.33~7.17 (m, 5H), 7.16~7.01 (m, 6H), 6.76 (d, $J=17.9$ Hz, 1H), 5.55 (d, $J=18.0$ Hz, 1H), 2.36 (s, 3H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ : 144.5, 137.9, 135.0, 1332.0 (d, $^2J_{\text{P,C}}=9.0$ Hz), 132.7 (d, $^2J_{\text{P,C}}=9.4$ Hz), 132.40 (d, $^3J_{\text{P,C}}=3.0$ Hz), 132.35 (d, $^3J_{\text{P,C}}=3.0$ Hz), 132.0, 130.2, 130.13, 130.10, 129.0, 128.4, 127.9 (d, $^1J_{\text{P,C}}=12.9$ Hz), 127.8 (d, $^1J_{\text{P,C}}=13.2$ Hz), 127.8, 127.3, 126.6, 86.3, 77.7 (d, $^1J_{\text{P,C}}=51.4$ Hz), 21.3; ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 1.8 (d, $J=135.9$ Hz, 2B), -1.0 (d, $J=157.6$ Hz, 2B), $-3.0\sim-16.0$ (m, 6B); ^{31}P NMR (162 MHz, Chloroform-*d*) δ : 18.2. HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{34}\text{B}_{10}\text{OP} [\text{M}+\text{H}]^+$: 537.3345; found 537.3345.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-(*m*-methylstyryl)-*o*-carborane (**7**): White solid, 34.60 mg, yield 43%. m.p. $227\sim 229$ °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.79~7.70 (m, 2H), 7.68~7.59 (m, 2H), 7.50~7.44 (m, 1H), 7.44~7.37 (m, 3H), 7.35~7.20 (m, 7H), 7.15~7.06 (m, 3H), 7.05~6.98 (m, 2H), 6.81 (d, $J=17.6$ Hz, 1H), 5.62 (d, $J=17.8$ Hz, 1H), 2.40 (s, 3H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ : 144.6, 137.8, 137.6, 133.0 (d, $^2J_{\text{P,C}}=8.9$ Hz), 132.7 (d, $^2J_{\text{P,C}}=9.1$ Hz), 132.4 (d, $^3J_{\text{P,C}}=3.3$ Hz), 132.3 (d, $^3J_{\text{P,C}}=3.3$ Hz), 132.0, 130.2, 130.11, 130.08, 129.0, 128.8, 128.2, 127.88 (d, $^1J_{\text{P,C}}=13.0$ Hz), 127.87 (d, $^1J_{\text{P,C}}=13.0$ Hz), 127.8, 127.4, 127.2, 124.0, 86.4, 77.9, 21.4; ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 1.8 (d, $J=144.0$ Hz, 2B), -1.0 (d, $J=109.4$ Hz, 2B), $-5.3\sim-15.3$ (m, 6B); ^{31}P NMR (162 MHz, Chloroform-*d*)

δ : 18.2. HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{34}\text{B}_{10}\text{OP} [\text{M}+\text{H}]^+$: 537.3345; found 537.3343.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-(2-(naphthalen-2-yl)vinyl)-*o*-carborane (**8**): White solid, 41.21 mg, yield 48%. m.p. $262\sim 264$ °C; ^1H NMR (500 MHz, Chloroform-*d*) δ : 7.82~7.77 (m, 2H), 7.75~7.68 (m, 3H), 7.58 (s, 1H), 7.59~7.53 (m, 2H), 7.48~7.42 (m, 2H), 7.39 (t, $J=7.4$ Hz, 1H), 7.35 (d, $J=7.7$ Hz, 2H), 7.31~7.27 (m, 2H), 7.25~7.18 (m, 3H), 7.17~7.11 (m, 2H), 7.04 (t, $J=7.6$ Hz, 2H), 6.92 (d, $J=17.9$ Hz, 1H), 5.73 (d, $J=17.9$ Hz, 1H); ^{13}C NMR (126 MHz, Chloroform-*d*) δ : 144.4, 135.2, 133.4, 133.2, 132.9 (d, $^2J_{\text{P,C}}=9.0$ Hz), 132.6 (d, $^2J_{\text{P,C}}=9.2$ Hz), 132.4 (d, $^3J_{\text{P,C}}=3.0$ Hz), 132.3 (d, $^3J_{\text{P,C}}=2.8$ Hz), 131.9, 130.3, 130.1, 129.9, 129.0, 128.6, 128.1, 127.92 (d, $^1J_{\text{P,C}}=12.8$ Hz), 127.90 (d, $^1J_{\text{P,C}}=12.8$ Hz), 127.81, 127.77, 127.6, 126.9, 126.2, 126.0, 123.7, 86.4, 77.7 (d, $^1J_{\text{P,C}}=51.4$ Hz); ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 2.0 (d, $J=165.0$ Hz, 2B), -1.1 (d, $J=141.3$ Hz, 2B), $-4.1\sim-15.2$ (m, 6B); ^{31}P NMR (162 MHz, Chloroform-*d*) δ : 18.0. HRMS (ESI) calcd for $\text{C}_{32}\text{H}_{34}\text{B}_{10}\text{OP} [\text{M}+\text{H}]^+$: 573.3345; found 573.3340.

1-Diphenylphosphinoyl-2-phenyl-(*E*)-4-(4-phenylstyrene)-*o*-carborane (**9**): White solid, 40.39 mg, yield 45%. m.p. $234\sim 236$ °C; ^1H NMR (500 MHz, Chloroform-*d*) δ : 7.74~7.67 (m, 2H), 7.63~7.58 (m, 2H), 7.58~7.56 (m, 2H), 7.55~7.51 (m, 2H), 7.46~7.37 (m, 3H), 7.36~7.30 (m, 4H), 7.28~7.16 (m, 7H), 7.03 (t, $J=7.2$ Hz, 2H), 6.78 (d, $J=17.9$ Hz, 1H), 5.64 (d, $J=17.9$ Hz, 1H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ : 143.9, 140.6, 140.5, 136.8, 133.0 (d, $^2J_{\text{P,C}}=9.0$ Hz), 132.6 (d, $^2J_{\text{P,C}}=9.2$ Hz), 132.45 (d, $^3J_{\text{P,C}}=2.8$ Hz), 132.35 (d, $^3J_{\text{P,C}}=2.8$ Hz), 131.9, 130.3, 130.1, 130.0, 129.0, 128.8, 128.6, 127.90 (d, $^1J_{\text{P,C}}=13.0$ Hz), 127.89 (d, $^1J_{\text{P,C}}=13.0$ Hz), 127.8, 127.5, 127.4, 127.1, 126.9, 86.3, 77.9; ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 1.8 (d, $J=135.9$ Hz, 2B), -1.1 (d, $J=90.0$ Hz, 2B), $-5.5\sim-15.0$ (m, 6B); ^{31}P NMR (162 MHz, Chloroform-*d*) δ : 18.0. HRMS (ESI) calcd for $\text{C}_{34}\text{H}_{36}\text{B}_{10}\text{OP} [\text{M}+\text{H}]^+$: 599.3501; found 599.3495.

1-Diphenylphosphinoyl-2-(*p*-tolyl)-(*E*)-4-(*p*-fluorostyryl)-*o*-carborane (**10**): White solid, 29.10 mg, yield 35%. m.p. $187\sim 190$ °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.79~7.69 (m, 2H), 7.60~7.51 (m, 2H), 7.43~7.37 (m, 1H), 7.37~7.31 (m, 1H), 7.25~7.18 (m, 5H), 7.16~7.06 (m, 3H), 7.03~6.94 (m, 2H), 6.83 (d, $J=7.6$ Hz, 2H), 6.68 (d, $J=18.0$ Hz, 1H), 5.56 (d, $J=18.0$ Hz, 1H), 2.31 (s, 3H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ : 162.5 (d, $^1J_{\text{F,C}}=247.4$ Hz), 142.9, 140.7, 134.1 (d, $^4J_{\text{F,C}}=3.1$ Hz), 133.3 (d, $^3J_{\text{F,C}}=9.4$ Hz), 132.9 (d, $^2J_{\text{P,C}}=9.0$ Hz), 132.5 (d, $^2J_{\text{P,C}}=9.2$ Hz), 132.4 (d, $^3J_{\text{P,C}}=2.8$ Hz), 132.1 (d, $^3J_{\text{P,C}}=2.8$ Hz), 131.7, 130.0, 129.2, 128.5, 128.8, 128.1, 127.8 (d, $^1J_{\text{P,C}}=12.8$ Hz), 127.7 (d, $^1J_{\text{P,C}}=12.8$ Hz), 115.1 (d, $^2J_{\text{F,C}}=21.5$ Hz), 86.6, 78.0, 21.0; ^{11}B NMR (128 MHz, Chloroform-*d*) δ : 1.7 (d, $J=157.6$ Hz, 2B), -1.5 (d, $J=117.4$ Hz, 2B), $-5.3\sim-17.8$ (m, 6B); ^{31}P NMR (162 MHz, Chloroform-*d*) δ : 17.4; ^{19}F NMR (376 MHz, Chloroform-*d*) δ : -113.8 . HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{33}\text{B}_{10}\text{FOP} [\text{M}+\text{H}]^+$: 555.3246; found 555.3251.

1-Diphenylphosphinoyl-2-(*p*-tolyl)-(E)-4-styryl-*o*-carborane (**11**): White solid, 31.38 mg, yield 39%. m.p. 162~163 °C; ¹H NMR (500 MHz, Chloroform-*d*) δ: 7.74~7.66 (m, 2H), 7.61~7.52 (m, 2H), 7.42~7.35 (m, 1H), 7.34~7.31 (m, 1H), 7.31~7.28 (m, 1H), 7.28~7.26 (m, 1H), 7.24~7.22 (m, 1H), 7.22~7.20 (m, 2H), 7.20~7.18 (m, 3H), 7.18~7.13 (m, 3H), 6.81 (d, *J*=7.7 Hz, 2H), 6.72 (d, *J*=18.0 Hz, 1H), 5.62 (d, *J*=17.8 Hz, 1H), 2.28 (s, 3H); ¹³C NMR (126 MHz, Chloroform-*d*) δ: 144.3, 140.6, 137.7, 132.9 (d, ²*J*_{P,C}=9.0 Hz), 132.6 (d, ²*J*_{P,C}=9.4 Hz), 132.4 (d, ³*J*_{P,C}=3.0 Hz), 132.2 (d, ³*J*_{P,C}=3.0 Hz), 131.8, 130.0, 129.1, 128.6, 128.5, 128.2, 127.9, 127.77 (d, ¹*J*_{P,C}=12.8 Hz), 127.76 (d, ¹*J*_{P,C}=12.8 Hz), 127.3, 126.6, 86.6, 77.75 (d, ¹*J*_{P,C}=51.6 Hz), 21.0; ¹¹B NMR (128 MHz, Chloroform-*d*) δ: 1.6 (d, *J*=175.7 Hz, 2B), -1.6 (d, *J*=112.1 Hz, 2B), -4.9~-18.0 (m, 6B); ³¹P NMR (202 MHz, Chloroform-*d*) δ: 18.3. HRMS (ESI) calcd for C₂₉H₃₃B₁₀OP [M+H]⁺: 537.3345; found 537.3349.

1-Diphenylphosphinoyl-2-(*p*-tolyl)-(E)-4-(*p*-chlorostyryl)-*o*-carborane (**12**): White solid, 23.10 mg, yield 27%. m.p. 182~184 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.86~7.75 (m, 2H), 7.65~7.54 (m, 2H), 7.54~7.41 (m, 1H), 7.41~7.35 (m, 1H), 7.35~7.20 (m, 8H), 7.17~7.10 (m, 2H), 6.95~6.84 (m, 2H), 6.71 (d, *J*=17.7 Hz, 1H), 5.68 (d, *J*=17.8 Hz, 1H), 2.35 (s, 3H); ¹³C NMR (101 MHz, Chloroform-*d*) δ: 142.7, 140.7, 136.3, 133.4, 132.9 (d, ¹*J*_{P,C}=8.9 Hz), 132.5 (d, ¹*J*_{P,C}=9.1 Hz), 132.4 (d, ¹*J*_{P,C}=2.4 Hz), 132.1 (d, ¹*J*_{P,C}=2.5 Hz), 131.7, 129.9, 129.1, 128.8, 128.5, 128.4, 127.81 (d, ¹*J*_{P,C}=12.6 Hz), 127.79, 127.80 (d, ¹*J*_{P,C}=12.6 Hz), 127.2, 86.6, 77.9, 21.0; ¹¹B NMR (128 MHz, Chloroform-*d*) δ: 1.8 (d, *J*=166.4 Hz, 2B), -1.4 (d, *J*=108.8 Hz, 2B), -4.6~-16.6 (m, 6B); ³¹P NMR (162 MHz, Chloroform-*d*) δ: 17.2. HRMS (ESI) calcd for C₂₉H₃₃B₁₀ClOP [M+H]⁺: 571.2955; found 571.2948.

1-Diphenylphosphinoyl-2-(*p*-tolyl)-(E)-4-(*p*-methylstyryl)-*o*-carborane (**13**): White solid, 34.67 mg, yield 42%. m.p. 109~111 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.75~7.67 (m, 2H), 7.64~7.56 (m, 2H), 7.46~7.40 (m, 1H), 7.39~7.32 (m, 1H), 7.30~7.17 (m, 6H), 7.16~7.03 (m, 4H), 6.84 (d, *J*=7.2 Hz, 2H), 6.74 (d, *J*=17.9 Hz, 1H), 5.57 (d, *J*=18.0 Hz, 1H), 2.36 (s, 1H), 2.31 (s, 1H); ¹³C NMR (101 MHz, Chloroform-*d*) δ: 144.4, 140.6, 137.8, 135.1, 133.0 (d, ²*J*_{P,C}=8.9 Hz), 132.7 (d, ²*J*_{P,C}=9.1 Hz), 132.4 (d, ³*J*_{P,C}=2.6 Hz), 132.2 (d, ³*J*_{P,C}=2.8 Hz), 131.8, 130.2, 129.1, 129.0, 128.4, 127.76 (d, ¹*J*_{P,C}=12.8 Hz), 127.75 (d, ¹*J*_{P,C}=12.8 Hz), 127.5, 127.4, 126.6, 86.7, 77.8 (d, ¹*J*_{P,C}=51.5 Hz), 21.3, 21.0; ¹¹B NMR (128 MHz, Chloroform-*d*) δ: 1.7 (d, *J*=136.5 Hz, 2B), -1.2 (d, *J*=111.5 Hz, 2B), -3.2~-14.3 (m, 6B); ³¹P NMR (162 MHz, Chloroform-*d*) δ: 18.0. HRMS (ESI) calcd for C₃₀H₃₆B₁₀OP [M+H]⁺: 551.3501; found 551.3497.

Supporting Information Crystal data of **3** and NMR spectra of **3**~**13** are available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Liu, L. J.; Mak T, C. W.; Zang, S. Q. *Chin. J. Chem.* **2021**, *39*, 81.
(b) Kolel-Veetil, M. K.; Beckham, H. W.; Keller, T. M. *Chem. Mater.* **2004**, *16*, 3162.
(c) Gao, Y.; Cui, P. F.; Aznarez, F.; Jin, G. X., *Chem.-Eur. J.* **2018**, *24*, 10357.
(d) Armstrong, A. F.; Valliant, J. F. *Dalton Trans.* **2007**, *38*, 4240.
- [2] (a) Cui, P.-F.; Lin, Y. J.; Li, Z. H.; Jin, G.-X. *J. Am. Chem. Soc.* **2020**, *142*, 8532.
(b) Guo, S.-T.; Cui, P.-F.; Yuan, R.-Z.; Jin, G.-X. *Chem. Commun.* **2021**, *57*, 2412.
- [3] (a) Xiong, H. J.; Li, Y. R.; Ye, H. H.; Huang, G.; Zhou, D. F.; Huang, Y. B. *J. Mater. Chem. B* **2020**, *8*, 10309.
(b) Zhang, J.; Liu, K.; Liu, Z. S.; Wang, Z. L.; Hua, C. X.; Liu, T. H.; Fang, Y. *ACS Appl. Mater. Inter.* **2021**, *13*, 5625.
(c) Yao, Z. J.; Jin, Y.-X.; Deng, W.; Liu, Z.-J. *Inorg. Chem.* **2021**, *60*, 2756.
- [4] Sarosi, M.-B.; Neumann, W.; Lybrand, T. P.; Hey-Hawkins, E. J. *Chem. Inf. Model.* **2017**, *57*, 2056.
- [5] (a) Zhang, J.; Xie, Z. W. *Chin. J. Chem.* **2018**, *36*, 1041.
(b) Chan, T. L.; Xie, Z. *Chem. Sci.* **2018**, *9*, 2284.
(c) Guo, J. X.; Liu, D. Q.; Zhang, J. H.; Zhang, J. J.; Miao, Q.; Xie, Z. W. *Chem. Commun.* **2015**, *51*, 12004.
(d) Tu, D. S.; Cai, S. Z.; Fernandez, C.; Ma, H. L.; Wang, X.; Wang, H.; Ma, C. Q.; Yan, H.; Lu, C. S.; An, Z. F. *Angew. Chem. Int. Ed.* **2019**, *58*, 9129.
(e) Wei, X.; Zhu, M. J.; Cheng, Z.; Lee, M.; Yan, H.; Lu, C. S.; Xu, J. J. *Angew. Chem. Int. Ed.* **2019**, *58*, 3162.
- [6] (a) Wang, Z. J.; Zhao, J. W.; Muddassir, M.; Guan, R. F.; Tao, S. L. *Inorg. Chem.* **2021**, *60*, 4705.
(b) Li, X.; Zhou, Q.; Zhu, M.; Chen, W.; Wang, B. B.; Sha, Y.; Yan, H. *Chem.-Asian J.* **2021**, *7*, 757.
(c) Ochi, J.; Tanaka, K.; Chujo, Y. *Angew. Chem. Int. Ed.* **2020**, *25*, 9841.
(d) Nishino, K.; Tanaka, K.; Chujo, Y. *Asian J. Org. Chem.* **2019**, *12*, 2228.
- [7] (a) Chen, Y.; Au, Y. K.; Quan, Y. J.; Xie, Z. W. *Sci. China Chem.* **2019**, *62*, 74.
(b) Zheng, F. R.; Leung, T. F.; Chan, K. W.; Sung, H. H. Y.; Williams, I. D.; Xie, Z. W.; Jia, G. C. *Chem. Commun.* **2016**, *52*, 10767.
(c) Lyu, H. R.; Quan, Y. J.; Xie, Z. W. *Angew. Chem. Int. Ed.* **2015**, *54*, 10623.
(d) Zhao, D.; Zhang, J. J.; Xie, Z. W. *Chem.-Eur. J.* **2015**, *21*, 10334.
- [8] (a) Kürti, L.; Czákó, B. *Strategic Applications of Named Reactions in Organic Synthesis: Background and Detailed Mechanisms*, Elsevier Academic Press, Burlington, **2007**, p. 196.
(b) Huang, X. L.; Teng, S. H.; Chi, Y. R.; Xu, W. Q.; Pu, M. P.; Wu, Y. D.; Zhou, J. S. *Angew. Chem. Int. Ed.* **2020**, *60*, 2828.
(c) Peng, X. P.; Yang, Y.; Luo, B. L.; Wen, S. J.; Huang, P. *Adv. Syn. Catal.* **2021**, *363*, 222.
(d) Ju, B. H.; Chen, S. G.; Kong, W. Q. *Org. Lett.* **2019**, *21*, 9343.
(e) Zhou, L. X.; Shi, Y. X.; Zhu, X. Y.; Zhang, P. *Tetrahedron Lett.* **2019**, *60*, 2005.
(f) Xie, J.-Q.; Liang, R.-X.; Jia, Y.-X. *Chin. J. Chem.* **2021**, *3*, 710.
(g) Yang, P.; Xu, R.-Q.; Zheng, C.; You, S.-L. *Chin. J. Chem.* **2020**, *3*, 235.
(h) Liu, C. X.; Liu, G. Y.; Zhao, H. K. *Chin. J. Chem.* **2016**, *10*, 1048.
- [9] (a) Wen, Z. K.; Zhao, Z. K.; Wang, N. J.; Chen, Z. L.; Chao, J. B.; Feng, L. H. *Org. Lett.* **2019**, *21*, 9545.
(b) Bettadapur, K. R.; Sherikar, M. S.; Lanke, V.; Prabhu, K. R. *Asian J. Org. Chem.* **2018**, *7*, 1338.
- [10] Grimes, R. N. *Carboranes*, 3rd ed., Elsevier, Oxford, **2016**.
- [11] (a) Wu, J.; Cao, K.; Xu, T.-T.; Zhang, X.-J.; Jiang, L.; Yang, J. X.;

- Huang, Y. W. *RSC Adv.* **2015**, *5*, 91683.
- (b) Lyu, H. R.; Quan, Y. J.; Xie, Z. W. *Angew. Chem. Int. Ed.* **2015**, *54*, 10623.
- (c) Zhang, C. Y.; Wang, Q.; Tian, S.; Zhang, J. W.; Li, J. Y.; Zhou, L.; L, J. *Org. Biomol. Chem.* **2020**, *18*, 4723.
- (d) Wang, Q.; Tian, S.; Zhang, C. Y.; Li, J. W.; Wang, Z. X.; Du, Y. M.; Zhou, L.; L, J. *Org. Lett.* **2019**, *21*, 8018.
- [12] (a) Quan, Y. J.; Xie, Z. W. *J. Am. Chem. Soc.* **2014**, *136*, 15513.
- (b) Quan, Y. J.; Xie, Z. W. *Angew. Chem. Int. Ed.* **2016**, *55*, 1295.
- (c) Lyu, H. R.; Quan, Y. J.; Xie, Z. W. *Angew. Chem. Int. Ed.* **2016**, *128*, 12019.
- (d) Lyu, H. R.; Quan, Y. J.; Xie, Z. W. *J. Am. Chem. Soc.* **2016**, *138*, 12727.
- (e) Lyu, H. R.; Quan, Y. J.; Xie, Z. W. *Chem.-Eur. J.* **2017**, *23*, 14866.
- (f) Baek, Y.; Kim, S. H.; Son, J.-Y.; Lee, K. Y.; Kim, D. W.; Lee, P. H. *ACS Catal.* **2019**, *9*, 10418.
- [13] (a) Xu, T.-T.; Cao, K.; Zhang, C.-Y.; Wu, J.; Ding, L.-F.; Yang, J. X. *Org. Lett.* **2019**, *21*, 9276.
- (b) Xu, T.-T.; Cao, K.; Zhang, C.-Y.; Wu, J.; Jiang, L. H.; Yang, J. X. *Chem. Commun.* **2018**, *54*, 13603.
- (c) Liang, Y.-F.; Yang, L.; Jei, B. B.; Kuniyila, R.; Ackermann, L. *Chem. Sci.* **2020**, *11*, 10764.
- (d) Lyu, H. R.; Zhang, J.; Yang, J. T.; Quan, Y. J.; Xie, Z. W. *J. Am. Chem. Soc.* **2019**, *141*, 4219.
- [14] Lian, L. X.; Lin, C. X.; Yu, Y.; Ye, K.-Y.; Yuan Y. F. *Tetrahedron Lett.* **2020**, *61*, 152625.
- [15] Popescu, A.-R.; Laromaine, A.; Teixidor, F.; Sillanpää, R.; Kivekäs, R.; Llambias, J. I.; Viñas, C. *Chem.-Eur. J.* **2011**, *17*, 4429.

(Fan, Y.)