无配体条件钯催化内炔的氢膦酰化反应合成(*E*)-烯基膦酸酯类化合物

许振丽^a 张宗雪^b 孟晨湘^b 张萧雅^b 许凯^{*,b} 刘澜涛^{b,c}
王涛^b 徐海云^b 毛国梁^{*,a}

(^a 东北石油大学化学化工学院 黑龙江大庆 163318)

(^b 商丘师范学院化学化工学院 药物绿色合成河南省工程实验室 河南商丘 476000)

(^c 郑州大学化学与分子工程学院 郑州 450001)

摘要 报道了在不同醇中无配体条件下钯催化二苯基 H-亚膦酸酯对内炔的氢膦酰化反应合成 *E*-烯基膦酸酯类化合物的方法。这种实用且通用的三组分反应涉及二苯基 H-亚膦酸酯对相对惰性内炔的顺式加成, 以及伴随的二苯基 H-亚膦酸酯的酯交换过程。反应具有高立体选择性、宽泛的底物范围和无需额外配体等显著特点。

关键词 加成反应; 三组分反应; 烯基膦酸酯; 钯; 内炔

Ligand-Free Pd-Catalyzed Hydrophosphorylation of Internal Alkynes for the Synthesis of *E*-Vinylphosphonates

Xu, Zhenli^a Zhang, Zongxue^b Meng, Chenxiang^b Zhang, Xiaoya^b Xu, Kai^{*,b}
Liu, Lantao^{b,c} Wang, Tao^b Xu, Haiyun^b Mao, Guoliang^{*,a}

(^a College of Chemistry and Chemical Engineering, Northeast Petroleum University, Daqing, Heilongjiang 163318)

(^b Henan Engineering Laboratory of Green Synthesis for Pharmaceuticals, College of Chemistry and Chemical Engineering, Shangqiu Normal University, Shangqiu, Henan 476000)

(^c College of Chemistry and Molecular Engineering, Zhengzhou University, Zhengzhou 450001)

Abstract A ligand-free Pd-catalyzed hydrophosphorylation of internal alkynes with diphenyl H-phosphonate in various alcohols for the synthesis of *E*-vinylphosphonates has been reported. This practical and versatile three-component reaction involves the *syn*-addition of diphenyl H-phosphonate towards relatively inactive internal alkynes, as well as the concomitant transesterification of diphenyl H-phosphonate. The remarkable features include high stereoselectivity, wide substrate scope and the absence of additional ligands.

Keywords addition reaction; three-component reaction; vinylphosphonate; palladium; internal alkyne

1 Introduction

Vinylphosphonates as an important class of organophosphorus compounds have wide application in the fields of organometallic chemistry, pharmaceuticals and agrochemistry.^[1] The typical approaches to construct C(sp²)-P bond in vinylphosphonates could be basically classified into Michaelis-Arbuzov- and Hirao-type coupling^[2] as well as Tanaka-type addition.^[3] In detail, although the former can

facilely deliver the stereospecific products by the starting materials, the latter has the advantage of high atomic economy and several brilliant works have emerged including phosphinic acid induced reversal of regioselectivity in Pd-catalyzed the formation of C(sp²)-P bond,^[3b] acid-free Ni/DPPE-catalyzed hydrophosphorylation of alkynes,^[3f] oxapalladacycle-mediated hydrophosphorylation of terminal alkynes,^[3g] Ni/DIBAL-catalyzed synthesis of mono- and bisphosphonates,^[3h] systematic and comprehensive palla-

* Corresponding authors. E-mail: liult05@iccas.ac.cn; xuk1008@163.com

Received March 22, 2021; revised April 13, 2021; published online May 7, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21572126, 21202095), the Technicians Troop Construction Projects of Henan Province (No. C20150030), the Program of Science and Technology Innovation Talents of Henan Province (No. 184100510011), the Natural Science Foundation of Henan Province (No. 212300410379), and the Key Scientific and Technological Project of Henan Province (No. 192102110222).

国家自然科学基金(Nos. 21572126, 21202095)、河南省技术人员队伍建设(No. C20150030)、河南省科技创新人才计划(No. 184100510011)、河南省自然科学基金(No. 212300410379)、河南省科技重大专项(No.192102110222)资助项目。

dium-catalyzed hydrophosphorylation of alkynes,^[3j] as well as solvent-free Ni-catalyzed hydrophosphonylation of alkynes with trialkyl phosphites^[3i] (Schemes 1a and 1b). Despite these significant advances, there are still limitations involving the low stereoselectivity, incompatibility to inactive internal alkynes and poor diversity for the desired vinylphosphonates. On the other hand, Stawinski and co-workers^[4] did pioneering work concerning transesterification of diphenyl H-phosphonate with an appropriate alcohol, which could easily provide an effective access for the generation of a wide range of dialkyl and alkyl phenyl H-phosphonates. The application and development of this method still have great potential and preparation significance^[5] on account of commercially available starting materials, simple operation and satisfactory efficiency.

Previous reports:

This work:

Scheme 1 Hydrophosphorylation of alkynes for vinylphosphonates

According to the mentioned above and our reports in related realm,^[6] we envisioned that transition-metal catalyzed three-component reaction involving inactive internal alkynes, diphenyl H-phosphonate and matching alcohols with higher stereoselectivity and better diversity could be achieved (Scheme 1c).

2 Results and discussion

Initially, the three-component reaction involving 1,2-diphenylethyne, diphenyl H-phosphonate and ethanol was performed as the model reaction. Especially, the ethanol served as both reactant and solvent. In the presence of Pd(OAc)₂ as the catalyst, both inorganic and organic bases were carefully screened^[7], and ⁿBu₃N gave the best results affording the stereospecific *E*-**4a** with an isolated yield of 85% (Table 1, Entries 1~7). Since the *E*-configuration of **4a** evidenced by the coupling constant of ³J(P,H)=24.8 Hz conforms to the previous reports^[3f] (see the Supporting Information for more details), it could be inferred that the *syn*-addition of diphenyl H-phosphonate to 1,2-diphenylethyne in ethanol was involved in this hydrophosphorylation process logically. In addition, either decrease of the reaction temperature or replacement of Pd(OAc)₂ with PdCl₂ or NiCl₂ could not further improve the efficiency

(Table 1, Entries 8~10). Unfortunately, when the solvent switched to the bulkier isopropanol, the reaction was significantly suppressed (Table 1, Entry 11). Therefore, it was still necessary to optimize the reaction condition again (Table 1, Entries 11~20). Upon a similar examination of various bases, K₂CO₃ showed higher reactivity and a yield of 79% was obtained (Table 1, Entries 11~17). Other metal salts such as PdCl₂ and NiCl₂ had a negative effect on this hydrophosphorylation again (Table 1, Entries 18~19). In order to improve the solubility of reactants, tetrabutylammonium bromide (TBAB) was added and the stereospecific *E*-**4l** was obtained with a yield of 83%.

Table 1 Optimization of reaction condition^a

Entry	Catalyst	Base	Additive	Solvent	Yield ^b /%
1	Pd(OAc) ₂	KOH	No	EtOH	43
2	Pd(OAc) ₂	K ₂ CO ₃	No	EtOH	18
3	Pd(OAc) ₂	K ₃ PO ₄	No	EtOH	63
4	Pd(OAc) ₂	KF	No	EtOH	67
5	Pd(OAc) ₂	Pyridine	No	EtOH	<5
6	Pd(OAc) ₂	NEt ₃	No	EtOH	68
7	Pd(OAc)₂	ⁿBu₃N	No	EtOH	85
8 ^c	Pd(OAc) ₂	ⁿ Bu ₃ N	No	EtOH	78
9	PdCl ₂	ⁿ Bu ₃ N	No	EtOH	43
10	NiCl ₂	ⁿ Bu ₃ N	No	EtOH	<5
11	Pd(OAc) ₂	ⁿ Bu ₃ N	No	ⁱ PrOH	17
12	Pd(OAc) ₂	DBU	No	ⁱ PrOH	<5
13	Pd(OAc) ₂	KOH	No	ⁱ PrOH	67
14	Pd(OAc) ₂	K ₂ CO ₃	No	ⁱ PrOH	79
15	Pd(OAc) ₂	KOAc	No	ⁱ PrOH	48
16	Pd(OAc) ₂	K ₃ PO ₄	No	ⁱ PrOH	64
17	Pd(OAc) ₂	KF	No	ⁱ PrOH	66
18	PdCl ₂	K ₂ CO ₃	No	ⁱ PrOH	24
19	NiCl ₂	K ₂ CO ₃	No	ⁱ PrOH	<5
20	Pd(OAc)₂	K₂CO₃	TBAB	ⁱPrOH	83

^a Reagents and conditions: diphenylacetylene (0.4 mmol), diphenyl H-phosphonate (0.8 mmol), catalyst (5 mol%), base (0.8 mmol) and additive (0.4 mmol) in ROH (1 mL) at 100 °C under air for 16 h. ^b Isolated yield based on the amount of diphenylacetylene. ^c At 80 °C.

The scope of internal alkynes and alcohols in this three-component reaction was evaluated under conditions A and B, respectively, and the selective results were shown in Table 2. As expected, various *E*-vinylphosphonates were afforded not only evidenced by the coupling constant of ³J(P,H) (21.2~25.2 Hz, see the Supporting Information for more details) but also unambiguously determined by

X-ray analysis of **4u**. In general, when the less bulkier alcohols served as the candidates, the *syn*-hydrophosphorylation was conducted under the standard condition A (**4a**~**4k**), while when the more bulkier alcohols served as the candidates, the *syn*-hydrophosphorylation was conducted under the standard condition B (**4l**~**4u**). In ethanol, symmetrical diarylacetylenes bearing electron-donating and electron-neutral groups readily afforded the corresponding (*E*)-diethyl phosphonates with yields ranging from 66% to 99% (**4a**~**4e**), while symmetrical diarylacetylenes containing electron-withdrawing groups displayed relatively lower reactivity (**4f**~**4h**). Unsymmetrical diarylacetylenes exemplified as prop-1-yn-1-ylbenzene could also survive in this reaction and a yield of 67% was obtained (**4i**). Surprisingly, 2-fluoroethanol and 2-methoxyethanol could also amenable to this transformation

(**4j**~**4k**). Meanwhile, the steric hindrance effect had little influence and comparable yields could be obtained (**4c** vs **4d**, **4a** vs **4e**). Similarly, in isopropanol, various (*E*)-diisopropyl phosphonates originated from symmetrical diarylacetylenes were obtained with yields ranging from 57% to 90% (**4l**~**4r**). The more bulkier alcohols such as 2-methylpropan-1-ol, cyclohexanol and cyclopentanol were also effective candidates for this *syn*-hydrophosphorylation and moderate to good yields were obtained (**4s**~**4u**). Meanwhile, the compatibility of other substrates involving both alkynes and alcohols was investigated individually (see the Supporting Information for more details). Moreover, the scale-up processes for the preparation of **4a** and **4l** were also performed, and comparable yield for **4a** was obtained, while a slightly lower yield for **4l** was observed (Scheme 2).

Table 2 Substrate scope

$\text{R}^1\text{---}\text{C}\equiv\text{C---Ar} \quad \text{1} + \quad \text{H---P(=O)(OR}^2\text{)}_2 \quad \text{2} \xrightarrow[\text{R}^2\text{OH (3), 100 }^\circ\text{C, air, 16 h}]{\text{Pd(OAc)}_2, \text{ base, additive}} \quad \text{O=P(OR}^2\text{)}_2\text{---C(R}^1\text{)=C(Ar)} \quad \text{4}$	
Standard condition A	
 4a 85%	 4b 90%
 4c 99%	 4d 93%
 4e 97%	
 4f 89%	 4g 66%
 4h 70%	 4i 67%
 4j 40%	 4k 40%
Standard condition B	
 4l 83%	 4m 78%
 4n 70%	 4o 90%
 4p 87%	 4q 81%
 4r 57%	 4s 78%
 4t 45%	 4u 81%
 (deposition number: 2063359)	

^a Standard condition A: internal alkyne (0.4 mmol), diphenyl H-phosphonate (0.8 mmol), Pd(OAc)₂ (5 mol%) and ^tBu₃N (0.8 mmol) in R²OH (1 mL) at 100 °C under air for 16 h. Yield of isolated product based on the amount of internal alkyne. ^b Reagents and condition B: with K₂CO₃ as base and TBAB as additive based on the above standard conditions.

Scheme 2 Scale-up experiments

Scheme 3 Proposed mechanism

Based on the aforementioned results and the related reports,^[4-5,3c,3i,3j] a possible mechanism for the *syn*-hydrophosphorylation was proposed as depicted in Scheme 3. Firstly, the transesterification of diphenyl H-phosphonate (**2**) with various alcohols (**3**) in the presence of appropriate base took place and resulted in the formation of the corresponding dialkyl H-phosphonate (**A**). The other tautomeric form **B** was readily generated and underwent the reduction of Pd^{II} to Pd⁰ species. The oxidative addition of **A** to Pd^{II}-H complex (**C**), which inserted into the diarylacetylenes in a *syn*-addition mode delivering the adduct **D**. Finally, the reductive elimination of **D** afforded the *E*-vinylphosphonates (**4**).

3 Conclusions

In conclusion, a practical and versatile method for the synthesis of *E*-vinylphosphonates via ligand-free Pd-catalyzed *syn*-hydrophosphorylation and transesterification of diphenyl H-phosphonate has been developed. The coupling constant of ³J(P,H) and the characterized X-ray analysis jointly confirmed the *E*-configuration of the desired products. This three-component reaction involving inactive

internal alkynes, diphenyl H-phosphonate and alcohols had the advantages of high stereoselectivity, good diversity, moderate to excellent yields. Further exploration of the substrate scope and synthetic applications of these methodologies are currently under way in our laboratory.

4 Experimental section

4.1 General information

¹H NMR, ¹³C NMR, ³¹P NMR and ¹⁹F NMR spectra were recorded on a Bruker DPX-400 spectrometer at room temperature with CDCl₃ as the solvent and TMS as an internal standard. High-resolution mass spectra were collected on a MALDI-FTMS. Melting points were measured using a XT4A microscopic apparatus. IR spectra were obtained on a Bruker VECTOR22 spectrophotometer in KBr pellets. X-ray structural analysis was conducted on the XtaLAB mini. Internal alkynes were synthesized according to the reported literature.^[8] Solvents and chemicals were bought from commercial sources and used directly unless otherwise stated.

4.2 General procedure for the synthesis of vinylphosphonates

Standard conditions A: A solution of internal alkyne (0.4 mmol), diphenyl H-phosphonate (0.8 mmol), Pd(OAc)₂ (5 mol%) and ⁿBu₃N (0.8 mmol) in R²OH (1 mL) was stirred in a 10 mL well-sealed thick-walled glass tube at 100 °C for 16 h. Then the reaction mixture was cooled down and filtered over a plug of silica gel (washed with EtOAc or CH₂Cl₂) directly. The resulting filtrate was concentrated and purified by flash chromatography over silica gel (ethyl acetate/hexane) to give the pure product.

Standard conditions B: A solution of internal alkyne (0.4 mmol), diphenyl H-phosphonate (0.8 mmol), Pd(OAc)₂ (5 mol%), K₂CO₃ (0.8 mmol) and TBAB (0.4 mmol) in R²OH (1 mL) was stirred in a 10 mL well-sealed thick-walled glass tube at 100 °C for 16 h. Then the reaction mixture was cooled down and filtered over a plug of silica gel (washed with EtOAc or CH₂Cl₂) directly. The resulting filtrate was concentrated and purified by flash chromatography over silica gel (ethyl acetate/hexane) to give the pure product.

(*E*)-Diethyl (1,2-diphenylvinyl)phosphonate (**4a**):^[9] Pale yellow oil, yield 85%. ¹H NMR (400 MHz, CDCl₃) δ : 7.63 (d, *J*=24.8 Hz, 1H), 7.39~7.24 (m, 5H), 7.22~7.12 (m, 3H), 7.07~7.04 (m, 2H), 4.17~4.03 (m, 4H), 1.30~1.25 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 143.7 (d, *J*=11.0 Hz), 135.6 (d, *J*=7.6 Hz), 134.7 (d, *J*=22.5 Hz), 131.0 (d, *J*=178.0 Hz), 130.4 (d, *J*=0.9 Hz), 129.2 (d, *J*=5.3 Hz), 129.0, 128.9 (d, *J*=1.4 Hz), 128.2, 127.8 (d, *J*=2.4 Hz), 62.2, 62.2, 16.3, 16.3; ³¹P NMR (CDCl₃, 163 MHz) δ : 18.1; IR (KBr) ν : 3430, 3055, 3023, 2975, 2927, 1611, 1486, 1444, 1386, 1236, 1159, 1015, 957, 791, 695, 618 cm⁻¹. HRMS (ESI) calcd for C₁₈H₂₂O₃P [M+H]⁺ 317.1301, found 317.1302.

(*E*)-Diethyl (1,2-di-*p*-tolylvinyl)phosphonate (**4b**): Pale yellow oil, yield 90%. ¹H NMR (400 MHz, CDCl₃) δ : 7.58 (d, *J*=24.4 Hz, 1H), 7.15 (s, 4H), 6.97 (s, 4H), 4.10 (s, 4H), 2.37 (s, 3H), 2.26 (s, 3H), 1.28 (s, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 143.6 (d, *J*=11.6 Hz), 139.2, 137.4 (d, *J*=2.6 Hz), 132.7 (d, *J*=7.8 Hz), 132.0 (d, *J*=22.6 Hz), 130.4, 129.7 (d, *J*=178.1 Hz), 129.6 (d, *J*=1.2 Hz), 129.0 (d, *J*=5.2 Hz), 128.9, 62.1, 62.0, 21.3, 21.3, 16.4, 16.3; ³¹P NMR (CDCl₃, 163 MHz) δ : 18.8; IR (KBr) 3443, 2985, 2917, 1611, 1505, 1447, 1390, 1236, 1025, 964, 813, 727, 589 cm⁻¹. HRMS (ESI) calcd for C₂₀H₂₆O₃P [M+H]⁺ 345.1614, found 345.1615.

(*E*)-Diethyl (1,2-bis(4-methoxyphenyl)vinyl)phosphonate (**4c**): Pale yellow oil, yield 99%. ¹H NMR (400 MHz, CDCl₃) δ : 7.54 (d, *J*=24.4 Hz, 1H), 7.19 (d, *J*=6.4 Hz, 2H), 7.04 (d, *J*=7.2 Hz, 2H), 6.91 (d, *J*=6.8 Hz, 2H), 6.69 (d, *J*=7.2 Hz, 2H), 4.09 (s, 4H), 3.84 (s, 3H), 3.75 (s, 3H), 1.30~1.26 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 160.1, 159.1 (d, *J*=2.3 Hz), 143.3 (d, *J*=12.2 Hz), 132.0, 130.5 (d, *J*=5.2 Hz), 127.9 (d, *J*=8.0 Hz), 127.6 (d, *J*=179.9 Hz), 127.5 (d, *J*=23.2 Hz), 114.4 (d, *J*=5.2 Hz), 113.6, 62.0, 62.0, 55.2, 55.2, 16.4, 16.3; ³¹P NMR (CDCl₃, 163 MHz) δ : 19.2; IR (KBr) ν : 3452, 2978, 2933, 2905, 2837, 1604, 1511, 1460, 1287, 1249, 1178, 1028, 983, 829, 768, 730, 592 cm⁻¹. HRMS (ESI) calcd for C₂₀H₂₆O₅P [M+H]⁺ 377.1512, found 377.1514.

(*E*)-Diethyl (1,2-bis(2-methoxyphenyl)vinyl)phosphonate (**4d**): Pale yellow oil, yield 93%. ¹H NMR (400 MHz, CDCl₃) δ : 8.00 (d, *J*=24.8 Hz, 1H), 7.30~7.25 (m, 1H), 7.22~7.13 (m, 2H), 6.96~6.91 (m, 1H), 6.86~6.77 (m, 3H), 6.58~6.53 (m, 1H), 4.15~4.06 (m, 4H), 3.84 (s, 3H), 3.61 (s, 3H), 1.29~1.25 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 157.8, 156.9 (d, *J*=5.5 Hz), 139.6 (d, *J*=12.3 Hz), 130.8 (d, *J*=4.6 Hz), 130.1, 129.2 (d, *J*=2.4 Hz), 128.9 (d, *J*=2.2 Hz), 126.2 (d, *J*=181.1 Hz), 124.7, 124.6 (d, *J*=31.3 Hz), 120.8 (d, *J*=2.0 Hz), 119.8, 111.0, 110.5, 62.0, 61.9, 55.5, 55.3, 16.3, 16.2; ³¹P NMR (CDCl₃, 163 MHz) δ : 19.0; IR (KBr) 3443, 3058, 2985, 2937, 2901, 2831, 1723, 1591, 1489, 1460, 1438, 1390, 1294, 1252, 1165, 1111, 1050, 1018, 970, 836, 752, 618, 521 cm⁻¹. HRMS (ESI) calcd for C₂₀H₂₆O₅P [M+H]⁺ 377.1512, found 377.1513.

(*E*)-Diethyl (1,2-di(naphthalen-1-yl)vinyl)phosphonate (**4e**): White solid, m.p. 111~112 °C, yield 97%. ¹H NMR

(400 MHz, CDCl₃) δ 8.63 (d, *J*=24.4 Hz, 1H), 8.29 (d, *J*=8.4 Hz, 1H), 7.91 (d, *J*=8.4 Hz, 1H), 7.77~7.72 (m, 3H), 7.62~7.22 (m, 7H), 6.94~6.85 (m, 2H), 4.19~4.01 (m, 4H), 1.28 (t, *J*=7.1 Hz, 3H), 1.11 (t, *J*=7.1 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ : 142.9 (d, *J*=10.9 Hz), 133.6 (d, *J*=1.0 Hz), 133.3, 132.8 (d, *J*=7.4 Hz), 131.7 (d, *J*=179.3 Hz), 131.6 (d, *J*=16.9 Hz), 131.6, 131.5, 129.1, 128.6, 128.2, 128.2, 127.6 (d, *J*=5.4 Hz), 126.7 (d, *J*=2.4 Hz), 126.6, 126.0, 125.9, 125.8, 125.8, 125.3 (d, *J*=2.4 Hz), 124.9, 123.8, 62.6 (d, *J*=6.4 Hz), 62.4 (d, *J*=6.1 Hz), 16.4 (d, *J*=6.2 Hz), 16.2 (d, *J*=6.3 Hz); ³¹P NMR (CDCl₃, 163 MHz) δ : 17.3; IR (KBr) ν : 3423, 3046, 2975, 2930, 2904, 1934, 1736, 1614, 1588, 1505, 1470, 1393, 1236, 1159, 1063, 1021, 957, 800, 772, 643 cm⁻¹. HRMS (ESI) calcd for C₂₆H₂₆O₃P [M+H]⁺ 417.1614, found 417.1616.

(*E*)-Diethyl (1,2-bis(4-chlorophenyl)vinyl)phosphonate (**4f**): Pale yellow oil, yield 89%. ¹H NMR (400 MHz, CDCl₃) δ : 7.58 (d, *J*=24.0 Hz, 1H), 7.35 (d, *J*=6.8 Hz, 2H), 7.17 (dd, *J*=14.6 Hz and 7.8 Hz, 4H), 6.99 (d, *J*=6.8 Hz, 2H), 4.11 (d, *J*=3.2 Hz, 4H), 1.30~1.27 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 142.7 (d, *J*=44.4 Hz), 135.1, 134.1 (d, *J*=2.8 Hz), 133.7 (d, *J*=7.8 Hz), 132.8 (d, *J*=22.3 Hz), 131.5, 130.7 (d, *J*=180.0 Hz), 130.6 (d, *J*=5.4 Hz), 129.3 (d, *J*=1.7 Hz), 128.6, 62.4, 62.3, 16.4, 16.3; ³¹P NMR (CDCl₃, 163 MHz) δ : 17.1; IR (KBr) ν : 3462, 2985, 2930, 2905, 1617, 1588, 1495, 1393, 1242, 1165, 1092, 1050, 1021, 983, 823, 746, 669 cm⁻¹. HRMS (ESI) calcd for C₁₈H₂₀Cl₂O₃P [M+H]⁺ 385.0522, found 385.0523.

(*E*)-Diethyl (1,2-bis(4-(trifluoromethyl)phenyl)vinyl)phosphonate (**4g**): Pale yellow oil, yield 66%. ¹H NMR (400 MHz, CDCl₃) δ : 7.73~7.62 (m, 3H), 7.46~7.37 (m, 4H), 7.13 (d, *J*=7.6 Hz, 2H), 4.16~4.13 (m, 4H), 1.33~1.28 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 142.6 (d, *J*=10.3 Hz), 139.0 (d, *J*=8.0 Hz), 137.6 (d, *J*=21.7 Hz), 132.9 (d, *J*=179.1 Hz), 130.8 (q, *J*=32.6 Hz), 130.3 (q, *J*=32.7 Hz), 130.3, 129.6 (d, *J*=5.2 Hz), 125.9 (q, *J*=2.2 Hz), 125.3 (q, *J*=3.9 Hz), 123.9 (q, *J*=272.8 Hz), 123.7 (q, *J*=271.8 Hz), 62.6, 62.6, 16.3, 16.3; ³¹P NMR (CDCl₃, 163 MHz) δ : 16.1; ¹⁹F NMR (CDCl₃, 376 MHz) δ : -62.6, -63.0; IR (KBr) ν : 3446, 2985, 2933, 2905, 1614, 1406, 1329, 1246, 1172, 1127, 1066, 1018, 983, 836, 647, 608 cm⁻¹. HRMS (ESI) calcd for C₂₀H₂₀F₆O₃P [M+H]⁺ 453.1049, found 453.1050.

(*E*)-diethyl (1,2-di(thiophen-2-yl)vinyl)phosphonate (**4h**): Pale yellow solid, mp 64~65 °C, yield 70%. ¹H NMR (400 MHz, CDCl₃) δ : 7.93 (d, *J*=21.2 Hz, 1H), 7.52~7.49 (m, 1H), 7.31~7.28 (m, 1H), 7.23~7.21 (m, 1H), 7.15~7.12 (m, 1H), 7.01~6.95 (m, 2H), 4.16~4.08 (m, 4H), 1.34~1.29 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ : 140.7 (d, *J*=14.9 Hz), 137.9 (d, *J*=23.2 Hz), 134.5 (d, *J*=8.0 Hz), 133.8, 130.9, 128.6 (d, *J*=6.4 Hz), 128.0 (d, *J*=3.3 Hz), 127.8 (d, *J*=2.1 Hz), 126.6 (d, *J*=1.0 Hz), 119.8 (d, *J*=189.5 Hz), 62.4, 62.3, 16.3, 16.3; ³¹P NMR (CDCl₃, 163 MHz) δ : 16.5; IR (KBr) ν : 3420, 3065, 2985, 2898, 1723, 1604, 1524, 1415, 1383, 1230, 1050, 1018, 966, 848, 749, 695, 627 cm⁻¹. HRMS (ESI) [M+

$\text{H}]^+$ calcd for $\text{C}_{14}\text{H}_{18}\text{O}_3\text{PS}_2^+$ 329.0429, found 329.0429.

(*E*)-Diethyl (1-phenylprop-1-en-2-yl)phosphonate (**4i**):^[2b] Pale yellow oil, yield 67%. ^1H NMR (400 MHz, CDCl_3) δ : 7.45~7.24 (m, 6H), 4.08~4.04 (m, 4H), 2.01 (d, $J=15.2$ Hz, 3H), 1.30~1.27 (m, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 142.6 (d, $J=11.6$ Hz), 135.5 (d, $J=23.8$ Hz), 129.4, 128.3, 128.3, 125.8 (d, $J=177.1$ Hz), 61.7, 61.7, 16.3, 16.3, 14.2 (d, $J=9.1$ Hz); ^{31}P NMR (CDCl_3 , 163 MHz) δ : 22.0; IR (KBr) ν : 3526, 3472, 3055, 2985, 2933, 2901, 1726, 1623, 1489, 1444, 1390, 1242, 1169, 1133, 1053, 1025, 964, 788, 752, 698, 618, 522 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{20}\text{O}_3\text{P}$ $[\text{M} + \text{H}]^+$ 255.1145, found 255.1145.

(*E*)-bis(2-Fluoroethyl) (1,2-diphenylvinyl)phosphonate (**4j**): Pale yellow oil, yield 40%. ^1H NMR (400 MHz, CDCl_3) δ : 7.65 (d, $J=25.2$ Hz, 1H), 7.37~7.05 (m, 10H), 4.61 (s, 2H), 4.49 (s, 2H), 4.31 (s, 2H), 4.24 (s, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 144.4 (d, $J=11.4$ Hz), 134.9 (d, $J=8.0$ Hz), 134.4 (d, $J=23.2$ Hz), 130.5, 129.8 (d, $J=181.3$ Hz), 129.3, 129.3, 129.0 (d, $J=1.5$ Hz), 128.3, 128.1 (d, $J=2.7$ Hz), 83.0 (d, $J=6.5$ Hz), 81.3 (d, $J=6.5$ Hz), 65.1 (d, $J=5.9$ Hz), 64.9 (d, $J=5.9$ Hz); ^{31}P NMR (CDCl_3 , 163 MHz) δ : 19.1; IR (KBr) ν : 3446, 3055, 2949, 1723, 1614, 1492, 1444, 1242, 1079, 1028, 983, 868, 797, 765, 698, 618 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{20}\text{F}_2\text{O}_3\text{P}$ $[\text{M} + \text{H}]^+$ 353.1113, found 353.1114.

(*E*)-bis(2-Methoxyethyl) (1,2-diphenylvinyl)phosphonate (**4k**): Pale yellow oil, yield 40%. ^1H NMR (400 MHz, CDCl_3) δ : 7.64 (d, $J=24.8$ Hz, 1H), 7.38~7.29 (m, 5H), 7.22~7.12 (m, 3H), 7.07~7.04 (m, 2H), 4.20~4.15 (m, 4H), 3.57~3.53 (m, 4H), 3.33 (s, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 143.8 (d, $J=11.1$ Hz), 135.3 (d, $J=7.9$ Hz), 134.6 (d, $J=22.7$ Hz), 130.5 (d, $J=180.4$ Hz), 130.4 (d, $J=10.0$ Hz), 129.4 (d, $J=10.0$ Hz), 129.0, 128.8 (d, $J=1.4$ Hz), 128.2, 127.9 (d, $J=2.2$ Hz), 71.5, 71.5, 65.1, 65.1, 58.9, 58.9; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 18.8; IR (KBr) ν : 3459, 3058, 2940, 2882, 2821, 1726, 1617, 1451, 1370, 1242, 1133, 1041, 967, 842, 759, 704, 615, 528 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{26}\text{O}_5\text{P}$ $[\text{M} + \text{H}]^+$ 377.1512, found 377.1514.

(*E*)-Diisopropyl (1,2-diphenylvinyl)phosphonate (**4l**):^[3h] Pale yellow oil, yield 83%. ^1H NMR (400 MHz, CDCl_3) δ : 7.62 (d, $J=24.4$ Hz, 1H), 7.36~7.27 (m, 5H), 7.21~7.04 (m, 5H), 4.76~4.64 (m, 2H), 1.30 (d, $J=6.4$ Hz, 6H), 1.23 (d, $J=6.4$ Hz, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 142.7 (d, $J=11.0$ Hz), 135.8 (d, $J=7.9$ Hz), 135.0 (d, $J=22.5$ Hz), 132.6 (d, $J=179.8$ Hz), 130.3 (d, $J=0.9$ Hz), 129.5 (d, $J=5.2$ Hz), 128.8, 128.6 (d, $J=1.4$ Hz), 128.1, 127.6 (d, $J=2.2$ Hz), 70.9, 70.9, 24.1, 24.0, 23.9, 23.8; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 15.9; IR (KBr) ν : 3462, 3058, 2975, 2933, 1614, 1489, 1451, 1380, 1239, 1178, 1105, 980, 887, 768, 698, 618 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{26}\text{O}_3\text{P}$ $[\text{M} + \text{H}]^+$ 345.1614, found 345.1615.

(*E*)-Diisopropyl (1,2-di-*p*-tolylvinyl)phosphonate (**4m**): Pale yellow oil, yield 78%. ^1H NMR (400 MHz, CDCl_3) δ : 7.56 (d, $J=24.8$ Hz, 1H), 7.19~7.12 (m, 4H), 6.99~6.94 (m, 4H), 4.74~4.62 (m, 2H), 2.36 (d, $J=0.3$ Hz, 3H),

2.26 (s, 3H), 1.30 (d, $J=6.0$ Hz, 6H), 1.22 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 142.6 (d, $J=11.5$ Hz), 138.9, 137.2 (d, $J=2.4$ Hz), 132.9 (d, $J=7.9$ Hz), 132.3 (d, $J=22.6$ Hz), 131.3 (d, $J=179.9$ Hz), 130.3, 129.4 (d, $J=1.6$ Hz), 129.3 (d, $J=5.6$ Hz), 128.9, 70.8, 70.7, 24.1, 24.1, 23.9, 23.8, 21.3, 21.3; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 16.5; IR (KBr) ν : 3440, 3030, 2978, 2924, 2866, 1614, 1508, 1451, 1380, 1242, 1175, 1108, 986, 887, 810, 768, 595 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{30}\text{O}_3\text{P}$ $[\text{M} + \text{H}]^+$ 373.1927, found 373.1930.

(*E*)-Diisopropyl (1,2-bis(4-(tert-butyl)phenyl)vinyl)phosphonate (**4n**): Colorless oil, yield 70%. ^1H NMR (400 MHz, CDCl_3) δ : 7.55 (d, $J=24.8$ Hz, 1H), 7.36 (d, $J=6.4$ Hz, 2H), 7.19 (dd, $J=22.0$ Hz and 6.8 Hz, 4H), 7.03 (d, $J=6.4$ Hz, 2H), 4.70~4.64 (m, 2H), 1.35~1.21 (m, 30H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 152.1, 150.6 (d, $J=2.4$ Hz), 142.3 (d, $J=11.5$ Hz), 132.9 (d, $J=7.8$ Hz), 132.1 (d, $J=6.5$ Hz), 131.4 (d, $J=195.8$ Hz), 130.2, 129.0 (d, $J=5.7$ Hz), 125.5 (d, $J=1.3$ Hz), 125.1, 70.8, 70.8, 34.7, 34.6, 31.4, 31.1, 24.1, 24.1, 23.8, 23.8; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 16.6; IR (KBr) ν : 3443, 2965, 2873, 1607, 1511, 1463, 1383, 1242, 1175, 1105, 980, 893, 829, 768, 589 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{42}\text{O}_3\text{P}$ $[\text{M} + \text{H}]^+$ 457.2866, found 457.2868.

(*E*)-Diisopropyl (1,2-bis(4-methoxyphenyl)vinyl)phosphonate (**4o**): Pale yellow oil, yield 90%. ^1H NMR (400 MHz, CDCl_3) δ : 7.54 (d, $J=24.0$ Hz, 1H), 7.22 (d, $J=6.0$ Hz, 2H), 7.05 (d, $J=6.8$ Hz, 2H), 6.89 (d, $J=6.0$ Hz, 2H), 6.68 (d, $J=6.4$ Hz, 2H), 4.70~4.64 (m, 2H), 3.82 (s, 3H), 3.73 (s, 3H), 1.31 (d, $J=4.4$ Hz, 6H), 1.24 (d, $J=3.2$ Hz, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 159.9, 159.0 (d, $J=2.1$ Hz), 142.2 (d, $J=12.1$ Hz), 131.9, 130.7 (d, $J=5.5$ Hz), 129.2 (d, $J=181.4$ Hz), 128.1 (d, $J=8.1$ Hz), 127.7 (d, $J=22.8$ Hz), 114.1 (d, $J=1.1$ Hz), 113.5, 70.6, 70.6, 55.1, 55.1, 24.1, 24.0, 23.9, 23.8; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 16.9; IR (KBr) ν : 3456, 2975, 2930, 2834, 1601, 1511, 1460, 1380, 1246, 1178, 1107, 978, 829, 775, 599 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{30}\text{O}_5\text{P}$ $[\text{M} + \text{H}]^+$ 405.1825, found 405.1827.

(*E*)-Diisopropyl (1,2-bis(4-chlorophenyl)vinyl)phosphonate (**4p**): Pale yellow oil, yield 87%. ^1H NMR (400 MHz, CDCl_3) δ : 7.57 (d, $J=24.0$ Hz, 1H), 7.34~7.31 (m, 2H), 7.23~7.12 (m, 4H), 7.00~6.97 (m, 2H), 4.76~4.64 (m, 2H), 1.31 (d, $J=6.0$ Hz, 6H), 1.24 (d, $J=6.4$ Hz, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 141.7 (d, $J=11.0$ Hz), 134.8, 134.0 (d, $J=8.0$ Hz), 133.9 (d, $J=2.6$ Hz), 133.1 (d, $J=22.5$ Hz), 132.2 (d, $J=181.2$ Hz), 131.4 (d, $J=1.0$ Hz), 130.8 (d, $J=5.2$ Hz), 129.0 (d, $J=1.3$ Hz), 128.6, 71.2, 71.1, 24.0, 24.0, 23.9, 23.8; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 14.9; IR (KBr) ν : 3452, 3065, 2981, 2933, 2869, 1617, 1588, 1486, 1380, 1246, 1178, 1101, 1018, 980, 820, 768, 699, 573 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{24}\text{Cl}_2\text{O}_3\text{P}$ $[\text{M} + \text{H}]^+$ 413.0835, found 413.0834.

(*E*)-Diisopropyl (1,2-bis(4-(trifluoromethyl)phenyl)vinyl)phosphonate (**4q**): Colorless oil, yield 81%. ^1H NMR (400 MHz, CDCl_3) δ : 7.70 (d, $J=24.0$ Hz, 1H), 7.62 (d, $J=8.4$ Hz, 2H), 7.45~7.39 (m, 4H), 7.14 (d, $J=8.0$ Hz,

2H), 4.80~4.68 (m, 2H), 1.32 (d, $J=6.4$ Hz, 6H), 1.26 (d, $J=3.2$ Hz, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 141.7 (d, $J=10.3$ Hz), 139.3 (d, $J=7.4$ Hz), 137.9 (d, $J=21.6$ Hz), 134.3 (d, $J=180.6$ Hz), 130.6 (q, $J=32.7$ Hz), 130.2 (d, $J=1.0$ Hz), 130.1 (q, $J=32.6$ Hz), 129.8 (d, $J=5.2$ Hz), 125.7 (q, $J=1.6$ Hz), 125.3 (q, $J=3.7$ Hz), 125.3 (q, $J=272.0$ Hz), 124.0 (q, $J=271.9$ Hz), 71.5, 71.4, 24.0, 23.9, 23.8, 23.8; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 13.9; ^{19}F NMR (CDCl_3 , 376 MHz) δ : -62.6, -63.0; IR (KBr) ν : 3456, 2985, 2933, 1617, 1457, 1380, 1326, 1242, 1172, 1127, 1063, 1015, 993, 836, 643, 560 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{24}\text{F}_6\text{O}_3\text{P}$ $[\text{M}+\text{H}]^+$ 481.1362, found 481.1365.

(*E*)-Diisopropyl (1,2-di(thiophen-2-yl)vinyl)phosphonate (**4r**): Pale yellow solid, m.p. 66~67 °C, yield 57%. ^1H NMR (400 MHz, CDCl_3) δ : 7.90 (d, $J=21.2$ Hz, 1H), 7.49~7.46 (m, 1H), 7.28~7.25 (m, 1H), 7.19 (d, $J=3.6$ Hz, 1H), 7.13~7.10 (m, 1H), 7.02~6.99 (m, 1H), 6.96~6.93 (m, 1H), 4.76~4.64 (m, 2H), 1.33~1.29 (m, 12H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 139.3 (d, $J=15.0$ Hz), 138.1 (d, $J=23.4$ Hz), 134.8 (d, $J=7.6$ Hz), 133.4, 130.5, 128.7 (d, $J=6.6$ Hz), 127.7 (d, $J=3.0$ Hz), 127.5 (d, $J=2.1$ Hz), 126.5 (d, $J=1.0$ Hz), 121.7 (d, $J=191.5$ Hz), 71.2, 71.1, 24.0, 24.0, 24.0, 23.9; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 14.1; IR (KBr) ν : 3433, 3071, 2984, 2930, 1601, 1524, 1444, 1377, 1342, 1226, 1105, 1012, 993, 848, 736, 704, 631, 541 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{22}\text{O}_3\text{PS}_2$ $[\text{M}+\text{H}]^+$ 357.0742, found 357.0743.

(*E*)-Diisobutyl (1,2-diphenylvinyl)phosphonate (**4s**): Colorless oil, yield 78%. ^1H NMR (400 MHz, CDCl_3) δ : 7.63 (d, $J=24.8$ Hz, 1H), 7.34~7.06 (m, 10H), 3.82~3.77 (m, 4H), 1.93~1.87 (m, 2H), 0.89~0.86 (m, 12H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 143.6 (d, $J=11.1$ Hz), 135.7 (d, $J=7.4$ Hz), 134.7 (d, $J=22.5$ Hz), 131.0 (d, $J=179.0$ Hz), 130.4, 129.2 (d, $J=5.4$ Hz), 129.0, 128.8 (d, $J=1.3$ Hz), 128.2, 127.8 (d, $J=2.4$ Hz), 72.0, 71.9, 29.2, 29.2, 18.7, 18.7, 18.7, 18.7; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 17.8; IR (KBr) ν : 3452, 3055, 2959, 2873, 1617, 1473, 1396, 1364, 1249, 1178, 1047, 864, 762, 698, 618, 528 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{30}\text{O}_3\text{P}$ $[\text{M}+\text{H}]^+$ 373.1927, found 373.1928.

(*E*)-Dicyclohexyl (1,2-diphenylvinyl)phosphonate (**4t**): Colorless oil, yield 45%. ^1H NMR (400 MHz, CDCl_3) δ : 7.61 (d, $J=24.8$ Hz, 1H), 7.31 (s, 5H), 7.16~7.06 (m, 5H), 4.45~4.41 (m, 2H), 2.07~1.65 (m, 9H), 1.49~1.45 (m, 6H), 1.30~1.25 (m, 5H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 142.4 (d, $J=11.0$ Hz), 136.0 (d, $J=7.5$ Hz), 135.0 (d, $J=22.5$ Hz), 132.8 (d, $J=180.7$ Hz), 130.3, 129.5 (d, $J=5.5$ Hz), 128.7, 128.6 (d, $J=1.4$ Hz), 128.1, 127.6 (d, $J=2.2$ Hz), 75.7, 75.6, 33.7, 33.7, 33.6, 33.5, 25.2, 25.2, 23.6, 23.6, 23.6, 23.6; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 15.9; IR (KBr) ν : 3440, 3062, 3020, 2933, 2857, 1614, 1489, 1444, 1370, 1249, 1047, 986, 893, 836, 756, 691, 611, 538 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{34}\text{O}_3\text{P}$ $[\text{M}+\text{H}]^+$ 425.2240, found 425.2242.

(*E*)-Dicyclopentyl (1,2-diphenylvinyl)phosphonate (**4u**): White solid, m.p. 75~76 °C, yield 81%. ^1H NMR (400 MHz, CDCl_3) δ : 7.60 (d, $J=24.4$ Hz, 1H), 7.32~7.06 (m,

10H), 4.94~4.90 (m, 2H), 1.79~1.53 (m, 16H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 142.8 (d, $J=11.1$ Hz), 135.9 (d, $J=7.6$ Hz), 134.9 (d, $J=22.3$ Hz), 132.3 (d, $J=179.5$ Hz), 130.3 (d, $J=0.9$ Hz), 129.4 (d, $J=5.3$ Hz), 128.8, 128.6 (d, $J=1.3$ Hz), 128.1, 127.6 (d, $J=2.5$ Hz), 79.6, 79.5, 34.2, 34.2, 34.0, 33.9, 23.1, 23.1, 23.0, 23.0; ^{31}P NMR (CDCl_3 , 163 MHz) δ : 16.2; IR (KBr) ν : 3427, 3055, 2965, 2869, 1726, 1607, 1569, 1492, 1441, 1361, 1319, 1246, 1172, 1073, 973, 919, 829, 772, 698, 605, 528 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{30}\text{O}_3\text{P}$ $[\text{M}+\text{H}]^+$ 397.1927, found 397.1929.

Supporting Information ^1H NMR, ^{13}C NMR and ^{31}P NMR spectra of products; X-ray analysis of **4u**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- For reviews: see:
 - Moonen, K.; Laureyn, O.; Stevens, C. V. *Chem. Rev.* **2004**, *104*, 6177.
 - Demmer, C. S.; Larsen, N. K.; Bunch, L. *Chem. Rev.* **2011**, *111*, 7981.
 - Queffelec, C.; Petit, M.; Janvier, P.; Knight, D. A.; Bujoli, B. *Chem. Rev.* **2012**, *112*, 3777.
 - Galezowska, J.; Gumienna-Kontecka, E. *Coord. Chem. Rev.* **2012**, *256*, 105.
 - Soller, B. S.; Salzinger, S.; Rieger, B. *Chem. Rev.* **2016**, *116*, 1993.
 - Horsman, G. P.; Zechel, D. L. *Chem. Rev.* **2017**, *117*, 5704.
 - Shi, B. C.; Fang, Y. W.; Zhang, L.; Jin, X. P.; Wu, Y. H.; Fang, M.; Yang, Y. F.; Chen, C. *Chin. J. Org. Chem.* **2016**, *36*, 673 (in Chinese).
 - 方烨汶, 张莉, 金小平, 武永辉, 方媚, 杨宇飞, 陈冲, 有机化学, **2016**, *36*, 673.)
 - Ye, X. Y.; Peng, L.; Bao, X. Z.; Tan, C. H.; Wang, H. *Green Synth. Catal.* **2021**, *2*, 6.
- Michaelis, A.; Kaehne, R. *Ber. Dtsch. Chem. Ges.* **1898**, *31*, 1048.
 - Arbuzov, A. E. *J. Russ. Phys. Chem. Soc.* **1906**, *38*, 687.
 - Hirao, T.; Masunaga, T.; Ohshiro, Y.; Agawa, T. *Tetrahedron Lett.* **1980**, *21*, 3595.
 - Bhattacharya, A. K.; Thyagarajan, G. *Chem. Rev.* **1981**, *81*, 415.
 - Hirao, T.; Masunaga, T.; Yamada, N.; Ohshiro, Y.; Agawa, T. *Bull. Chem. Soc. Jpn.* **1982**, *55*, 909.
 - Russell, G. A.; Yao, C. F. *J. Org. Chem.* **1992**, *57*, 6508.
 - Zhong, P.; Huang, X.; Xiong, Z. X. *Synlett* **1999**, 721.
 - Zhong, P.; Xiong, Z. X.; Huang, X. *Synth. Commun.* **2000**, *30*, 273.
 - Kabalka, G. K.; Guchhait, S. K. *Org. Lett.* **2003**, *5*, 729.
 - Thielges, S.; Bissere, P.; Eustache, J. *Org. Lett.* **2005**, *7*, 681.
 - Zhuang, R. Q.; Xu, J.; Cai, Z. S.; Tang, G.; Fang, M. J.; Zhao, Y. F. *Org. Lett.* **2011**, *13*, 2110.
 - Evano, G.; Tadiparthi, K.; Couty, F. *Chem. Commun.* **2011**, *47*, 179.
 - Liu, L.; Wang, Y. L.; Zeng, Z. P.; Xu, P. X.; Gao, Y. X.; Yin, Y. W.; Zhao, Y. F. *Adv. Synth. Catal.* **2013**, *355*, 659.
 - Wu, Y. L.; Liu, L. L.; Yan, K. L.; Xu, P. X.; Gao, Y. X.; Zhao, Y. F. *J. Org. Chem.* **2014**, *79*, 8118.
 - Liu, L.; Lv, Y.; Wu, Y. L.; Gao, X.; Zeng, Z. P.; Gao, Y. X.; Tang, G.; Zhao, Y. F. *RSC Adv.* **2014**, *4*, 2322.
 - Gui, Q. W.; Hu, L.; Chen, X.; Liu, J. D.; Tan, Z. *Chem. Commun.* **2015**, *51*, 13922.
 - Xue, J. F.; Zhou, S. F.; Liu, Y. Y.; Pan, X. Q.; Zou, J. P.; Asekun,

- O. T. *Org. Biomol. Chem.* **2015**, *13*, 4896.
- (r) Yuan, J. W.; Yang, L. R.; Mao, P.; Qu, L. B. *RSC Adv.* **2016**, *6*, 87058.
- (s) Liaom L. L.; Gui, Y. Y.; Zhang, X. B.; Shen, G.; Liu, H. D.; Zhou, W. J.; Li, J.; Yu, D. G. *Org. Lett.* **2017**, *19*, 3735.
- (t) Gu, J.; Cai, C. *Org. Biomol. Chem.* **2017**, *15*, 4226.
- (u) Liu, L. X.; Zhou, D.; Dong, J. Y.; Zhou, Y. B.; Yin, S. F.; Han, L. B. *J. Org. Chem.* **2018**, *83*, 4190.
- (v) Wang, L.; Yang, Z.; Zhu, H. J.; Liu, H. T.; Lv, S. P.; Xu, Y. *Eur. J. Org. Chem.* **2019**, *2019*, 2138.
- (w) Ghasemzadeh, M. S.; Akhlaghinia, B. *New J. Chem.* **2019**, *43*, 5341.
- (x) Liu, C. W.; Ji, C. L.; Zhou, T. L.; Hong, X.; Szostak, M. *Org. Lett.* **2019**, *21*, 9256.
- (y) Qiao, B. K.; Cao, H. Q.; Huang, Y. J.; Zhang, Y.; Nie, J.; Zhang, F. G.; Ma, J. A. *Chin. J. Chem.* **2018**, *36*, 809.
- (z) Ren, L. J.; Ran, M. G.; Fang, X. H.; Zhao, L.; Yao, Q. L. *Chin. J. Org. Chem.* **2018**, *38*, 2791 (in Chinese).
(任林静, 冉茂刚, 方学红, 赵玲, 姚秋丽, 有机化学, **2018**, *38*, 2791.)
- [3] (a) Han, L. B.; Tanaka, M. *J. Am. Chem. Soc.* **1996**, *118*, 1571.
(b) Han, L. B.; Hua, R. M.; Tanaka, M. *Angew. Chem. Int. Ed.* **1998**, *37*, 94.
(c) Lai, C. B.; Xi, C. J.; Chen, C.; Ma, M. M.; Hong, X. Y. *Chem. Commun.* **2003**, 2736.
(d) Trostyanskaya, I. G.; Beletskaya, I. P. *Tetrahedron* **2005**, *61*, 6315.
(e) Ananikov, V. P.; Khemchyan, L. L.; Beletskaya, I. P. *Russ. J. Org. Chem.* **2010**, *46*, 1269.
(f) Ananikov, V. P.; Khemchyan, L. L.; Beletskaya, I. P.; Starikova, Z. A. *Adv. Synth. Catal.* **2010**, *352*, 2979.
(g) Xu, Q.; Shen, R. W.; Ono, Y.; Nagahata, R.; Shimada, S.; Goto, M.; Han, L. B. *Chem. Commun.* **2011**, *47*, 2333.
(h) Khemchyan, L. L.; Ivanova, J. V.; Zaleskiy, S. S.; Ananikov, V. P.; Beletskaya, I. P.; Starikova, Z. A. *Adv. Synth. Catal.* **2014**, *356*, 771.
(i) Islas, R. E.; García, J. J. *ChemCatChem* **2017**, *9*, 4125.
(j) Chen, T. Q.; Zhao, C. Q.; Han, L. B. *J. Am. Chem. Soc.* **2018**, *140*, 3139.
(k) Wei X. H.; Bai, C. Y.; Zhao, L. B.; Zhang, P.; Li, Z. H.; Wang, Y. B.; Su, Q. *Chin. J. Chem.* **2021**, *39*, 1855.
- [4] Kers, A.; Kers, I.; Stawinski, J.; Sobkowski, M.; Kraszewski, A. *Synthesis* **1995**, 427.
- [5] (a) Huang, H.; Denne, J.; Yang, C. H.; Wang, H. B.; Kang, J. Y. *Angew. Chem. Int. Ed.* **2018**, *57*, 6624.
(b) Adler, P.; Pons, A.; Li, J.; Heider, J.; Brutiu, B. R.; Maulide, N. *Angew. Chem. Int. Ed.* **2018**, *57*, 13330.
- [6] (a) Xu, K.; Hu, H.; Yang, F.; Wu, Y. J. *Eur. J. Org. Chem.* **2013**, *2013*, 319.
(b) Xu, K.; Yang, F.; Zhang, G. D.; Wu, Y. J. *Green Chem.* **2013**, *15*, 1055.
- [7] Ouyang, K. B.; Xi, Z. F. *Acta Chim. Sinica* **2013**, *71*, 13.
- [8] Park, K.; Bae, G.; Moon, J.; Choe, J.; Song, K.; Lee, S. J. *Org. Chem.* **2010**, *75*, 6244.
- [9] Kobayashi, Y.; William, A. D. *Adv. Synth. Catal.* **2004**, *346*, 1749.

(Li, L.; Fan, Y.)