新型拟天然芪类拓扑异构酶 I 抑制剂的设计与合成

陆 棋^a 叶飞霞^a 孙晓彤^a 翁建全^{*,a} 余 茜^{*,b} 胡德玄^c^(a) 浙江工业大学化学工程学院 杭州 310014)^(b) 广东药科大学临床药学院 广州 510006)^(c) 中山大学药学院 广州 510006)

摘要 为了寻找新型抗肿瘤药物先导, 设计合成了 23 个含噻唑环的天然芪类似物, 其结构均经 NMR 和 ESI-HRMS 表征确证. 通过 Top1-介导的松散实验筛选了目标化合物的拓扑异构酶 I (Top1) 抑制活性, 其中 (*E*)-5-溴-2-(2-氯苯乙烯基)-4-对氟苯基噻唑(**6k**)显示出了良好的 Top1 抑制活性. 同时通过分子对接研究其构效关系, 结果表明化合物的 Top1 抑制活性与分子对接研究之间具有良好的相关性. 此外, 采用 3-(4,5-二甲基噻唑-2-基)-2,5-二苯基四氮唑溴盐(MTT)法测定了化合物对人乳腺癌细胞(MCF-7)和人结肠癌细胞(HCT116)的体外抗肿瘤活性, 结果表明(*E*)-5-溴-4-对氟苯基-2-(4-三氟甲基苯乙烯基)噻唑(**6e**)、(*E*)-5-溴-2-(2-氯苯乙烯基)-4-对氟苯基噻唑(**6k**)和(*E*)-5-溴-2-(4-氯苯乙烯基)-4-对氟苯基噻唑(**6l**)在低摩尔浓度下表现出较高的细胞毒活性.

关键词 天然芪; 噻唑; 拓扑异构酶 I 抑制剂; 合成; 细胞毒性

Design and Synthesis of Novel Nature-Inspired Stilbene Analogues as Potential Topoisomerase 1 Inhibitors

Lu, Qi^a Ye, Feixia^a Sun, Xiaotong^a Weng, Jianquan^{*,a} Yu, Qian^{*,b} Hu, Dexuan^c^(a) College of Chemical Engineering, Zhejiang University of Technology, Hangzhou 310014)^(b) School of Clinical Pharmacy, Guangdong Pharmaceutical University, Guangzhou 510006)^(c) School of Pharmaceutical Sciences, Sun Yat-sen University, Guangzhou 510006)

Abstract In order to find novel antitumor drug leads, twenty-three nature-inspired stilbene analogues containing thiazole moiety were designed and synthesized, and their structures were confirmed by NMR and ESI-HRMS. These compounds were screened for their topoisomerase I (Top1) inhibitory activity using Top1-mediated relaxation assay, and (*E*)-5-bromo-2-(2-chlorostyryl)-4-(4-fluorophenyl)thiazole (**6k**) possessed promising Top1 inhibitory activity. Molecular docking was also established to study the structure-activity relationship and a good correlation was observed between Top1 inhibitory activity and molecular docking study. Furthermore, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay against human breast cancer (MCF-7) and human colon cancer (HCT116) cell lines indicated that (*E*)-5-bromo-4-(4-fluorophenyl)-2-(4-(trifluoromethyl)styryl)thiazole (**6e**), (*E*)-5-bromo-2-(2-chlorostyryl)-4-(4-fluorophenyl)thiazole (**6k**), and (*E*)-5-bromo-2-(4-chlorostyryl)-4-(4-fluorophenyl)thiazole (**6l**) showed high cytotoxicity at low micromolar concentrations.

Keywords natural stilbene; thiazole; topoisomerase 1 inhibitor; synthesis; cytotoxicity

1 Introduction

DNA topoisomerase IB (Top1) is an essential nuclear enzyme that regulates DNA topology structure in several cellular metabolic processes including replicated and transcribed.^[1-3] Top1 can be used as a molecular target to screen antitumor agents. Although the classical Top1 inhibitors of camptothecin (CPT) derivatives (topotecan and irinotecan,

Figure 1A) have been developed and applied clinically, their several shortcomings remain a serious problem.^[4-5] Therefore, in the last decades many investigations have focused on the discovery of novel non-CPT Top1 inhibitors. The indenoisoquinolines LMP400 (indotecan), LMP776 (indimitecan), and LMP744 are novel non-CPT Top1 inhibitors (Figure 1B) in clinical trials.^[6]

Natural products have a long history of being used as or

* Corresponding authors. E-mail: jqweng@zjut.edu.cn; yuxi024@163.com

Received February 19, 2021; revised April 16, 2021; published online May 13, 2021.

Project supported by the Natural Science Foundation of Zhejiang Province (No. LY17C140003).

浙江省自然科学基金(No. LY17C140003)资助项目.

serving as inspiration for novel pharmaceutical and agrochemical agents.^[7-9] Natural stilbenes are phytoalexins produced by some plants in response to pathogen attack and other environmental stresses. The stilbene scaffold, structurally characterized by a 1,2-diphenylethylene nucleus, is proved to hold a wide range of biological activities, such as anticancer,^[10] anti-inflammatory,^[11] antioxidative,^[12] anti-

(A) Clinically used CPT derivatives

(B) Non-camptothecin TOP1 inhibitors

Figure 1 Structures of the camptothecins derivatives in clinical uses (A), indenoisoquinolines in clinical trials and the stilbene analogue Top1 inhibitor in laboratory (B)

bacterial,^[13] antifungal,^[14] insecticidal^[15] and algicidal^[16] activities. In our previous works, some novel biologically active stilbenes^[17] and stilbene-based analogues^[18] were discovered, and accordingly we found a new class of Top1 inhibitors, the stilbene analogues (Figure 2). Amongst them, (*E*)-2-(3-bromostyryl)-4-(2,6-difluorophenyl)thiazole showed strong (+++) Top1 inhibition (Figure 1B). These findings suggest that the stilbenes could be potential lead compounds for the development of Top1 inhibitors, which inspired us to further investigate Top1 inhibitory activity of substituted styrylthiazole derivatives.

Thus far, chemical scaffold hopping is a common approach for the discovery of new drugs. Derivatives of thiazole have been shown to exhibit various activities, such as anticancer,^[19] anti-inflammatory,^[20] analgesic,^[21] antimicrobial,^[22] antifungal^[23] and insecticidal activities,^[24] which are widely used as a scaffold in development of medicinal and agricultural active agents. In line with our continuous efforts to synthesize bioactive thiazole derivatives^[25-28] and to discover novel stilbene analogues as new Top1 inhibitors,

Figure 2 Design strategy of title compounds

the title stilbene analogues were designed by replacing one phenyl ring of stilbene scaffold with 4-(4-halophenyl)-thiazole moiety, and totally 23 novel styrylthiazole derivatives were synthesized. The design strategy of title compounds was shown in Figure 2. Their Top1 inhibition and cytotoxicity were also evaluated. Meanwhile, the preliminary relationship between structure and activity, and their molecular docking study were discussed.

2 Results and discussion

2.1 Synthesis

The synthetic route to the title stilbene-inspired substituted styrylthiazole derivatives **6a** ~ **6w** was shown in Scheme 1.

Scheme 1 Synthetic route to title compounds **6**

2-Bromo-1-(4-halophenyl)ethan-1-ones (**1**) were prepared from the corresponding 1-(4-halophenyl)ethan-1-ones according to the procedure reported in literature.^[29] The reaction of **1** with thioacetamide in *N,N*-dimethyl-

formamide (DMF) at room temperature gave the intermediate 4-(4-halophenyl)-2-methylthiazoles (**2**). Then the 5-position and benzyl position of thiazole ring in intermediate **2** were sequentially brominated under different conditions to give 5-bromo-2-(bromomethyl)-4-(4-halophenyl)thiazole **4**. According to our experimental results, the single benzyl-brominated product was almost impossible to be obtained in the process of preparing intermediate **4** because the proton on 5-position has higher reactivity than that on the benzyl position, and thus 5-position will be brominated preferentially. Meanwhile, the attempt to prepare intermediate **4** by a single step using excess *N*-bromo-succinimide (NBS) only gave dibromo product **4** in low yield. Therefore, the bromination was carried out in two steps, the 5-position was firstly brominated by reacting NBS in nucleophilic substitution mechanism, and then the obtained 5-bromo-4-(4-halophenyl)-2-methylthiazole (**3**) further underwent the free radical benzyl bromination by treatment with NBS under light irradiation to get intermediate **4**. Next, the phosphonate **5** was prepared by Arbuzov reaction procedure and directly used to next step without further purification. Finally, the title compounds **6a**~**6w** were obtained in 47.6%~71.5% yields by reacting the above crude phosphonate with various benzaldehydes *via* Wittig-Horner reaction. All the title compounds were identified and characterized by ^1H NMR, ^{13}C NMR and ESI-HRMS. In the ^1H NMR spectra of title compounds, the peaks of the vinylic CH proton connected to the thiazole ring appeared in the δ 7.14~7.38 range and the signals of the vinylic CH proton connected to the benzene ring were observed around δ 7.31~7.72, and they generally appeared as a doublet with a coupling constant $J=16.0$ Hz or 16.5 Hz which proved the title compounds are of *E*-configuration.^[17] Meanwhile, all the title compounds exhibited the $[\text{M}+\text{H}]^+$ peak in the ESI-MS results.

2.2 Top1 inhibitory activities

Top1 inhibitory activity of the title styrylthiazole derivatives **6a**~**6w** were assessed by Top1-mediated relaxation assay with CPT as a positive control. All com-

pounds were tested at $50\text{ }\mu\text{mol}\cdot\text{L}^{-1}$. The compounds were semiquantitatively expressed relative to CPT at $50\text{ }\mu\text{mol}\cdot\text{L}^{-1}$ as follows: +++, more than 80% of the activity; ++, between 40% and 79% of the activity; +, between 10% and 39% of the activity; +, less than 10% of the activity. The Top1 inhibitory activity is summarized in Table 1. The results showed that the compounds with an electron-withdrawing group (halogen and trifluoromethyl) in the benzene ring generally displayed better Top1 inhibition than those with an electron-donating group (alkyl and methoxyl). The compound **6k** exhibited moderate Top1 inhibition of +++. The representative Top1-mediated relaxation assay gels are shown in Figure 3. They show that compound **6k** exhibited dose-dependent manner inhibit of Top1 at the concentrations of 0.2, 1, 5, 25 and 125 $\mu\text{mol}\cdot\text{L}^{-1}$.

Figure 3 Representative gels for Top1-mediated relaxation assay. Lane 1, pBR322 DNA alone; Lane 2, pBR322 DNA and Top1 without compound; Lanes 3~7, pBR322 DNA, Top1 and compounds at 0.2, 1, 5, 25, 125 $\mu\text{mol/L}$, respectively. R, relaxed DNA; Sc, supercoiled DNA

2.3 Molecular docking study

To investigate the binding modes of our inhibitors, representative compound **6k** was docked into the topotecan binding pocket of the Top1-DNA-topotecan ternary complex (PDB ID:1K4T). The energy-minimized, top-ranked

Table 1 Top1 inhibitory activities of stilbene analogues **6a**~**6w**

Compd.	X	R	Relaxation assay ^a	Compd.	X	R	Relaxation assay ^a
6a	F	H	+	6m	F	2,4-Cl ₂	++
6b	F	2-Me	+	6n	F	2-Br	++
6c	F	3-Me	+	6o	F	3-Br	+
6d	F	4-Me	+	6p	Cl	4-CF ₃	++
6e	F	4-CF ₃	++	6q	Cl	4- <i>tert</i> -Bu	+
6f	F	4- <i>tert</i> -Bu	+	6r	Cl	3-OMe	+
6g	F	2-OMe	+	6s	Cl	3,4,5-(OMe) ₃	+
6h	F	3-OMe	+	6t	Cl	2-F	+
6i	F	4-OMe	+	6u	Cl	3-Br	+
6j	F	4-F	++	6v	Br	2-F	++
6k	F	2-Cl	+++	6w	Br	4-F	++
6l	F	4-Cl	++				

^a Top1 inhibitory activity was semiquantitatively expressed relative to CPT at 50 $\mu\text{mol/L}$ as follows: +++, more than 80% of the activity; ++, between 40% and 79% of the activity; +, between 10% and 39% of the activity; +, less than 10% of the activity. Every experiment was repeated at least twice independently.

pose of **6k** in the ternary complex is displayed in Figure 4. The fluorobenzene scaffold turns to a suitable pocket which binds to the DNA cleavage site, and stacks with the DNA (T and A) via a strong π - π stack along with the thiazole scaffold (Figure 4, A and B). The N atom of the thiazole forms a hydrogen bond to Arg364 with 3.2 Å (Figure 4, C), which come in contact with Top1 and DNA, and will be attributed to Top1 inhibition. Moreover, because of the strongly polar region along with Lys532, Asp533 and the DNA pairs (T and G), an electron-drawing group substituted in the benzene ring, like X and CF₃, show a better Top1 inhibition than an electron-donating group, like OMe and Me. The simulated results generated a reasonable binding mode of **6k**, further strengthening our concept of this rational design.

Figure 4 Hypothetical binding mode of **6k** in a ternary complex with DNA and Top1 (derived from PDB ID:1K4T)

All distances are measured from heavy atom to heavy atom. The carbon atoms of **6k** are shown in yellow, and the base pairs are displayed in blue lines.

2.4 Cytotoxicity

The *in vitro* cytotoxicities of **6a**~**6w** were also evaluated through MTT assay against two human cancer cell lines, including MCF-7 and HCT116. The etoposide (vp-16) and CPT were used as positive controls. The compounds **6a**~**6w** were incubated with cells for 72 h in five-dose assay ranging from 0.01 $\mu\text{mol}\cdot\text{L}^{-1}$ to 100 $\mu\text{mol}\cdot\text{L}^{-1}$. The IC₅₀ values defined as the inhibitor concentration which inhibits growth by 50%, are summarized in Table 2. It was found that compounds showed good correlation between their Top1 inhibitory activity and cytotoxicity toward two cancer

cell lines. Generally, the compounds which showed strong Top1 inhibition displayed high cytotoxicity. The potent Top1 inhibitor **6k** (+++) showed significant cytotoxicity against MCF-7 and HCT116 cell lines with IC₅₀ values of 2.29 and 7.37 $\mu\text{mol}\cdot\text{L}^{-1}$, respectively. The compounds **6e**, **6j**, **6l**, and **6m** displayed high cytotoxicity against HCT116 cell lines with IC₅₀ values ranging from 0.72 $\mu\text{mol}\cdot\text{L}^{-1}$ to 8.06 $\mu\text{mol}\cdot\text{L}^{-1}$ with Top1 inhibition of ++.

3 Conclusions

In summary, a series of novel nature-inspired stilbene analogues containing thiazole moiety were designed and synthesized, and their Top1 inhibitory and cytotoxicity were evaluated. The results showed that all compounds possessed certain Top1 inhibitory activity and some of them exhibited high cytotoxicity. Amongst them, compound **6k** displayed promising Top1 inhibitory activity and compound **6k** exhibited the highest cytotoxicity against MCF-7 with IC₅₀ value of 2.29 $\mu\text{mol}\cdot\text{L}^{-1}$. Compound **6e** showed the highest cytotoxicity against HCT116 with IC₅₀ value of 0.72 $\mu\text{mol}\cdot\text{L}^{-1}$. Furthermore, the molecular docking was studied and the preliminary structure-activity relationships were discussed. These results are useful for the design of more potent natural stilbene-based anticancer agents in the future.

4 Experimental section

4.1 General

Melting points were determined using an X-4 apparatus without correction. ¹H NMR and ¹³C NMR spectra were measured on a Bruker ADVANCE III instrument (500 MHz for ¹H NMR and 125 MHz for ¹³C NMR) using TMS as an internal standard and CDCl₃ as the solvents. HRMS was determined on a Shimadzu LCMS-IT-TOF mass spectrometer equipped with electrospray ionization (ESI) ion source. All reactions were carried out under air atmosphere in dried glassware. The reaction progress was monitored by TLC on silica gel GF254, and spots were visualized by exposure to UV light (254 nm). All chemical reagents and solvents used in this study were commercial and were used without further purification.

4.2 Synthesis

4.2.1 Synthesis of 4-(4-halophenyl)-2-methylthiazole (2)

2-Bromo-1-(4-halophenyl)ethanone (**1**) (10 mmol) and the thioacetamide (0.83 g, 11 mmol) were dissolved in DMF (20 mL). Then the mixture was stirred at room temperature for 5 h. After the reaction was completed, 50 mL of H₂O was added and stirred for 30 min, the resulting precipitate was filtered and washed with water for 2 times to give 4-(4-halophenyl)-2-methylthiazole (**2**).

4-(4-Fluorophenyl)-2-methylthiazole (**2a**): White solid, 1.78 g, yield 92.0%. m.p. 183~184 °C (lit.^[30] 184~185 °C); ¹H NMR (500 MHz, CDCl₃) δ : 7.86 (dd, *J*=8.9, 5.4 Hz, 2H), 7.25 (s, 1H), 7.11 (t, *J*=8.7 Hz, 2H), 2.78 (s,

Table 2 Cytotoxic activities of compounds **6a**~**6w** against two cancer cell lines

Compd.	IC ₅₀ /(μmol·L ⁻¹)		Compd.	IC ₅₀ /(μmol·L ⁻¹)	
	MCF-7	HCT116		MCF-7	HCT116
6a	22.16±5.01	15.78±1.89	6n	20.95±1.41	58.38±3.49
6b	>100	>100	6o	>100	>100
6c	>100	>100	6p	>100	>100
6d	95.52±3.11	31.87±4.29	6q	>100	>100
6e	6.31±0.51	0.72±0.09	6r	61.92±8.05	5.26±0.35
6f	59.42±4.21	6.47±0.62	6s	>100	>100
6g	49.53±9.08	17.8±1.16	6t	>100	>100
6h	29.85±5.47	4.99±0.80	6u	>100	62.91±2.34
6i	57.20±1.79	17.32±1.05	6v	64.57±1.45	37.18±6.65
6j	49.64±3.58	8.06±0.92	6w	>100	63.25±6.17
6k	2.29±1.08	7.37±1.15	vp-16 ^a	24.68±3.12	18.95±1.35
6l	13.45±2.46	3.07±0.53	CPT ^a	0.34	0.012
6m	72.43±2.15	4.79±0.73			

^a Positive control.

3H).

4-(4-Chlorophenyl)-2-methylthiazole (**2b**): White solid, 2.01 g, yield 95.8%. m.p. 121~123 °C (lit.^[31] 122~123 °C); ¹H NMR (500 MHz, CDCl₃) δ: 7.73 (d, *J*=7.5 Hz, 2H), 7.64 (s, 1H), 7.54 (d, *J*=7.5 Hz, 2H), 2.85 (s, 3H).

4-(4-Bromophenyl)-2-methylthiazole (**2c**): White solid, 2.37 g, yield 93.3%. m.p. 132~133 °C (lit.^[32] 134 °C); ¹H NMR (500 MHz, CDCl₃) δ: 7.70 (d, *J*=7.5 Hz, 2H), 7.66 (s, 1H), 7.55 (d, *J*=7.5 Hz, 2H), 2.85 (s, 3H).

4.2.2 Synthesis of 5-bromo-4-(4-halophenyl)-2-methylthiazole (**3**)

4-(4-Halophenyl)-2-methylthiazole (**2**) (10 mmol) and *N*-bromosuccinimide (NBS) (1.96 g, 11 mmol) were dissolved in ethyl acetate (20 mL), and the resulting solution was heated to reflux for 3 h. The reaction mixture was cooled to room temperature and washed with water. The organic layer was dried over anhydrous MgSO₄ and evaporated *in vacuo*. Then, the residual solid was recrystallized from anhydrous ethanol and dried to give pure 5-bromo-4-(4-halophenyl)-2-methylthiazole (**3**).

5-Bromo-4-(4-fluorophenyl)-2-methylthiazole (**3a**): Yellow solid, 2.18 g, yield 80.0%. m.p. 56~57 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.89 (dd, *J*=10.0, 5.0 Hz, 2H), 7.12 (t, *J*=8.7 Hz, 2H), 2.70 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.7, 163.6 (*J*=247.5 Hz), 151.0, 130.5 (*J*=21.2 Hz), 129.6, 115.4 (dd, *J*=21.2, 3.7 Hz), 101.9, 19.5. HRMS (ESI) calcd C₁₀H₈BrFNS [M + H]⁺ 271.9467, found 271.9471.

5-Bromo-4-(4-chlorophenyl)-2-methylthiazole (**3b**): Yellow solid, 2.33 g, yield 80.8%. m.p. 79~80 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.88~7.84 (m, 2H), 7.42~7.39 (m, 2H), 2.69 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.8, 150.7, 134.3, 131.9, 129.8, 128.5, 102.4, 19.6. HRMS (ESI) calcd C₁₀H₈BrClNS [M + H]⁺ 287.9171, found 287.9177.

5-Bromo-4-(4-bromophenyl)-2-methylthiazole (**3c**): light yellow solid, 2.71 g, yield 81.5 %. m.p. 80~82 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.82 (d, *J*=8.5 Hz, 2H),

7.58 (d, *J*=8.5 Hz, 2H), 2.71 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.9, 150.8, 132.3, 131.5, 130.1, 122.6, 102.5, 19.6. HRMS (ESI) calcd C₁₀H₈Br₂NS [M + H]⁺ 331.8666, found 331.8672.

4.2.3 Synthesis of 5-bromo-2-(bromomethyl)-4-(4-halophenyl)thiazole (**4**)

5-Bromo-4-(4-halophenyl)-2-methylthiazole (**3**) (10 mmol) and NBS (1.96 g, 11 mmol) were dissolved in CH₃CN (30 mL). The reaction mixture was stirred and irradiated by 15 W LED for 4 h at room temperature. Ethyl acetate (50 mL) was added, and washed with saturated sodium bicarbonate solution and water. Then, the solvent was removed and the residue was recrystallized from anhydrous ethanol and dried to give intermediates **4**.

5-Bromo-2-(bromomethyl)-4-(4-fluorophenyl)thiazole (**4a**): Yellow solid, 1.90 g, yield 54.0%. m.p. 35~36 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.89 (dd, *J*=10.0, 5.5 Hz, 2H), 7.13 (t, *J*=8.7 Hz, 2H), 4.69 (s, 2H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.5, 163.8 (*J*=247.5 Hz), 151.6, 130.5 (*J*=7.5 Hz), 129.0, 115.4 (*J*=38.7 Hz), 105.6, 29.7. HRMS (ESI) calcd C₁₀H₇Br₂FNS [M + H]⁺ 349.8572, found 349.8581.

5-Bromo-2-(bromomethyl)-4-(4-chlorophenyl)thiazole (**4b**): Yellow solid, 2.40 g, yield 65.2%. m.p. 83~84 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.86 (d, *J*=8.6 Hz, 2H), 7.41 (d, *J*=8.6 Hz, 2H), 4.67 (s, 2H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.4, 151.1, 134.5, 131.1, 129.6, 128.4, 105.9, 26.3. HRMS (ESI) calcd C₁₀H₇Br₂ClNS [M + H]⁺ 365.8276, found 365.8269.

5-Bromo-2-(bromomethyl)-4-(4-bromophenyl)thiazole (**4c**): Yellow solid, 2.49 g, yield 60.4%. m.p. 80~81 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.81 (d, *J*=8.6 Hz, 2H), 7.58 (d, *J*=8.6 Hz, 2H), 4.69 (s, 2H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.7, 151.4, 131.8, 131.6, 130.1, 123.0, 106.2, 26.4. HRMS (ESI) calcd C₁₀H₇Br₃NS [M + H]⁺ 409.7771, found 409.7766.

4.2.4 Synthesis of diethyl ((4-(4-halophenyl)thiazol-2-yl)methyl)phosphonate (**5**)

5-Bromo-2-(bromomethyl)-4-(4-halophenyl)thiazole (**4**)

(10 mmol) and triethyl phosphite (3.34 g, 20 mmol) were dissolved in toluene (20 mL), and the resulting solution was heated to reflux for 2 h. After the reaction is completed, the solvent and excess triethyl phosphite were evaporated to give phosphonate **5** as a brown oil that was used without further purification.

4.2.5 General synthetic procedure for title compounds **6a~6o**

The resulting phosphonate **5**, substituted benzaldehyde (12 mmol) and NaOH (0.48 g, 12 mmol) were dissolved in 20 mL of anhydrous DMF. Then the mixture was stirred at room temperature for 5 h. After the reaction is completed, 50 mL of H₂O was added and continued to stir for 30 min, and then extracted with CHCl₃ (30 mL×3), the organic layer was combined and dried by anhydrous MgSO₄. Then the solvent was removed and the residue was purified by column chromatography with petroleum ether-EtOAc (*V*: *V*=5:1) to obtain (*E*)-5-bromo-4-(4-halophenyl)-2-(substituted-styryl)thiazoles **6a~6w**.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-styrylthiazole (**6a**): Yellow solid, yield 63.2%. m.p. 109~111 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (dd, *J*=8.5, 5.5 Hz, 2H), 7.55 (d, *J*=8.5 Hz, 2H), 7.46~7.33 (m, 4H), 7.26 (d, *J*=16.0 Hz, 1H), 7.17 (t, *J*=8.5 Hz, 2H); ¹³C NMR (125 MHz, CDCl₃) δ: 166.3, 162.8 (d, *J*=248.7 Hz), 152.2, 135.3, 130.5 (d, *J*=8.2 Hz), 129.4 (d, *J*=3.3 Hz), 128.9, 127.2, 121.0, 115.3 (d, *J*=21.6 Hz), 102.6. HRMS (ESI) calcd C₁₇H₁₂BrFNS [M+H]⁺ 359.9852, found 359.9836.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(2-methylstyryl)thiazole (**6b**): Yellow solid, yield 61.6%. m.p. 99~100 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (dd, *J*=9.0, 5.5 Hz, 2H), 7.68~7.55 (m, 2H), 7.31~7.21 (m, 3H), 7.20~7.11 (m, 3H), 2.47 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 166.6, 162.8 (d, *J*=249.7 Hz), 152.2, 136.7, 134.3, 133.0, 130.8, 130.5 (d, *J*=8.4 Hz), 129.5 (d, *J*=3.2 Hz), 129.1, 126.4, 125.7, 122.1, 115.3 (d, *J*=21.6 Hz), 102.5, 19.9. HRMS (ESI) calcd C₁₈H₁₄BrFNS [M+H]⁺ 374.0009, found 373.9994.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(3-methylstyryl)thiazole (**6c**): Yellow solid, yield 64.6%. m.p. 101~103 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (dd, *J*=8.5, 5.5 Hz, 2H), 7.41~7.27 (m, 4H), 7.24 (d, *J*=16.5 Hz, 1H), 7.17 (t, *J*=8.5 Hz, 3H), 2.40 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 166.4, 162.7 (d, *J*=248.7 Hz), 152.1, 138.5, 135.5, 135.3, 130.5 (d, *J*=8.2 Hz), 130.1, 129.5 (d, *J*=3.1 Hz), 128.8, 127.9, 124.4, 120.8, 115.3 (d, *J*=21.7 Hz), 102.4, 21.4. HRMS (ESI) calcd C₁₈H₁₄BrFNS [M+H]⁺ 374.0009, found 373.9995.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(4-methylstyryl)thiazole (**6d**): Yellow solid, yield 64.4%. m.p. 148~150 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.95 (s, 2H), 7.44 (d, *J*=7.0 Hz, 2H), 7.33 (d, *J*=16.0 Hz, 1H), 7.26~7.10 (m, 5H), 2.39 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 166.6, 162.8 (d, *J*=248.7 Hz), 152.1, 139.5, 135.4, 132.6, 130.5 (d, *J*=8.3 Hz), 129.7, 129.5 (d, *J*=3.0 Hz), 120.1, 115.3 (d, *J*=21.6 Hz), 102.2, 21.4. HRMS (ESI) calcd C₁₈H₁₄Br-

FNS [M+H]⁺ 374.0009, found 373.9972.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(4-(trifluoromethyl)styryl)thiazole (**6e**): Yellow solid, yield 70.8%. m.p. 123~125 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (dd, *J*=9.0, 5.5 Hz, 2H), 7.65 (dd, *J*=12.5, 3.5 Hz, 4H), 7.39 (d, *J*=16.0 Hz, 1H), 7.30 (d, *J*=16.0 Hz, 1H), 7.17 (t, *J*=8.5 Hz, 2H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.3, 162.8 (d, *J*=249.1 Hz), 152.5, 138.8, 133.3, 130.5 (d, *J*=8.3 Hz), 129.3 (d, *J*=3.3 Hz), 127.3, 127.2, 125.9 (q, *J*=3.8 Hz), 125.0, 123.2, 122.9, 120.7, 115.4 (d, *J*=21.6 Hz), 103.5. HRMS (ESI) calcd C₁₈H₁₁BrF₄NS [M+H]⁺ 427.9726, found 427.9716.

(*E*)-5-Bromo-2-(4-(*tert*-butyl)styryl)-4-(4-fluorophenyl)thiazole (**6f**): Yellow solid, yield 55.5%. m.p. 98~99 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.98 (dd, *J*=9.0, 5.5 Hz, 2H), 7.50 (d, *J*=8.5 Hz, 2H), 7.44 (d, *J*=8.0 Hz, 2H), 7.35 (d, *J*=16.5 Hz, 1H), 7.24 (d, *J*=16.5 Hz, 1H), 7.17 (t, *J*=8.5 Hz, 2H), 1.37 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ: 166.5, 162.7 (d, *J*=248.7 Hz), 152.6, 152.0, 135.2, 132.5, 130.5 (d, *J*=8.2 Hz), 129.5 (d, *J*=3.2 Hz), 127.0, 125.8, 120.2, 115.2 (d, *J*=21.6 Hz), 102.2, 34.8, 31.2. HRMS (ESI) calcd C₂₁H₂₀BrFNS [M+H]⁺ 416.0478, found 416.0468.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(2-methoxystyryl)thiazole (**6g**): Yellow solid, yield 71.5%. m.p. 111~113 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (dd, *J*=8.5, 5.5 Hz, 2H), 7.66 (d, *J*=16.5 Hz, 1H), 7.56 (dd, *J*=7.5, 1.5 Hz, 1H), 7.41~7.31 (m, 2H), 7.16 (t, *J*=8.5 Hz, 2H), 7.00 (t, *J*=7.5 Hz, 1H), 6.94 (d, *J*=8.5 Hz, 1H), 3.93 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 167.4, 162.7 (d, *J*=248.5 Hz), 157.6, 152.0, 130.9, 130.5 (d, *J*=8.3 Hz), 130.4, 129.6 (d, *J*=3.1 Hz), 127.8, 124.3, 121.8, 120.9, 115.3 (d, *J*=21.3 Hz), 111.1, 102.2. HRMS (ESI) calcd C₁₈H₁₄BrFNOS [M+H]⁺ 389.9958, found 389.9945.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(3-methoxystyryl)thiazole (**6h**): Yellow solid, yield 64.5%. m.p. 167~168 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (dd, *J*=8.5, 5.5 Hz, 2H), 7.64 (d, *J*=16.0 Hz, 1H), 7.49 (d, *J*=8.5 Hz, 1H), 7.26~7.08 (m, 5H), 6.79 (dd, *J*=9.0, 3.0 Hz, 1H), 3.84 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 165.8, 162.80 (d, *J*=248.8 Hz), 159.1, 152.4, 135.8, 134.0, 133.9, 130.5 (d, *J*=8.3 Hz), 129.4 (d, *J*=3.2 Hz), 123.8, 117.0, 115.4 (m), 111.7, 103.3, 55.6. HRMS (ESI) calcd C₁₈H₁₄BrFNOS [M+H]⁺ 389.9958, found 389.9767.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(4-methoxystyryl)thiazole (**6i**): Yellow solid, yield 71.3%. m.p. 145~146 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.95 (dd, *J*=8.5, 5.5 Hz, 2H), 7.49 (d, *J*=8.5 Hz, 2H), 7.31 (d, *J*=16.0 Hz, 1H), 7.20~7.05 (m, 3H), 6.93 (d, *J*=8.5 Hz, 2H), 3.86 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 166.8, 162.7 (d, *J*=248.6 Hz), 160.6, 152.0, 135.0, 130.5 (d, *J*=8.2 Hz), 129.5 (d, *J*=3.3 Hz), 128.7, 128.1, 118.9, 115.3 (d, *J*=21.6 Hz), 114.4, 101.9, 55.4. HRMS (ESI) calcd C₁₈H₁₄BrFONS [M+H]⁺ 389.9958, found 389.9940.

(*E*)-5-Bromo-4-(4-fluorophenyl)-2-(4-fluorostyryl)thiazole (**6j**): Yellow solid, yield 58.7%. m.p. 132~134 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.95 (dd, *J*=8.5, 5.5 Hz,

2H), 7.52 (dd, $J=8.5, 5.5$ Hz, 2H), 7.32 (d, $J=16.0$ Hz, 1H), 7.20~7.12 (m, 3H), 7.10 (t, $J=8.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.0, 164.7~161.5 (m), 152.2, 134.0, 131.6 (d, $J=3.4$ Hz), 130.5 (d, $J=8.4$ Hz), 129.4 (d, $J=3.4$ Hz), 128.9 (d, $J=8.2$ Hz), 120.8 (d, $J=2.4$ Hz), 116.1 (d, $J=21.9$ Hz), 115.3 (d, $J=21.7$ Hz), 102.6. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{BrF}_2\text{NS}$ $[\text{M} + \text{H}]^+$ 377.9758, found 377.9802.

(*E*)-5-Bromo-2-(2-chlorostyryl)-4-(4-fluorophenyl)thiazole (**6k**): Yellow solid, yield 57.3%. m.p. 124~126 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.96 (dd, $J=9.0, 5.5$ Hz, 2H), 7.74 (d, $J=16.0$ Hz, 1H), 7.68 (dd, $J=7.5, 2.5$ Hz, 1H), 7.43 (dd, $J=7.5, 2.5$ Hz, 1H), 7.34~7.23 (m, 3H), 7.17 (t, $J=8.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.0, 162.8 (d, $J=248.9$ Hz), 152.3, 134.1, 133.5, 131.2, 130.6, 130.5, 130.1, 130.1, 129.3 (d, $J=3.2$ Hz), 127.2, 126.8, 123.5, 115.4 (d, $J=21.6$ Hz), 103.2. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{BrClFNS}$ $[\text{M} + \text{H}]^+$ 393.9463, found 393.9451.

(*E*)-5-Bromo-2-(4-chlorostyryl)-4-(4-fluorophenyl)thiazole (**6l**): Yellow solid, yield 71.0%. m.p. 149~151 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.94 (dd, $J=9.0, 5.5$ Hz, 2H), 7.47 (d, $J=8.5$ Hz, 2H), 7.37 (d, $J=8.5$ Hz, 2H), 7.31 (d, $J=16.5$ Hz, 1H), 7.20 (d, $J=16.5$ Hz, 1H), 7.16 (t, $J=9.0$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 173.0, 165.8, 162.8 (d, $J=247.9$ Hz), 152.3, 133.9, 133.8, 130.5 (d, $J=8.4$ Hz), 129.3 (d, $J=3.1$ Hz), 129.2, 128.3, 121.5, 115.4 (d, $J=21.6$ Hz), 102.9. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{BrClFNS}$ $[\text{M} + \text{H}]^+$ 393.9463, found 393.9439.

(*E*)-5-Bromo-2-(2,4-dichlorostyryl)-4-(4-fluorophenyl)thiazole (**6m**): Yellow solid, yield 66.2%. m.p. 170~171 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.95 (dd, $J=9.0, 5.5$ Hz, 2H), 7.66 (d, $J=16.5$ Hz, 1H), 7.61 (d, $J=8.5$ Hz, 1H), 7.45 (d, $J=2.5$ Hz, 1H), 7.32~7.26 (m, 1H), 7.23 (d, $J=16.5$ Hz, 1H), 7.17 (t, $J=9.0$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 165.5, 162.8 (d, $J=249.1$ Hz), 152.5, 135.2, 134.6, 132.2, 130.5 (d, $J=8.3$ Hz), 129.9, 129.8, 129.3 (d, $J=3.3$ Hz), 127.7, 127.5, 123.9, 115.4 (d, $J=21.7$ Hz), 103.5. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{10}\text{BrCl}_2\text{FNS}$ $[\text{M} + \text{H}]^+$ 427.9073, found 427.9057.

(*E*)-5-Bromo-2-(2-bromostyryl)-4-(4-fluorophenyl)thiazole (**6n**): Yellow solid, yield 50.9%. m.p. 137~138 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.96 (dd, $J=9.0, 5.5$ Hz, 2H), 7.71 (d, $J=16.0$ Hz, 1H), 7.68 (dd, $J=8.0, 1.5$ Hz, 1H), 7.63 (dd, $J=8.0, 1.5$ Hz, 1H), 7.35 (t, $J=7.0$ Hz, 1H), 7.25~7.12 (m, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ : 165.9, 162.8 (d, $J=248.7$ Hz), 152.4, 135.3, 133.8, 133.4, 130.5 (d, $J=8.4$ Hz), 130.3, 129.4 (d, $J=3.3$ Hz), 127.8, 127.0, 124.6, 123.7, 115.4 (d, $J=21.6$ Hz), 103.3. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{Br}_2\text{FNS}$ $[\text{M} + \text{H}]^+$ 437.8957, found 437.8947.

(*E*)-5-Bromo-2-(3-bromostyryl)-4-(4-fluorophenyl)thiazole (**6o**): Yellow solid, yield 55.7%. m.p. 160~161 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.95 (dd, $J=8.5, 5.5$ Hz, 2H), 7.69 (s, 1H), 7.46 (t, $J=9.0$ Hz, 2H), 7.34~7.25 (m, 2H), 7.22 (d, $J=16.0$ Hz, 1H), 7.17 (t, $J=8.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 165.5, 162.8 (d, $J=248.8$ Hz),

152.4, 137.5, 133.4, 132.0, 130.5 (d, $J=8.3$ Hz), 130.4, 130.0, 129.3 (d, $J=3.3$ Hz), 125.7, 123.1, 122.2, 115.4 (d, $J=21.3$ Hz), 103.1. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{Br}_2\text{FNS}$ $[\text{M} + \text{H}]^+$ 437.8958, found 437.8968.

(*E*)-5-bromo-4-(4-chlorophenyl)-2-(4-(trifluoromethyl)styryl)thiazole (**6p**): Light yellow solid, yield 61.4%. m.p. 121~123 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.95~7.91 (m, 2H), 7.68~7.61 (m, 4H), 7.48~7.43 (m, 2H), 7.39 (d, $J=16.5$ Hz, 1H), 7.30 (d, $J=16.0$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 165.4, 152.2, 138.8, 134.7, 133.4, 131.6, 130.7 (q, $J=32.6$ Hz), 129.9, 128.6, 127.3, 125.9 (q, $J=3.8$ Hz), 125.0, 123.1, 103.9. HRMS (ESI) calcd $\text{C}_{18}\text{H}_{11}\text{BrClF}_3\text{NS}$ $[\text{M} + \text{H}]^+$ 443.9431, found 443.9428.

(*E*)-5-Bromo-2-(4-(tert-butyl)styryl)-4-(4-chlorophenyl)thiazole (**6q**): Yellow solid, yield 60.8%. m.p. 77~79 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.95~7.91 (m, 2H), 7.49 (d, $J=8.5$ Hz, 2H), 7.47~7.41 (m, 4H), 7.34 (d, $J=16.0$ Hz, 1H), 7.23 (d, $J=16.0$ Hz, 1H), 1.36 (s, 9H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.7, 152.8, 151.8, 135.4, 134.5, 132.6, 131.8, 129.9, 128.5, 127.1, 125.9, 120.2, 102.8, 34.8, 31.2. HRMS (ESI) calcd $\text{C}_{21}\text{H}_{20}\text{BrClNS}$ $[\text{M} + \text{H}]^+$ 432.0183, found 432.0185.

(*E*)-5-Bromo-4-(4-chlorophenyl)-2-(3-methoxystyryl)thiazole (**6r**): Yellow solid, yield 50.9%. m.p. 103~104 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.94~7.91 (m, 2H), 7.47~7.43 (m, 2H), 7.35~7.30 (m, 1H), 7.34 (d, $J=16.5$ Hz, 1H), 7.24 (d, $J=16.5$ Hz, 1H), 7.14 (d, $J=7.5$ Hz, 1H), 7.08~7.07 (m, 1H), 6.92 (ddd, $J=8.0, 2.5, 0.5$ Hz, 1H), 3.86 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.4, 160.0, 152.0, 136.8, 135.4, 134.6, 131.8, 130.0, 128.6, 121.3, 120.0, 115.3, 112.1, 103.2, 55.4. HRMS (ESI) calcd $\text{C}_{18}\text{H}_{14}\text{BrClNOS}$ $[\text{M} + \text{H}]^+$ 405.9663, found 405.9665.

(*E*)-5-Bromo-4-(4-chlorophenyl)-2-(3,4,5-trimethoxystyryl)thiazole (**6s**): Yellow solid, yield 50.6%. m.p. 162~163 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.92 (d, $J=8.5$ Hz, 2H), 7.44 (d, $J=8.5$ Hz, 2H), 7.27 (d, $J=16.0$ Hz, 1H), 7.17 (d, $J=16.0$ Hz, 1H), 6.75 (m, 2H), 3.92 (s, 9H). ^{13}C NMR (125 MHz, CDCl_3) δ : 166.3, 163.7, 153.8, 153.6, 151.9, 150.6, 139.4, 138.7, 135.9, 135.4, 134.5, 131.8, 131.0, 129.9, 128.6, 123.9, 120.4, 105.8, 105.0, 104.4, 103.0, 61.1, 56.3, 29.7. HRMS (ESI) calcd $\text{C}_{20}\text{H}_{18}\text{BrClNO}_3\text{S}$ $[\text{M} + \text{H}]^+$ 465.9874, found 465.9874.

(*E*)-5-Bromo-4-(4-chlorophenyl)-2-(2-fluorostyryl)thiazole (**6t**): Light yellow solid, yield 52.4%. m.p. 149~151 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.96~7.91 (m, 1H), 7.89~7.83 (m, 1H), 7.50 (d, $J=16.5$ Hz, 1H), 7.48~7.39 (m, 3H), 7.34 (d, $J=16.5$ Hz, 1H), 7.24~7.16 (m, 2H), 6.95 (d, $J=7.0$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.2, 163.2, 161.8 (d, $J=250.8$ Hz), 152.0, 150.7, 134.5, 131.7, 130.6, 129.8, 128.5, 128.0, 127.8, 126.3, 124.5, 123.4 (d, $J=23.0$ Hz), 123.3, 116.3 (d, $J=21.4$ Hz), 103.6, 100.0, 29.7. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{BrClFNS}$ $[\text{M} + \text{H}]^+$ 393.9463, found 393.9467.

(*E*)-5-Bromo-2-(3-bromostyryl)-4-(4-chlorophenyl)thiazole (**6u**): Yellow solid, yield 57.9%. m.p. 169~170 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.95~7.91 (m, 2H), 7.70

(t, $J=1.5$ Hz, 1H), 7.50~7.44 (m, 4H), 7.30~7.27 (m, 2H), 7.22 (d, $J=16.0$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 165.6, 152.1, 137.5, 134.6, 133.9, 133.5, 132.1, 131.7(m), 130.5, 130.0, 129.9, 128.6, 127.5, 125.8, 124.5, 123.1, 122.2, 103.6, 29.7. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{Br}_2\text{-CINS}$ $[\text{M}+\text{H}]^+$ 453.8662, found 453.8659.

(*E*)-5-Bromo-4-(4-bromophenyl)-2-(2-fluorostyryl)thiazole (**6v**): Light yellow solid, yield 47.6%. m.p. 140~141 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.89~7.85 (m, 2H), 7.63~7.59 (m, 2H), 7.57 (d, $J=8.6$ Hz, 1H), 7.50 (d, $J=16.5$ Hz, 1H), 7.36 (d, $J=16.5$ Hz, 1H), 7.34 (m, 1H), 7.19 (t, $J=7.5$ Hz, 1H), 7.15~7.10 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.3, 161.8 (d, $J=250.8$ Hz), 152.1, 132.2, 131.5, 130.6, 130.1, 128.0, 128.9, 126.3, 124.5, 123.4 (d, $J=23.5$ Hz), 122.9, 116.3 (d, $J=21.4$ Hz), 103.6, 29.7. HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{Br}_2\text{FNS}$ $[\text{M}+\text{H}]^+$ 437.8958, found 437.8958.

(*E*)-5-Bromo-4-(4-bromophenyl)-2-(4-fluorostyryl)thiazole (**6w**): Yellow solid, yield 50.1%. m.p. 179~180 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.88~7.84 (m, 2H), 7.63~7.59 (m, 2H), 7.56~7.50 (m, 2H), 7.33 (d, $J=16.5$ Hz, 1H), 7.16 (d, $J=16.5$ Hz, 1H), 7.14~7.07 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 166.2, 164.2 (d, $J=251.8$ Hz), 152.0, 145.0, 137.8, 134.1, 132.2, 131.5, 130.1, 128.9 (d, $J=8.4$ Hz), 127.5, 125.0, 122.8, 120.7, 116.1 (d, $J=21.9$ Hz). HRMS (ESI) calcd $\text{C}_{17}\text{H}_{11}\text{Br}_2\text{FNS}$ $[\text{M}+\text{H}]^+$ 437.8958, found 437.8955.

4.3 Bioassay of Top1-mediated relaxation assay

Compounds **6a**~**6w** were screened for their Top1 inhibitory activity using Top1-mediated relaxation assay.^[33] Briefly, reaction mixture (20 μL) with 0.5 μg supercoiled pBR322 DNA in Relaxation Buffer (10 $\text{mmol}\cdot\text{L}^{-1}$ Tris-HCl, pH 7.5, 50 $\text{mmol}\cdot\text{L}^{-1}$ KCl, 5 $\text{mmol}\cdot\text{L}^{-1}$ MgCl_2 , 15 $\mu\text{g}/\text{mL}$ BSA, 40 $\mu\text{g}/\text{mL}$ DTT) was incubated with 1 unit of calf thymus Top1 in the absence or in the presence of compound for 0.5 h at 37 °C. Then the reaction solution was added with 6 $\times$ loading buffer (4 μL), and was analyzed using a 0.8 % agarose gel in TBE buffer at 4.6 V/cm for 1.5 h. Gel was stained with 1 $\times$ gel red for 0.5 h and subsequently visualized with a UV transilluminator.

4.4 Molecular modelling

The PDB file for the X-ray crystal structure of Top1 was obtained using the protein data bank code 1K4T.^[34] The ternary complex ligand centroid coordinates for docking were defined using the ligand in the Top1-DNA-Topotecan crystal structure as the centre of the binding pocket. The protein structure was cleaned, inspected for errors and missing residues, hydrogens were added, and the water molecules and the ligand were deleted. The **6k** to be modeled was constructed and optimized using ChemDraw and saved in SDF file formats and was corrected using MOE software. Hydrogens were added, and the ligand was minimized by the conjugate gradient method using the MMFF94x force field with MMFF94 charges, a distance-dependent dielectric function, and a 418 $\text{J}\cdot\text{mol}^{-1}\cdot\text{nm}^{-1}$ energy gradient convergence criterion. Induced fit was

used for docking with the default parameters. The top 30 docking poses of the ligand were inspected visually following the docking runs. The highest-ranked pose was merged into the crystal structure. Energy minimizations were performed for the highest-ranked pose for the ligand. The AMBER forcefield was utilized within the MOE software for energy minimization.

4.5 Cell culture and MTT assay

The cells were cultured on RPMI-1640 medium at 37 °C in a humidified atmosphere with 5% CO_2 . The cytotoxicity of compounds **6a**~**6w** was evaluated through MTT assay against two human cancer cell lines (human breast cancer MCF-7, human colon cancer HCT116). For the compound treatment experiments, the cancer cells were treated with the compounds **6a**~**6w** (predissolved in DMSO) at a five-dose assay ranging from 10^{-8} to 10^{-4} $\text{mol}\cdot\text{L}^{-1}$. After incubation for 72 h at 37 °C, MTT solution (20 μL , 2.5 mg/mL) in PBS (PBS without MTT as the blank) was added to each well of the culture plate. After being incubated for 4 h, the formazan crystal formed in the well was dissolved in 100 μL of DMSO for optical density at 570 nm. The IC_{50} value was calculated by nonlinear regression analysis (GraphPad Prism). Every experiment was conducted for three independent times.

Supporting Information The ^1H NMR, ^{13}C NMR and ESI-HRMS spectra of title compounds **6a**~**6w**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Champoux, J. J. *Annu. Rev. Biochem.* **2001**, 70, 369.
- [2] Stewart, L.; Redinbo, M. R.; Qui, X.; Hol, W. G. J.; Champoux, J. *J. Science* **1998**, 279, 1534.
- [3] Berger, J. M.; Gamblin, S. J.; Harrison, S. C.; Wang, J. C. *Nature* **1996**, 379, 225.
- [4] Lian, Q.; Xu, J.; Yan, S.; Huang, M.; Ding, H.; Sun, X.; Bi, A.; Ding, J.; Sun, B.; Geng, M. *Cell Res.* **2017**, 27, 784.
- [5] Pommier, Y. *Nat. Rev. Cancer* **2006**, 6, 789.
- [6] Thomas, A.; Pommier, Y. *Clin. Cancer Res.* **2019**, 25, 6581.
- [7] Newman, D. J.; Cragg, G. M. *J. Nat. Prod.* **2020**, 83, 770.
- [8] Loiseleur, O. *Chimia* **2017**, 71, 810.
- [9] Lin, Z. Q.; Xia, W. L.; Liu, R. Y.; Jiang, S. H.; Ma, Z. Q. *Chin. J. Org. Chem.* **2020**, 40, 2980 (in Chinese). (林芷晴, 夏婉铃, 刘仁义, 姜少华, 马志强, 有机化学, **2020**, 40, 2980.)
- [10] Rimando, A. M.; Cuendet, M.; Desmarchelier, C.; Mehta, R. G.; Pezzuto, J. M.; Duke, S. O. *J. Agric. Food Chem.* **2002**, 50, 3453.
- [11] Kimura, Y.; Okuda, H.; Arichi, S. *Biochim. Biophys. Acta* **1985**, 834, 275.
- [12] Stivala, L. A.; Savio, M.; Carafoli, F.; Perucca, P.; Bianchi, L.; Maga, G.; Forti, L.; Pagnoni, U. M.; Albini, A.; Prosperi, E.; Vannini, V. *J. Biol. Chem.* **2001**, 276, 22586.
- [13] Mahady, G. B.; Pendland, S. L.; Chadwick, L. R. *Am. J. Gastroenterol.* **2003**, 98, 1440.
- [14] Shi, D.; An, R.; Zhang, W. B.; Zhang, G. L.; Yu, Z. G. *J. Agric. Food Chem.* **2017**, 65, 60.
- [15] Kronenwerth, M.; Dauth, C.; Kaiser, M.; Pemberton, I.; Bode, H. B. *Eur. J. Org. Chem.* **2014**, 36, 8026.
- [16] Mizuno, C. S.; Schrader, K. K.; Rimando, A. M. *J. Agric. Food Chem.* **2008**, 56, 9140.

- [17] Weng, J. Q.; Ali, A.; Estep, A.; Becnel, J.; Meyer, S. L. F.; Wedge, D. E.; Jacob, M.; Rimando, A. M. *Chem. Biodiversity* **2016**, *13*, 1165.
- [18] Lu, Q.; Yu, Q.; Zhu, Y. B.; Weng, J. Q.; Yuan, J.; Hu, D. X.; Chen, J.; Liu, X. H.; Tan, C. X. *J. Mol. Struct.* **2019**, *1180*, 780.
- [19] Mabkhot, Y. N.; Alharbi, M. M.; Al-Showiman, S. S.; Ghabbour, H. A.; Kheder, N. A.; Soliman, S. M.; Frey, W. *Chem. Cent. J.* **2018**, *12*, 56.
- [20] Kouatly, O.; Geronikaki, A.; Kamoutsis, C.; Hadjipavlou-Litina, D.; Eleftheriou, P. *Eur. J. Med. Chem.* **2009**, *44*, 1198.
- [21] Pember, S. O.; Mejia, G. L.; Price, T. J.; Pasteris, R. J. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 2965.
- [22] Koti, R. S.; Kolavi, G. D.; Hegde, V. S.; Khazi, I. M. *Indian J. Chem. B* **2006**, *45*, 1900.
- [23] Weng, J. Q.; Tan, C. X.; Liu, X. H. *J. Pestic. Sci.* **2012**, *37*, 164.
- [24] Yu, H. B.; Qin, Z. F.; Dai, H.; Zhang, X.; Qin, X.; Wang, T. T.; Fang, J. X. *J. Agric. Food Chem.* **2018**, *56*, 11356.
- [25] Weng, J. Q.; Liu, X. H.; Huang, H.; Tan, C. X.; Chen, J. *Molecules* **2012**, *17*, 989.
- [26] Zhang, J. H.; Zhu, Y. B.; Weng, J. Q.; Yu, Q.; Yuan, J.; Chen, J. *Chin. J. Org. Chem.* **2020**, *40*, 1055 (in Chinese). (章俊辉, 朱亚波, 翁建全, 余茜, 袁静, 陈杰, 有机化学, **2020**, *40*, 1055.)
- [27] Kong, Y. L.; Xu, W. X.; Liu, X. H.; Weng, J. Q. *Chin. Chem. Lett.* **2020**, *31*, 3245.
- [28] Xu, W. X.; Ye, F. X.; Liu, X. H.; Weng, J. Q. *Tetrahedron Lett.* **2020**, *61*, 151807.
- [29] Dou, D. F.; He, G. J.; Li, Y.; Lai, Z.; Wei, L. Q.; Alliston, K. R.; Lushington, G. H.; Eichhorn, D. M.; Groutas, W. C. *Bioorg. Med. Chem.* **2010**, *18*, 1093.
- [30] Kocabas, E.; Sariguney, A. B.; Coskun, A. *Heterocycles* **2010**, *81*, 2849.
- [31] Zhu, D. J.; Chen, J. X.; Wu, D. Z.; Liu, M. C.; Ding, J. C.; Wu, H. Y. *J. Chem. Res.* **2009**, (2), 84.
- [32] Wetherill, J. P.; Hann, R. M. *J. Am. Chem. Soc.* **1934**, *56*, 970.
- [33] Pommier, Y.; Covey, J. M.; Kerngan, D.; Markovits, J.; Pham, R. *Nucleic Acids Res.* **1987**, *15*, 6713.
- [34] Staker, B. L.; Hjerrild, K.; Feese, M. D.; Behnke, C. A.; Burgin, A. B.; Jr.; Stewart, L. P. *Natl. Acad. Sci. U. S. A.* **2002**, *99*, 15387.

(Li, L.; Fan, Y.)