无配体参与的铁催化分子内 C(sp³)—H 键胺化合成咪唑啉酮

邬林洋 钟大猷 刘文博*

(武汉大学化学与分子科学学院 索维奇国际分子科学研究中心
有机硅化合物及材料教育部工程研究中心 武汉 430072)

摘要 金属氮宾的分子内碳氢键插入反应是合成氮杂环最有效的方法之一。以 2-叠氮基-*N,N*-二苄基乙酰胺为底物, 发展了无配体、无外加氧化剂条件下的铁催化分子内 sp³-碳氢键胺化反应, 以中等到良好的收率合成了一系列咪唑啉酮类化合物。

关键词 铁催化; 氮宾插入反应; 咪唑啉酮; 氮杂环合成

Ligand-Free Iron-Catalyzed Intramolecular Amination of C(sp³)—H Bond for the Synthesis of Imidazolinones

Wu, Linyang Zhong, Dayou Liu, Wenbo*

(Savage Center for Molecular Sciences, Engineering Research Center of Organosilicon Compounds & Materials, Ministry of Education, College of Chemistry and Molecular Sciences, Wuhan University, Wuhan 430072)

Abstract Transition-metal-nitrenoids intramolecular C—H insertion is one of the most effective methods to synthesize nitrogen-containing heterocycles. Using 2-azido-*N,N*-dibenzylacetamides as substrates, an iron-catalyzed intramolecular C(sp³)—H bond amination reaction under ligand-free and external oxidant-free conditions has been developed. A series of imidazolinones were synthesized in moderate to good yields.

Keywords iron catalysis; nitrene insertion; imidazolinone; synthesis of *N*-containing heterocycles

1 Introduction

It is of great importance to construct C—N bond due to the wide existence of nitrogen-containing structures in natural products and drug molecules.^[1-2] Conventionally, transition-metal catalyzed cross-coupling reaction was widely used for C—N bond formation albeit necessitating pre-functionalization of substrates (e.g. Buchwald-Hartwig coupling^[3-4] and Ullmann-Ma coupling reaction^[5-6]). In past few decades, iron-catalyzed C—H bond amination has been received much attention due to its high atom efficiency (Scheme 1a).^[7] Tremendously remarkable results were achieved with both activated and inert C(sp³)—H bond-containing substrates.^[8-23]

The first iron-catalyzed C—H nitrene insertion was reported by Breslow in 1982 using [Fe(TPP)Cl] as a catalyst.^[24] After that, many heme-type and non-heme-type catalysts were developed, such as porphyrin-based iron

complex,^[22] phthalocyanine-based iron complex,^[13] and iron-dipyrinato complex,^[25] greatly promoting the advances of the field. In addition, intra- and inter-molecular C—H aminations were also realized using iron salts as catalysts.^[26-29] However, only limited examples were released to date.

In another hand, sulfamates were among the most investigated substrates which were used as the nitrene precursors in Fe-catalyzed C—H amination, but those reactions required stoichiometric external oxidants.^[13,15,24] In comparison, readily available organoazides were superior nitrene precursors with N₂ as the only by-product under oxidant-free conditions, as evidenced by significant achievements made by a number of groups (Scheme 1b). For instance, Che group^[30] successfully realized an aliphatic C—H amination reaction with [Fe^{III}(F₂₀TPP)Cl] as a catalyst. Recently, Yu group also reported the C—H amination reactions of organoazides catalyzed by Fe(II)-diketi-

* Corresponding author. E-mail: wenbolu@whu.edu.cn

Received April 26, 2021; revised May 9, 2021; published online May 13, 2021.

Dedicated to the 100th anniversary of Chemistry at Nankai University.

Project supported by the National Natural Science Foundation of China (Nos. 21971198, 21772148).

国家自然科学基金(Nos. 21971198, 21772148)资助项目.

electron-donating group (OMe) delivered the amination product in 55% yield. However, an electron-withdrawing group (CF₃)-containing substrate led to the formation of product **2h** in higher yield (72%). Heteroaromatic substrate, 2-azido-*N,N*-bis(thiophen-2-ylmethyl)acetamide, was also compatible by this protocol, delivering the corresponding product **2i** in 75% yield. For the substrate **1j** bearing both electron-donating group (OMe) and electron-withdrawing group (NO₂), a 1.6 : 1 mixture of product **2j** and **2j'** was obtained in 60% yield with the electron-rich C—H bond slightly favored. This trend is consistent with Ru-catalyzed amination of aliphatic azide as reported by Meggers.^[40] Surprisingly, the reaction was sensitive to the steric hindrance since no product was detected using 2-azido-*N,N*-dibenzyl-2-methylpropanamide (**1k**) as the substrate.

Table 2 Substrate scope^a

Entry	R ¹	R ²	R ³	Time/h	Yield ^b / % of 2
1	Ph	Ph	H	1	59 (2a)
2	4-MeC ₆ H ₄	4-MeC ₆ H ₄	H	6	52 (2b)
3	4- ^t BuC ₆ H ₄	4- ^t BuC ₆ H ₄	H	6	63 (2c)
4	4-FC ₆ H ₄	4-FC ₆ H ₄	H	5	47 (2d)
5	4-ClC ₆ H ₄	4-ClC ₆ H ₄	H	5	60 (2e)
6	4-BrC ₆ H ₄	4-BrC ₆ H ₄	H	5	60 (2f)
7	4-MeOC ₆ H ₄	4-MeOC ₆ H ₄	H	4	55 (2g)
8	4-CF ₃ C ₆ H ₄	4-CF ₃ C ₆ H ₄	H	2	72 (2h)
9	2-Thienyl	2-Thienyl	H	1	75 (2i)
10 ^c	4-MeOC ₆ H ₄	4-O ₂ NC ₆ H ₄	H	2	39 (2j)
	4-O ₂ NC ₆ H ₄	4-MeOC ₆ H ₄	H		21 (2j')
11	Ph	Ph	Me	6	NR

^a Reaction conditions: FeBr₂ (5 mol%), **1** (0.4 mmol), Boc₂O (0.48 mmol, 1.2 equiv.), DCE (2 mL), 1~6 h. ^b Isolated yield. ^c The reaction was carried out with substrate **1j** and a 1.6 : 1 mixture of **2j** and **2j'** were obtained.

3 Conclusions

In summary, an iron-catalyzed intramolecular amination of aliphatic azide for the synthesis of imidazolinones has been developed. A series of imidazolinones were synthesized in moderate yields under ligand- and oxidant-free condition. This research provides a convenient method for the C(sp³)—H amination to deliver *N*-containing heterocycles.

4 Experimental section

4.1 Instruments and reagents

Reactions were monitored by thin-layer chromatography (TLC). TLC was performed using Huanghai (8±0.2) μm precoated glass plates (0.25 mm) and visualized by UV fluorescence quenching and KMnO₄. Huanghai silica gel

(particle size 200~300 mesh) was used for column chromatography. ¹H NMR spectra were recorded at room temperature on a Bruker ADVANCE III 400 MHz spectrometer and were reported relative to residual CDCl₃ (δ 7.26). ¹³C NMR spectra were recorded on a Bruker ADVANCE III 400 MHz spectrometer (100 MHz) and were reported relative to CDCl₃ (δ 77.16). High resolution mass spectra (HRMS) were obtained by use of a Bruker Compact TOF mass spectrometer in electrospray ionization mode (ESI⁺).

Unless otherwise noted, all reagents were purchased commercially and used without further purification. Petroleum ester (PE, 60~90 °C), ethyl acetate (EA) and dichloromethane (CH₂Cl₂) were used as eluent for silica gel chromatography. Dry solvents were purchased commercially or were dried by passage through an activated alumina column under argon.

4.2 General procedure for iron-catalyzed intramolecular amination of C(sp³)—H bond

To an oven-dried 10 mL sealed tube equipped with a stirring bar were added FeBr₂ (4.4 mg, 5 mol%), substrate **1** (0.4 mmol, 1.0 equiv.), Boc₂O (104.6 mg, 0.48 mmol, 1.2 equiv.) and DCE (2 mL) under an argon atmosphere. The reaction mixture was stirred at 65 °C as monitored by TLC. After the completion of the reaction, the mixture was filtered. After removing the solvent, the crude mixture was purified by flash column chromatography (200~300 mesh SiO₂) to give the desired product **2**.

tert-Butyl 3-benzyl-4-oxo-2-phenylimidazolidine-1-carboxylate (**2a**): White solid (83.1 mg, 59% yield). m.p. 138~140 °C (Ref.^[40] 136~138 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.43~7.37 (m, 2H), 7.35~7.31 (m, 3H), 7.23~7.12 (m, 5H), 5.76~5.46 (m, 1H), 5.09 (d, *J*=14.9 Hz, 1H), 4.28~4.17 (m, 2H), 3.44~3.26 (m, 1H), 1.43~1.09 (m, 9H).

tert-Butyl 3-(4-methylbenzyl)-4-oxo-2-(*p*-tolyl)imidazolidine-1-carboxylate (**2b**):^[41] White solid (79.3 mg, 52% yield). m.p. 134~136 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.24~7.01 (m, 8H), 5.76~5.40 (m, 1H), 5.05 (d, *J*=14.7 Hz, 1H), 4.29~4.13 (m, 2H), 3.37~3.19 (m, 1H), 2.38 (s, 3H), 2.35 (s, 3H), 1.44~1.09 (m, 9H).

tert-Butyl 3-(4-(*tert*-butyl)benzyl)-2-(4-(*tert*-butyl)phenyl)-4-oxoimidazolidine-1-carboxylate (**2c**): White solid (117.7 mg, 63% yield). m.p. 136~138 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.40 (d, *J*=7.5 Hz, 2H), 7.37~7.30 (m, 2H), 7.24~7.06 (m, 4H), 5.79~5.49 (m, 1H), 5.08~4.97 (m, 1H), 4.31~4.10 (m, 2H), 3.43~3.30 (m, 1H), 1.33 (s, 9H), 1.32 (s, 9H), 1.39~1.15 (m, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.7/167.5, 152.8/152.5, 151.1, 135.1, 134.4, 132.5/132.3, 128.6/128.4, 127.3/127.1, 126.0, 125.8/125.6, 81.2, 74.6/74.2, 49.1/48.5, 43.1, 34.9/ 34.7, 31.5/31.4, 28.4/28.1; HRMS (ESI⁺) calcd for C₂₉H₄₁N₂O₃ [M+H]⁺ 465.3112, found 465.3119.

tert-Butyl 3-(4-fluorobenzyl)-2-(4-fluorophenyl)-4-oxoimidazolidine-1-carboxylate (**2d**):^[41] White solid (73.5 mg, 47% yield). m.p. 134~136 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.31~6.92 (m, 8H), 5.73~5.44 (m, 1H), 4.99

(m, 1H), 4.29~4.12 (m, 2H), 3.48~3.31 (m, 1H), 1.43~1.11 (m, 9H).

tert-Butyl 3-(4-chlorobenzyl)-2-(4-chlorophenyl)-4-oxoimidazolidine-1-carboxylate (**2e**):^[41] White solid (100.7 mg, 60% yield). m.p. 110~112 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.37 (d, *J*=8.4 Hz, 2H), 7.34~7.09 (m, 4H), 7.06 (d, *J*=8.2 Hz, 2H), 5.71~5.41 (m, 1H), 5.00 (d, *J*=15.0 Hz, 1H), 4.32~4.10 (m, 2H), 3.45~3.29 (m, 1H), 1.43~1.10 (m, 9H).

tert-Butyl 3-(4-bromobenzyl)-2-(4-bromophenyl)-4-oxoimidazolidine-1-carboxylate (**2f**): White solid (122.4 mg, 60% yield). m.p. 128~130 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.53 (d, *J*=8.3 Hz, 2H), 7.49~7.42 (m, 2H), 7.17~7.02 (m, 2H), 6.99 (d, *J*=8.2 Hz, 2H), 5.68~5.37 (m, 1H), 4.98 (d, *J*=15.0 Hz, 1H), 4.29~4.10 (m, 2H), 3.47~3.25 (m, 1H), 1.47~1.12 (m, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.7/167.4, 152.6/152.1, 136.9, 136.2, 134.0/133.9, 132.2/132.1, 130.2/130.1, 129.0, 123.7, 122.2, 81.8/81.6, 74.1/73.9, 48.7/48.2, 42.9, 28.2/28.1; HRMS (ESI⁺) calcd for C₂₁H₂₃Br₂N₂O₃ [M+H]⁺ 511.0051, found 511.0069.

tert-Butyl 3-(4-methoxybenzyl)-2-(4-methoxyphenyl)-4-oxoimidazolidine-1-carboxylate (**2g**):^[41] White solid (88.7 mg, 55% yield). m.p. 76~78 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.23~7.09 (m, 2H), 7.07 (d, *J*=8.3 Hz, 2H), 6.91 (d, *J*=8.4 Hz, 2H), 6.88~6.81 (m, 2H), 5.77~5.38 (m, 1H), 5.02 (d, *J*=14.8 Hz, 1H), 4.24~4.13 (m, 2H), 3.84 (s, 3H), 3.81 (s, 3H), 3.42~3.19 (m, 1H), 1.43~1.11 (m, 9H).

tert-Butyl 4-oxo-3-(4-(trifluoromethyl)benzyl)-2-(4-(trifluoromethyl)phenyl)imidazolidine-1-carboxylate (**2h**):^[41] White solid (139.8 mg, 72% yield). m.p. 72~74 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.64 (d, *J*=8.1 Hz, 2H), 7.57 (d, *J*=8.0 Hz, 2H), 7.40~7.27 (m, 2H), 7.23 (d, *J*=8.0 Hz, 2H), 5.79~5.55 (m, 1H), 5.05~4.95 (m, 1H), 4.33~4.17 (m, 2H), 3.58 (d, *J*=15.2 Hz, 1H), 1.43~1.11 (m, 9H).

tert-Butyl 4-oxo-2-(thiophen-2-yl)-3-(thiophen-2-ylmethyl)imidazolidine-1-carboxylate (**2i**):^[41] Yellow solid (109.0 mg, 75% yield). m.p. 90~92 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.38 (d, *J*=4.6 Hz, 1H), 7.31~7.25 (m, 1H), 7.23~7.07 (m, 1H), 7.02 (dd, *J*=5.0, 3.5 Hz, 1H), 6.99~6.89 (m, 2H), 6.24~5.90 (m, 1H), 5.19 (d, *J*=15.5 Hz, 1H), 4.20~4.04 (m, 2H), 3.90~3.77 (m, 1H), 1.49~1.20 (m, 9H).

tert-Butyl 2-(4-methoxyphenyl)-3-(4-nitrobenzyl)-4-oxoimidazolidine-1-carboxylate (**2j**): White solid (66.5 mg, 39% yield). m.p. 38~40 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.17 (d, *J*=8.4 Hz, 2H), 7.33~7.26 (m, 2H), 7.20~7.03 (m, 2H), 6.92~6.86 (m, 2H), 5.73~5.42 (m, 1H), 4.99 (d, *J*=15.3 Hz, 1H), 4.32~4.16 (m, 2H), 3.83 (s, 3H), 3.68~3.60 (m, 1H), 1.44~1.14 (m, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.9, 160.7, 152.3, 147.8, 142.9, 129.3, 128.7, 124.2, 114.5, 114.3, 81.6, 74.9/74.6, 55.5, 48.8/48.3, 43.1, 28.4/28.2; HRMS (ESI⁺) calcd for C₂₂H₂₆N₃O₆ [M+H]⁺ 428.1816, found 428.1816.

tert-Butyl 3-(4-methoxybenzyl)-2-(4-nitrophenyl)-4-

oxoimidazolidine-1-carboxylate (**2j'**): White solid (35.7 mg, 21% yield). m.p. 54~56 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.25 (d, *J*=8.6 Hz, 2H), 7.51~7.30 (m, 2H), 7.05~6.99 (m, 2H), 6.87~6.83 (m, 2H), 5.81~5.57 (m, 1H), 5.02 (d, *J*=15.0 Hz, 1H), 4.30~4.12 (m, 2H), 3.81 (s, 3H), 3.45~3.25 (m, 1H), 1.43~1.08 (m, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.7/167.4, 159.7, 152.7/152.0, 148.7, 145.4/144.6, 130.0/129.9, 128.6, 126.6/126.5, 124.3/124.1, 114.5, 82.2/82.0, 73.7/73.4, 55.5, 49.0/48.5, 43.3, 28.4/28.2; HRMS (ESI⁺) calcd for C₂₂H₂₆N₃O₆ [M+H]⁺ 428.1816, found 428.1819.

Supporting Information Synthetic information of substrates **1**, and ¹H NMR, ¹³C NMR spectra of compound **1c**, **1f**, **1j**, **2c**, **2f**, **2j** and **2j'**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Trowbridge, A.; Walton, S. M.; Gaunt, M. J. *Chem. Rev.* **2020**, *120*, 2613.
- [2] Hazelard, D.; Nocquet, P.; Compain, P. *Org. Chem. Front.* **2017**, *4*, 2500.
- [3] Ruiz-Castillo, P.; Buchwald, S. L. *Chem. Rev.* **2016**, *116*, 12564.
- [4] Baumgartner, L. M.; Dennis, J. M.; White, N. A.; Buchwald, S. L.; Jensen, K. F. *Org. Process Res. Dev.* **2019**, *23*, 1594.
- [5] Dai, L. *Prog. Chem.* **2018**, *30*, 1257 (in Chinese). (戴立信, 化学进展, **2018**, *30*, 1257.)
- [6] Cai, Q.; Zhou, W. *Chin. J. Chem.* **2020**, *38*, 879.
- [7] Park, Y.; Kim, Y.; Chang, S. *Chem. Rev.* **2017**, *117*, 9247.
- [8] Wang, P.; Deng, L. *Chin. J. Chem.* **2018**, *36*, 1222.
- [9] Plietker, B.; Röske, A. *Catal. Sci. Technol.* **2019**, *9*, 4188.
- [10] Liu, Y.; You, T.; Wang, H. X.; Tang, Z.; Zhou, C. Y.; Che, C.-M. *Chem. Soc. Rev.* **2020**, *49*, 5310.
- [11] Shi, D.; Bao, H.; Xu, Z.; Liu, Y. *Chin. J. Org. Chem.* **2017**, *37*, 1290 (in Chinese). (施冬冬, 鲍汉扬, 徐峥, 刘运奎, 有机化学, **2017**, *37*, 1290.)
- [12] Xu, L.; Xu, H.; Lin, H.; Dai, H. *Chin. J. Org. Chem.* **2018**, *38*, 1940 (in Chinese). (徐琳琳, 徐辉, 林海霞, 戴辉雄, 有机化学, **2018**, *38*, 1940.)
- [13] Paradine, S. M.; White, M. C. *J. Am. Chem. Soc.* **2012**, *134*, 2036.
- [14] Liang, L.; Lü, H.; Yu, Y.; Wang, P.; Zhang, J. *Dalton Trans.* **2012**, *41*, 1457.
- [15] Liu, Y.; Guan, X.; Wong, E. L.; Liu, P.; Huang, J.; Che, C.-M. *J. Am. Chem. Soc.* **2013**, *135*, 7194.
- [16] Wang, H.; Li, Y.; Wang, Z.; Lou, J.; Xiao, Y.; Qiu, G.; Hu, X.; Altenbach, H.; Liu, P. *RSC Adv.* **2014**, *4*, 25287.
- [17] Tseberlidis, G.; Zardi, P.; Caselli, A.; Cancogni, D.; Fusari, M.; Lay, L.; Gallo, E. *Organometallics* **2015**, *34*, 3774.
- [18] Thacker, N. C.; Lin, Z.; Zhang, T.; Gilhula, J. C.; Abney, C. W.; Lin, W.-B. *J. Am. Chem. Soc.* **2016**, *138*, 3501.
- [19] Alt, I. T.; Plietker, B. *Angew. Chem., Int. Ed.* **2016**, *55*, 1519.
- [20] Thacker, N. C.; Ji, P.; Lin, Z.; Urban, A.; Lin, W.-B. *Faraday Discuss.* **2017**, *21*, 303.
- [21] Shing, K.; Liu, Y.; Cao, B.; Chang, X.; You, T.; Che, C.-M. *Angew. Chem., Int. Ed.* **2018**, *57*, 11947.
- [22] Liu, Y.; You, T.; Wang, T.; Che, C.-M. *Tetrahedron* **2019**, *75*, 130607.
- [23] Du, Y.; Xu, Z.; Zhou, C.; Che, C.-M. *Org. Lett.* **2019**, *21*, 895.
- [24] Breslow, R.; Gellman, S. H. *J. Chem. Soc., Chem. Commun.* **1982**, 1400.
- [25] King, E. R.; Betley, T. A. *Inorg. Chem.* **2009**, *48*, 2361.
- [26] Wang, Z.; Zhang, Y.; Fu, H.; Jiang, Y.; Zhao, Y. *Org. Lett.* **2008**, *10*, 1863.

- [27] Shen, M.; Driver, T. G. *Org. Lett.* **2008**, *10*, 3367.
- [28] Bonnamour, J.; Bolm, C. *Org. Lett.* **2011**, *13*, 2012.
- [29] Nguyen, Q.; Nguyen, T.; Driver, T. G. *J. Am. Chem. Soc.* **2013**, *135*, 620.
- [30] Liu, Y.; Che, C.-M. *Chem.-Eur. J.* **2010**, *16*, 10494.
- [31] Zhao, X.; Liang, S.; Fan, X.; Yang, T.; Yu, W. *Org. Lett.* **2019**, *21*, 1559.
- [32] Liang, S.; Zhao, X.; Yang, T.; Yu, W. *Org. Lett.* **2020**, *22*, 1961.
- [33] Kweon, J.; Chang, S. *Angew. Chem., Int. Ed.* **2021**, *60*, 2909.
- [34] Prasanthi, A. V. G.; Begum, S.; Srivastava, H. K.; Tiwari, S. K.; Singh, R. *ACS Catal.* **2018**, *8*, 8369.
- [35] Zhou, Z.; Tan, Y.; Yamahira, T.; Ivlev, S.; Xie, X.; Riedel, R.; Hemming, M.; Kimura, M.; Meggers, E. *Chem* **2020**, *6*, 2024.
- [36] Zhong, D.; Wu, D.; Zhang, Y.; Lu, Z.; Usman, M.; Liu, W.; Lu, X.; Liu, W.-B. *Org. Lett.* **2019**, *21*, 5808.
- [37] Liu, W.; Zhong, D.; Yu, C.; Zhang, Y.; Wu, D.; Feng, Y.; Cong, H.; Lu, X.; Liu, W.-B. *Org. Lett.* **2019**, *21*, 2673.
- [38] Zhang, Y.; Zhong, D.; Usman, M.; Xue, P.; Liu, W.-B. *Chin. J. Chem.* **2020**, *38*, 1651.
- [39] Zhong, D.; Wu, L.-Y.; Wang, X.-Z.; Liu, W.-B. *Chin. J. Chem.* **2021**, *39*, 855.
- [40] Liang, S.; Zhao, X.; Yang, T.; Yu, W. *Org. Lett.* **2020**, *22*, 1961.
- [41] Zhou, Z.; Chen, S.; Qin, J.; Nie, X.; Zheng, X.; Harms, K.; Riedel, R.; Houk, K. N.; Meggers, E. *Angew. Chem., Int. Ed.* **2019**, *58*, 1088.

(Cheng, F.)