t-BuOK 促进的还原脱磺/脱氢反应：选择性合成 2-取代的 1,3-共轭二烯及其应用

孙佳兵^a 苗涛^{*,a} 李品华^a 王磊^{*,a,b,c}

(^a 淮北师范大学化学与材料科学学院 绿色和精准合成化学及应用教育部重点实验室 安徽淮北 235000)

(^b 台州学院医药化工与材料工程学院 浙江台州 318000)

(^c 中国科学院上海有机化学研究所金属有机化学国家重点实验室 上海 200032)

摘要 发展了一种叔丁醇钾促进的(*Z*)-C6-乙烯基磺基菲啶的脱磺、脱氢串联反应，用于选择性地合成 1,3-共轭二烯。该反应无需过渡金属催化剂，底物适用范围广，以良好的收率得到相应的目标产物，为菲啶官能化的 1,3-共轭二烯的合成提供了新颖和有效的方法。此外，通过与富马腈的[4+2]环加成反应，实现了 1,3-共轭二烯的合成应用。

关键词 叔丁醇钾；脱磺反应；1,3-共轭二烯；环加成反应

t-BuOK-Mediated Reductive Desulfonylation/Dehydrogenation for the Synthesis of 2-Substituted 1,3-Dienes and Their [4+2] Cycloaddition Reactions

Sun, Jiabing^a Miao, Tao^{*,a} Li, Pinhua^a Wang, Lei^{*,a,b,c}

(^a Key Laboratory of Green and Precise Synthetic Chemistry and Applications, Ministry of Education; School of Chemistry and Materials Science, Huabei Normal University, Huabei, Anhui 235000)

(^b Advanced Research Institute and Department of Chemistry, Taizhou University, Taizhou, Zhejiang 318000)

(^c State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032)

Abstract A *t*-BuOK-mediated desulfonylation/dehydrogenation cascade reaction of (*Z*)-C6-(vinyl sulfone)phenanthridines is described. This process provides a novel and efficient strategy for the synthesis of 1,3-dienes with phenanthridinyl units in good yields under transition-metal-free conditions. Furthermore, the application of the generated 1,3-dienes in the [4+2] cycloaddition has been demonstrated by reacting with dienophiles such as fumaronitrile in a two-step procedure.

Keywords *t*-BuOK; desulfonylation; 1,3-conjugated dienes; cycloaddition reactions

1 Introduction

Conjugated dienes are of great importance since they not only are key structural motifs in various bioactive natural products and pharmaceuticals,^[1] but also are recognized to be highly valuable synthetic precursors in organic synthesis,^[2] and material science.^[3] Therefore, great efforts from synthetic organic chemists have been devoted to the developing efficient and practical methods for synthesis of conjugated dienes. Classical procedures for the preparation of 1,3-dienes mainly rely on the conventional elimination

reactions from allyl bromide, allyl alcohol or dihalogenated compound,^[4] and the carbonyl olefinations represented by the Wittig reaction^[5] (Scheme 1a) and its variants such as Horner-Wadsworth-Emmons reaction,^[6] Julia olefination,^[7] and Peterson reaction.^[8] Over the past decades, the transition-metal catalyzed C(sp²)—C(sp²) cross-couplings including Suzuki, Negishi, Stille, or Heck reactions have emerged as privileged and powerful strategies for the assembly of dienes moieties (Scheme 1b).^[9] Recently, several alternative and direct methods *via* oxidative cross-coupling

* Corresponding authors. E-mail: leiwang88@hotmail.com; taomiao@chnu.edu.cn

Received February 9, 2021; revised May 7, 2021; published online May 23, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21772062, 22071171, 21602072) and the Anhui University Natural Science Research Project (No. KJ2019ZD66).

国家自然科学基金(Nos. 21772062, 22071171, 21602072)和安徽省高校自然科学基金(No. KJ2019ZD66)资助项目。

of alkenes with acrylates and styrenes have also been developed by Loh,^[10] Yu,^[11] and others^[12] (Scheme 1c). Furthermore, the catalytic ene-yne cross-metathesis has provided a useful and atom-economical cross-coupling to access the 1,3-diene motif (Scheme 1d).^[13] Despite the effectiveness of the existing methods, the development of additional complementary processes with organic phosphine- and transition-metal-free conditions for the construction of substituted conjugated dienes remains challenging and highly desirable.

Previous works

a) Wittig reaction

b) Transition-metal catalyzed Heck/Suzuki couplings

c) Transition-metal catalyzed oxidative dehydrogenative couplings

d) Ru catalyzed eneyne cross-metathesis

This work

Scheme 1 Synthetic strategies for the construction of conjugated dienes

Vinylsulfones are important building blocks in synthetic organic chemistry due to their versatility as substrates and intermediates in numerous transformations, and have been extensively exploited in the synthesis of complex molecules including natural products.^[14] The most common transformations of vinylsulfones are Michael addition reactions with various nucleophiles,^[15] the substitution of the sulfone group by hydrogen^[16] and variety of metal-catalyzed desulfonylative cross-couplings.^[17] However, to the best of our knowledge, there have been no reports for the straightforward conversion of vinylsulfones into conjugated dienes *via* reductive process. Herein, we disclose a new, simple and efficient method for the preparation of 1,3-dienes with phenanthridinyl units by *t*-BuOK-mediated desulfonylation/dehydrogenation cascade reaction of (*Z*)-C6-(vinyl sulfone) phenanthridines and their subsequent transformations in [4+2] cycloaddition reactions (Scheme 1). It should be noted that the phenanthridine skeleton is an important substructure widely existing in natural and pharmaceutical compounds.^[18]

2 Results and discussion

In our initial investigation, (*Z*)-C6-(1-tosylhept-1-en-2-yl)phenanthridine (**1a**) was chosen as the model substrate to find the optimal reaction conditions, and the results are summarized in Table 1. To our delight, 40% yield of 6-(hepta-1,3-dien-2-yl)phenanthridine (**2a**) was obtained when 2.0 equiv of *t*-BuOK was added to a tetrahydrofuran (THF) solution of (*Z*)-C6-(1-tosylhept-1-en-2-yl)phenanthridine (**1a**) and the mixture was stirred at 70 °C for 18 h (Entry 1). Further survey of solvents showed that THF was the ideal solvent for this reaction, providing the highest yield of **2a** at 87% (Table 1, Entry 2). In addition, *p*-methyl phenylsulfonic acid was detected by HPLC-HRMS analysis after the acidification of reaction solution (for detail see Supporting Information, Scheme S1). Other solvents such as 1,4-dioxane, CH₃CN, toluene, *N,N*-dimethyl-formamide (DMF), acetone was less effective and gave the inferior yields of the desired products (Table 1, Entries 3~7). Dimethyl sulfoxide (DMSO) and 1,2-dichloroethane (DCE) were not suitable solvents and the reaction was prohibited completely (Table 1, Entries 8, 9). Furthermore, *t*-BuOK was an essential base in the formation of **2a**, *t*-BuONa was found to be harmful, and gave **2a** in 36% yield (Table 1,

Table 1 Optimization of reaction conditions^a

Entry	Base	Solvent	Yield ^b /%
1	<i>t</i> -BuOK	CH ₂ Cl ₂	40
2	<i>t</i>-BuOK	THF	87
3	<i>t</i> -BuOK	1,4-Dioxane	58
4	<i>t</i> -BuOK	CH ₃ CN	41
5	<i>t</i> -BuOK	Toluene	17
6	<i>t</i> -BuOK	DMF	30
7	<i>t</i> -BuOK	Acetone	41
8	<i>t</i> -BuOK	DMSO	0
9	<i>t</i> -BuOK	DCE	0
10	<i>t</i> -BuONa	THF	36
11	K ₂ CO ₃	THF	0
12	Na ₂ CO ₃	THF	0
13	Cs ₂ CO ₃	THF	0
14	Et ₃ N	THF	0
15	Pyridine	THF	0
16	<i>t</i> -BuOK	THF	86 ^c
17	<i>t</i> -BuOK	THF	85 ^d
18	<i>t</i> -BuOK	THF	58 ^e
19	<i>t</i> -BuOK	THF	60 ^f

^a Reaction conditions: (*Z*)-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1a**, 0.25 mmol), base (2.0 equiv.), solvent (3.0 mL), air, 70 °C oil bathing, 18 h.

^b Isolated yield based on **1a**. ^c In the N₂ atmosphere. ^d In the O₂ atmosphere.

^e Reaction time 12 hours. ^f 1.0 equiv. of *t*-BuOK was used.

Entry 10). Other bases such as K_2CO_3 , Na_2CO_3 , Cs_2CO_3 , Et_3N and pyridine instead of t -BuOK did not work in this reaction (Table 1, Entries 11~15). It should be noted that this transformation also proceeded well in comparable efficiency and yield when the reaction was carried out under O_2 or N_2 atmosphere (Table 1, Entries 16, 17). Shortening the reaction time to 12 h led to 65% yield of **2a** (Table 1, Entry 18). Meanwhile, the decreased yield was also observed when the amount of t -BuOK was reduced (Table 1, Entry 19).

With the optimized reaction conditions in hand, the above method was applied to the synthesis of various terminal 1,3-dienes (**2**) from (Z)-C6-(vinyl sulfone)phenanthridines (**1**). First, the substituent effects on the phenanthridinyl ring were investigated, and the results were depicted in Table 2. It was found that a variety of substrates bearing electron-donating (Me, t -Bu, MeO) or electron-withdrawing groups (F, Cl) on the phenanthridinyl ring were all viable, affording the terminal 1,3-dienes **2b**~**2m** in 72%~87% yields, so the electronic effects of the phenanthridinyl moiety are not pronounced. Subsequently, the effects of the

Table 2 Scope of phenanthridine moiety in the substrates^a

^a Reaction conditions: (Z)-C6-(vinyl sulfone)phenanthridine (**1**, 0.25 mmol), t -BuOK (2.0 equiv.) and THF (3.0 mL) at 70 °C oil bathing for 18 h in the air; isolated yield based on **1**. ^b 4.0 mmol **1a** was used.

substituted position were also studied. When the phenanthridinyl ring was substituted on the 2-position, the products **2b**~**2d** were achieved with 76%~80% yields. 3-Position substituted substrates generated the targeted products **2e**~**2g** in 77%~85% yields, the 9- or 10-position substituted substrates also worked well, and terminal 1,3-dienes **2h**~**2l** were isolated with good yields. In the case of disubstituted (Z)-C6-(vinyl sulfone)phenanthridine, the reaction underwent smoothly affording product **2m** in 80% yield. Furthermore, a gram-scale reaction for the synthesis of 1,3-diene was carried out under the optimal reaction conditions, and desired product **2a** was obtained with 79% yield (0.86 g).

The scope of (Z)-6-(1-tosylalkyl-1-en-2-yl)phenanthridines is shown in Table 3. Installation of longer alkyl chain with R^3 furnished the corresponding products **2n** and **2o** in 80% and 88% yields, respectively.

Table 3 Scope of (Z)-C6-(vinyl sulfone)phenanthridines with alkyl chain^a

^a Reaction conditions: (Z)-C6-(vinyl sulfone)phenanthridine (**1**, 0.25 mmol), t -BuOK (2.0 equiv.) and THF (3.0 mL) at 70 °C oil bathing for 18 h in the air; isolated yield based on **1**.

When ethyl or methyl group instead of propyl, the tosyl moiety was also smoothly cleaved in the presence of t -BuOK, leading to the desired products **2p** and **2q** with good yields. Moreover, (Z)-6-(1-tosyloct-1-en-2-yl)phenanthridine was successfully transformed into the product **2r** in 80% yield.

(Z)-6-(1-Tosylhex-1-en-2-yl)phenanthridines with Me, F or Cl substituents on phenanthridine ring were suitable substrates and gave the terminal 1,3-dienes **2s**~**2w** in 70%~84% yields.

An important synthetic application of 1,3-dienes leading to a variety of complex products is the Diels-Alder reaction.^[19] Hence, to demonstrate the usefulness of the cascade desulfonylation/dehydrogenation, the dienes **2** were tested in [4+2] cycloaddition with fumaronitrile by a two-step procedure. As shown in Table 4, a number of polysubstituted cyclohexenes (**4a~4n**) were observed with good yields. The reaction was also not sensitive to the electronegativity of the substituents on phenanthridine ring, and Me-, MeO-, F-, Cl-, and CF₃-substituted substrates ran smoothly to deliver the final products in 68%~84% yields. Fortunately, F, Cl and CN groups were inert under these reaction conditions and would allow further functionalization of the polysubstituted cyclohexenes. Furthermore, the

Table 4 Applications of 1,3-dienes by reacting with fumaronitrile^a

method was shown to be useful in the construction of polysubstituted cyclohexenes containing biologically relevant moieties such as F and CF₃, as in the case of **4c**, **4e**, **4g** and **4i~4k**.

To gain a good understanding of the reaction mechanism, several control experiments were conducted, as shown in Scheme 2. First, (*E*)-1-methyl-4-((2-phenylhex-1-en-1-yl)sulfonyl)benzene (**5**) was subjected to the reaction under the standard conditions, and no desired diene **6** was formed, yet the other isomer **5'** was isolated with 73% yield (Scheme 2a). Second, a well-known radical scavenger, 2,2,6,6-tetramethyl-piperidine-1-oxyl (TEMPO, 2.0 equiv.) was added in the reaction system of **1a**, the present reaction was not inhibited, and the diene **2a** was observed in 84% yield (Scheme 2b). This result implied that radical process was not involved in this transformation. Thirdly, the D₂O experiment indicated that the terminal hydrogen of product **2a** probably originated from H₂O by reacting with carbanion intermediate (Scheme 2c).

Scheme 2 Control experiments

Based on the above observations and previous reports,^[4] a proposed reaction pathway for the formation of products **2** is assumed, which agrees with an ionic reaction process (Scheme 3). Initially, *t*-BuOK abstracts a proton from (Z)-C6-(vinyl sulfone)phenanthridine (**1**) to form the anionic intermediate **A**, followed by an isomerization process to form intermediate **A'**. Then, the generated **A'** can react with H₂O by abstraction of hydrogen to give the compound **B**, which undergoes a *t*-BuOK-mediated C—S bond cleavage to give *p*-toluenesulfinic acid anion and cationic intermediate **C** with the assistance of phenanthridine group. Specifically, *p*-toluenesulfinic acid is successfully detected

by ESI-MS to support this proposed pathway. Subsequently, the allylic cation **C'** is produced from **C** through a resonance. Finally, losing of proton from **C'** affords 1,3-diene product **2**.

Scheme 3 Proposed mechanism

3 Conclusions

In summary, a novel *t*-BuOK-mediated desulfonylation/dehydrogenation cascade reaction of (Z)-C6-(vinyl sulfone)phenanthridines has been successfully realized under organic phosphine- and transition-metal-free conditions. This olefination provides an efficient method for the selective synthesis of phenanthridine substituted 1,3-dienes in good yields. Furthermore, the generated 1,3-dienes show good reactivity in the [4 + 2] cycloaddition, and have proven to be powerful for the synthesis of polysubstituted cyclohexenes by reacting with dienophiles such as fumaronitrile in a two-step procedure. Further investigation on utilizing *in situ* generated 1,3-dienes in synthetic transformations will be considered.

4 Experimental section

4.1 Instruments and reagents

The chemicals and solvents were purchased from commercial suppliers without further purification. All reactions were set up in air unless otherwise specified. ¹H NMR and ¹³C NMR spectra were measured on a Bruker Avance NMR spectrometer (600 MHz or 400 MHz, respectively) in

CDCl₃ or DMSO-*d*₆ solvent and recorded relative to internal standard tetramethylsilane. Flash column chromatography was performed using silica gel of 200~300 mesh. High resolution mass spectroscopic data of the products were obtained on an Agilent Technologies 6540 UHD Accurate-Mass Q-TOF LC/MS using ESI.

4.2 Typical procedure for the synthesis of (Z)-C6-(vinyl sulfone)phenanthridines

All (Z)-C6-(vinyl sulfone)phenanthridines were prepared according to the reported method.^[20] A representative procedure for the preparation of (Z)-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1a**) is shown as following: 1-heptyne (48.0 mg, 0.50 mmol), *p*-toluenesulfonic acid (117.0 mg, 0.75 mmol), pyridine (6.0 equiv.) and 2-isocyanobiphenyl (44.8 mg, 0.25 mmol) were added into an air-tight reaction tube (10.0 mL) with 4.5 mL of DMSO/H₂O (*V* : *V* = 8 : 1). The tube containing the reactants and solvent was evacuated by pump and back-filled with oxygen, then an oxygen balloon was connected to the reaction flask. The reaction was stirred at room temperature under photo-irradiation with UV LED (365 ± 5 nm, 3 W) for 24 h. After the reaction was completed, the solution was diluted with ethyl acetate (10 mL), washed with water (5 mL), and extracted with ethyl acetate (5 mL × 2). The organic layers were combined, dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. the resulting crude mixture was purified by flash chromatography on silica gel (petroleum ether/EtOAc, *V* : *V* = 5 : 1) to provide the desired product **1a** in 73% yield (78.3 mg).

The characterization data of (Z)-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1a**), (Z)-2-methyl-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1b**), (Z)-2-fluoro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1c**), (Z)-2-chloro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1d**), (Z)-8-methyl-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1i**), (Z)-8-chloro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1l**), (Z)-8,10-dichloro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1m**), (Z)-6-(1-tosyloct-1-en-2-yl)phenanthridine (**1o**), (Z)-6-(1-tosylhex-1-en-2-yl)phenanthridine (**1p**), (Z)-6-(1-tosylpent-1-en-2-yl)phenanthridine (**1q**) are consistent with the reference [20].

(Z)-3-Methyl-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1e**): 78.6 mg, 71% yield. Colorless solid, m.p. 113.6~115.4 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.53 (d, *J* = 8.3 Hz, 1H), 8.42 (d, *J* = 8.3 Hz, 1H), 7.91 (d, *J* = 8.1 Hz, 1H), 7.87 (s, 1H), 7.77 (t, *J* = 7.6 Hz, 1H), 7.59 (t, *J* = 7.5 Hz, 1H), 7.50~7.46 (m, 3H), 7.02 (d, *J* = 7.9 Hz, 2H), 6.70 (s, 1H), 2.61~2.59 (m, 5H), 2.24 (s, 3H), 1.57~1.54 (m, 2H), 1.30~1.22 (m, 4H), 0.82 (t, *J* = 6.8 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 158.3, 154.7, 143.8, 143.3, 138.7, 137.6, 132.3, 130.6, 129.3, 129.2, 129.1, 128.8, 127.9, 127.2, 126.8, 124.2, 121.8, 121.5, 38.7, 31.1, 26.2, 22.2, 21.4, 21.3, 13.8. HRMS (ESI) calcd for C₂₈H₃₀NO₂S [M + H]⁺: 444.1992, found 444.1994.

(Z)-3-Fluoro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1f**): 78.2 mg, 70% yield. Colorless solid, m.p. 170.1~171.8 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.53~8.52 (m, 2H), 7.96 (d, *J* = 8.0 Hz, 1H), 7.82~7.81 (m, 1H), 7.74~

7.72 (m, 1H), 7.65 (t, $J=7.5$ Hz, 1H), 7.54 (t, $J=7.8$ Hz, 2H), 7.43~7.40 (m, 1H), 7.12 (d, $J=8.0$ Hz, 2H), 6.70 (s, 1H), 2.60 (t, $J=7.8$ Hz, 2H), 2.32 (s, 3H), 1.58~1.55 (m, 2H), 1.33~1.23 (m, 4H), 0.83 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.6 (d, $J=246.9$ Hz), 160.0, 154.4, 144.5 (d, $J=11.6$ Hz), 144.1, 137.7, 132.2, 131.1, 129.3, 129.2, 127.9, 127.5, 127.3, 124.3, 124.2 (d, $J=9.5$ Hz), 121.9, 120.6 (d, $J=1.6$ Hz), 116.1 (d, $J=23.6$ Hz), 114.3 (d, $J=20.5$ Hz), 38.7, 31.1, 26.2, 22.2, 21.4, 13.8. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{27}\text{FNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 448.1741, found 448.1743.

(Z)-3-Chloro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1g**): 83.3 mg, 72% yield. Colorless solid, m.p. 184.8~186.1 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.54 (d, $J=8.2$ Hz, 1H), 8.47 (d, $J=8.8$ Hz, 1H), 8.06 (d, $J=2.2$ Hz, 1H), 7.97 (d, $J=7.9$ Hz, 1H), 7.84 (td, $J=7.6, 1.1$ Hz, 1H), 7.68 (t, $J=7.5$ Hz, 1H), 7.61 (dd, $J=8.8, 2.2$ Hz, 1H), 7.53 (d, $J=8.2$ Hz, 2H), 7.12 (d, $J=8.0$ Hz, 2H), 6.69 (s, 1H), 2.60 (t, $J=7.8$ Hz, 2H), 2.33 (s, 3H), 1.58~1.53 (m, 2H), 1.32~1.24 (m, 4H), 0.83 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 159.9, 154.3, 144.2, 143.9, 137.7, 134.3, 132.0, 131.2, 129.4, 129.3, 129.0, 127.9, 127.8, 127.6, 127.5, 124.7, 123.6, 122.4, 122.1, 38.7, 31.1, 26.3, 22.2, 21.5, 13.8. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{27}^{35}\text{ClNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 464.1446, found 464.1450; $\text{C}_{27}\text{H}_{27}^{37}\text{ClNO}_2\text{S}$: 466.1416, found 464.1419.

(Z)-10-Fluoro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1h**): 69.3 mg, 62% yield. White solid, m.p. 134.7~136.1 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.98 (d, $J=8.3$ Hz, 1H), 8.11 (d, $J=8.0$ Hz, 1H), 7.81~7.77 (m, 2H), 7.70 (t, $J=7.5$ Hz, 1H), 7.64~7.60 (m, 1H), 7.57~7.51 (m, 3H), 7.10 (d, $J=8.0$ Hz, 2H), 6.69 (s, 1H), 2.60 (t, $J=5.4$ Hz, 2H), 2.30 (s, 3H), 1.58~1.53 (m, 2H), 1.32~1.24 (m, 4H), 0.83 (t, $J=6.9$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.5 (d, $J=253.6$ Hz), 157.9 (d, $J=1.8$ Hz), 154.5, 144.1, 143.6, 137.6, 129.7, 129.4, 129.3, 129.0, 127.9, 127.8, 127.7 (d, $J=1.8$ Hz), 127.1, 127.0 (d, $J=2.7$ Hz), 123.4 (d, $J=3.4$ Hz), 121.7 (d, $J=18.6$ Hz), 121.6 (d, $J=13.1$ Hz), 117.3 (d, $J=23.3$ Hz), 38.6, 31.1, 26.2, 22.2, 21.4, 13.8. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{27}\text{FNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 448.1741, found 448.1739.

(Z)-8-Methoxy-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1j**): 80.3 mg, 70% yield. Colorless solid, m.p. 161.6~164.2 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.50 (d, $J=9.0$ Hz, 1H), 8.46 (d, $J=8.0$ Hz, 1H), 8.04 (d, $J=8.0$ Hz, 1H), 7.67 (t, $J=7.4$ Hz, 1H), 7.62 (t, $J=7.4$ Hz, 1H), 7.47~7.44 (m, 3H), 7.25 (d, $J=0.9$ Hz, 1H), 7.02 (d, $J=8.0$ Hz, 2H), 6.72 (s, 1H), 3.91 (s, 3H), 2.63 (t, $J=7.7$ Hz, 2H), 2.24 (s, 3H), 1.58~1.54 (m, 2H), 1.32~1.24 (m, 4H), 0.82 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 158.5, 157.4, 154.4, 144.0, 142.4, 137.4, 129.7, 129.3, 129.2, 127.9, 127.6, 127.1, 126.7, 125.9, 123.9, 123.8, 121.6, 121.5, 107.2, 55.4, 38.7, 31.1, 26.3, 22.2, 21.4, 13.8. HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{30}\text{NO}_3\text{S}$ $[\text{M}+\text{H}]^+$: 460.1941, found 460.1939.

(Z)-8-Fluoro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1k**): 79.3 mg, 71% yield. Colorless solid, m.p. 152.8~

155.2 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.62~8.60 (m, 1H), 8.50 (d, $J=7.3$ Hz, 1H), 8.10 (d, $J=8.1$ Hz, 1H), 7.74 (t, $J=7.5$ Hz, 1H), 7.69~7.66 (m, 1H), 7.57~7.54 (m, 4H), 7.13 (d, $J=8.0$ Hz, 2H), 6.71 (d, $J=0.8$ Hz, 1H), 2.61~2.59 (m, 2H), 2.32 (s, 3H), 1.57~1.55 (m, 2H), 1.32~1.25 (m, 4H), 0.83 (t, $J=6.9$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.2 (d, $J=247.4$ Hz), 157.7 (d, $J=3.7$ Hz), 154.3, 144.2, 142.9, 137.7, 129.9, 129.5, 129.4, 129.1, 128.6, 127.9, 127.5, 125.9 (d, $J=7.8$ Hz), 124.8 (d, $J=8.4$ Hz), 123.5, 122.0, 120.0 (d, $J=23.7$ Hz), 111.7 (d, $J=21.6$ Hz), 38.6, 31.1, 26.2, 22.2, 21.5, 13.8. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{27}\text{FNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 448.1741, found 448.1745.

(Z)-6-(1-Tosyldec-1-en-2-yl)phenanthridine (**1n**): 70.7 mg, 60% yield. Yellow liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.60 (d, $J=8.2$ Hz, 1H), 8.56 (d, $J=8.2$ Hz, 1H), 8.08 (d, $J=8.1$ Hz, 1H), 7.96 (d, $J=8.0$ Hz, 1H), 7.82 (t, $J=7.6$ Hz, 1H), 7.75~7.72 (m, 1H), 7.68~7.64 (m, 2H), 7.50 (d, $J=7.6$ Hz, 2H), 7.05 (d, $J=8.2$ Hz, 2H), 6.71 (d, $J=1.1$ Hz, 1H), 2.62~2.60 (m, 2H), 2.26 (s, 3H), 1.59~1.53 (m, 2H), 1.32~1.18 (m, 10H), 0.84 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 158.5, 154.7, 143.9, 143.2, 137.6, 132.3, 130.7, 129.8, 129.3, 129.2, 128.6, 127.9, 127.4, 127.3, 127.1, 124.7, 123.9, 122.12, 122.07, 38.8, 31.6, 29.1, 29.0, 28.9, 26.5, 22.5, 21.4, 14.0. HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{34}\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 472.2305, found 472.2306.

(Z)-3-Chloro-6-(1-tosyloct-1-en-2-yl)phenanthridine (**1r**): 78.7 mg, 66% yield. Colorless solid, m.p. 190.4~193.0 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.57 (d, $J=8.3$ Hz, 1H), 8.50 (d, $J=8.8$ Hz, 1H), 8.06 (d, $J=2.2$ Hz, 1H), 7.98 (d, $J=8.0$ Hz, 1H), 7.88~7.85 (m, 1H), 7.71~7.68 (m, 1H), 7.64 (dd, $J=8.7, 2.2$ Hz, 1H), 7.52 (d, $J=8.2$ Hz, 2H), 7.13 (d, $J=8.0$ Hz, 2H), 6.69 (t, $J=1.4$ Hz, 1H), 2.60 (t, $J=7.8$ Hz, 2H), 2.34 (s, 3H), 1.58~1.52 (m, 2H), 1.33~1.20 (m, 6H), 0.83 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.0, 154.4, 144.2, 140.0, 137.7, 134.3, 132.0, 131.2, 129.41, 129.37, 129.1, 128.0, 127.8, 127.7, 127.6, 124.8, 123.6, 122.5, 122.1, 38.8, 31.4, 28.7, 26.6, 22.4, 21.5, 13.9. HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{29}\text{ClNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 478.1602, found 478.1601; $\text{C}_{28}\text{H}_{29}^{37}\text{ClNO}_2\text{S}$: 480.1573, found 480.1577.

(Z)-3-Chloro-6-(1-tosylhex-1-en-2-yl)phenanthridine (**1s**): 79.7 mg, 71% yield. Colorless solid, m.p. 173.1~175.2 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.52~8.51 (m, 1H), 8.46~8.43 (m, 1H), 8.06 (s, 1H), 7.96 (d, $J=8.1$ Hz, 1H), 7.83~7.80 (m, 1H), 7.68~7.65 (m, 1H), 7.60~7.58 (m, 1H), 7.53 (d, $J=8.0$ Hz, 2H), 7.12 (d, $J=8.0$ Hz, 2H), 6.70 (d, $J=1.0$ Hz, 1H), 2.60 (t, $J=7.5$ Hz, 2H), 2.32 (s, 3H), 1.57~1.50 (m, 2H), 1.38~1.29 (m, 2H), 0.85 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 159.9, 154.3, 144.1, 143.9, 137.6, 134.2, 131.9, 131.2, 129.35, 129.28, 129.0, 127.9, 127.7, 127.6, 127.4, 124.6, 123.6, 122.4, 122.1, 38.4, 28.7, 22.1, 21.4, 13.7. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{25}\text{ClNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 450.1289, found 450.1288; $\text{C}_{26}\text{H}_{25}^{37}\text{ClNO}_2\text{S}$: 452.1260, found 452.1256.

(Z)-2-Methyl-6-(1-tosylhex-1-en-2-yl)phenanthridine (**1t**): 75.1 mg, 70% yield. Colorless solid, m.p. 123.1~

123.9 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.57 (d, $J=8.3$ Hz, 1H), 8.33 (s, 1H), 7.96 (d, $J=8.3$ Hz, 1H), 7.92 (d, $J=8.1$ Hz, 1H), 7.78 (t, $J=7.3$ Hz, 1H), 7.62 (t, $J=7.6$ Hz, 1H), 7.55 (d, $J=8.3$ Hz, 1H), 7.48 (d, $J=8.0$ Hz, 2H), 7.03 (d, $J=7.9$ Hz, 2H), 6.71 (s, 1H), 2.62~2.60 (m, 5H), 2.25 (s, 3H), 1.55~1.51 (m, 2H), 1.35~1.30 (m, 2H), 0.84 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 157.4, 154.8, 143.8, 141.5, 137.6, 137.0, 132.0, 130.5, 130.3, 129.4, 129.23, 129.16, 127.9, 127.21, 127.19, 124.7, 123.7, 122.0, 121.7, 38.5, 28.6, 22.1, 21.8, 21.3, 13.6. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{28}\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 430.1835, found 430.1837.

(Z)-2-Chloro-6-(1-tosylhex-1-en-2-yl)phenanthridine (**1u**): 76.3 mg, 68% yield. Colorless solid, m.p. 193.3~193.8 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.52~8.51 (m, 2H), 8.01 (d, $J=8.6$ Hz, 1H), 7.96 (d, $J=8.1$ Hz, 1H), 7.83 (t, $J=7.6$ Hz, 1H), 7.70~7.67 (m, 2H), 7.51 (d, $J=8.1$ Hz, 2H), 7.10 (d, $J=8.0$ Hz, 2H), 6.70 (s, 1H), 2.61 (t, $J=7.8$ Hz, 2H), 2.30 (s, 3H), 1.57~1.51 (m, 2H), 1.37~1.31 (m, 2H), 0.85 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 158.9, 154.3, 144.1, 141.6, 137.6, 133.0, 131.3, 131.2, 131.0, 129.3, 129.2, 128.1, 127.9, 127.4, 125.0, 124.9, 122.2, 121.8, 38.4, 28.7, 22.2, 21.4, 13.7. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{25}\text{ClNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 450.1289, found 450.1289; $\text{C}_{26}\text{H}_{25}^{37}\text{ClNO}_2\text{S}$: 452.1260, found 452.1264.

(Z)-10-Fluoro-6-(1-tosylhex-1-en-2-yl)phenanthridine (**1v**): 70.4 mg, 65% yield. Colorless solid, m.p. 169.5~170.9 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.95 (d, $J=7.2$ Hz, 1H), 8.14 (d, $J=8.2$ Hz, 1H), 7.98 (d, $J=8.0$ Hz, 1H), 7.92 (d, $J=8.0$ Hz, 2H), 7.76 (t, $J=7.4$ Hz, 1H), 7.69 (t, $J=6.9$ Hz, 1H), 7.64~7.61 (m, 1H), 7.59~7.55 (m, 1H), 7.38 (d, $J=8.0$ Hz, 2H), 6.51 (s, 1H), 3.31 (t, $J=7.5$ Hz, 2H), 2.45 (s, 3H), 1.41~1.34 (m, 4H), 0.81 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.5 (d, $J=254.3$ Hz), 158.3, 155.3, 144.5, 143.3, 138.7, 132.0, 130.1, 129.9, 129.2, 128.0, 127.9 (d, $J=9.4$ Hz), 127.5, 126.8 (d, $J=22.9$ Hz), 126.5 (d, $J=3.4$ Hz), 123.3 (d, $J=3.5$ Hz), 122.5 (d, $J=10.0$ Hz), 121.3 (d, $J=4.7$ Hz), 117.5 (d, $J=23.5$ Hz), 32.0, 29.9, 22.9, 21.6, 13.7, 13.8. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{25}\text{FNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 434.1585, found 434.1587.

(Z)-8,10-Dichloro-6-(1-tosylhex-1-en-2-yl)phenanthridine (**1w**): 79.7 mg, 66% yield. Colorless solid, m.p. 213.8~215.4 °C; ^1H NMR (600 MHz, CDCl_3) δ : 9.75 (d, $J=8.6$ Hz, 1H), 8.13 (d, $J=8.0$ Hz, 1H), 7.91 (s, 1H), 7.83~7.80 (m, 2H), 7.72 (t, $J=7.7$ Hz, 1H), 7.52 (d, $J=7.9$ Hz, 2H), 7.16 (d, $J=7.9$ Hz, 2H), 6.70 (s, 1H), 2.60~2.58 (m, 2H), 2.36 (s, 3H), 1.56~1.52 (m, 2H), 1.40~1.31 (m, 2H), 0.87 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 157.4, 154.0, 144.4, 144.3, 137.5, 134.3, 132.6, 132.3, 130.3, 129.9, 129.5, 129.4, 128.1, 127.9, 127.6, 127.2, 126.2, 126.0, 122.6, 38.3, 28.7, 22.2, 21.5, 13.7. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{24}\text{Cl}_2\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 484.0899, found 484.0898; $\text{C}_{26}\text{H}_{24}^{35}\text{Cl}^{37}\text{ClNO}_2\text{S}$: 486.0870, found 486.0872; $\text{C}_{26}\text{H}_{24}^{37}\text{Cl}_2\text{NO}_2\text{S}$: 488.0840, found 488.0845.

(Z)-2-Fluoro-8-methyl-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1x**): 79.5 mg, 69% yield. Colorless solid, m.p. 198.7~200.2 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.36~

8.34 (m, 1H), 8.12~8.10 (m, 1H), 8.06~8.04 (m, 1H), 7.65 (s, 1H), 7.63 (t, $J=7.9$ Hz, 1H), 7.51 (d, $J=8.0$ Hz, 2H), 7.44 (td, $J=8.4$, 2.2 Hz, 1H), 7.10 (d, $J=8.0$ Hz, 2H), 6.70 (s, 1H), 2.60 (t, $J=7.7$ Hz, 2H), 2.55 (s, 3H), 2.31 (s, 3H), 1.60~1.55 (m, 2H), 1.33~1.25 (m, 4H), 0.83 (t, $J=6.9$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.3 (d, $J=245.7$ Hz), 157.4 (d, $J=2.6$ Hz), 154.8, 144.0, 139.8, 138.1, 137.7, 132.6, 131.9 (d, $J=9.0$ Hz), 129.8 (d, $J=4.1$ Hz), 129.3, 129.2, 127.9, 126.6, 125.4 (d, $J=9.3$ Hz), 124.7, 122.2, 117.0 (d, $J=24.2$ Hz), 106.9 (d, $J=23.3$ Hz), 38.6, 31.1, 26.2, 22.2, 21.7, 21.4, 13.8. HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{29}\text{FNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 462.1898, found 462.1895.

(Z)-2-Chloro-8-fluoro-6-(1-tosylhept-1-en-2-yl)phenanthridine (**1y**): 81.8 mg, 68% yield. Colorless solid, m.p. 224.2~225.8 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.50~8.47 (m, 1H), 8.41 (d, $J=1.2$ Hz, 1H), 8.03 (d, $J=8.6$ Hz, 1H), 7.67 (dd, $J=8.6$, 1.5 Hz, 1H), 7.55 (t, $J=7.8$ Hz, 4H), 7.17 (d, $J=8.0$ Hz, 2H), 6.70 (s, 1H), 2.59 (t, $J=7.7$ Hz, 2H), 2.35 (s, 3H), 1.56~1.54 (m, 2H), 1.32~1.25 (m, 4H), 0.83 (t, $J=6.8$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.6 (d, $J=248.9$ Hz), 158.1 (d, $J=3.5$ Hz), 153.9, 144.3, 141.3, 137.6, 133.4, 131.4, 129.6, 129.5, 129.1, 128.1, 127.9, 126.2 (d, $J=7.8$ Hz), 125.0 (d, $J=8.5$ Hz), 124.6, 121.6, 120.3 (d, $J=23.8$ Hz), 111.8 (d, $J=21.7$ Hz), 38.5, 31.1, 26.2, 22.2, 21.5, 13.8. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{26}^{35}\text{ClFNO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 482.1351, found 482.1349; $\text{C}_{27}\text{H}_{26}^{37}\text{ClFNO}_2\text{S}$: 484.1322, found 484.1314.

4.3 Typical procedure for the synthesis of 1,3-diene (**2a**)

(Z)-6-(1-Tosylhept-1-en-2-yl)phenanthridine (**1a**, 0.25 mmol) and *t*-BuOK (0.50 mmol) were added into an air-tight reaction tube (10 mL) with 3.0 mL of THF. Then, the reaction was stirred at 70 °C for 18 h. After the reaction was completed, the mixture was diluted with water (10 mL) and extracted with ethyl acetate (5 mL $\times$ 3). The organic layers were combined, dried over anhydrous Na_2SO_4 , and filtered. The solvents were concentrated *in vacuo*, and the resulting crude mixture was purified by flash chromatography on silica gel (petroleum ether/EtOAc, $V:V=15:1$) to provide the desired products **2a** in 87% yield (59.4 mg).

4.4 Typical procedure for the synthesis of **4a**

(Z)-6-(1-Tosylhept-1-en-2-yl)phenanthridine (**1a**, 0.25 mmol) and *t*-BuOK (0.5 mmol) were added into an air-tight reaction tube (10 mL) with 3.0 mL of THF. The reaction was stirred at 70 °C for 18 h. After the reaction was completed, the mixture was diluted with water (10 mL) and extracted with ethyl acetate (5 mL $\times$ 3). The organic layers were combined, and concentrated *in vacuo*, generating the resulting crude mixture. The crude mixture and fumaronitrile (**3**, 0.50 mmol) were added into an air-tight reaction tube (10 mL) with 3.0 mL of toluene, and the reaction was stirred at 120 °C for 12 h. After the reaction was completed, the reaction mixture was concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel (petroleum ether/EtOAc, $V:V=3:1$) to provide the desired product **4a** in 83% yield (72.8 mg).

6-(Hepta-1,3-dien-2-yl)phenanthridine (**2a**): 59.3 mg, 87% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.60 (d, $J=8.2$ Hz, 1H), 8.55 (d, $J=8.2$ Hz, 1H), 8.26 (d, $J=8.2$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.80~7.77 (m, 1H), 7.74~7.71 (m, 1H), 7.65~7.63 (m, 1H), 7.59 (t, $J=7.6$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.30 (s, 1H), 5.26~5.21 (m, 1H), 2.00 (q, $J=7.1$ Hz, 2H), 1.29~1.23 (m, 2H), 0.78 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.1, 146.6, 143.7, 135.8, 132.8, 131.6, 130.5, 130.2, 128.7, 128.6, 127.0, 126.8, 125.3, 123.7, 121.82, 121.79, 117.6, 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{20}\text{N}$ $[\text{M}+\text{H}]^+$: 274.1590, found 274.1592.

6-(Hexa-1,3-dien-2-yl)-2-methylphenanthridine (**2b**): 56.0 mg, 78% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.60 (d, $J=8.3$ Hz, 1H), 8.35 (s, 1H), 8.14 (d, $J=8.3$ Hz, 2H), 7.79~7.76 (m, 1H), 7.59~7.57 (m, 1H), 7.55 (dd, $J=8.3$, 1.7 Hz, 1H), 6.52 (d, $J=15.6$ Hz, 1H), 5.62 (s, 1H), 5.60 (s, 1H), 5.25~5.20 (m, 1H), 2.61 (s, 3H), 2.02~1.98 (m, 2H), 1.30~1.23 (m, 2H), 0.78 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.1, 146.7, 142.0, 136.7, 135.7, 132.5, 131.7, 130.3, 130.0, 128.7, 126.8, 125.4, 123.6, 121.8, 121.5, 117.5, 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{20}\text{N}$ $[\text{M}+\text{H}]^+$: 288.1747, found 288.1746.

2-Fluoro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2c**): 58.2 mg, 80% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.49 (d, $J=8.2$ Hz, 1H), 8.22 (t, $J=6.9$ Hz, 1H), 8.17 (d, $J=9.2$ Hz, 2H), 7.82 (t, $J=7.4$ Hz, 1H), 7.65 (t, $J=7.5$ Hz, 1H), 7.47 (t, $J=8.4$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.30 (s, 1H), 5.25~5.20 (m, 1H), 2.01 (q, $J=7.0$ Hz, 2H), 1.31~1.25 (m, 2H), 0.79 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.3 (d, $J=245.5$ Hz), 160.4 (d, $J=2.1$ Hz), 146.4, 140.6, 135.9, 132.5 (d, $J=9.2$ Hz), 132.3 (d, $J=3.9$ Hz), 131.5, 130.6, 128.9, 127.7, 125.4, 125.1 (d, $J=8.9$ Hz), 122.0, 117.7, 117.5 (d, $J=24.1$ Hz), 106.8 (d, $J=22.9$ Hz), 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{FN}$ $[\text{M}+\text{H}]^+$: 292.1496, found 292.1496.

2-Chloro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2d**): 58.3 mg, 76% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.54~8.53 (m, 2H), 8.18~8.15 (m, 2H), 7.83 (t, $J=7.59$ Hz, 1H), 7.68~7.63 (m, 2H), 6.52 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.30 (s, 1H), 5.24~5.19 (m, 1H), 2.01 (q, $J=7.2$ Hz, 2H), 1.31~1.25 (m, 2H), 0.79 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.4, 146.4, 142.2, 135.9, 132.7, 131.9, 131.8, 131.5, 130.9, 129.1, 128.9, 127.7, 125.5, 124.9, 121.9, 121.6, 117.8, 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{ClN}$ $[\text{M}+\text{H}]^+$: 308.1201, found 308.1198; $\text{C}_{20}\text{H}_{19}^{37}\text{ClN}^+$: 310.1171, found 310.1175.

6-(Hepta-1,3-dien-2-yl)-3-methylphenanthridine (**2e**): 60.3 mg, 84% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.54 (d, $J=8.2$ Hz, 1H), 8.42 (d, $J=8.2$ Hz, 1H), 8.13 (d, $J=8.1$ Hz, 1H), 8.05 (s, 1H), 7.75 (t, $J=7.4$ Hz, 1H), 7.55 (t, $J=7.4$ Hz, 1H), 7.46 (d, $J=8.2$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.61 (s, 1H), 5.29 (s, 1H), 5.25~5.21 (m, 1H), 2.57 (s, 3H), 2.00 (q, $J=6.8$ Hz, 2H), 1.30~1.23 (m, 2H), 0.78 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz,

CDCl_3) δ : 161.1, 146.7, 143.8, 138.7, 135.7, 132.8, 131.6, 130.4, 129.8, 128.7, 128.5, 126.5, 125.0, 121.6, 121.4, 117.4, 34.8, 21.9, 21.4, 13.6. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{N}$ $[\text{M}+\text{H}]^+$: 288.1747, found 288.1749.

3-Fluoro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2f**): 56.0 mg, 77% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.55 (d, $J=8.4$ Hz, 2H), 8.16 (d, $J=8.2$ Hz, 1H), 7.89 (dd, $J=9.8$, 2.0 Hz, 1H), 7.82 (t, $J=7.6$ Hz, 1H), 7.61 (t, $J=7.6$ Hz, 1H), 7.42 (td, $J=8.4$, 1.8 Hz, 1H), 6.52 (d, $J=7.9$ Hz, 1H), 5.64 (s, 1H), 5.30 (s, 1H), 5.25~5.20 (m, 1H), 2.02 (q, $J=7.0$ Hz, 2H), 1.30~1.25 (m, 2H), 0.79 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.6 (d, $J=246.7$ Hz), 162.5, 146.4, 145.1 (d, $J=11.8$ Hz), 135.9, 132.7, 131.5, 131.0, 129.0, 126.9, 124.9, 123.8 (d, $J=9.4$ Hz), 121.7, 120.5, 117.7, 115.9 (d, $J=23.7$ Hz), 114.8 (d, $J=20.4$ Hz), 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{FN}$ $[\text{M}+\text{H}]^+$: 292.1496, found 292.1496.

3-Chloro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2g**): 65.2 mg, 85% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.53 (d, $J=8.2$ Hz, 1H), 8.45 (d, $J=8.7$ Hz, 1H), 8.23 (s, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.80 (t, $J=7.6$ Hz, 1H), 7.62 (t, $J=7.6$ Hz, 1H), 7.58 (d, $J=8.7$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.29 (s, 1H), 5.25~5.20 (m, 1H), 2.01 (q, $J=7.2$ Hz, 2H), 1.30~1.24 (m, 2H), 0.79 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.4, 146.3, 144.4, 135.9, 134.2, 132.4, 131.4, 130.9, 129.4, 128.9, 127.3, 125.2, 123.2, 122.3, 121.7, 117.7, 34.8, 21.9, 13.5. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{ClN}$ $[\text{M}+\text{H}]^+$: 308.1201, found 308.1200; $\text{C}_{20}\text{H}_{19}^{37}\text{ClN}^+$: 310.1171, found 310.1171.

10-Fluoro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2h**): 52.3 mg, 72% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.98 (d, $J=8.4$ Hz, 1H), 8.27 (d, $J=8.1$ Hz, 1H), 8.00 (d, $J=7.9$ Hz, 1H), 7.76 (t, $J=7.5$ Hz, 1H), 7.68 (t, $J=7.7$ Hz, 1H), 7.57~7.49 (m, 2H), 6.51 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.31 (s, 1H), 5.23~5.18 (m, 1H), 2.01 (q, $J=7.2$ Hz, 2H), 1.30~1.24 (m, 2H), 0.78 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.4 (d, $J=253.3$ Hz), 160.4 (d, $J=1.8$ Hz), 146.6, 144.0, 136.0, 131.4, 130.2, 128.9, 127.6 (d, $J=3.4$ Hz), 127.4 (d, $J=1.7$ Hz), 127.2 (d, $J=9.3$ Hz), 126.8 (d, $J=22.6$ Hz), 124.8 (d, $J=3.5$ Hz), 122.1 (d, $J=10.3$ Hz), 121.5 (d, $J=4.5$ Hz), 117.8, 117.0 (d, $J=23.1$ Hz), 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{FN}$ $[\text{M}+\text{H}]^+$: 292.1496, found 292.1495.

6-(Hepta-1,3-dien-2-yl)-8-methylphenanthridine (**2i**): 55.2 mg, 77% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.53 (d, $J=8.1$ Hz, 1H), 8.50 (d, $J=8.3$ Hz, 1H), 8.23 (d, $J=8.1$ Hz, 1H), 7.93 (s, 1H), 7.70 (t, $J=7.5$ Hz, 1H), 7.63 (d, $J=7.7$ Hz, 2H), 6.52 (d, $J=15.8$ Hz, 1H), 5.62 (s, 1H), 5.29 (s, 1H), 5.26~5.21 (m, 1H), 2.53 (s, 3H), 2.02 (q, $J=7.1$ Hz, 2H), 1.31~1.25 (m, 2H), 0.79 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.8, 146.7, 143.4, 136.9, 135.7, 132.3, 131.6, 130.7, 130.2, 128.1, 126.7, 125.5, 123.9, 121.8, 121.7, 117.5, 34.8, 21.9, 21.6, 13.5. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{N}$ $[\text{M}+\text{H}]^+$: 288.1747, found 288.1748.

6-(Hepta-1,3-dien-2-yl)-8-methoxyphenanthridine (**2j**):

56.0 mg, 74% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.52 (d, $J=8.9$ Hz, 1H), 8.48 (d, $J=7.9$ Hz, 1H), 8.22 (d, $J=8.0$ Hz, 1H), 7.67 (t, $J=7.5$ Hz, 1H), 7.62 (t, $J=7.5$ Hz, 1H), 7.50 (d, $J=2.6$ Hz, 1H), 7.43 (d, $J=9.0$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.32 (s, 1H), 5.29~5.24 (m, 1H), 3.90 (s, 3H), 2.02 (q, $J=7.2$ Hz, 2H), 1.31~1.25 (m, 2H), 0.80 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.3, 158.4, 146.8, 142.9, 135.7, 131.3, 130.2, 127.6, 127.2, 126.9, 126.6, 123.9, 123.5, 121.4, 117.5, 108.4, 55.4, 34.8, 22.0, 13.6. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{NO}$ $[\text{M}+\text{H}]^+$: 304.1696, found 304.1698.

8-Fluoro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2k**): 58.9 mg, 81% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.62~8.59 (m, 1H), 8.50 (d, $J=7.6$ Hz, 1H), 8.25 (dd, $J=8.2$, 1.0 Hz, 1H), 7.80 (dd, $J=9.7$, 2.7 Hz, 1H), 7.74~7.72 (m, 1H), 7.68~7.65 (m, 1H), 7.57~7.54 (m, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.31 (s, 1H), 5.26~5.21 (m, 1H), 2.02 (q, $J=7.0$ Hz, 2H), 1.32~1.26 (m, 2H), 0.80 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.2 (d, $J=246.8$ Hz), 160.2 (d, $J=3.5$ Hz), 146.1, 143.4, 136.0, 131.3, 130.4, 129.5 (d, $J=1.6$ Hz), 128.5, 127.2, 126.6 (d, $J=7.7$ Hz), 124.4 (d, $J=8.0$ Hz), 123.4, 121.6, 119.8 (d, $J=23.4$ Hz), 117.9, 113.0 (d, $J=21.7$ Hz), 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{F-N}$ $[\text{M}+\text{H}]^+$: 292.1496, found 292.1495.

8-Chloro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2l**): 63.7 mg, 83% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.55 (d, $J=8.8$ Hz, 1H), 8.16 (d, $J=8.8$ Hz, 1H), 8.24 (dd, $J=8.2$, 0.9 Hz, 1H), 8.13 (d, $J=2.2$ Hz, 1H), 7.76~7.74 (m, 2H), 7.68~7.66 (m, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.65 (s, 1H), 5.31 (s, 1H), 5.25~5.20 (m, 1H), 2.03 (q, $J=7.0$ Hz, 2H), 1.33~1.26 (m, 2H), 0.81 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.0, 146.0, 143.7, 136.1, 133.0, 131.4, 131.2, 131.1, 130.4, 129.0, 127.9, 127.3, 126.3, 123.7, 123.2, 121.8, 118.1, 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{Cl-N}$ $[\text{M}+\text{H}]^+$: 308.1201, found 308.1200; $\text{C}_{20}\text{H}_{19}^{37}\text{Cl-N}^+$: 310.1171, found 310.1171.

8,10-Dichloro-6-(hepta-1,3-dien-2-yl)phenanthridine (**2m**): 68.2 mg, 80% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 9.73 (d, $J=8.6$ Hz, 1H), 8.26 (dd, $J=8.1$, 1.4 Hz, 1H), 8.13 (d, $J=2.3$ Hz, 1H), 7.87 (t, $J=2.0$ Hz, 1H), 7.79 (td, $J=7.1$, 1.0 Hz, 1H), 7.70~7.67 (m, 1H), 6.51 (d, $J=15.8$ Hz, 1H), 5.66 (s, 1H), 5.31 (s, 1H), 5.20~5.15 (m, 1H), 2.05~2.01 (m, 2H), 1.32~1.26 (m, 2H), 0.80 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 159.8, 146.2, 144.7, 136.5, 134.1, 132.1, 132.0, 131.4, 130.7, 129.3, 128.4, 128.3, 127.6, 126.9, 125.9, 122.5, 118.4, 34.8, 21.9, 13.6. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{18}\text{Cl}_2\text{N}$ $[\text{M}+\text{H}]^+$: 342.0811, found 342.0808; $\text{C}_{20}\text{H}_{18}^{35}\text{Cl}^{37}\text{Cl-N}^+$: 344.0781, found 344.0776; $\text{C}_{20}\text{H}_{18}^{37}\text{Cl}_2\text{N}^+$: 346.0751, found 346.0751.

6-(Deca-1,3-dien-2-yl)phenanthridine (**2n**): 63.0 mg, 80% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.61 (d, $J=8.2$ Hz, 1H), 8.57 (d, $J=8.2$ Hz, 1H), 8.25 (d, $J=8.2$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.80 (t, $J=7.6$ Hz, 1H), 7.74~7.72 (m, 1H), 7.65 (t, $J=7.5$ Hz, 1H), 7.60 (t, $J=7.6$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.30

(s, 1H), 5.26~5.21 (m, 1H), 2.02 (q, $J=7.0$ Hz, 2H), 1.24~1.15 (m, 8H), 0.82 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.1, 146.6, 143.7, 136.1, 132.8, 131.4, 130.5, 130.3, 128.8, 128.6, 127.0, 126.8, 125.3, 123.8, 121.85, 121.82, 117.5, 32.8, 31.5, 28.71, 28.66, 22.4, 14.0. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{26}\text{N}$ $[\text{M}+\text{H}]^+$: 316.2060, found 316.2061.

6-(Octa-1,3-dien-2-yl)phenanthridine (**2o**): 63 mg, 88% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.63 (d, $J=8.3$ Hz, 1H), 8.58 (d, $J=8.1$ Hz, 1H), 8.25 (d, $J=8.1$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.81 (t, $J=7.6$ Hz, 1H), 7.74 (t, $J=7.1$ Hz, 1H), 7.66 (t, $J=7.4$ Hz, 1H), 7.61 (t, $J=7.6$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.30 (s, 1H), 5.26~5.21 (m, 1H), 2.03 (q, $J=6.8$ Hz, 2H), 1.25~1.17 (m, 4H), 0.80 (t, $J=6.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.2, 146.7, 143.8, 136.0, 132.8, 131.4, 130.5, 130.3, 128.8, 128.6, 127.0, 126.8, 125.4, 123.8, 121.88, 121.85, 117.6, 32.4, 30.9, 22.1, 13.8. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{N}$ $[\text{M}+\text{H}]^+$: 288.1747, found 288.1747.

6-(Hexa-1,3-dien-2-yl)phenanthridine (**2p**): 56.3 mg, 87% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.59 (d, $J=8.3$ Hz, 1H), 8.55 (d, $J=8.1$ Hz, 1H), 8.26 (d, $J=8.2$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.78 (t, $J=7.6$ Hz, 1H), 7.72 (t, $J=7.5$ Hz, 1H), 7.63 (t, $J=7.6$ Hz, 1H), 7.59 (t, $J=7.6$ Hz, 1H), 6.52 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.32~5.27 (m, 2H), 2.05~2.00 (m, 2H), 0.89 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.1, 146.5, 143.7, 137.2, 132.8, 130.48, 130.46, 130.2, 128.7, 128.5, 127.0, 126.7, 125.3, 123.7, 121.81, 121.78, 117.5, 25.6, 12.8. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{18}\text{N}$ $[\text{M}+\text{H}]^+$: 260.1434, found 260.1433.

6-(Penta-1,3-dien-2-yl)phenanthridine (**2q**): 45.3 mg, 74% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.61 (d, $J=8.3$ Hz, 1H), 8.56 (d, $J=8.2$ Hz, 1H), 8.25 (d, $J=8.2$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.80 (t, $J=7.6$ Hz, 1H), 7.73 (t, $J=7.5$ Hz, 1H), 7.65 (t, $J=7.5$ Hz, 1H), 7.61 (t, $J=7.6$ Hz, 1H), 6.53 (d, $J=15.7$ Hz, 1H), 5.62 (s, 1H), 5.28~5.22 (m, 2H), 1.68 (d, $J=6.7$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.1, 146.4, 143.7, 132.84, 132.81, 130.6, 130.5, 130.2, 128.7, 128.6, 127.1, 126.8, 125.3, 123.7, 121.843, 121.835, 117.3, 18.2. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{N}$ $[\text{M}+\text{H}]^+$: 246.1277, found 246.1276.

3-Chloro-6-(octa-1,3-dien-2-yl)phenanthridine (**2r**): 64.2 mg, 80% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.55 (d, $J=8.3$ Hz, 1H), 8.47 (d, $J=8.7$ Hz, 1H), 8.23 (d, $J=1.4$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.82 (t, $J=7.6$ Hz, 1H), 7.62 (t, $J=7.8$ Hz, 1H), 7.60 (dd, $J=8.9$, 1.7 Hz, 1H), 6.51 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.28 (s, 1H), 5.25~5.20 (m, 1H), 2.03 (q, $J=6.5$ Hz, 2H), 1.25~1.18 (m, 4H), 0.80 (t, $J=6.9$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.4, 146.3, 144.5, 136.1, 134.2, 132.4, 131.2, 131.0, 129.5, 129.0, 127.34, 127.32, 125.3, 123.3, 122.3, 121.8, 117.7, 32.4, 30.9, 22.1, 13.8. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{20}\text{Cl-N}$ $[\text{M}+\text{H}]^+$: 322.1357, found 322.1357; $\text{C}_{21}\text{H}_{21}^{37}\text{Cl-N}^+$: 324.1328, found 324.1328.

3-Chloro-6-(hexa-1,3-dien-2-yl)phenanthridine (**2s**): 59.3 mg, 81% yield. Colorless liquid. ^1H NMR (600 MHz,

CDCl_3) δ : 8.56 (d, $J=8.3$ Hz, 1H), 8.46 (d, $J=8.8$ Hz, 1H), 8.23 (d, $J=2.2$ Hz, 1H), 8.16 (d, $J=8.2$ Hz, 1H), 7.83 (t, $J=7.6$ Hz, 1H), 7.63 (t, $J=7.6$ Hz, 1H), 7.61 (dd, $J=8.7$, 2.1 Hz, 1H), 6.51 (d, $J=15.9$ Hz, 1H), 5.65 (s, 1H), 5.31~5.26 (m, 2H), 2.07~2.02 (m, 2H), 0.91 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.4, 146.2, 144.4, 137.4, 134.2, 132.4, 131.0, 130.3, 129.5, 128.9, 127.37, 127.36, 125.3, 123.3, 122.3, 121.8, 117.8, 25.7, 12.9. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{17}\text{ClN}$ $[\text{M}+\text{H}]^+$: 294.1044, found 294.1042; $\text{C}_{19}\text{H}_{17}^{37}\text{ClN}^+$: 296.1015, found 296.1016.

6-(Hexa-1,3-dien-2-yl)-2-methylphenanthridine (**2t**): 51.2 mg, 75% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.60 (d, $J=8.3$ Hz, 1H), 8.35 (s, 1H), 8.14 (d, $J=8.3$ Hz, 2H), 7.77 (t, $J=7.6$ Hz, 1H), 7.58 (t, $J=7.6$ Hz, 1H), 7.55 (dd, $J=8.3$, 1.5 Hz, 1H), 6.51 (d, $J=15.8$ Hz, 1H), 5.63 (s, 1H), 5.31~5.26 (m, 2H), 2.61 (s, 3H), 2.06~2.01 (m, 2H), 0.89 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.1, 146.6, 142.0, 137.1, 136.6, 132.5, 130.6, 130.3, 130.0, 128.7, 126.8, 125.4, 123.6, 121.8, 121.4, 117.5, 25.7, 21.9, 12.9. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{20}\text{N}$ $[\text{M}+\text{H}]^+$: 274.1590, found 274.1593.

2-Chloro-6-(hexa-1,3-dien-2-yl)phenanthridine (**2u**): 51.3 mg, 70% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.50~8.49 (m, 2H), 8.16 (d, $J=8.6$ Hz, 2H), 7.79 (t, $J=7.2$ Hz, 1H), 7.66~7.61 (m, 2H), 6.51 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.30~5.26 (m, 2H), 2.06~2.01 (m, 2H), 0.90 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 161.3, 146.2, 142.0, 137.3, 132.6, 131.75, 131.65, 130.8, 130.3, 129.0, 128.7, 127.7, 125.4, 124.8, 121.8, 121.5, 117.7, 25.6, 12.8. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{17}\text{ClN}$ $[\text{M}+\text{H}]^+$: 294.1044, found 294.1048; $\text{C}_{19}\text{H}_{17}^{37}\text{ClN}^+$: 296.1015, found 296.1018.

10-Fluoro-6-(hexa-1,3-dien-2-yl)phenanthridine (**2v**): 50.4 mg, 73% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 8.99 (d, $J=8.4$ Hz, 1H), 8.27 (d, $J=8.1$ Hz, 1H), 8.01 (d, $J=7.8$ Hz, 1H), 7.78 (t, $J=7.5$ Hz, 1H), 7.70 (t, $J=7.7$ Hz, 1H), 7.59~7.52 (m, 2H), 6.51 (d, $J=15.8$ Hz, 1H), 5.64 (s, 1H), 5.30 (s, 1H), 5.29~5.24 (m, 1H), 2.07~2.02 (m, 2H), 0.91 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.5 (d, $J=253.3$ Hz), 160.4 (d, $J=1.7$ Hz), 146.6, 144.1, 137.5, 130.4, 130.3, 128.9, 127.7 (d, $J=3.4$ Hz), 127.4 (d, $J=1.6$ Hz), 127.3 (d, $J=9.2$ Hz), 126.9 (d, $J=22.6$ Hz), 124.8 (d, $J=3.9$ Hz), 122.2 (d, $J=9.9$ Hz), 121.5 (d, $J=4.6$ Hz), 117.8, 117.0 (d, $J=23.2$ Hz), 25.7, 12.9. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{17}\text{FN}$ $[\text{M}+\text{H}]^+$: 277.3354, found 277.3352.

8,10-Dichloro-6-(hexa-1,3-dien-2-yl)phenanthridine (**2w**): 68.7 mg, 84% yield. Colorless liquid. ^1H NMR (600 MHz, CDCl_3) δ : 9.73 (d, $J=8.6$ Hz, 1H), 8.26 (dd, $J=8.1$, 1.4 Hz, 1H), 8.13 (d, $J=2.3$ Hz, 1H), 7.87 (t, $J=1.8$ Hz, 1H), 7.80~7.78 (m, 1H), 7.70~7.67 (m, 1H), 6.51 (d, $J=15.8$ Hz, 1H), 5.67 (s, 1H), 5.30 (s, 1H), 5.26~5.22 (m, 1H), 2.09~2.04 (m, 2H), 0.92 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 159.8, 146.1, 144.7, 137.9, 134.1, 132.1, 132.0, 130.7, 130.3, 129.3, 128.43, 128.35, 127.5, 126.9, 125.9, 122.4, 118.4, 25.7, 12.9. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{16}\text{Cl}_2\text{N}$ $[\text{M}+\text{H}]^+$: 328.0654, found 328.0648;

$\text{C}_{19}\text{H}_{16}^{35}\text{Cl}^{37}\text{ClN}^+$: 330.0625, found 330.0625; $\text{C}_{19}\text{H}_{16}^{37}\text{Cl}_2\text{N}^+$: 332.0595, found 332.0597.

5-(Phenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4a**): 72.8 mg, 83% yield. White solid, m.p. 169.7~170.9 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.64 (d, $J=8.2$ Hz, 1H), 8.55 (d, $J=8.1$ Hz, 1H), 8.29 (d, $J=8.2$ Hz, 1H), 8.16 (d, $J=8.0$ Hz, 1H), 7.85 (t, $J=7.5$ Hz, 1H), 7.74 (t, $J=7.4$ Hz, 1H), 7.70~7.65 (m, 2H), 6.01 (s, 1H), 3.60~3.58 (m, 1H), 3.45 (t, $J=5.0$ Hz, 1H), 3.31~3.27 (m, 1H), 3.15 (d, $J=18.4$ Hz, 1H), 3.03 (d, $J=2.5$ Hz, 1H), 1.85~1.71 (m, 2H), 1.56~1.49 (m, 2H), 1.01 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.3, 143.2, 133.4, 133.3, 130.7, 130.4, 130.0, 128.8, 127.4, 127.3, 127.1, 124.2, 123.5, 122.3, 121.8, 118.9, 117.2, 34.8, 34.3, 32.2, 29.0, 27.2, 19.9, 13.9. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{22}\text{N}_3$ $[\text{M}+\text{H}]^+$: 352.1808, found 352.1812.

5-(3-Methylphenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4b**): 65.7 mg, 72% yield. White solid, m.p. 147.9~149.6 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.59 (d, $J=8.2$ Hz, 1H), 8.42 (d, $J=8.4$ Hz, 1H), 8.26 (d, $J=8.2$ Hz, 1H), 7.95 (s, 1H), 7.83~7.79 (m, 1H), 7.64 (t, $J=7.3$ Hz, 1H), 7.48 (dd, $J=8.3$, 1.1 Hz, 1H), 5.99 (d, $J=1.0$ Hz, 1H), 3.59~3.55 (m, 1H), 3.43 (t, $J=5.0$ Hz, 1H), 3.31~3.25 (m, 1H), 3.13 (d, $J=18.3$ Hz, 1H), 3.03~3.01 (m, 1H), 2.58 (s, 3H), 1.84~1.71 (m, 2H), 1.57~1.48 (m, 2H), 1.01 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.4, 143.4, 139.0, 133.5, 133.4, 130.6, 130.3, 129.6, 128.9, 127.3, 126.9, 123.9, 122.2, 121.6, 121.2, 118.9, 117.2, 34.9, 34.3, 32.2, 29.0, 27.3, 21.4, 19.9, 13.9. HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{24}\text{N}_3$ $[\text{M}+\text{H}]^+$: 366.1965, found 366.1966.

5-(3-Fluorophenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4c**): 69.2 mg, 75% yield. White solid, m.p. 169.6~171.0 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.58 (d, $J=8.0$ Hz, 1H), 8.53 (dd, $J=9.0$, 5.9 Hz, 1H), 8.30 (d, $J=7.8$ Hz, 1H), 7.87 (td, $J=7.6$, 1.0 Hz, 1H), 7.80 (dd, $J=9.7$, 2.6 Hz, 1H), 7.69 (td, $J=8.1$, 1.0 Hz, 1H), 7.42 (td, $J=8.5$, 2.7 Hz, 1H), 6.01 (d, $J=1.6$ Hz, 1H), 3.61~3.59 (m, 1H), 3.46 (t, $J=5.0$ Hz, 1H), 3.33~3.28 (m, 1H), 3.16 (d, $J=18.5$ Hz, 1H), 3.07~3.04 (m, 1H), 1.85~1.74 (m, 2H), 1.58~1.52 (m, 2H), 1.03 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.7 (d, $J=247.2$ Hz), 160.7, 144.6 (d, $J=11.8$ Hz), 133.4, 133.2, 131.1, 130.9, 127.5, 127.3, 123.8 (d, $J=9.5$ Hz), 122.2, 120.3 (d, $J=1.7$ Hz), 118.8, 117.1, 116.3 (d, $J=23.7$ Hz), 114.6 (d, $J=20.3$ Hz), 34.9, 34.3, 32.2, 28.9, 27.4, 20.0, 13.9. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{FN}_3$ $[\text{M}+\text{H}]^+$: 370.1714, found 370.1715.

5-(3-Chlorophenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4d**): 74.1 mg, 77% yield. White solid, m.p. 193.7~194.8 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.55 (d, $J=8.2$ Hz, 1H), 8.42 (d, $J=8.8$ Hz, 1H), 8.29 (d, $J=8.2$ Hz, 1H), 8.11 (d, $J=2.0$ Hz, 1H), 7.85 (t, $J=7.5$ Hz, 1H), 7.70 (t, $J=7.6$ Hz, 1H), 7.57 (dd, $J=8.7$, 2.1 Hz, 1H), 6.00 (s, 1H), 3.61~3.59 (m, 1H), 3.45 (t, $J=4.9$ Hz, 1H), 3.30~3.26 (m, 1H), 3.15 (d, $J=18.4$ Hz, 1H), 3.05~3.04 (m, 1H), 1.84~1.73 (m, 2H), 1.57~1.51 (m, 2H), 1.02 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 160.5, 143.9,

134.4, 133.1, 133.0, 131.1, 131.0, 129.2, 127.7, 127.6, 127.5, 124.1, 123.2, 122.3, 122.0, 118.8, 117.0, 34.8, 34.3, 32.1, 28.8, 27.3, 19.9, 13.9. HRMS (ESI) calcd for $C_{24}H_{20}ClN_3$ $[M+H]^+$: 386.1419, found 386.1415; $C_{24}H_{21}-^{37}ClN_3$: 388.1389, found 388.1393.

5-(2-Fluorophenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4e**): 74.7 mg, 81% yield. White solid, m.p. 158.9~160.1 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.49 (d, $J=8.3$ Hz, 1H), 8.30 (d, $J=8.1$ Hz, 1H), 8.15~8.09 (m, 2H), 7.87~7.83 (m, 1H), 7.72 (t, $J=7.6$ Hz, 1H), 7.45 (td, $J=8.4, 2.6$ Hz, 1H), 6.00 (d, $J=1.2$ Hz, 1H), 3.63~3.59 (m, 1H), 3.46 (t, $J=4.9$ Hz, 1H), 3.33~3.26 (m, 1H), 3.15 (d, $J=18.3$ Hz, 1H), 3.06~3.00 (m, 1H), 1.87~1.71 (m, 2H), 1.59~1.50 (m, 2H), 1.02 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 161.4 (d, $J=245.8$ Hz), 158.6 (d, $J=2.7$ Hz), 140.1, 133.2, 132.9 (d, $J=4.3$ Hz), 132.4 (d, $J=9.0$ Hz), 130.7 (d, $J=6.1$ Hz), 128.1, 127.4, 124.9 (d, $J=9.0$ Hz), 124.3, 122.5, 118.8, 117.7 (d, $J=24.2$ Hz), 117.1, 106.8 (d, $J=23.2$ Hz), 34.9, 34.3, 32.2, 28.9, 27.4, 19.9, 13.9. HRMS (ESI) calcd for $C_{24}H_{21}FN_3$ $[M+H]^+$: 370.1714, found 370.1713.

5-(8-*tert*-Butylphenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4f**): 81.4 mg, 80% yield. White solid, m.p. 156.1~157.4 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 8.57 (d, $J=8.7$ Hz, 1H), 8.52 (d, $J=7.6$ Hz, 1H), 8.31 (d, $J=1.9$ Hz, 1H), 8.14 (dd, $J=8.1, 0.8$ Hz, 1H), 7.93 (dd, $J=8.7, 2.0$ Hz, 1H), 7.71~7.68 (m, 1H), 7.64~7.62 (m, 1H), 6.03 (d, $J=1.6$ Hz, 1H), 3.60~3.58 (m, 1H), 3.44 (t, $J=5.0$ Hz, 1H), 3.31 (d, $J=18.4$ Hz, 1H), 3.22~3.16 (m, 1H), 3.03~3.01 (m, 1H), 1.85~1.75 (m, 2H), 1.57~1.53 (m, 2H), 1.45 (s, 9H), 1.02 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 159.5, 150.4, 143.1, 133.4, 131.3, 130.3, 130.0, 129.0, 128.4, 127.0, 124.3, 123.5, 123.1, 122.1, 121.7, 118.9, 117.1, 35.1, 35.0, 34.2, 32.3, 31.2, 28.8, 27.4, 19.9, 13.8. HRMS (ESI) calcd for $C_{28}H_{29}N_3$ $[M+H]^+$: 408.2434, found 408.2435.

5-(8-Fluorophenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4g**): 70.1 mg, 76% yield. White solid, m.p. 172.3~174.4 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.67~8.63 (m, 1H), 8.51 (t, $J=7.6$ Hz, 1H), 8.16 (t, $J=7.4$ Hz, 1H), 7.94 (d, $J=9.8, 2.5$ Hz, 1H), 7.77~7.67 (m, 2H), 7.64~7.59 (m, 1H), 6.04 (s, 1H), 3.64~3.60 (m, 1H), 3.48 (t, $J=5.0$ Hz, 1H), 3.36~3.30 (m, 1H), 3.17 (d, $J=18.2$ Hz, 1H), 3.08~3.05 (m, 1H), 1.86~1.75 (m, 2H), 1.60~1.54 (m, 2H), 1.05 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 161.3 (d, $J=247.2$ Hz), 158.3 (d, $J=3.4$ Hz), 142.9, 133.1, 130.9, 130.3, 130.2, 128.8, 127.6, 125.5 (d, $J=7.7$ Hz), 125.1 (d, $J=8.1$ Hz), 123.2, 121.7, 119.9 (d, $J=24.0$ Hz), 118.8, 117.1, 111.9 (d, $J=22.0$ Hz), 34.9, 34.3, 32.2, 28.9, 27.3, 20.0, 13.9. HRMS (ESI) calcd for $C_{24}H_{21}FN_3$ $[M+H]^+$: 370.1714, found 370.1714.

5-(8-Chlorophenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4h**): 72.2 mg, 75% yield. White solid, m.p. 181.3~184.2 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.56 (d, $J=8.8$ Hz, 1H), 8.48 (d, $J=8.0$ Hz, 1H), 8.25 (d, $J=1.6$ Hz, 1H), 8.14 (d, $J=8.0$ Hz, 1H), 7.79~7.73 (m, 1H), 7.67 (t, $J=7.4$ Hz, 1H), 6.02 (s, 1H), 3.61~3.57 (m, 1H), 3.46 (t,

$J=5.1$ Hz, 1H), 3.33~3.27 (m, 1H), 3.15 (d, $J=18.4$ Hz, 1H), 3.05 (d, $J=2.5$ Hz, 1H), 1.87~1.73 (m, 2H), 1.60~1.51 (m, 2H), 1.03 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 158.1, 143.2, 133.4, 133.0, 131.9, 131.22, 131.16, 130.3, 129.2, 127.6, 126.6, 125.1, 124.2, 123.0, 121.8, 118.8, 117.1, 34.9, 34.4, 32.3, 29.0, 27.3, 20.0, 13.9. HRMS (ESI) calcd for $C_{24}H_{21}ClN_3$ $[M+H]^+$: 386.1419, found 386.1419; $C_{24}H_{21}-^{37}ClN_3$: 388.1389, found 388.1396.

3-Propyl-5-(8-(trifluoromethyl)phenanthridin-6-yl)cyclohex-4-ene-1,2-dicarbonitrile (**4i**): 71.2 mg, 68% yield. White solid, m.p. 171.1~173.2 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 8.77 (d, $J=8.6$ Hz, 1H), 8.62 (s, 1H), 8.57 (d, $J=8.2$ Hz, 1H), 8.19 (d, $J=8.0$ Hz, 1H), 8.04 (dd, $J=8.7, 1.1$ Hz, 1H), 7.82 (t, $J=7.2$ Hz, 1H), 7.73 (t, $J=7.6$ Hz, 1H), 6.05 (d, $J=1.3$ Hz, 1H), 3.61 (q, $J=5.2$ Hz, 1H), 3.47 (t, $J=5.2$ Hz, 1H), 3.32~3.24 (m, 2H), 3.08~3.03 (m, 1H), 1.90~1.75 (m, 2H), 1.60~1.54 (m, 2H), 1.04 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 159.0, 143.9, 135.7, 132.8, 131.9, 130.4, 130.1, 129.2 (q, $J=32.7$ Hz), 127.8, 126.5, 124.9 (q, $J=4.1$ Hz), 123.9 (q, $J=270.9$ Hz), 123.59, 123.56, 122.6, 122.3, 118.7, 117.0, 34.9, 34.5, 32.3, 28.9, 27.3, 19.9, 13.9. HRMS (ESI) calcd for $C_{25}H_{21}F_3N_3$ $[M+H]^+$: 420.1682, found 420.1680.

5-(2-Fluoro-8-methylphenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4j**): 80.4 mg, 84% yield. White solid, m.p. 161.6~164.0 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.36 (d, $J=8.4$ Hz, 1H), 8.11~8.05 (m, 3H), 7.67 (d, $J=8.4$ Hz, 1H), 7.44~7.39 (m, 1H), 5.99 (s, 1H), 3.62~3.58 (m, 1H), 3.46 (t, $J=4.9$ Hz, 1H), 3.29~3.23 (m, 1H), 3.12~3.05 (m, 2H), 2.58 (s, 3H), 1.86~1.73 (m, 2H), 1.60~1.51 (m, 2H), 1.03 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 161.3 (d, $J=245.6$ Hz), 158.4, 139.8, 138.2, 133.4, 132.4, 132.2 (d, $J=9.1$ Hz), 130.7 (d, $J=4.3$ Hz), 130.2, 126.8, 125.0 (d, $J=9.2$ Hz), 124.4, 122.4, 118.9, 117.21 (d, $J=24.2$ Hz), 117.19, 106.6 (d, $J=23.0$ Hz), 34.9, 34.3, 32.2, 29.0, 27.3, 21.7, 19.9, 13.9. HRMS (ESI) calcd for $C_{25}H_{23}FN_3$ $[M+H]^+$: 384.1871, found 384.1874.

5-(2-Chloro-8-fluorophenanthridin-6-yl)-3-propylcyclohex-4-ene-1,2-dicarbonitrile (**4k**): 68.5 mg, 68% yield. White solid, m.p. 193.8~195.0 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 8.54~8.52 (m, 1H), 8.40 (d, $J=2.0$ Hz, 1H), 8.04 (d, $J=8.7$ Hz, 1H), 7.93 (dd, $J=9.7, 2.5$ Hz, 1H), 7.64 (dd, $J=8.7, 2.1$ Hz, 1H), 7.62~7.59 (m, 1H), 6.02 (s, 1H), 3.63~3.61 (m, 1H), 3.46 (t, $J=4.9$ Hz, 1H), 3.32~3.29 (m, 1H), 3.16 (d, $J=18.4$ Hz, 1H), 3.06~3.06 (m, 1H), 1.84~1.76 (m, 2H), 1.58~1.53 (m, 2H), 1.04 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 161.7 (d, $J=248.6$ Hz), 158.5 (d, $J=3.5$ Hz), 141.3, 133.5, 132.8, 131.7, 131.3, 129.3, 129.2, 125.6 (d, $J=7.7$ Hz), 125.1 (d, $J=8.5$ Hz), 124.1, 121.3, 120.2 (d, $J=23.8$ Hz), 118.7, 117.0, 112.0 (d, $J=22.5$ Hz), 34.8, 34.4, 32.1, 28.8, 27.3, 19.9, 13.9. HRMS (ESI) calcd for $C_{24}H_{20}ClFN_3$ $[M+H]^+$: 404.1324, found 404.1328; $C_{24}H_{20}-^{37}ClFN_3$: 406.1295, found 406.1299.

3-Ethyl-5-(phenanthridin-6-yl)cyclohex-4-ene-1,2-dicarbonitrile (**4l**): 59.0 mg, 70% yield. White solid, m.p.

180.2~183.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.63 (d, $J=8.3$ Hz, 1H), 8.54 (d, $J=7.8$ Hz, 1H), 8.29 (d, $J=8.1$ Hz, 1H), 8.16~8.14 (m, 1H), 7.86~7.82 (m, 1H), 7.75~7.63 (m, 3H), 6.02 (d, $J=1.1$ Hz, 1H), 3.60~3.57 (m, 1H), 3.46 (t, $J=5.0$ Hz, 1H), 3.33~3.25 (m, 1H), 3.14 (d, $J=18.4$ Hz, 1H), 2.94~2.92 (m, 1H), 1.91~1.77 (m, 2H), 1.12 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.4, 143.2, 133.4, 130.7, 130.2, 130.1, 128.8, 127.4, 127.3, 127.1, 124.2, 123.5, 122.3, 121.8, 118.9, 117.1, 36.1, 32.0, 29.0, 27.3, 25.9, 11.3. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{20}\text{N}_3$ $[\text{M}+\text{H}]^+$: 338.1652, found 338.1653.

5-(3-Chlorophenanthridin-6-yl)-3-ethylcyclohex-4-ene-1,2-dicarbonitrile (**4m**): 70.5 mg, 76% yield. White solid, m.p. 192.0~196.3 °C; ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ : 8.81 (d, $J=8.3$ Hz, 1H), 8.74 (d, $J=8.8$ Hz, 1H), 8.27 (d, $J=8.2$ Hz, 1H), 8.07 (d, $J=2.2$ Hz, 1H), 7.95 (t, $J=7.6$ Hz, 1H), 7.79 (t, $J=7.6$ Hz, 1H), 7.69 (dd, $J=8.8, 2.2$ Hz, 1H), 6.01 (s, 1H), 4.04~4.00 (m, 1H), 3.10~3.06 (m, 1H), 2.97 (d, $J=18.7$ Hz, 1H), 2.81~2.80 (m, 1H), 1.75~1.70 (m, 2H), 1.06 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$) δ : 161.8, 143.9, 133.9, 133.2, 133.0, 132.1, 131.6, 128.7, 128.6, 127.84, 127.76, 125.2, 124.3, 123.4, 122.4, 120.5, 118.8, 36.1, 31.3, 29.5, 26.6, 25.8, 11.6. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{19}^{35}\text{ClN}_3$ $[\text{M}+\text{H}]^+$: 372.1262, found 372.1260; $\text{C}_{23}\text{H}_{19}^{37}\text{ClN}_3$: 374.1233, found 374.1235.

3-Methyl-5-(phenanthridin-6-yl)cyclohex-4-ene-1,2-dicarbonitrile (**4n**): 59.8 mg, 74% yield. White solid, m.p. 134.2~136.8 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.65 (d, $J=8.3$ Hz, 1H), 8.55 (d, $J=8.1$ Hz, 1H), 8.27 (d, $J=8.0$ Hz, 1H), 8.14 (dd, $J=8.1, 1.0$ Hz, 1H), 7.85 (t, $J=7.6$ Hz, 1H), 7.75~7.72 (m, 1H), 7.69~7.65 (m, 2H), 5.97 (d, $J=2.2$ Hz, 1H), 3.55~3.52 (m, 1H), 3.41 (t, $J=5.8$ Hz, 1H), 3.32~3.27 (m, 1H), 3.19~3.16 (m, 1H), 3.13~3.09 (m, 1H), 1.45 (d, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 159.2, 143.2, 133.5, 133.2, 131.7, 130.7, 130.1, 128.9, 127.4, 127.3, 127.2, 124.2, 123.6, 122.4, 121.9, 118.8, 117.3, 33.6, 29.8, 29.2, 27.0, 17.9. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{N}_3$ $[\text{M}+\text{H}]^+$: 324.1495, found 324.1496.

Supporting Information Mechanism investigation and ^1H NMR and ^{13}C NMR spectra for compounds **2** and **4**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- (a) Devonn, T. K.; Scott, A. I. *Handbook of Naturally Occurring Compounds*, Academic, New York, **1972**, Vol. II.
(b) Pereira, A. R.; Cabezas, J. A. *J. Org. Chem.* **2005**, *70*, 2594.
(c) Figueiredo, R. M.; Berner, R.; Julis, J.; Liu, T.; Turp, D.; Christmann, M. *J. Org. Chem.* **2007**, *72*, 640.
(d) Nicolaou, K. C.; Daines, R. A.; Ogawa, Y.; Chakraborty, T. K. *J. Am. Soc. Chem.* **1988**, *110*, 4696.
(e) Negishi, E.-I.; Wang, G. In *Science of Synthesis*, Eds.: Jacobsen, E. N.; Meijere, A.; Thieme, Stuttgart, **2010**, Vol. 47, p. 909.
- (a) Rappoport, Z. *The Chemistry of Dienes and Polyenes*, John Wiley & Sons, Chichester, **1997**, Vol. 1; **2001**, Vol. 2.
(b) Nicolaou, K. C.; Snyder, S. S.; Montagnon, T.; Vassilikogiannakis, G. *Angew. Chem., Int. Ed.* **2002**, *41*, 1668.
(c) Negishi, E.-I.; Huang, Z.; Wang, G.; Mohan, S.; Wang, C.; Hat-
tori, H. *Acc. Chem. Res.* **2008**, *41*, 1474.
- (a) Metzker, M. L.; Lu, J.; Gibbs, R. A. *Science* **1996**, *271*, 1420.
(b) Valente, A.; Mortreux, A.; Visseaux, M.; Zinck, P. *Chem. Rev.* **2013**, *113*, 3836.
- (a) Normant, J. F. *Modern Synthetic Methods*, Ed.: Scheffold, R., Wiley, Chichester, **1983**, Vol. 3, p. 139.
(b) Larock, R. C. *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, **1989**, p. 241.
(c) Sellén, M.; Bäckvall, J.-E.; Helquist, P. *J. Org. Chem.*, **1991**, *56*, 835.
- (a) Wittig, G.; Schöllkopf, U. *Chem. Ber.* **1954**, *87*, 1318.
(b) Dong, D. J.; Li, H. H.; Tian, S. K. *J. Am. Soc. Chem.* **2010**, *132*, 5018.
(c) Hampton, C. S.; Harmata, M. *Org. Lett.* **2014**, *16*, 1256.
(d) Jacobsen, M. J.; Funder, E. D.; Cramer, J. R.; Gothelf, K. V. *Org. Lett.* **2011**, *13*, 3418.
(e) Xu, S.; Zhou, L.; Zeng, S.; Ma, R.; Wang, Z.; He, Z. *Org. Lett.* **2009**, *11*, 3498.
- (a) Horner, L.; Hoffmann, H.; Wippel, H. G.; Klahre, G. *Chem. Ber.* **1959**, *92*, 2499.
(b) Wadsworth, W. S.; Emmons, W. D., Jr. *J. Am. Soc. Chem.* **1961**, *83*, 1733.
(c) Wadsworth, W. S. *Org. React.* **1977**, *25*, 73.
- (a) Julia, M.; Paris, J. M. *Tetrahedron Lett.* **1973**, *14*, 4833.
(b) Blakemore, P. R. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2563.
(c) Aïssa, C. *Eur. J. Org. Chem.* **2009**, 1831.
(d) Markó, I. E.; Pospíšil, J. In *Science of Synthesis*, Vol. 47a, Eds.: Alkenes; de Meijere, A., George Thieme Verlag, Stuttgart, **2009**, 105.
- Peterson, D. J. *J. Org. Chem.* **1968**, *33*, 780.
- (a) Wojtkiewicz, A. *Curr. Org. Synth.* **2013**, *10*, 43.
(b) Cornil, J.; Echeverria, P.-G.; Phansavath, P.; Ratovelomanana-Vidal, V.; Guérinot, A.; Cossy, J. *Org. Lett.* **2015**, *17*, 948.
(c) Stille, J. K. *Angew. Chem., Int. Ed.* **1986**, *25*, 508.
(d) Heck, R. F. *Org. React.* **1982**, *27*, 345.
(e) Yoon, C. H.; Yoo, K. S.; Yi, S. W.; Mishra, R. K.; Jung, K. W. *Org. Lett.* **2004**, *6*, 4037.
(f) Molander, G. A.; Yokoyama, Y. *J. Org. Chem.* **2006**, *71*, 2493.
(g) Nguyen, V. T.; Dang, H. T.; Pham, H. H.; Nguyen, V. D.; Flores-Hansen, C.; Arman, H. D.; Larionov, O. V. *J. Am. Chem. Soc.* **2018**, *140*, 8434.
(h) Fiorito, D.; Folliet, S.; Liu, Y.; Mazet, C. *ACS Catal.* **2018**, *8*, 1392.
(i) He R.; Deng, M.-Z. *Org. Lett.* **2002**, *4*, 2759.
- (a) Xu, Y.-H.; Lu, J.; Loh, T.-P. *J. Am. Soc. Chem.* **2009**, *131*, 1372.
(b) Xu, Y.-H.; Chok, Y. K.; Loh, T.-P. *Chem. Sci.* **2011**, *2*, 1822.
- Yu, H.; Jin, W.; Sun, C.; Chen, J.; Du, W.; He, S.; Yu, Z. *Angew. Chem., Int. Ed.* **2010**, *49*, 5792.
- (a) Hatamoto, Y.; Sakaguchi, S.; Ishii, Y. *Org. Lett.* **2004**, *6*, 4623.
(b) Besset, T.; Kuhl, N.; Patureau, F. W.; Glorius, F. *Chem.-Eur. J.* **2011**, *17*, 7167.
(c) Zhang, Y.; Cui, Z.; Li, Z.; Liu, Z.-Q. *Org. Lett.* **2012**, *14*, 1838.
(d) Meng, K.; Sun, Y.; Zhang, J.; Zhang, K.; Ji, X.; Ding, L.; Zhong, G. *Org. Lett.* **2019**, *21*, 8219.
(e) Gong, T.-J.; Su, W.; Liu, Z.-J.; Cheng, W.-M.; Xiao, B.; Fu, Y. *Org. Lett.* **2014**, *16*, 330.
- (a) Poulsen, C. S.; Madsen, R. *Synthesis* **2003**, 1.
(b) Mori, M. Ene-Yne Metathesis. In *Handbook of Metathesis*, Ed.: Grubbs, R. H., Wiley-VCH, Weinheim, Germany, **2003**, Vol. 2, p. 176.
(c) Diver, S. T.; Giessert, A. J. *Chem. Rev.* **2004**, *104*, 1317.
(d) Diver, S. T. *Sci. Synth.* **2009**, *46*, 97.
(e) Clark, J. R.; Diver, S. T. *Org. Lett.* **2011**, *13*, 2896.
(f) Alcaide, B.; Almendros, P.; Luna, A. *Chem. Rev.* **2009**, *109*, 3817.
(g) Clark, J. R.; Griffiths, J. R.; Diver, S. T. *J. Am. Chem. Soc.* **2013**, *135*, 3327.
- (a) Nakata, T. *Chem. Soc. Rev.* **2010**, *39*, 1955.

- (b) Pathak, T. *Tetrahedron* **2008**, *64*, 3605.
 (c) Back, T. G.; Clary, K. N.; Gao, D. *Chem. Rev.* **2010**, *110*, 4498.
 (d) Bäckvall, J.-E.; Chinchilla, R.; Nájera, C.; Yus, M. *Chem. Rev.* **1998**, *98*, 2291.
 (e) Alba, A. R.; Companyo, X.; Rios, R. *Chem. Soc. Rev.* **2010**, *39*, 2018.
 (f) Caturla, F.; Nájera, C. *Tetrahedron* **1998**, *54*, 11255.
 (g) Tsogoeva, S. B. *Eur. J. Org. Chem.* **2007**, 1701.
 (h) El-Awa, A.; Noshi, M. N.; Jourdin, X. M.; Fuchs, P. L. *Chem. Rev.* **2009**, *109*, 2315.
 (i) Legarda, P.; García-Rubia, D. A.; Gómez Arrayás, R.; Carretero, J. C. *Adv. Synth. Catal.* **2016**, *358*, 1065.
 (j) Tang, Y.; Ran, S.; Wang, P.; Chen, P. *Chin. J. Org. Chem.* **2019**, *39*, 1116 (in Chinese).
 (唐裕才, 冉书童, 王萍, 陈飘, 有机化学, **2019**, *39*, 1116.)
 (k) Chen, D.; Sun, Y.; Dong, D.; Han, Q.; Wang, Z. *Chin. J. Org. Chem.* **2020**, *40*, 4267 (in Chinese).
 (陈德茂, 孙媛媛, 董道青, 韩晴晴, 王祖利, 有机化学, **2020**, *40*, 4267.)
- [15] (a) Qiao, B.; Liu, Q.; Liu, H.; Yan, L.; Jiang, Z. *Chem.-Asian J.* **2014**, *9*, 1252.
 (b) Zhu, Q.; Lu, Y. *Org. Lett.* **2008**, *10*, 4803.
 (c) Landa, A.; Maestro, M.; Masdeu, C.; Puente, Á.; Vera, S.; Oiarbide, M.; Palomo, C. *Chem.-Eur. J.* **2009**, *15*, 1562.
 (d) Dou, X.; Lu, Y. *Org. Biomol. Chem.* **2013**, *11*, 5217.
 (e) Chepyshev, S. V.; Lujan-Montelongo, J. A.; Chao, A.; Fleming, F. F. *Angew. Chem., Int. Ed.* **2017**, *56*, 4310.
 (f) Bos, P. H.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2008**, *10*, 4219.
- [16] (a) Nájera, C.; Yus, M. *Tetrahedron* **1999**, *55*, 10547.
 (b) Braun, M.-G.; Quiclet-Sire, B.; Zard, S. Z. *J. Am. Chem. Soc.* **2011**, *133*, 15954.
 (c) Chang, M.; Wu, Y.; Tsai, Y.; Chen, H. *J. Org. Chem.* **2019**, *84*, 11699.
- [17] (a) Ma, S.; Ren, H.; Wei, Q. *J. Am. Chem. Soc.* **2003**, *125*, 4817.
 (b) Chen, G.; Yu, Y.; Cai, M. *Synth. Commun.* **2009**, *39*, 1478.
 (c) Clayden, J.; Julia, M. J. *Chem. Soc., Chem. Commun.* **1993**, 1682.
 (d) Fabre, J. L.; Julia, M.; Verpeaux, J. N. *Tetrahedron Lett.* **1982**, *23*, 2469.
 (e) Alvarez, E.; Cuvigny, T.; Hervé du Penhoat, C.; Julia, M. *Tetrahedron*, **1988**, *44*, 111.
 (f) Alvarez, E.; Cuvigny, T.; du Penhoat, C. H.; Julia, M. *Tetrahedron* **1988**, *44*, 119.
 (g) Fukuda, J.; Nogi, K.; Yorimitsu, H. *Asian J. Org. Chem.* **2018**, *7*, 2049.
- [18] (a) Theobald, R. S.; Schofield, K. *Chem. Rev.* **1950**, *46*, 170.
 (b) Simeon, S.; Rios, J. L.; Villar, A. *Pharmazie* **1989**, *44*, 593.
 (c) Janin, Y. L.; Croisy, A.; Riou, J.-F.; Bisagni, E. *J. Med. Chem.* **1993**, *36*, 3686.
 (d) Nakanishi, T.; Suzuki, M.; Saimoto, A.; Kabasawa, T. *J. Nat. Prod.* **1999**, *62*, 864.
 (e) Abdel-Halim, O. B.; Morikawa, T.; Ando, S.; Matsuda, H.; Yoshikawa, M. *J. Nat. Prod.* **2004**, *67*, 1119.
- [19] (a) Nicolaou, K. C.; Snyder, S. A.; Montagnon, T.; Vassilikogiannakis, G. *Angew. Chem., Int. Ed.* **2002**, *41*, 1668.
 (b) Yang, B.; Gao, S. *Chem. Soc. Rev.* **2018**, *47*, 7926.
 (c) Ju, C.; Wu, Z.; Li, Y.; Zhang, W. *Chin. J. Org. Chem.* **2020**, *40*, 3925 (in Chinese).
 (居辰阳, 吴正兴, 李云艺, 张万斌, 有机化学, **2020**, *40*, 3925.)
 (d) Sun, M.; Zhang, J.; Zhang, Y.; Li, P.; Wang, L. *Chin. J. Org. Chem.* **2020**, *40*, 2419 (in Chinese).
 (孙明利, 张家俊, 张义成, 李品华, 王磊, 有机化学, **2020**, *40*, 2419.)
- [20] Li, Y.; Miao, T.; Li, P.; Wang, L. *Org. Lett.* **2018**, *20*, 1735.

(Li, L.; Fan, Y.)