烯胺调控下和对甲苯磺酰叠氮在纯水介质中的无金属环化反应 合成 1,2,3-三氮唑

郑茜茜^a 刘云云^a 万结平^{*,a}

(^a江西师范大学 化学化工学院 南昌 330022)

摘要 通过使用简单的 Et₃N 催化容易获得的 β -取代 NH-烯胺酯和对甲苯磺酰叠氮的反应, 高效实现了全取代的 1,2,3-三氮唑的合成. 在该方法中, 水用作反应的唯一介质, 以高底物适用性和中等至优秀的产率合成了 1,2,3-三氮唑产物. 对照实验表明, 使用稳定的 NH-烯胺作为底物, 可能是通过 NH 基团与水之间的氢键作用实现水介导反应的关键因素. 研究还发现, 在相同条件下经由相应的 *N*-烷基烯胺酯和对甲苯磺酰叠氮的反应, 选择性地生成 *N*-烷基磺酰胺.

关键词 烯胺; 环化反应; 无金属; 纯水介质; 1,2,3-三氮唑

Metal-Free Synthesis of 1,2,3-Triazoles in Pure Water via the Enamine Modified Annulation Reactions with Tosyl Azide

Zheng, Xixi^a Liu, Yunyun^a Wan, Jie-Ping^{*}

(College of Chemistry and Chemical Engineering, Jiangxi Normal University, Nanchang 330022)

Abstract The synthesis of full substituted 1,2,3-triazoles has been accomplished with high efficiency through the reactions of readily available β -substituted NH-enaminoesters and tosyl azide with Et₃N catalysis. In this method, water was used as the sole medium for the reactions which provide 1,2,3-triazole products with broad scope and moderate to excellent yield. Control experiments disclose that the employment of stable NH-enamines as substrates is the key factor enabling the water mediated synthesis probably via the hydrogen bonding effect between NH group and water. In addition, the novel and selective production of *N*-alkyl sulfonamides via the reactions of corresponding *N*-alkyl enaminoesters and tosyl azide has been observed under identical conditions.

Keywords enamine; annulation; metal-free; pure water medium; 1,2,3-triazole

1 Introduction

In the process of developing green organic synthesis, replacing organic solvent with water as the reaction medium is inarguably the ultimate aim for those reactions requiring a medium.^[1] In this context, revolutionary progress has been gained in the area of water mediated sustainable synthesis. Besides simply acting as alternative reaction medium, water has also been found to be capable of accelerating reaction processes and/or switching reaction selectivity in many cases, thus demonstrating the high merits of water in organic synthesis.^[2] To date, a number of organic reactions featuring the construction of C—C,^[3] C—heteroatom,^[4] heteroatom—heteroatom,^[5] chemical bond as well as other important transformation in water medium have been dis-

closed.^[6] However, comparing with huge number of organic reactions employing organic solvent(s) as medium, the overall availability of practical water-mediated organic reactions is yet scarce. Therefore, devising more practical and sustainable organic reactions using pure water as medium is yet highly desirable work.

As valuable heterocyclic scaffolds with widespread applications in multidisciplinary fields, 1,2,3-triazoles have received extensive and everlasting attention from different scientific communities,^[7] which has driven accordingly the splendid advances in the area of 1,2,3-triazole synthesis. Particularly, the transition metal-free annulation reactions have been proved as powerful tools in the synthesis of 1,2,3-triazoles with enriched substructure diversity.^[8] Representatively, the annulations of azide with aldehydes,^[9]

* Corresponding author. E-mail: wanjieping@jxnu.edu.cn

Received April 26, 2021; revised May 10, 2021; published online May 24, 2021.

Project supported by the National Natural Science Foundation of China (No. 21861019) and the Natural Science Foundation of Jiangxi Province (No. 20202ACBL203006).

国家自然科学基金(No. 21861019)和江西省自然科学基金(No. 20202ACBL203006)资助项目.

ketones,^[10] electron-deficient alkenes,^[11] cycloalkynes or alkynes,^[12] enamines,^[13] hydrazones,^[14] *etc.* have been successfully developed as practical routes to access various 1,2,3-triazoles under metal-free conditions. Moreover, alternative synthetic protocols without using azide as build block have also been established as sustainable complement to the common azide-based synthesis.^[15] Regardless these significant advances taking place in the synthesis of 1,2,3-triazoles, pure water mediated synthesis of 1,2,3-triazoles without transition metal catalysis is yet hardly available. Considering the incomparable advantages of greenness and inherent safety of synthetic methods in pure water, devising metal-free methods for the practical synthesis of 1,2,3-triazoles with pure water as medium is inarguably highly desirable.

Over the past few years, our sustaining efforts in novel synthetic and/or catalytic methods on enamines have disclosed that the hydrophilic effect of the enamines could facilitate their synthetic transformations in different reactions providing sulfonyl pyrazoles, *N*-sulfonyl amidines, α -diazoketones *etc.*^[16] Being inspired by these results, we envision that the synthesis of 1,4,5-trisubstituted 1,2,3-triazoles which have been previously synthesized via annulation reactions in volatile organic solvents may be alternatively achieved in pure water medium. To further disclose the magic application of stable enamines in the synthesis of diverse organic products,^[17] herein, we report our results on the green synthesis of these polyfunctionalized 1,2,3-triazoles via the annulation reactions of *NH*-enaminones and tosyl azide in pure water using Et₃N (10 mol%) as the only catalyst. In addition, the novel formation of sulfonamides via the reactions of tosyl azide and *N*-alkyl enaminones are also reported. Despite the splendid advances being achieved in the organocatalytic 1,2,3-triazole synthesis, to the best of our knowledge, the present work is the first example on the organocatalytic synthesis of 5-ester functionalized 1,2,3-triazoles in pure water medium.

2 Results and discussion

In an initial effort, the reaction of β -substituted *NH*-enaminoester **1a** and tosyl azide **2** (75% solution in ethyl acetate) was tentatively run in water by heating at 80 °C for 12 h in the presence of 0.4 equiv. of Et₃N, which provided 1,4,5-trisubstituted 1,2,3-triazole product **3a** with 80% yield (Entry 1, Table 1). A blank entry without using any catalyst gave only trace **3a** (Entry 2, Table 1). In addition, the comparison using different base species such as K₂CO₃, Cs₂CO₃ and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) proved that Et₃N was the ideal catalyst for this reaction (Entries 3~5, Table 1). Later, the same reaction conducted at different temperature led to the observation of excellent 92% excellent yield by heating at 70 °C (Entries 6~8, Table 1). Notably, the parallel entry in organic medium, including EtOH, dimethyl sulfoxide (DMSO), *N,N*-dimethylformamide (DMF) and toluene all gave evidently inferior results to the reaction in water (Entries 9~12, Ta-

ble 1). Moreover, shortening the reaction time of the in-water entry to 40 min gave yet identically excellent yield of the product, indicating that water could efficiently accelerate the target reaction (Entry 13, Table 1). Finally, varying the loading of Et₃N confirmed that only 10 mol% catalyst was abundant for the synthesis of **3a** with excellent yield (Entries 14~15, Table 1). However, product **3a** could not be afforded by using sodium azide and enaminoester **1a** under the optimized conditions.

Table 1 Optimization on reaction conditions^a

Entry	Solvent	Base	T/°C	Yield ^b /%
1	H ₂ O	Et ₃ N	80	80
2	H ₂ O	—	80	Trace
3	H ₂ O	K ₂ CO ₃	80	39
4	H ₂ O	Cs ₂ CO ₃	80	50
5	H ₂ O	DBU	80	46
6	H ₂ O	Et ₃ N	60	76
7	H ₂ O	Et ₃ N	70	92
8	H ₂ O	Et ₃ N	90	78
9	EtOH	Et ₃ N	70	46
10	DMSO	Et ₃ N	70	78
11	DMF	Et ₃ N	70	70
12	Toluene	Et ₃ N	70	Trace
13 ^c	H ₂ O	Et ₃ N	70	92
14 ^{c,d}	H ₂ O	Et ₃ N	70	91
15 ^{c,e}	H ₂ O	Et ₃ N	70	91

^a General conditions: **1a** (0.5 mmol), **2** (1.25 mmol), base (0.2 mmol), stirred for 12 h in 2.0 mL of solvent. ^b Yield of isolated product based on **1a**. ^c The reaction time was 40 min. ^d Et₃N (0.1 mmol). ^e Et₃N (0.05 mmol).

Having the efficient and green reaction conditions in hand, the scope of the green method on the synthesis of 1,2,3-triazoles was then investigated by employing enaminoesters **1** containing different substructures. The reactions of enaminoesters derived from ethyl acetates and different anilines were first performed (Table 2). As indicated by the results (**3a**~**3i**), these reactions all provided the target product with excellent yield. On other hand, the enaminoesters functionalized with β -aryl structure were found to be much less reactive toward the 1,2,3-triazole formation. Prolonging the reaction time to 5 h could give products with moderate to good yields (**3j**~**3r**). Electron donating group in the *N*-phenyl ring turned out to provide corresponding products with better yields (**3p**~**3r**). Subsequently, the reactions employing enaminoesters containing various alkyls in ester fragment also displayed fine tolerance to the present protocol by affording 1,2,3-triazoles with excellent yield (**3s**~**3v**). The alkyl group in this position showed no evident effect to the reaction result. Moreover, when substrates with different alkyl substituents in the β -site of **1** were used, the products were also provided with excellent yield (**3w**~**3x**). When an enaminone ((*Z*)-1-phenyl-3-(phenylamino)but-2-en-1-one) was used to react with tosyl

azide under the standard conditions, no such reaction forming 1,2,3-triazole was observed. The reaction using NH_2 -based enaminoester of type **1** (H instead of Ar, $\text{R}^1 = \text{Et}$ and $\text{R}^2 = \text{Me}$) with tosyl azide was also conducted, but the expected NH-1,2,3-triazole was not produced therein, either.

Table 2 Scope on the 1,2,3-triazole synthesis in pure water^{a,b}

^a General conditions: enaminone **1** (0.5 mmol), tosyl azide **2** (1.25 mmol), and Et₃N (0.05 mmol) in H₂O (2.0 mL), stirred for 40 min at 70 °C. ^b Isolated yield based on **1**. ^c The yield resulting from a 5 mmol scale reaction. ^d The reaction time was 5 h.

To testify the role of enamine structure for this pure water-mediated synthesis, the reaction directly using ethyl acetoacetate, aniline and tosyl azide was performed under the standard conditions in pure water. In this entry, no formation of product **3a** was observed, confirming that the modification using enaminone as starting materials was crucial for the present method toward clean synthesis of 1,2,3-triazoles in pure water medium. The state-of-the-art of green synthesis by means of substrate designation was thus reflected in the work (Eq. 1). The hydrogen bond between the enaminone substrate and water was crucial to the reaction, the general mechanism of the present reaction was similar with the reaction we previously reported in organic medium involving a key Regitz diazo transfer.^[13b]

Based on the good results, the reaction was carried out with *N*-alkyl enaminone, our attention was then turned to the potential synthesis of *N*-alkyl 1,2,3-triazoles by using *N*-alkyl enaminone. Amazingly, *N*-alkyl sulfonamides **5** were selectively afforded when the substrates were run under the standard conditions. The expected 1,2,3-triazoles, however, were not observed (Eq. 2). Such results indicated the additional application potential of this water mediated green system in the synthesis of sulfonamides.

3 Conclusions

In conclusion, by making use of the featured hydrophilic site in enaminone, we have developed a new and green catalytic approach for 1,2,3-triazole synthesis via the annulation of enaminone and tosyl azide. Most of the reactions proceed efficiently in pure water medium by using only low loading Et₃N catalyst (10 mol%) and provide target products with excellent yield. In addition, the novel synthesis of *N*-alkyl sulfonamides via the reactions of *N*-alkyl enaminone and tosyl azide has been discovered for the first time. The clean operation as well as the low-cost reagents make these methods highly attractive in the synthesis of these triazole heterocycles and sulfonamides.

4 Experimental section

4.1 General experimental information

All experiments were carried out under air atmosphere. The NH-enaminone **1** and **4** were synthesized following literature process.^[18] All other chemicals and solvents used in the experiments were obtained from commercial sources and used directly without further treatment. The ¹H NMR and ¹³C NMR spectra were recorded in 400 MHz apparatus at the frequencies of 400 MHz and 100 MHz, respectively. The chemical shifts were reported with TMS as internal standard. Melting points were tested in an X-4A instrument without correcting temperature and the HRMS were obtained under ESI model in a mass spectrometer equipped with TOF analyzer.

4.2 General procedure for the synthesis of 1,2,3-triazoles **3** and sulfonamides **5**

To a 25 mL round-bottom flask were added enaminone **1** or **4** (0.5 mmol), tosyl azide **2** (1.25 mmol) and pure water (2.0 mL). Then the mixture was stirred at 70 °C for 40 min or 5 h. Upon completion (TLC), the reaction mixture was allowed to cool to room temperature and diluted with water (5 mL). The resulting suspension was extracted with ethyl acetate (8 mL × 3). The combined organic solution was dried over anhydrous Na₂SO₄. After filtration, the solvent was removed at reduced pressure. Purification of the residue by flash column chromatography using mixed ethyl acetate (EA)/petroleum ether (PE) (*V*: *V* = 1:10) as eluent afforded analytically pure products **3** and **5**.

Ethyl 5-methyl-1-phenyl-1*H*-1,2,3-triazole-4-carboxylate (**3a**):^[10d] Yield 91%, 105 mg, white solid, m.p. 56~57 °C (lit. 46~48 °C). ¹H NMR (400 MHz, CDCl₃) δ:

7.64~7.51 (m, 3H), 7.48~7.39 (m, 2 H), 4.47 (q, $J=7.1$ Hz, 2H), 2.60 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.7, 138.8, 136.7, 135.4, 130.1, 129.7, 125.4, 61.0, 14.4, 10.0.

Ethyl 5-methyl-1-(*p*-tolyl)-1*H*-1,2,3-triazole-4-carboxylate (**3b**):^[10e] Yield 97%, 119 mg, white solid, m.p. 137~138 °C (lit. 130~132 °C). ^1H NMR (400 MHz, CDCl_3) δ : 7.34 (q, $J=8.0$ Hz, 4H), 4.46 (q, $J=7.1$ Hz, 2H), 2.57 (s, 3H), 2.46 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.8, 140.3, 138.8, 136.6, 133.0, 130.2, 125.2, 61.0, 21.3, 14.4, 10.0.

Ethyl 1-(4-fluorophenyl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3c**): Yield 94%, 117 mg, white solid, m.p. 126~127 °C. ^1H NMR (400 MHz, CDCl_3) δ : 7.53~7.41 (m, 2H), 7.34~7.23 (m, 2H), 4.46 (q, $J=7.1$ Hz, 1H), 2.59 (s, 1H), 1.45 (t, $J=7.1$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.2 (d, $^1J_{\text{C-F}}=249.7$ Hz), 161.6, 138.9, 136.7, 131.5 (d, $^4J_{\text{C-F}}=3.1$ Hz), 127.4 (d, $^3J_{\text{C-F}}=8.9$ Hz), 116.7 (d, $^2J_{\text{C-F}}=23.2$ Hz), 61.1, 14.3, 9.9; ESI-HRMS calcd for $\text{C}_{12}\text{H}_{13}\text{FN}_3\text{O}_2$ [$\text{M}+\text{H}$]⁺ 250.0986, found 250.1000.

Ethyl 1-(4-chlorophenyl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3d**):^[10d] Yield 94%, 125 mg, white solid, m.p. 165~166 °C (lit. 168~170 °C). ^1H NMR (400 MHz, CDCl_3) δ : 7.56 (d, $J=8.6$ Hz, 2H), 7.42 (d, $J=8.7$ Hz, 2H), 4.47 (q, $J=7.1$ Hz, 2H), 2.60 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.6, 138.8, 136.9, 136.2, 133.9, 129.9, 126.6, 61.1, 14.4, 10.0.

Ethyl 1-(4-bromophenyl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3e**):^[10e] Yield 93%, 144 mg, white solid, m.p. 179~180 °C (lit. 176~178 °C). ^1H NMR (400 MHz, CDCl_3) δ : 7.72 (d, $J=8.7$ Hz, 2H), 7.36 (d, $J=8.7$ Hz, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 2.60 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.6, 138.8, 136.9, 134.4, 132.9, 126.8, 124.2, 61.2, 14.4, 10.0.

Ethyl 1-(4-methoxyphenyl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3f**):^[10e] Yield 98%, 128 mg, white solid, m.p. 140~141 °C (lit. 140~142 °C). ^1H NMR (400 MHz, CDCl_3) δ : 7.36 (d, $J=8.9$ Hz, 2H), 7.06 (d, $J=8.9$ Hz, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 3.89 (s, 3H), 2.55 (s, 3H), 1.44 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.8, 160.7, 139.0, 136.5, 128.2, 126.7, 114.7, 61.0, 55.7, 14.4, 9.9.

Ethyl 1-([1,1'-biphenyl]-2-yl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3g**): Yield 95%, 146 mg, colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.65 (t, $J=7.5$ Hz, 1H), 7.60~7.52 (m, 2H), 7.45 (d, $J=7.3$ Hz, 1H), 7.27~7.20 (m, 3H), 7.07 (dd, $J=6.4, 3.0$ Hz, 2H), 4.37 (q, $J=7.1$ Hz, 2H), 1.98 (s, 3H), 1.39 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.5, 140.0, 139.1, 136.8, 136.0, 132.8, 131.1, 130.9, 128.7, 128.6, 128.2, 128.2, 128.1, 60.8, 14.2, 9.1; ESI-HRMS calcd for $\text{C}_{18}\text{H}_{18}\text{N}_3\text{O}_2$ [$\text{M}+\text{H}$]⁺ 308.1394, found 308.1409.

Ethyl 1-(3-methoxyphenyl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3h**): Yield 94%, 123 mg, colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.47 (t, $J=8.3$ Hz, 1H), 7.08 (d, $J=8.5$ Hz, 1H), 7.01 (d, $J=7.7$ Hz, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 3.87 (s, 3H), 2.60 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H);

^{13}C NMR (100 MHz, CDCl_3) δ : 161.7, 160.4, 138.8, 136.6, 136.3, 130.4, 117.3, 115.8, 111.2, 61.0, 55.6, 14.3, 10.0; ESI-HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_3\text{O}_3$ [$\text{M}+\text{H}$]⁺ 262.1186, found 262.1206.

Ethyl 1-(2-methoxyphenyl)-5-methyl-1*H*-1,2,3-triazole-4-carboxylate (**3i**): Yield 90%, 117 mg, colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.54 (td, $J=8.2, 1.6$ Hz, 1H), 7.36 (dd, $J=7.7, 1.5$ Hz, 1H), 7.12 (t, $J=8.3$ Hz, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 3.80 (s, 3H), 2.43 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.9, 153.9, 140.9, 135.9, 132.1, 128.4, 124.0, 121.0, 112.2, 60.9, 55.8, 14.4, 9.4; ESI-HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_3\text{O}_3$ [$\text{M}+\text{H}$]⁺ 262.1186, found 262.1198.

Ethyl 1,5-diphenyl-1*H*-1,2,3-triazole-4-carboxylate (**3j**):^[10d] Yield 40%, 59 mg, white solid, m.p. 127~128 °C (lit. 95~97 °C). ^1H NMR (400 MHz, CDCl_3) δ : 7.45~7.35 (m, 6H), 7.32~7.26 (m, 4H), 4.37 (q, $J=7.1$ Hz, 2H), 1.33 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.0, 140.9, 137.0, 135.8, 130.3, 129.9, 129.5, 129.3, 128.4, 125.8, 125.2, 61.2, 14.2.

Ethyl 5-phenyl-1-(*p*-tolyl)-1*H*-1,2,3-triazole-4-carboxylate (**3k**): Yield 45%, 69 mg, white solid, m.p. 119~120 °C. ^1H NMR (400 MHz, CDCl_3) δ : 7.44~7.34 (m, 3H), 7.29 (d, $J=7.5$ Hz, 2H), 7.15 (s, 4H), 4.36 (q, $J=7.1$ Hz, 2H), 2.35 (s, 3H), 1.32 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.0, 140.8, 139.7, 136.9, 133.4, 130.3, 129.8, 129.8, 128.3, 125.9, 125.0, 61.1, 21.2, 14.2; ESI-HRMS calcd for $\text{C}_{18}\text{H}_{18}\text{N}_3\text{O}_2$ [$\text{M}+\text{H}$]⁺ 308.1394, found 308.1414.

Ethyl 1-(4-fluorophenyl)-5-phenyl-1*H*-1,2,3-triazole-4-carboxylate (**3l**): Yield 46%, 72 mg, white solid, m.p. 101~102 °C. ^1H NMR (400 MHz, CDCl_3) δ : 7.40 (dt, $J=14.4, 7.0$ Hz, 3H), 7.32~7.24 (m, 4H), 7.07 (t, $J=8.4$ Hz, 2H), 4.36 (q, $J=7.1$ Hz, 2H), 1.31 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.7 (d, $^1J_{\text{C-F}}=249.2$ Hz), 160.8, 140.9, 137.0, 131.9 (d, $^4J_{\text{C-F}}=2.9$ Hz), 130.2, 130.0, 128.4, 127.2 (d, $^3J_{\text{C-F}}=8.9$ Hz), 125.6, 116.4 (d, $^2J_{\text{C-F}}=23.0$ Hz), 61.2, 14.1; ESI-HRMS calcd for $\text{C}_{17}\text{H}_{15}\text{FN}_3\text{O}_2$ [$\text{M}+\text{H}$]⁺ 312.1143, found 312.1155.

Ethyl 1-(4-chlorophenyl)-5-phenyl-1*H*-1,2,3-triazole-4-carboxylate (**3m**): Yield 43%, 70 mg, white solid, m.p. 96~97 °C. ^1H NMR (400 MHz, CDCl_3) δ : 7.47~7.34 (m, 5H), 7.30~7.26 (m, 2H), 7.22 (d, $J=8.3$ Hz, 2H), 4.37 (q, $J=7.1$ Hz, 2H), 1.32 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.8, 140.8, 137.2, 135.6, 134.3, 130.2, 130.1, 129.6, 128.5, 126.3, 125.5, 61.3, 14.2; ESI-HRMS calcd for $\text{C}_{17}\text{H}_{15}\text{ClN}_3\text{O}_2$ [$\text{M}+\text{H}$]⁺ 328.0847, found 328.0864.

Ethyl 1-(4-bromophenyl)-5-phenyl-1*H*-1,2,3-triazole-4-carboxylate (**3n**): Yield 38%, 70 mg, white solid, m.p. 106~107 °C. ^1H NMR (400 MHz, CDCl_3) δ : 7.57~7.38 (m, 5H), 7.29 (d, $J=7.4$ Hz, 2H), 7.16 (d, $J=8.4$ Hz, 2H), 4.36 (q, $J=6.7$ Hz, 2H), 1.31 (t, $J=6.9$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.8, 140.8, 137.2, 134.8, 132.6, 130.2, 130.1, 128.5, 126.6, 125.5, 123.6, 61.3, 14.1; ESI-HRMS calcd for $\text{C}_{17}\text{H}_{15}\text{BrN}_3\text{O}_2$ [$\text{M}+\text{H}$]⁺ 372.0342, found 372.0354.

Ethyl 1-([1,1'-biphenyl]-2-yl)-5-phenyl-1*H*-1,2,3-triazole-4-carboxylate (**3o**): Yield 55%, 101 mg, colorless liquid.

^1H NMR (400 MHz, CDCl_3) δ : 7.59~7.52 (m, 1H), 7.46~7.37 (m, 2H), 7.20~7.16 (m, 1H), 7.13~7.06 (m, 2H), 6.97 (dt, $J=15.6, 7.6$ Hz, 4H), 6.56~6.41 (m, 4H), 4.19 (q, $J=7.0$ Hz, 2H), 1.16 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.9, 142.2, 138.6, 136.8, 136.0, 133.4, 130.8, 130.7, 129.6, 129.2, 128.5, 128.1, 127.6, 127.5, 124.8, 61.0, 14.1; ESI-HRMS calcd for $\text{C}_{23}\text{H}_{20}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 370.1550, found 370.1568.

Ethyl 1-(4-methoxyphenyl)-5-phenyl-1H-1,2,3-triazole-4-carboxylate (**3p**):^[9b] Yield 68%, 110 mg, white solid, m.p. 112~113 °C (lit. 83~85 °C). ^1H NMR (400 MHz, CDCl_3) δ : 7.42~7.34 (m, 3H), 7.29 (d, $J=6.5$ Hz, 2H), 7.18 (d, $J=8.9$ Hz, 2H), 6.86 (d, $J=8.9$ Hz, 2H), 4.36 (q, $J=7.1$ Hz, 2H), 3.80 (s, 3H), 1.31 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.0, 160.2, 140.8, 136.8, 130.3, 129.8, 128.8, 128.3, 126.6, 125.9, 114.4, 61.1, 55.5, 14.1.

Ethyl 1-(3-methoxyphenyl)-5-phenyl-1H-1,2,3-triazole-4-carboxylate (**3q**): Yield 62%, 100 mg, white solid, m.p. 126~127 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.45~7.36 (m, 3H), 7.31 (d, $J=7.0$ Hz, 2H), 7.24 (t, $J=8.1$ Hz, 1H), 6.92 (d, $J=9.0$ Hz, 1H), 6.82 (d, $J=7.3$ Hz, 2H), 4.36 (q, $J=7.1$ Hz, 2H), 3.69 (s, 3H), 1.31 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.9, 160.0, 140.8, 137.0, 136.7, 130.2, 130.0, 129.9, 128.3, 125.9, 117.3, 115.6, 110.7, 61.1, 55.5, 14.1; ESI-HRMS calcd for $\text{C}_{18}\text{H}_{18}\text{N}_3\text{O}_3$ $[\text{M} + \text{H}]^+$ 324.1343, found 324.1358.

Ethyl 1-(2-methoxyphenyl)-5-phenyl-1H-1,2,3-triazole-4-carboxylate (**3r**): Yield 59%, 95 mg, white solid, m.p. 123~124 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.38 (t, $J=7.7$ Hz, 2H), 7.35~7.25 (m, 5H), 7.01 (t, $J=7.6$ Hz, 1H), 6.85 (d, $J=8.2$ Hz, 1H), 4.36 (q, $J=7.1$ Hz, 2H), 3.46 (s, 3H), 1.32 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.2, 153.6, 142.7, 136.0, 131.8, 129.6, 129.5, 128.5, 127.8, 126.2, 124.7, 120.8, 112.2, 61.0, 55.4, 14.2; ESI-HRMS calcd for $\text{C}_{18}\text{H}_{18}\text{N}_3\text{O}_3$ $[\text{M} + \text{H}]^+$ 324.1343, found 324.1359.

Methyl 5-methyl-1-phenyl-1H-1,2,3-triazole-4-carboxylate (**3s**):^[10d] Yield 92%, 100 mg, white solid, m.p. 50~51 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.58 (d, $J=7.1$ Hz, 3H), 7.46 (dd, $J=7.6, 1.8$ Hz, 2H), 3.98 (s, 3H), 2.60 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.0, 139.0, 136.4, 135.3, 130.1, 129.7, 125.3, 51.9, 9.9.

Propyl 5-methyl-1-phenyl-1H-1,2,3-triazole-4-carboxylate (**3t**): Yield 86%, 105 mg, colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.50~7.42 (m, 3H), 7.38~7.31 (m, 2H), 4.28~4.19 (m, 2H), 2.50~2.44 (m, 3H), 1.78~1.67 (m, 2H), 0.98~0.88 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.7, 138.7, 136.7, 135.4, 130.0, 129.6, 125.3, 66.5, 22.0, 10.4, 10.0; ESI-HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 246.1237, found 246.1255.

Isobutyl 5-methyl-1-phenyl-1H-1,2,3-triazole-4-carboxylate (**3u**): Yield 84%, 109 mg, yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.46 (t, $J=6.3$ Hz, 3H), 7.35 (t, $J=6.3$ Hz, 2H), 4.07 (t, $J=6.2$ Hz, 2H), 2.48 (d, $J=5.2$ Hz, 3H), 2.03 (dq, $J=12.6, 6.5$ Hz, 1H), 0.92 (t, $J=6.0$ Hz, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.7, 138.6, 136.7, 135.4, 130.0, 129.6, 125.3, 70.9, 27.8, 19.1, 10.0; ESI-

HRMS calcd for $\text{C}_{14}\text{H}_{18}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 260.1394, found 260.1410.

tert-Butyl 5-methyl-1-phenyl-1H-1,2,3-triazole-4-carboxylate (**3v**): Yield 81%, 105 mg, white solid, m.p. 77~78 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.50~7.41 (m, 3H), 7.33 (td, $J=5.1, 3.1$ Hz, 2H), 2.45 (d, $J=3.4$ Hz, 3H), 1.54 (d, $J=3.7$ Hz, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.9, 138.1, 137.8, 135.6, 129.9, 129.6, 125.3, 82.0, 28.3, 10.1; ESI-HRMS calcd for $\text{C}_{14}\text{H}_{18}\text{N}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 260.1394, found 260.1408.

Methyl 5-ethyl-1-phenyl-1H-1,2,3-triazole-4-carboxylate (**3w**):^[11a] Yield 83%, 96 mg, yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.62~7.56 (m, 3H), 7.45 (dd, $J=6.8, 2.9$ Hz, 2H), 4.00 (s, 3H), 3.01 (q, $J=7.5$ Hz, 2H), 1.17 (t, $J=7.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.9, 144.6, 135.7, 135.4, 130.3, 129.7, 125.6, 51.9, 16.9, 13.1.

Methyl 1-phenyl-5-propyl-1H-1,2,3-triazole-4-carboxylate (**3x**):^[11a] Yield 80%, 98 mg, white solid, m.p. 72~73 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.61~7.56 (m, 3H), 7.43 (dd, $J=6.5, 2.9$ Hz, 2H), 4.00 (s, 3H), 2.97 (t, $J=7.8$ Hz, 2H), 1.57 (dt, $J=15.2, 7.5$ Hz, 2H), 0.86 (t, $J=7.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.0, 143.4, 136.1, 135.5, 130.3, 129.7, 125.7, 51.9, 25.1, 22.1, 13.7.

N-Butyl-4-methylbenzenesulfonamide (**5a**):^[19] Yield 42%, 48 mg, colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.75 (d, $J=8.2$ Hz, 2H), 7.31 (d, $J=8.1$ Hz, 2H), 4.62 (s, 1H), 2.96~2.89 (m, 2H), 2.43 (s, 3H), 1.47~1.40 (m, 2H), 1.33~1.25 (m, 2H), 0.84 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 143.3, 137.1, 129.7, 127.1, 42.9, 31.59, 21.5, 19.7, 13.5.

N-Hexyl-4-methylbenzenesulfonamide (**5b**):^[19] Yield 52%, 66 mg, colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (d, $J=8.1$ Hz, 2H), 7.30 (d, $J=8.0$ Hz, 2H), 4.94 (s, 1H), 2.91 (q, $J=6.5$ Hz, 2H), 2.42 (s, 3H), 1.43 (q, $J=7.0$ Hz, 2H), 1.27~1.16 (m, 6H), 0.83 (t, $J=6.7$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 143.2, 137.1, 129.6, 127.1, 43.2, 31.2, 29.5, 26.2, 22.4, 21.5, 13.9.

4-Methyl-*N*-propylbenzenesulfonamide (**5c**):^[19] Yield 56%, 95 mg, white solid, m.p. 70~71 °C (lit. 73 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.69 (d, $J=8.2$ Hz, 2H), 7.21 (d, $J=8.1$ Hz, 2H), 5.05 (s, 1H), 2.89~2.76 (m, 2H), 2.33 (s, 3H), 1.34 (q, $J=7.0$ Hz, 2H), 1.13 (d, $J=21.4$ Hz, 18 H), 0.79 (t, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 143.1, 137.1, 129.6, 127.1, 43.2, 31.9, 29.6, 29.6, 29.5, 29.5, 29.4, 29.3, 29.1, 26.5, 22.7, 21.4, 14.1.

4.3 5 mmol scale reaction for the synthesis of **3a**

To a 100 mL round-bottom flask were charged with enaminoester **1a** (5 mmol), tosyl azide **2** (12.5 mmol), Et_3N (0.5 mmol) and H_2O (20 mL). The resulting mixture was stirred at 70 °C for 40 min (TLC) under air atmosphere. After cooling down to room temperature, the suspension was extracted with ethyl acetate (80 mL $\times$ 3). The combined organic solution was dried over anhydrous Na_2SO_4 . After filtration, the solvent was removed from the solution at reduced pressure. Purification of the residue by flash column chromatography using mixed ethyl acetate (EA)/pe-

troleum ether (PE) ($V:V=1:10$) as eluent provided pure product **3a**.

Supporting Information The ^1H NMR and ^{13}C NMR spectra of all products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Lipshutz, B. H.; Ghorai, S.; Cortes-Clerget, M. *Chem.-Eur. J.* **2018**, *24*, 6672.
(b) Gawande, M. B.; Bonifácio, V. D. B.; Luque, R.; Branco, P. S.; Varma, R. S. *Chem. Soc. Rev.* **2013**, *42*, 5522.
(c) Anastas, P.; Eghbali, N. *Chem. Soc. Rev.* **2010**, *39*, 301.
- [2] (a) Butler, R. N.; Goynes, A. G. *Chem. Rev.* **2010**, *110*, 6302.
(b) Chanda, A.; Fokin, V. V. *Chem. Rev.* **2009**, *109*, 725.
(c) Sun, K.; Lv, Q.-Y.; Chen, X.-L.; Qu, L.-B.; Yu, B. *Green Chem.* **2021**, *23*, 232.
(d) Chen, D.; Liu, J.; Zhang, X.; Jiang, H.; Li, J. *Chin. J. Org. Chem.* **2019**, *39*, 3353 (in Chinese).
(陈丹, 刘剑沉, 张馨元, 蒋合众, 李加洪, 有机化学. **2019**, *39*, 3353.)
- [3] (a) Li, Y.; Huang, Y.; Gui, Y.; Sun, J.; Li, J.; Zha, Z.; Wang, Z. *Org. Lett.* **2017**, *19*, 6416.
(b) Zhang, F.; Tian, Y.; Li, G.; Qu, J. *J. Org. Chem.* **2015**, *80*, 1107.
(c) Álvarez, M.; Gava, R.; Rodríguez, M. R.; Rull, S. G.; Pérez, P. *J. ACS Catal.* **2017**, *7*, 3707.
(d) Zhang, N.; Yang, D.; Wei, W.; Yuan, L.; Nie, F.; Tian, L.; Wang, H. *J. Org. Chem.* **2015**, *80*, 3258.
(e) Tang, S.; Li, L.; Ren, X.; Li, J.; Yang, G.; Li, H.; Yuan, B. *Green Chem.* **2019**, *21*, 2899.
- [4] (a) Peng, S.; Song, Y.-X.; He, J.-Y.; Tang, S.-S.; Tan, J.-X.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, *30*, 2287.
(b) Wu, Y.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2020**, *31*, 2999.
(c) Shukla, P.; Asati, A.; Bhardiya, S. R.; Singh, M.; Rai, V. K.; Rai, A. *J. Org. Chem.* **2020**, *85*, 15552.
- [5] (a) Tang, L.; Yang, Y.; Wen, L.; Yang, X.; Wang, Z. *Green Chem.* **2016**, *18*, 1224.
(b) Lin, Y.; Lu, G.; Wang, G.; Yi, W. *J. Org. Chem.* **2017**, *82*, 382.
- [6] (a) Chen, L.; Huang, R.; Li, K.; Yun, X.-H.; Yang, C.-L.; Yan, S.-J. *Green Chem.* **2020**, *22*, 6943.
(b) Husain, A. A.; Bisht, K. *J. Org. Chem.* **2020**, *85*, 9928.
(c) Köhling, S.; Exner, M. P.; Nojoudi, S.; Schiller, J.; Budisa, N.; Rademann, J. *Angew. Chem. Int. Ed.* **2016**, *55*, 15510.
(d) Yang, J.; Mei, F.; Fu, S.; Gu, Y. *Green Chem.* **2018**, *20*, 1367.
(e) Reddy, G. T.; Kumar, G.; Reddy, N. C. G. *Adv. Synth. Catal.* **2018**, *360*, 995.
(f) Yang, L.; Wu, Y.; Yang, Y.; Wen, C.; Wan, J.-P. *Beilstein J. Org. Chem.* **2018**, *14*, 2348.
(g) Chen, X.; Xia, F.; Zhao, Y.; Ma, J.; Zhang, D.; Yang, L. *Sun, P. Chin. J. Chem.* **2020**, *38*, 1239.
(h) Sun, G.; He, Y.; Tian, C.; Borzov, M.; Hu, Q.; Nie, W. *Acta Chem. Sinica* **2019**, *77*, 166 (in Chinese).
(孙国峰, 何云清, 田冲, Borzov Maxim, 胡启山, 聂万丽, 化学学报. **2019**, *77*, 166.)
- [7] (a) Meldal, M.; Tomøe, C. W. *Chem. Rev.* **2008**, *108*, 2952.
(b) Thirumurugan, P.; Matosiuk, D.; Jozwiak, K. *Chem. Rev.* **2013**, *113*, 4905.
(c) Jiang, Y.; Sun, R.; Tang, X.-Y.; Shi, M. *Chem.-Eur. J.* **2016**, *22*, 17910.
- [8] For selected reviews, see:
(a) Opsomer, T.; Dehaen, W. *Chem. Commun.* **2021**, *57*, 1568.
(b) Ramasastry, S. S. V. *Angew. Chem. Int. Ed.* **2014**, *53*, 14310.
(c) Lima, C. G. S.; Ali, A.; Berkel, S. S.; Westermann, B.; Paixão, M. W. *Chem. Commun.* **2015**, *51*, 10784.
- [9] (a) Ramachary, D. B.; Shashank, A. B.; Karthik, S. *Angew. Chem. Int. Ed.* **2014**, *53*, 10420.
(b) Thomas, J.; John, J.; Parekh, N.; Dehaen, W. *Angew. Chem. Int. Ed.* **2014**, *53*, 10155.
(c) Zhang, D.; Fan, Y.; Yan, Z.; Nie, Y.; Xiong, X.; Gao, L. *Green Chem.* **2019**, *21*, 4211.
(d) Reddy, G. S.; Reddy, L. M.; Kumar, A. S.; Ramachary, D. B. *J. Org. Chem.* **2020**, *85*, 15488.
- [10] (a) Belkheira, M.; Abed, D. E.; Pons, J.-M.; Bressy, C. *Chem.-Eur. J.* **2011**, *17*, 12917.
(b) Ramachary, D. B.; Krishna, P. M.; Gujral, J.; Reddy, G. S. *Chem.-Eur. J.* **2015**, *21*, 16775.
(c) Guo, N.; Liu, X.; Xu, H.; Zhou, X.; Zhao, H. *Org. Biomol. Chem.* **2019**, *17*, 6148.
(d) Danence, L. J. T.; Gao, Y.; Li, M.; Huang, Y.; Wang, J. *Chem.-Eur. J.* **2011**, *17*, 3584.
(e) Nelsona, R.; Kesternicha, V.; Pérez-Fehrmanna, M.; Jaldina, S.; Marcourt, L.; Christen, P. *J. Chem. Res.* **2016**, *40*, 453.
- [11] (a) Li, W.; Wang, J. *Angew. Chem. Int. Ed.* **2014**, *53*, 14186.
(b) Ramachary, D. B.; Ramakumar, K.; Narayana, V. V. *Chem.-Eur. J.* **2008**, *14*, 9143.
(c) Das, J.; Dey, S.; Pathak, T. *J. Org. Chem.* **2019**, *84*, 15437.
- [12] (a) Agard, N. J.; Prescher, J. A.; Bertozzi, C. R. *J. Am. Chem. Soc.* **2004**, *126*, 15046.
(b) Kwok, S. W.; Fotsing, J. R.; Fraser, R. J.; Rodionov, V. O.; Fokin, V. V. *Org. Lett.* **2010**, *12*, 4217.
- [13] (a) Cheng, G.; Zeng, X.; Shen, J.; Wang, X.; Cui, X. *Angew. Chem. Int. Ed.* **2013**, *52*, 13265.
(b) Wan, J.-P.; Cao, S.; Liu, Y. *Org. Lett.* **2016**, *18*, 6034.
(c) Deng, L.; Cao, X.; Liu, Y.; Wan, J.-P. *J. Org. Chem.* **2019**, *84*, 14179.
(d) De Nino, A.; Algieri, V.; Tallarida, M. A.; Constanzo, P.; Pedron, M.; Tejero, T.; Merino, P.; Maiuolo, L. *Eur. J. Org. Chem.* **2019**, *33*, 5725.
(e) Huang, W.; Zhu, C.; Li, M.; Yu, Y.; Wu, W.; Tu, Z.; Jiang, H. *Adv. Synth. Catal.* **2018**, *360*, 3117.
(f) Thomas, J.; Goyvaerts, V.; Liekens, S.; Dehaen, W. *Chem.-Eur. J.* **2016**, *22*, 9966.
(g) Cao, S.; Liu, Y.; Hu, C.; Wen, C.; Wan, J.-P. *ChemCatChem* **2018**, *10*, 5007.
- [14] (a) Shu, W.-M.; Zhang, X.-F.; Zhang, X.-X.; Li, M.; Wang, A.-J.; Wu, A.-X. *J. Org. Chem.* **2019**, *84*, 14919.
(b) Berkel, S. S.; Brauch, S.; Gabriel, L.; Henze, M.; Stark, S.; Vassilev, D.; Wessjohann, L. A.; Abbas, M.; Westermann, B. *Angew. Chem. Int. Ed.* **2012**, *51*, 5343.
- [15] For reviews and selected examples, see:
(a) Wan, J.-P.; Hu, D.; Liu, Y.; Sheng, S. *ChemCatChem* **2015**, *7*, 901.
(b) Chen, Z.; Gao, G.; Song, J.; Ren, H. *Chin. J. Chem.* **2017**, *35*, 1797.
(c) Shang, Z.-H.; Zhang, Z.-X.; Weng, W.-Z.; Wang, Y.-F.; Cheng, T.-W.; Zhang, Q.-Y.; Song, L.-Q.; Shao, T.-Q.; Liu, K.-X.; Zhu, Y.-P. *Adv. Synth. Catal.* **2020**, *363*, 490.
(d) Wan, J.-P.; Cao, S.; Liu, Y. *J. Org. Chem.* **2015**, *80*, 9028.
(e) Panda, S.; Maity, P.; Manna, D. *Org. Lett.* **2017**, *19*, 1534.
(f) Cai, Z.-J.; Lu, X.-M.; Zi, Y.; Yang, C.; Shen, L.-J.; Li, J.; Wang, S.-Y.; Ji, S.-J. *Org. Lett.* **2014**, *16*, 5108.
(g) Wu, P.; He, Y.; Wang, H.; Zhou, Y.-G.; Yu, Z. *Org. Lett.* **2020**, *22*, 310.
(h) Chen, Z.; Yan, Q.; Liu, Z.; Zhang, Y. *Chem.-Eur. J.* **2014**, *20*, 17635.
- [16] (a) Guo, Y.; Wang, G.; Wei, L.; Wan, J.-P. *J. Org. Chem.* **2019**, *84*, 2984.
(b) Zheng, X.; Wan, J.-P. *Adv. Synth. Catal.* **2019**, *361*, 5690.
(c) Gan, L.; Wei, L.; Wan, J.-P. *ChemistrySelect* **2020**, *5*, 7822.
(d) Fu, L.; Cao, X.; Wan, J.-P. *Chin. J. Chem.* **2020**, *38*, 254.
- [17] For a few recent examples, see
(a) Gan, L.; Yu, Q.; Liu, Y.; Wan, J.-P. *J. Org. Chem.* **2021**, *86*,

1231.
(b) Fu, L.; Xu, Z.; Wan, J.-P.; Liu, Y. *Org. Lett.* **2020**, *24*, 9518.
(c) Hu, D.; Yang, L.; Wan, J.-P. *Green Chem.* **2020**, *22*, 6773.
(d) Klintworth, R.; de Koning, C. B.; Opatz, T.; Michael, J. P. *J. Org. Chem.* **2019**, *84*, 11025.
(e) Chen, J.; Guo, P.; Zhang, J.; Rong, J.; Sun, W.; Jiang, Y.; Loh, T.-P. *Angew. Chem. Int. Ed.* **2019**, *58*, 12674.
(f) Gao, Y.; Liu, Y.; Wan, J.-P. *J. Org. Chem.* **2019**, *84*, 2243.
(g) Shang, Z.; Chen, Q.; Xing, L.; Zhang, Y.; Wait, L.; Du, Y. *Adv. Synth. Catal.* **2019**, *361*, 4926.
(h) Zhou, P.; Hu, B.; Li, L.; Rao, K.; Yang, J.; Yu, F. *J. Org. Chem.* **2017**, *82*, 13268.
(i) Gu, F.; Yao, W. *Chin. J. Org. Chem.* **2020**, *40*, 4384 (in Chinese).
(谷枫, 姚伟军, 有机化学, **2020**, *40*, 4384.)
(j) Wang, G.; Guo, Y.; Wan, J.-P. *Chin. J. Org. Chem.* **2020**, *40*, 645 (in Chinese).
(王国栋, 郭艳辉, 万结平, 有机化学, **2020**, *40*, 645.)
(k) Yu, Q.; Liu, Y.; Wan, J.-P. *Chin. Chem. Lett.* **2021**, DOI: 10.1016/j.ccl.2021.04.037.
[18] (a) Nitin, A. R.; Varun K.; Vipin, A. N. *Monatsh. Chem.* **2010**, *141*, 1329.
(b) Nishiwaki, N.; Nishimoto, T.; Tamura, M.; Ariga, M. *Synlett* **2006**, 1437.
[19] Tang, X.; Huang, L.; Qi, C.; Wu, X.; Wu, W.; Jiang, H. *Chem. Commun.* **2013**, *49*, 6102.

(Li, L.; Fan, Y.)