钯催化的芳酰氧基氨基甲酸酯的分子内脱羧烯丙基胺化反应

李晓颖 李培贺 王 峰 付 辉 戴启谱*

(北京理工大学化学与化工学院 原子分子簇科学教育部重点实验室 光电材料北京市重点实验室 北京 100081)

摘要 烯丙基胺化合物是有机合成化学和药物化学的重要化合物之一, 其合成方法备受关注. 以芳酰氧基氨基甲酸酯为原料通过分子内脱羧一步合成了烯丙基胺化合物, 而原料芳酰氧基氨基甲酸酯可由廉价易得的羧酸和羟胺原位制备. 反应条件温和, 底物普适性好, 并且可以实现克级和吲哚的合成.

关键词 钯催化; 芳酰氧基氨基甲酸酯; 脱羧; 烯丙基胺化; 烯丙胺

Palladium-Catalyzed Intramolecular Decarboxylative Allylic Amination of Aroyloxycarbamates

Li, Xiaoying Li, Peihe Wang, Zheng Fu, Hui Dai, Qipu*

(Beijing Key Laboratory of Photoelectronic/Electrophotonic, Key Laboratory of Cluster Science of Ministry of Education, School of Chemistry and Chemical Engineering, Beijing Institute of Technology, Beijing 100081)

Abstract Allylamine compounds are one of the important compounds in organic synthetic chemistry and pharmaceutical chemistry, the synthetic methods of which are concerned. Using aroyloxycarbamates as start material, which was prepared *in situ* by cheap and easy-available carboxylic acid and hydroxylamine, allylamine compounds were synthesized by decarboxylation in one step. The reaction conditions are mild and the substrate is generally applicable. Moreover, the reaction realized the synthesis of indole.

Keywords palladium-catalysis; aroyloxycarbamate; decarboxylation; allylic amination; allylamine

1 Introduction

As a valuable building block in chemistry, allylamines widely exist in active natural products, so many elegant methods have been developed for the synthesis of allylamines.^[1] In general, the allylamines were synthesized through the nucleophilic allylic substitution reaction employing allyl alcohols, allyl halides, acetates and carbonates as the electrophilic reagents.^[2] For instance, palladium,^[3] gold,^[4] iridium,^[5] nickel,^[6] oxovanadium^[7] and iron^[8] have been reported being employed as catalysts for the nucleophilic allylic substitution reaction (Scheme 1, a). What's more, metal-catalyzed reductive coupling of dienes, allenes with amines is an efficient synthetic method for allylamine.^[9] (Scheme 1, b). Despite of the above-mentioned advances, an intramolecular decarboxylation method was employed for the direct synthesis of allylic amines^[10] (Scheme 1, c). A palladium-catalyzed cross-coupling reaction is the most powerful tool for the formation of carbon-

carbon^[11] and carbon-heteroatom bonds.^[12] Although there are so many ways to obtain allylamines, there is no method to synthesize allylamines from carboxylic acid. Based on a long-time development of intramolecular decarboxylation coupling in our laboratory,^[13] we proposed an alternative, cheap, stable and environment-friendly carboxylic acids and hydroxylamine as raw materials to synthesize aroyloxycarbamates. Allylamines were obtained in one pot by palladium-catalyzed intramolecular decarboxylation process with two molecules of CO₂ releasing (Scheme 1, d).

2 Results and discussion

The optimization of the reaction was carried out using allyl ((2-methylbenzoyl)oxy)carbamate (**1a**) as the model substrate. Based on our previous study, we got Cs₂CO₃ as the optimal base. As shown in Table 1, after screening different catalysts, such as [Ir(cod)Cl]₂, Pd(OAc)₂, Pd(TFA)₂ and PdCl₂ with the Xantphos ligand or Pd(DPPF)Cl₂•

* Corresponding author. E-mail: daiqipu@bit.edu.cn

Received April 15, 2021; revised May 15, 2021; published online June 1, 2021.

Project supported by the Excellent Young Scholars Research Fund of Beijing Institute of Technology (Nos. 3190012211808, 3190012331518, 3190012331523) and the National Natural Science Foundation of China (No. 21871026).

北京理工大学优秀青年学者研究基金(Nos. 3190012211808, 3190012331518, 3190012331523)和国家自然科学基金(No. 21871026)资助项目.

Previous work:

(a) Nucleophilic allylic substitution reaction

(b) Metal-catalyzed reductive coupling reaction.

(c) Decarboxylative rearrangement of allylic carbamate

This work:

(d) Decarboxylation coupling reaction of alkanoyloxycarbamates

Scheme 1 Overview of synthetic allylamines

Table 1 Optimization of the reaction conditions^a

Entry	Catalyst	Ligand	Solvent	T/°C	Yield ^b /%
1	[Ir(cod)Cl] ₂	XantPhos	MeCN	80	17
2	Pd(OAc) ₂	XantPhos	MeCN	80	43
3	Pd(TFA) ₂	XantPhos	MeCN	80	69
4	PdCl ₂	XantPhos	MeCN	80	80
5	Pd(DPPF)Cl ₂ •CH ₂ Cl ₂		MeCN	80	21
6	PdCl ₂	L1	MeCN	80	12
7	PdCl ₂	L2	MeCN	80	Trace
8	PdCl ₂	XantPhos	Hexane	80	49
9	PdCl ₂	XantPhos	Toluene	80	52
10	PdCl ₂	XantPhos	THF	80	64
11	PdCl ₂	XantPhos	DMF	80	75
12	PdCl ₂	XantPhos	MeCN	100	65
13	PdCl ₂	XantPhos	MeCN	90	70
14	PdCl ₂	XantPhos	MeCN	70	65
15	PdCl ₂	XantPhos	MeCN	60	62

^a Reaction conditions: **1a** (0.20 mmol), base (0.20 mmol) in MeCN (2.0 mL) at 80 °C until **1a** was fully consumed (monitored by TLC) to get intermediate. Then the reaction mixture was filtrated and transferred to a nitrogen-filled Schlenk tube, then catalyst (2.5 mmol%) and ligand (5.0 mmol%) were added. The reaction mixture was stirred at *t* °C until intermediate was fully consumed (monitored by TLC) to get **2a**. ^b Unless otherwise noted, the yields were determined by GC analysis using biphenyl as internal standard. **L1** = 1,10-phenanthroline, **L2** = tri-*tert*-butyl phosphine tetrafluoroborate.

CH₂Cl₂, the PdCl₂ was selected as the efficient catalyst (Table 1, Entries 1~5). And the screening result of different phosphine and nitrogen ligands, such as Xantphos, 1,10-phenanthroline (**L1**), tri-*tert*-butyl phosphine tetrafluoroborate (**L2**), showed that Xantphos provided the best result (Table 1, Entries 4, 6, 7). Screening of solvent (Table 1, Entries 4, 8~11) and temperature (Table 1, Entries 4, 12~15) showed that MeCN and 80 °C were most suitable for this reaction. Thus, the highlighted condition in Entry 4 in Table 1 was selected as the optimal condition to explore the generality of this transformation.

With the optimum reaction conditions in hand, the scope of the reaction was investigated. As shown in Table 2, many functional groups, such as nitro (**2h**, **2i**, **2k**, **2o**), alkoxy (**2b**, **2j**) and halide groups (**2d**, **2e**, **2f**, **2l**, **2m**, **2n**, **2o**, **2p**, **2q**, **2r** and **2s**) were applicable to this reaction. *ortho*-Substituted allyl carbamates (**2b**) gave higher yield than that of *para*-substituted (**2j**). The yields of electron-donating groups were higher than those of the electron-withdrawing groups (**2a**, **2b**, **2d**, **2e**, **2f**, **2g**). To further investigate the property of the decarboxylation coupling reaction, different substrates were explored. As shown in Table 3, all of them had good yields. Compounds **1aa**, **1ae**, **1af**, **1ag** and **1ah** reacted smoothly and converted to the corresponding product in moderate yields. Introduction of a methyl group at the α-position of the aryloxycarbamates (**1ad**) got product in 63% yield. In 2010, Floch group^[14] reported that the introduction of a methyl group at the β-position led a mixture of S_N2 and S_N2' products in a 1 : 1 ratio such as **1ac**. The reaction systems of **1aa**, **1ad**~**1ah** were detected by NMR, and almost only one product of attack mode was obtained.

To evaluate the practical potential of this novel method on a large scale, 5.0 mmol of allyl ((2-methylbenzoyl)-oxy)carbamate (**1a**) was used to perform the decarboxylative allylic amination, and the product **2a** was obtained in 75% yield (Scheme 2, a).

The potential utility of decarboxylation coupling reaction was tested. Indoles with carbon substituents in the 3-position are of pivotal importance in central-nervous-system (CNS) drugs, particularly for antipsychotics.^[15] Although many indole syntheses are known, another truly general approach has yet to be discovered. Herein, we designed a route to synthesize 3-substituted indole from *N*-allyl-2-bromoaniline as shown in (Scheme 2, b), and the target product was obtained in 72% yield.

According to the proposed possible mechanism, in the presence of base, the proton was removed from nitrogen atom with the formation of intermediate **IV**. Then, the intermediate **II** was obtained via intramolecular nucleophilic reaction. Subsequently, the intermediate **III** was formed after the 1,3-migration and rearrangement. Simultaneously, the dioxide carbon was released. At last, the intermediate product **IV** was generated.^[16] Then, the external nucleophilic **B** attack of **A** occurs with concomitant decarboxylation to afford the product and regenerated Pd(0) (Scheme 3).^[10b]

Table 2 Substrate scope^a

^a Reaction conditions: **1** (0.70 mmol), Cs₂CO₃ (0.7 mmol) in MeCN (3 mL) at 80 °C until the starting material was fully consumed (monitored by TLC) to get intermediate. Then the reaction mixture was filtrated and transferred to a nitrogen-filled Schlenk tube, then PdCl₂ (2.5 mmol%) and XantPhos (5.0 mmol%) were added. The reaction mixture was stirred at 80 °C (or higher temperature) until intermediate was fully consumed (monitored by TLC) to get product.

Table 3 Substrate scope^a

^a Reaction conditions: **1** (0.70 mmol), Cs₂CO₃ (0.7 mmol) in MeCN (3 mL) at 80 °C until the starting material was fully consumed (monitored by TLC) to get intermediate. Then the reaction mixture was filtrated and transferred to a nitrogen-filled Schlenk tube, then PdCl₂ (2.5 mmol%) and XantPhos (5.0 mmol%) were added. The reaction mixture was stirred at 80 °C (or higher temperature) until intermediate was fully consumed (monitored by TLC) to get product. ^b 100 °C.

Scheme 2 Application of reaction

Scheme 3 Mechanism for allylamines

3 Conclusions

In conclusion, an efficient synthesis of allylic amines via palladium-catalyzed allylic amination that works through decarboxylation coupling of aryloxycarbamates, which generated *in situ* from inexpensive, readily available and nontoxic carboxylic acids and hydroxylamine, was newly developed. This reaction enables reliable access to a broad range of functional allylic amines. Furthermore, the products can be easily transformed into 3-methyl-indole that possess great potential in drug discovery. It is believed that this new method will possess wide application in organic synthesis and medicinal chemistry.

4 Experimental section

4.1 Instruments and reagents

^1H NMR spectra were recorded on a Bruker 400 MHz NMR spectrometer. ^{13}C NMR spectra were recorded on a 101 MHz NMR spectrometer. IR spectra were obtained on a Bruker ALPHA FT-IR spectrometer. Melting points were determined using an X-4 apparatus and uncorrected. GC analyses were performed on a Shimadzu GC-2014 gas chromatography equipped with a capillary column (HP-5 30 m $\times$ 0.25 μm) using a flame ionization detector. HRMS was performed on a TOF LC-MS in ESI mode. HPLC analyses were performed using a Shimadzu SPD-20A high

performance liquid chromatography. Rotation polarity: Krüss P8000. For product purification by flash column chromatography, silica gel (200~300 mesh) was used. All commercially available reagents were used without further purification. All solvents were purified and dried according to standard methods prior to use.

4.2 General procedure for the synthesis of allylamines

To a dried glass bottle (5 mL) were added allyl ether (0.7 mmol, 1.0 equiv.), Cs_2CO_3 (0.7 mmol, 1.0 equiv.) and acetonitrile (3 mL). The reaction mixture was allowed to heat at 80 °C until **1a** was fully consumed (monitored by TLC) to get intermediate. Then the reaction mixture was filtrated and transferred to a nitrogen-filled Schlenk tube (10 mL), then PdCl_2 (2.5 mol%) and Xantphos (5.0 mol%) were added.^[17] The reaction mixture was stirred at 80 °C (or higher temperature) until intermediate was fully consumed [monitored by thin layer chromatography (TLC)] to get **2a**. Then the reaction mixture was filtrated and purified carefully by flash chromatography (Hexane/EtOAc, $V:V=100:1$) to provide *N*-allyl-2-methylaniline (**2a**).^[18] Oil (0.082 g, 80% yield). $R_f=0.25$ (Hexane/EtOAc, $V:V=100:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 7.16~7.05 (m, 2H), 6.72~6.60 (m, 2H), 5.98~6.02 (m, 1H), 5.31 (dd, $J=17.2, 1.5$ Hz, 1H), 5.19 (dd, $J=10.3, 1.3$ Hz, 1H), 3.84 (d, $J=4.6$ Hz, 2H), 3.63 (s, 1H), 2.17 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.93, 135.56, 130.05, 127.07, 121.95, 117.09, 116.18, 110.01, 46.49, 17.43.

N-Allyl-2-methoxyaniline (**2b**).^[18] Oil (0.091 g, 80% yield). $R_f=0.5$ (Hexane/EtOAc, $V:V=80:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 6.86 (dd, $J=7.6, 1.3$ Hz, 1H), 6.79 (dd, $J=7.9, 1.2$ Hz, 1H), 6.69 (dd, $J=7.7, 1.4$ Hz, 1H), 6.62 (dd, $J=7.8, 1.4$ Hz, 1H), 6.05~5.94 (m, 1H), 5.34~5.26 (m, 1H), 5.16~5.19 (m, 1H), 4.37 (s, 1H), 3.86 (s, 3H), 3.81 (d, $J=4.9$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 146.84, 138.00, 135.57, 121.22, 116.56, 116.01, 110.15, 109.42, 55.40, 46.28.

N-Allylaniline (**2c**).^[19] Oil (0.037 g, 40% yield). $R_f=0.3$ (Hexane/EtOAc, $V:V=100:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 7.18 (t, $J=7.9$ Hz, 2H), 6.72 (t, $J=7.3$ Hz, 1H), 6.63 (d, $J=7.8$ Hz, 2H), 5.93~6.00 (m, 1H), 5.29 (dd, $J=17.2, 1.0$ Hz, 1H), 5.17 (d, $J=10.3$ Hz, 1H), 3.78 (d, $J=5.1$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 148.02, 135.43, 129.18, 117.49, 116.18, 112.95, 46.52.

N-Allyl-2-fluoroaniline (**2d**).^[20] Oil (0.049 g, 46% yield). $R_f=0.3$ (Hexane, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 7.03~6.92 (m, 2H), 6.73~6.58 (m, 2H), 5.96~6.01 (m, 1H), 5.36~5.26 (m, 1H), 5.23~5.13 (m, 1H), 4.05 (s, 1H), 3.81 (dd, $J=11.6, 6.2$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.77 (s), 150.40 (s), 136.44 (s), 135.00 (s), 124.52 (d, $J=3.5$ Hz), 116.72 (d, $J=7.0$ Hz), 116.41 (s), 114.38 (d, $J=18.4$ Hz), 112.36 (d, $J=3.3$ Hz), 46.09 (s); ^{19}F NMR (377 MHz, CDCl_3) δ : -122.72 (s), -136.68 (s).

N-Allyl-2-bromoaniline (**2e**).^[21] Oil (0.101 g, 46% yield). $R_f=0.3$ (Hexane, add 1% Et_3N); ^1H NMR (400 MHz,

CDCl_3) δ : 7.43 (dd, $J=7.9, 1.5$ Hz, 1H), 7.21~7.11 (m, 1H), 6.68~6.52 (m, 2H), 5.93~5.97 (m, 1H), 5.28~5.32 (m, 1H), 5.19~5.22 (m, 1H), 4.48 (s, 1H), 3.99~3.75 (m, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 144.74, 134.67, 132.36, 128.39, 117.84, 116.40, 111.58, 109.72, 46.22.

N-Allyl-2-iodoaniline (**2f**).^[22] Oil (0.056 g, 31% yield). $R_f=0.5$ (Hexane/EtOAc, $V:V=30:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 7.67 (d, $J=7.8$ Hz, 1H), 7.20 (t, $J=7.7$ Hz, 1H), 6.57 (d, $J=8.2$ Hz, 1H), 6.45 (t, $J=7.5$ Hz, 1H), 5.93~5.97 (m, 1H), 5.30 (d, $J=17.2$ Hz, 1H), 5.21 (d, $J=10.3$ Hz, 1H), 4.33 (s, 1H), 3.84 (t, $J=5.3$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 147.02, 139.00, 134.61, 129.35, 118.74, 116.43, 110.92, 85.38, 46.57.

N-Allyl-2-(trifluoromethyl)aniline (**2g**).^[23] Oil (0.087 g, 62% yield). $R_f=0.3$ (Hexane, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 7.44 (d, $J=7.3$ Hz, 1H), 7.35 (t, $J=7.8$ Hz, 1H), 6.72 (t, $J=7.6$ Hz, 2H), 5.91~5.98 (m, 1H), 5.30~5.32 (m, 1H), 5.19~5.22 (m, 1H), 4.51 (s, 1H), 3.90~3.82 (m, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.39, 134.31, 133.00, 126.59, 126.54, 123.85, 116.56, 116.01, 113.58, 113.29, 112.08, 45.84; ^{13}C NMR (101 MHz, CDCl_3) δ : 145.39 (s), 134.31 (s), 133.00 (s), 126.57 (d, $J=5.5$ Hz), 123.85 (s), 116.56 (s), 116.01 (s), 113.58 (s), 113.29 (s), 112.08 (s), 45.84 (s); ^{19}F NMR (377 MHz, CDCl_3) δ : -62.49 (s).

N-Allyl-2-nitroaniline (**2h**).^[24] Oil (0.060 g, 48% yield). $R_f=0.5$ (Hexane/EtOAc, $V:V=80:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 8.21~8.17 (m, 1H), 7.41~7.43 (m, 1H), 6.83 (d, $J=8.7$ Hz, 1H), 6.66~6.67 (m, 1H), 5.92~5.96 (m, 1H), 5.36~5.22 (m, 2H), 4.03~3.95 (m, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.23, 136.02, 133.13, 132.02, 126.76, 116.98, 115.42, 113.99, 45.18.

N-Allyl-3-nitroaniline (**2i**): Yellowish-brown solid (0.062 g, 50% yield). m.p. 60~61 °C (lit.^[25] m.p. 61~61 °C); $R_f=0.5$ (Hexane/EtOAc, $V:V=80:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 7.52 (dd, $J=8.1, 2.1$ Hz, 1H), 7.40 (t, $J=2.2$ Hz, 1H), 7.28 (d, $J=8.1$ Hz, 1H), 6.88 (dd, $J=8.2, 2.4$ Hz, 1H), 5.90~5.97 (m, 1H), 5.21~5.33 (m, 2H), 4.18 (s, 1H), 3.84 (t, $J=5.2$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 148.72, 134.05, 129.66, 118.82, 116.98, 112.07, 106.55, 46.17.

N-Allyl-4-methoxyaniline (**2j**).^[26] Oil (0.042 g, 37% yield). $R_f=0.5$ (Hexane/EtOAc, $V:V=100:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 6.84~6.75 (m, 2H), 6.61 (t, $J=6.0$ Hz, 2H), 5.26~5.30 (m, 1H), 5.28 (dd, $J=17.2, 1.4$ Hz, 1H), 5.16 (dd, $J=10.3, 1.1$ Hz, 1H), 3.78~3.70 (m, 5H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.21, 142.28, 135.79, 116.11, 114.86, 114.30, 55.79, 47.54.

N-Allyl-4-nitroaniline (**2k**).^[19,25] Yellowish-brown solid (0.064 g, 51% yield). m.p. 66~67 °C (lit.^[25] 65~66 °C); $R_f=0.3$ (Hexane/EtOAc, $V:V=9:1$, add 1% Et_3N); ^1H NMR (400 MHz, CDCl_3) δ : 8.12~8.05 (m, 2H), 6.61~6.49 (m, 2H), 5.88~5.92 (m, 1H), 5.38~5.21 (m, 2H), 4.69 (s, 1H), 3.94~3.82 (m, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 153.10, 138.25, 133.35, 126.33, 117.32, 111.28, 45.76.

N-Allyl-2-bromo-3-methylaniline (**2l**): Oil (0.106 g, 67% yield). $R_f=0.3$ (Hexane/EtOAc, $V:V=100:1$, add 1%

Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.07 (t, *J*=7.8 Hz, 1H), 6.61 (d, *J*=7.1 Hz, 1H), 6.49 (d, *J*=8.1 Hz, 1H), 5.94~5.98 (m, 1H), 5.30 (dd, *J*=17.2, 1.5 Hz, 1H), 5.20 (dd, *J*=10.3, 1.4 Hz, 1H), 4.59 (s, 1H), 3.90~3.79 (m, 2H), 2.39 (s, 3H); ¹³C NMR (176 MHz, CDCl₃) δ: 144.88, 138.38, 134.82, 127.52, 119.05, 116.29, 112.41, 108.97, 46.45, 23.76; IR (film) ν: 3420, 1592, 1498, 1470, 1320, 1019 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₀H₁₂NBr [M+H]⁺ 228.0166, found 228.0194.

N-Allyl-2-bromo-4-chloroaniline (**2m**): Oil (0.114 g, 66% yield). *R*_f=0.3 (Hexane/EtOAc, *V*:*V*=80:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.42 (s, 1H), 7.13 (d, *J*=8.7 Hz, 1H), 6.53 (d, *J*=8.7 Hz, 1H), 6.01~5.84 (m, 1H), 5.24 (dd, *J*=27.6, 13.7 Hz, 2H), 4.46 (s, 1H), 3.81 (d, *J*=5.2 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 143.50, 134.20, 131.68, 128.28, 121.54, 116.64, 112.00, 109.44, 46.27; IR (film) ν: 3420, 1595, 1505, 1317 cm⁻¹; HRMS (ESI-TOF) calcd for C₉H₉NBrCl [M+H]⁺ 247.9620, found 247.9646.

N-Allyl-2-bromo-4-fluoroaniline (**2n**): Oil (0.106 g, 67% yield). *R*_f=0.3 (Hexane/EtOAc, *V*:*V*=100:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.26~7.16 (m, 1H), 7.07~6.83 (m, 1H), 6.56~6.58 (m, 1H), 5.93~6.00 (m, 1H), 5.40~5.11 (m, 2H), 4.27 (s, 1H), 3.79 (d, *J*=5.1 Hz, 2H); ¹³C NMR (176 MHz, CDCl₃) δ: 155.18 (s), 153.82 (s), 141.53 (d, *J*=2.0 Hz), 134.57 (s), 119.44 (s), 119.30 (s), 116.53 (s), 114.98 (s), 114.85 (s), 111.56 (d, *J*=7.7 Hz), 108.81 (d, *J*=9.7 Hz), 46.71 (s); ¹⁹F NMR (377 MHz, CDCl₃) δ: -127.17 (s); IR (film) ν: 3428, 1513, 1317, 1266, 1188 cm⁻¹; HRMS (ESI-TOF) calcd for C₉H₉NFBr [M+H]⁺ 229.9936, found 229.9962.

N-Allyl-2-bromo-4-nitroaniline (**2o**): Yellow solid (0.079 g, 44% yield). m.p. 65~66 °C; *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=50:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 8.38 (d, *J*=2.6 Hz, 1H), 8.14~8.05 (m, 1H), 6.58 (d, *J*=9.2 Hz, 1H), 5.99~5.86 (m, 1H), 5.35~5.20 (m, 3H), 4.02~3.90 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 149.61, 132.64, 128.74, 125.24, 117.58, 109.23, 107.77, 45.90; IR (film) ν: 3390, 1590, 1527, 1525, 1323, 1300 cm⁻¹; HRMS (ESI-TOF) calcd for C₉H₉N₂BrO₂ [M+H]⁺ 258.9860, found 258.9890.

N-Allyl-2-bromo-5-methylaniline (**2p**):^[27] Oil (0.125 g, 79% yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=100:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.29 (d, *J*=8.0 Hz, 1H), 6.49~6.37 (m, 2H), 5.94~5.99 (m, 1H), 5.30 (d, *J*=17.2 Hz, 1H), 5.20 (d, *J*=10.3 Hz, 1H), 4.40 (s, 1H), 3.83 (t, *J*=4.9 Hz, 2H), 2.26 (s, 3H); ¹³C NMR (176 MHz, CDCl₃) δ: 144.41, 138.40, 134.73, 131.96, 118.78, 116.34, 112.36, 106.54, 46.22, 21.47.

N-Allyl-2-bromo-5-methoxyaniline (**2q**): Oil (0.134 g, 79% yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=100:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.30 (d, *J*=8.6 Hz, 1H), 6.19 (dd, *J*=13.9, 5.7 Hz, 2H), 5.92~5.99 (m, 1H), 5.30 (dd, *J*=17.2, 1.4 Hz, 1H), 5.21 (dd, *J*=10.3, 1.3 Hz, 1H), 4.45 (s, 1H), 3.88~3.73 (m, 5H); ¹³C NMR (176 MHz, CDCl₃) δ: 160.19, 145.52, 134.52, 132.43, 116.49, 102.71, 100.92, 98.37, 55.29, 46.22; IR (film) ν: 3418,

1600, 1509, 1313, 1213 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₀H₁₂NBrO [M+H]⁺ 244.0115, found 244.0135.

N-Allyl-2-bromo-5-chloroaniline (**2r**): Oil (0.123 g, 71% yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=80:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.32 (d, *J*=8.3 Hz, 1H), 6.64~6.51 (m, 2H), 6.02~5.87 (m, 1H), 5.26 (dd, *J*=27.9, 13.7 Hz, 2H), 4.54 (s, 1H), 3.82 (t, *J*=5.4 Hz, 2H); ¹³C NMR (176 MHz, CDCl₃) δ: 145.52, 134.30, 133.86, 132.89, 117.54, 116.81, 111.37, 107.32, 46.03; IR (film) ν: 3415, 1590, 1505, 1415, 1095, 1017 cm⁻¹; HRMS (ESI-TOF) calcd for C₉H₉BrClN [M+H]⁺ 247.9620, found 247.9651.

N-Allyl-2-bromo-5-fluoroaniline (**2s**): Oil (0.139 g, 86% yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=50:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.34 (dd, *J*=8.6, 6.0 Hz, 1H), 6.30~6.35 (m, 2H), 6.02~5.84 (m, 1H), 5.21~5.32 (m, 2H), 4.58 (s, 1H), 3.81 (dd, *J*=7.3, 3.6 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 164.53 (s), 162.11 (s), 146.06 (d, *J*=11.4 Hz), 133.95 (s), 132.81 (d, *J*=9.9 Hz), 116.78 (s), 104.44 (s), 104.21 (s), 103.59 (d, *J*=2.8 Hz), 99.05 (s), 98.78 (s), 46.12 (s); ¹⁹F NMR (377 MHz, CDCl₃) δ: -113.13 (s); IR (film) ν: 3420, 1611, 1510, 1435, 1305, 1175, 1023 cm⁻¹; HRMS (ESI-TOF) calcd for C₉H₉BrFN [M+H]⁺ 231.9915, found 231.9958.

(*E*)-*N*-Cinnamyl-2-methylaniline (**2aa**):^[28] Oil (0.0754 g, 86% yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=100:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.29 (d, *J*=7.4 Hz, 2H), 7.22 (t, *J*=7.5 Hz, 2H), 7.14 (s, 1H), 7.01 (d, *J*=26.0 Hz, 2H), 6.58 (d, *J*=7.7 Hz, 2H), 6.55 (d, *J*=16.0 Hz, 1H), 6.30 (s, 1H), 3.89 (dd, *J*=5.8, 1.1 Hz, 2H), 3.59 (s, 1H), 2.08 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 145.94, 136.85, 131.60, 130.09, 128.55, 127.53, 127.15, 127.13, 126.33, 122.04, 117.19, 110.07, 46.18, 17.49; HRMS (ESI-TOF) calcd for C₁₆H₁₇N [M+H]⁺ 224.1395, found 224.1274.

N-(But-3-en-2-yl)-2-methylaniline (**2ab**) and (*E*)-*N*-(but-2-en-1-yl)-2-methylaniline (**2ac**):^[29] Oil (45% or 86% total yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=100:1, add 1% Et₃N). **2ab**: ¹H NMR (400 MHz, CDCl₃) δ: 7.16~7.03 (m, 2H), 6.64 (dd, *J*=13.2, 7.9 Hz, 2H), 5.94~5.90 (m, 1H), 5.17 (dd, *J*=48.9, 13.8 Hz, 2H), 4.09~3.99 (m, 1H), 3.47 (s, 1H), 2.17 (s, 3H), 1.37 (d, *J*=6.6 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 145.28, 141.40, 130.08, 126.95, 121.65, 116.76, 113.96, 110.80, 50.85, 21.82, 17.52; IR (film) ν: 3445, 2920, 1607, 1513, 1450, 1317, 1258 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₁H₁₅N [M+H]⁺ 162.1238, found 162.1274. **2ac**: ¹H NMR (400 MHz, CDCl₃) δ: 7.18~7.09 (m, 1H), 7.06 (d, *J*=7.2 Hz, 1H), 6.71~6.59 (m, 2H), 5.80~5.57 (m, 2H), 3.78 (dd, *J*=33.6, 6.0 Hz, 2H), 3.61 (d, *J*=52.9 Hz, 1H), 2.15 (s, 3H), 1.73 (dt, *J*=6.2, 3.1 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 146.11, 129.99, 128.12, 127.95, 127.05, 121.93, 116.91, 109.91, 45.97, 17.78, 17.46; IR (film) ν: 3445, 2977, 1607, 1511, 1315, 1258 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₁H₁₅N [M+H]⁺ 162.1238, found 162.1274.

2-Methyl-*N*-(2-methylallyl)aniline (**2ad**): Oil (0.0610 g, 63% yield). *R*_f=0.5 (Hexane/EtOAc, *V*:*V*=100:1, add

1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.14~7.01 (m, 2H), 6.65 (t, *J*=7.4 Hz, 1H), 6.57 (d, *J*=8.0 Hz, 1H), 4.94 (d, *J*=28.5 Hz, 2H), 3.74 (s, 3H), 2.17 (s, 3H), 1.81 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 146.09, 142.78, 129.99, 127.05, 121.70, 116.89, 110.80, 109.94, 49.83, 20.48, 17.44; IR (film) ν: 3445, 2920, 1607, 1513, 1450, 1317, 1258 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₁H₁₅N [M+H]⁺ 162.1238, found 162.1276.

(*E*)-*N*-cinnamyl-2-methoxyaniline (**2ae**):^[30] Oil (0.1390 g, 83% yield). *R*_f=0.5 (Hexane/EtOAc, *V*: *V*=50:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.38 (d, *J*=7.4 Hz, 2H), 7.31 (s, 2H), 7.23 (t, *J*=7.2 Hz, 1H), 6.87 (dd, *J*=7.6, 1.1 Hz, 1H), 6.80 (d, *J*=7.8 Hz, 1H), 6.72~6.60 (m, 3H), 6.19~6.40 (m, 1H), 4.44 (s, 1H), 3.99~3.94 (m, 2H), 3.87 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 146.89, 137.98, 136.96, 131.35, 128.53, 127.43, 127.24, 126.32, 121.29, 116.69, 110.23, 109.42, 55.41, 45.93, 30.31; IR (film) ν: 3436, 1513, 1456, 1246, 1221, 1127, 1029, 736 cm⁻¹.

(*E*)-2-Bromo-*N*-cinnamyl-5-methoxyaniline (**2af**): Light yellow solid (0.1671 g, 75% yield). m.p. 86.3~87.5 °C; *R*_f=0.3 (Hexane/EtOAc, *V*: *V*=50:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.39 (d, *J*=7.3 Hz, 2H), 7.35~7.29 (m, 3H), 7.25 (d, *J*=6.3 Hz, 1H), 6.64 (d, *J*=15.9 Hz, 1H), 6.29 (dd, *J*=16.1, 9.4 Hz, 2H), 6.18 (dd, *J*=8.7, 2.8 Hz, 1H), 4.52 (s, 1H), 3.98 (s, 2H), 3.76 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 160.26, 145.54, 136.66, 132.50, 131.86, 128.57, 127.63, 126.38, 126.08, 102.83, 101.00, 98.47, 55.32, 45.95; IR (film) ν: 3405, 1598, 1596, 1511, 1315, 1205, 1172 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₆H₁₆BrNO [M+H]⁺ 218.0449, found 218.0475.

(*E*)-2-Bromo-*N*-cinnamylaniline (**2ag**): Oil (0.1493 g, 74% yield). *R*_f=0.3 (Hexane/EtOAc, *V*: *V*=50:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.46 (dd, *J*=7.9, 1.3 Hz, 1H), 7.40 (d, *J*=7.3 Hz, 2H), 7.34 (t, *J*=7.5 Hz, 2H), 7.27 (t, *J*=3.5 Hz, 1H), 7.19 (s, 1H), 6.72 (dd, *J*=8.1, 0.9 Hz, 1H), 6.64 (t, *J*=11.9 Hz, 2H), 6.34 (d, *J*=15.9 Hz, 1H), 4.58 (s, 1H), 4.02 (d, *J*=5.1 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 144.73, 136.67, 132.39, 131.73, 128.57, 128.48, 127.61, 126.36, 126.24, 117.94, 111.63, 109.77, 45.94; IR (film) ν: 3407, 1598, 1507, 1321, 1019, 966, 742 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₅H₁₄BrN [M+H]⁺ 290.0323, found 290.0356.

(*E*)-2-Bromo-5-chloro-*N*-cinnamylaniline (**2ah**):^[31] Light yellow solid (0.1536 g, 68% yield). m.p. 56.6~57.6 °C; *R*_f=0.5 (Hexane/EtOAc, *V*: *V*=50:1, add 1% Et₃N); ¹H NMR (400 MHz, CDCl₃) δ: 7.40 (d, *J*=7.3 Hz, 2H), 7.33 (dd, *J*=8.0, 6.5 Hz, 3H), 7.27 (d, *J*=7.3 Hz, 1H), 6.64 (t, *J*=8.6 Hz, 2H), 6.57 (dd, *J*=8.4, 2.3 Hz, 1H), 6.29 (d, *J*=15.9 Hz, 1H), 4.60 (s, 1H), 3.98 (dd, *J*=5.6, 4.5 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 145.54, 136.48, 134.39, 132.95, 132.19, 128.61, 127.77, 126.42, 125.32, 117.68, 111.39, 107.41, 45.78; IR (film) ν: 3407, 1590, 1505, 1445, 1360, 1299, 1017, 736 cm⁻¹.

3-Methyl-1*H*-indole (**3b**): White solid (0.0450 g, 72% yield). m.p. 92~93 °C (lit.^[21] 92~93 °C), *R*_f=0.5 (Hexane/EtOAc, *V*: *V*=10:1); ¹H NMR (400 MHz,

CDCl₃) δ: 7.85 (s, 1H), 7.60 (d, *J*=7.8 Hz, 1H), 7.35 (d, *J*=8.0 Hz, 1H), 7.21 (s, 1H), 7.17~7.10 (m, 1H), 6.98 (d, *J*=1.0 Hz, 1H), 2.36 (d, *J*=1.0 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 136.22, 128.25, 121.83, 121.53, 119.08, 118.79, 111.68, 110.90, 9.63.

Supporting Information ¹H NMR and ¹³C NMR spectra of compounds **2a**~**2s** and **2aa**~**2ah**, HRMS spectra of some materials and products, ¹⁹F NMR spectra of compounds **2d**, **2g**, **2n**, **2s**, the results of some reaction system NMR and experimental procedures of raw materials. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Johannsen, M.; Jørgensen, K. A. *Chem. Rev.* **1998**, *98*, 1689.
(b) Sweeney, J. B.; Ball, A. K.; Lawrence, P. A.; Sinclair, M. C.; Smith, L. J. *Angew. Chem., Int. Ed.* **2018**, *57*, 10202.
(c) Wang, D.; Yu, X.; Ge, B.; Miao, H.; Ding, Y. *Chin. J. Org. Chem.* **2015**, *35*, 676 (in Chinese).
(d) Hu, X.; Yang, B.; Yao, W.; Wang, D. *Chin. J. Org. Chem.* **2018**, *38*, 3296 (in Chinese).
(e) 胡昕宇, 杨伯斌, 姚玮, 王大伟, 有机化学, **2018**, *38*, 3296.
- [2] (a) Shekhar, S.; Trantow, B.; Leitner, A. J. *Am. Chem. Soc.* **2006**, *128*, 11770.
(b) Trost, B. M.; Malhotra, S.; Olson, D. E.; Maruniak, A.; Du Bois, J. J. *Am. Chem. Soc.* **2009**, *131*, 4190.
(c) Uozumi, Y.; Tanaka, H.; Shibatomi, K. *Org. Lett.* **2004**, *6*, 281.
(d) Wang, X.; Guo, P.; Han, Z.; Wang, X.; Wang, Z.; Ding, K. J. *Am. Chem. Soc.* **2014**, *136*, 405.
(e) You, S. L.; Zhu, X. Z.; Luo, Y. M.; Hou, X. L.; Dai, L. X. *J. Am. Chem. Soc.* **2001**, *123*, 7471.
(f) Shi, Y.; Wu, H.; Huang, G. *Org. Chem. Front.* **2021**, *8*, 3320.
- [3] (a) Hirata, G.; Satomura, H.; Kumagai, H.; Shimizu, A.; Onodera, G.; Kimura, M. *Org. Lett.* **2017**, *19*, 6148.
(b) Jing, J.; Huo, X.; Shen, J.; Fu, J.; Meng, Q.; Zhang, W. *Chem. Commun.* **2017**, *53*, 5151.
(c) Kleij, A. W.; Guo, W.; Cai, A.; Xie, J. *Angew. Chem., Int. Ed.* **2017**, *56*, 11797.
(d) Ozawa, F.; Okamoto, H.; Kawagishi, S.; Yamamoto, S.; Minami, T.; Yoshifuji, M. *J. Am. Chem. Soc.* **2002**, *124*, 10968.
(e) Wang, X.; Wang, X.; Han, Z.; Wang, Z.; Ding, K. *Org. Chem. Front.* **2017**, *4*, 271.
- [4] Mukherjee, P.; Widenhoefer, R. A. *Org. Lett.* **2010**, *41*, 1184.
- [5] (a) Lafrance, M.; Roggen, M.; Carreira, P. D. E. M. *Angew. Chem., Int. Ed.* **2012**, *51*, 3470.
(b) Roggen, M.; Carreira, R. M. *J. Am. Chem. Soc.* **2010**, *132*, 11917.
- [6] (a) Kita, Y.; Sakaguchi, H.; Hoshimoto, Y.; Nakauchi, D.; Nakahara, Y.; Carpentier, J.-F.; Ogoshi, S.; Mashima, K. *Chem.-Eur. J.* **2015**, *21*, 14571.
(b) Sweeney, J. B.; Anthony, B.; Philippa, L.; Mackenzie, S.; Luke, S. *Angew. Chem., Int. Ed.* **2018**, *57*, 10202.
- [7] Sakuramoto, T.; Hirao, T.; Tobisu, M.; Moriuchi, T. *ChemCatChem* **2019**, *11*, 1175.
- [8] Emayavaramban, B.; Roy, M.; Sundararaju, B. *Chem.-Eur. J.* **2016**, *22*, 3952.
- [9] (a) Patel, S. J.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2004**, *43*, 3941.
(b) Zhou, C. Y.; Zhu, S. F.; Wang, L. X.; Zhou, Q. L. *J. Am. Chem. Soc.* **2010**, *132*, 10955.
(c) Holmes, M.; Schwartz, L. A.; Krische, M. J. *Chem. Rev.* **2018**,

- 118, 6026.
- (d) Liu, L.; Chen, Y.; Zhang, A.; Liu, X.; Zhang, L.; Bai, J.; Li, H.; Mao, G. *Chin. J. Org. Chem.* **2019**, *39*, 475 (in Chinese). (刘澜涛, 陈莹莹, 张安安, 刘雪, 张丽, 白静茹, 李恒, 毛国梁, 有机化学, **2019**, *39*, 475.)
- (e) Banerjee, D.; Junge, K.; Beller, M. *Org. Chem. Front.* **2014**, *1*, 368.
- [10] (a) Doekal, V.; Imek, M.; Draínsk, M.; Vesel, J. *Chem.-Eur. J.* **2018**, *24*, 13441.
- (b) Kondoh, A.; Kamata, Y.; Terada, M. *Org. Lett.* **2017**, *19*, 1682.
- (c) Mellegaard-Waetzig, S. R.; Rayabarapu, D. K.; Tunge, J. A. *Synlett* **2005**, 2759.
- [11] (a) Li, J.; Yang, S.; Wu, W.; Jiang, H. *Chem.-Asian. J.* **2019**, *14*, 4114.
- (b) Ngamnithiporn, A.; Iwayama, T.; Bartberger, M.; Stoltz, B. M. *Chem. Sci.* **2020**, *11*, 11068.
- (c) Zhou, Q.; Zhang, B.; Chen, D.; Chen, R.; Jiang, H. *Chin. J. Org. Chem.* **2011**, *31*, 2181 (in Chinese). (周其忠, 张斌, 陈丹, 陈仁尔, 蒋华江, 有机化学, **2011**, *31*, 2181.)
- (d) Dai, J.; Wang, G.; Xu, X.; Xu, H. *Chin. J. Org. Chem.* **2013**, *33*, 2460 (in Chinese). (戴建军, 王光祖, 徐小岚, 许华建, 有机化学, **2013**, *33*, 2460.)
- [12] (a) Li, J.; Yang, S.; Wu, W.; Jiang, H. *Org. Chem. Front.* **2020**, *7*, 1395.
- (b) Mellegaard-Waetzig, S. R.; Rayabarapu, D. K.; Tunge, J. A. *Synlett* **2005**, 2759.
- [13] (a) Dai, Q.; Li, P.; Ma, N.; Hu, C. *Org. Lett.* **2016**, *18*, 5560.
- (b) Li, P.; Ma, N.; Wang, Z.; Dai, Q.; Hu, C. *W. J. Org. Chem.* **2018**, *83*, 8233.
- [14] Thoumazet, C.; Grützmacher, H.; Deschamps, B.; Ricard, L.; Floch, P. L. *Eur. J. Inorg. Chem.* **2006**, 3911.
- [15] Jensen, T.; Pedersen, H.; Bang-Andersen, B.; Madsen, R.; Jrgensen, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 888.
- [16] Fu, H.; Li, P.; Wang, Z.; Li, X.; Hu, C. *Org. Biomol. Chem.* **2020**, *18*, 4439.
- [17] Alexy, E. J.; Zhang, H.; Stoltz, B. M. *J. Am. Chem. Soc.* **2018**, *140*, 10109.
- [18] Jing, J.; Huo, X.; Shen, J.; Fu, J.; Meng, Q.; Zhang, W. *Chem. Commun.* **2017**, *53*, 5151.
- [19] Tao, Y.; Wang, B.; Wang, B.; Qu, L.; Qu, J. *Org. Lett.* **2010**, *12*, 2726.
- [20] Chanda, K.; ReJ, S.; Liu, S.-Y.; Huang, M. H. *ChemCatChem* **2015**, *7*, 1813.
- [21] Jensen, T.; Pedersen, H.; Bang-Andersen, B.; Madsen, R.; Jorgensen, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 888.
- [22] Chen, Y.-Z.; Jiang, H.-L. *Chem. Mater.* **2016**, *28*, 6698.
- [23] Lam, J. K.; Schmidt, Y.; Vanderwal, C. D. *Org. Lett.* **2012**, *14*, 5566.
- [24] Kommi, D. N.; Jadhavar, P. S.; Kumar, D.; Chakraborti, A. K. *Green Chem.* **2013**, *15*, 798.
- [25] Du, Z.; Yan, Y.; Fu, Y.; Wang, K. *Asian J. Org. Chem.* **2016**, *5*, 812.
- [26] Sweeney, J. B.; Ball, A. K.; Lawrence, P. A.; Sinclair, M. C.; Smith, L. J. *Angew. Chem., Int. Ed.* **2018**, *57*, 10202.
- [27] Repka, L. M.; Ni, R.; Reisman, R. E. *J. Am. Chem. Soc.* **2010**, *132*, 14418.
- [28] Wang, D.; Peng, X.; Wang, G.; Zhang, Y.; Guo, T.; Zhang, P. *Asian J. Org. Chem.* **2018**, *7*, 875.
- [29] Yang, S. C.; Tsai, Y. C. *Organometallics* **2001**, *20*, 763.
- [30] Everett, R. K.; Wolfe, J. P. *J. Org. Chem.* **2015**, *80*, 9041.
- [31] Yang, S. C.; Hsu, Y. C.; Gan, K. H. *Tetrahedron* **2006**, *62*, 3949.

(Zhao, C.)