5-苯基-1,3,4-噻二唑衍生物的合成及含 SH2 结构域 蛋白酪氨酸磷酸酶 1 (SHP1)抑制活性研究

于丽杰^{†,a} 冯 勃^{†,b,c} 王智佳^a 高立信^{a,b} 张 纯^{a,d}
Rajendran, Satheeshkumar^a 李 佳^b 周宇波^{*,b,c} 王文龙^{*,a}

(^a 江南大学药学院 江苏无锡 214122)

(^b 中国科学院上海药物研究所 国家新药筛选中心 上海 201203)

(^c 中国科学院药物创新研究院中山研究院 广东中山 528400)

(^d 南京大学化学化工学院 南京 210023)

摘要 作为细胞信号转导通路的关键节点分子,含 SH2 结构域蛋白酪氨酸磷酸酶 1 (SHP1)是潜在的抗肿瘤靶点. 已知的 SHP1 抑制剂屈指可数. 设计并合成了 11 个 5-苯基-1,3,4-噻二唑衍生物. 活性测试结果表明,部分衍生物对 SHP1 显示了一定强度的抑制活性. 其中,化合物 **5b** [$IC_{50} = (1.33 \pm 0.16) \mu\text{mol/L}$]对 SHP1 显示了中等强度的抑制活性,对 PTP1B 和 TCPTP 不显示抑制活性,对 SHP2 显示了 2 倍的选择性,为发现新型 SHP1 抑制剂提供了新的骨架类型.

关键词 5-苯基-1,3,4-噻二唑衍生物; 含 SH2 结构域蛋白酪氨酸磷酸酶 1 (SHP1); 抑制剂; 构效关系

Synthesis of 5-Phenyl-1,3,4-thiadiazole Derivatives and Their Biochemical Evaluation against Src Homology 2 Domain-Containing Protein Tyrosine Phosphatase 1 (SHP1)

Yu, Lijie^{†,a} Feng, Bo^{†,b,c} Wang, Zhijia^a Gao, Lixin^{a,b} Zhang, Chun^{a,d}
Satheeshkumar, Rajendran^a Li, Jia^b Zhou, Yubo^{*,b,c} Wang, Wenlong^{*,a}

(^a School of Pharmaceutical Sciences, Jiangnan University, Wuxi, Jiangsu 214122)

(^b National Center for Drug Screening, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203)

(^c Zhongshan Institute for Drug Discovery, Institutes of Drug Discovery and Development, Chinese Academy of Sciences, Zhongshan, Guangdong 528400)

(^d School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210023)

Abstract The Src homology 2 domain-containing protein tyrosine phosphatase 1 (SHP1) is a convergent node for oncogenic cell-signaling cascades. Consequently, SHP1 represents a potential target for drug development in cancer treatment. Meanwhile only countable SHP1 inhibitors have been reported. A new type of SHP1 inhibitors with 5-phenyl-1,3,4-thiadiazole scaffold was developed. The representative compound **5b** exhibited SHP1 inhibitory activity with IC_{50} of $(1.33 \pm 0.16) \mu\text{mol/L}$, exhibited about 2-fold selectivity for SHP1 over SHP2, and had no detectable activity against PTP1B and TCPTP, and offered a novel scaffold to develop new SHP1 inhibitors.

Keywords 5-phenyl-1,3,4-thiadiazole derivatives; Src homology 2 domain-containing protein tyrosine phosphatase 1 (SHP1); inhibitors; structure-activity relationships

* Corresponding authors. E-mail: wwenlong2011@163.com; ybzhou@simmm.ac.cn

Received April 18, 2021; revised May 19, 2021; published online May 28, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21772068, 81773779), the Postgraduate Research & Practice Innovation Program of Jiangsu Province (No. KYCX20-1965), and the Natural Science Foundation of Jiangsu Province (No. BK20190608).

国家自然科学基金(Nos. 21772068, 81773779)、江苏省研究生科研与实践创新计划(No. KYCX20-1965)和江苏省自然科学基金(No. BK20190608)资助项目.

[†] 共同第一作者(These authors contributed equally to this work).

1 Introduction

Reversible protein tyrosine phosphorylation, catalyzed by the opposing actions of protein tyrosine kinases (PTKs) and protein tyrosine phosphatases (PTPs), is a key regulatory mechanism in eukaryotic cell physiology.^[1] Dysregulation of PTKs and PTPs is highly correlated with numerous human diseases, including diabetes, obesity, infection, autoimmune, and neuropsychiatric disorders and cancer.^[2-3] Two important non-receptor PTPs, SHP1 and Src homology 2 (SH2) domain-containing protein tyrosine phosphatase 2 (SHP2), have high similarity in basic structure and work as key hubs to connect several intracellular oncogenic signaling pathways, such as Jak/STAT, PI3K/AKT, and PD-1/PD-L1.^[4-5] Malfunctions of SHP1 and SHP2 have been associated with hematological malignancies and solid tumors, and viewed as valuable targets for cancer therapy.^[6] Compared to significant progress in identifying chemotypes of SHP2 inhibitors,^[7] developing potent and selective SHP1 inhibitors is unsatisfied.^[8] Only countable SHP1 inhibitors have been reported, including NSC-87877,^[9] tyrosine phosphatase inhibitor 1 (TPI-1),^[10] α -bromo-4'-hydroxyacetophenone (PI-1),^[11] sodium stibogluconate (SSG),^[12] oxovanadium complex,^[13] copper complexes,^[14] and luminescent Tb(III) complex^[15] (Figure 1), and their potency and selectivity are disappointed. Therefore, the imminent development of potent and specific SHP1 inhibitor remains necessary.

The 1,2,4-thiadiazole scaffold represents an important class of pharmacophores in medicinal chemistry due to its myriad bioactivities^[16] and 1,2,4-thiadiazole derivatives are well documented with antifungal, anti-inflammatory, antibacterial, antioxidant, anticancer activities.^[17] Our efforts to develop modulators of protein tyrosine phosphatases (PTPs) started from 1*H*-2,3-dihydroperimidines.^[18-23] Using the scaffold hopping strategy, we developed 3-aryl-1-oxa-2,8-diazaspiro[4.5]dec-2-enes as PTP1B inhibitors,^[19] bis-

aryl amides^[20] and benzo[*c*][1,2,5]thiadiazoles as SHP2 inhibitors.^[21] To explore novel scaffolds for modulators of PTPs, we designed and synthesized a series of 1,3,4-thiadiazole derivatives (Figure 2), evaluated their inhibitory activities against SHP1. Among them, several 1,3,4-thiadiazole derivatives exhibited potent inhibitory activities against SHP1. The representative compound **5b** showed good potency against SHP1 with IC₅₀ of (1.33 ± 0.16) μ mol/L and was also subjected to selectivity analyses to determine whether its biological properties made it suitable for further development.

2 Results and discussion

2.1 Chemistry

Eleven 5-phenyl-1,3,4-thiadiazole derivatives (**2a~2e**, **4** and **5a~5e**, Table 1) were designed and synthesized. A series of aromatic carboxylic acids (**1a~1e**) were condensed with thiosemicarbazide to give 1,3,4-thiadiazoles (**2a~2e**)^[24] (Scheme 1). By coupling of 3-((*tert*-butoxycarbonyl)amino)benzoic acid with compound **2a** in the presence of 2-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) and *N,N*-diisopropylethylamine (DIPEA), compound **3** was obtained in 40% isolated yield. The compounds **5a~5e** were obtained from compound **3** in two-step sequential reactions with deprotection of Boc group and coupling reaction with corresponding carboxylic acids in *N,N*-dimethylformamide (DMF). The compounds **5d~5e** were yielded using similar protocol to compound **3**.

2.2 Biochemical activities

2.2.1 SHP1 inhibitory activities and structure-activity relationships

The inhibitory activities of all the synthesized compounds against SHP1 were measured using 6,8-difluoro-4-thylumbelliferyl phosphate (DiFMUP) as substrate,^[25-26]

Figure 1 Chemical structures of reported SHP1 inhibitors

Figure 2 Structures of modulators of PTPs

Scheme 1 Synthetic route of compounds **2a**~**2e**, **4** and **5a**~**5e**

and the results were detailed in Table 1. Among amine compounds **2a**~**2e**, compounds with nitro group (**2a**~**2c**) showed slightly better potency than compounds without nitro group (**2d**~**2e**). Among compound **4** and its derivatives **5a**~**5c**, compound **5a** with furan-2-carbonyl group, compound **5b** with 5-methylfuran-2-carbonyl group showed inhibitory activities against SHP1 in low micromolar range, especially compound **5b** with IC₅₀ = (1.33 ± 0.16) μmol/L, significantly improving the potency compared to compound **4** and NSC-87877. Compound **5c** with tetrahydrofuran-2-carbonyl group exhibited similar activities against SHP1 compared to compound **4**. These results indicated that introduction of furan-carbonyl group

to the amino group on compound **4** benefited for the improvement of inhibitory activities against SHP1. By replacing the methylamino group with isopropylamino group and methoxy group on R² position respectively, the inhibitory activities of compounds **5d** and **5e** dropped significantly compared to compound **5a**.

2.2.2 Selectivity against other PTPs

In addition to potency improvements, we investigated the selectivity of compound **5b** against the other PTPs (PTP1B, TCPTP).^[27] As shown in Table 2, compound **5b** showed about 2-fold selectivity for SHP1 over SHP2 and no visible activity against PTP1B and TCPTP at the dose of 50 μmol/

Table 1 SHP1 inhibitory activities of compounds **2a**~**2e**, **4** and **5a**~**5e**

Compd.	R ¹	R ²	R ³	SHP1	
				Inhibition/% at 50 μmol/L	IC ₅₀ /(μmol·L ⁻¹)
2a	3-NO ₂	4-NHMe	NH ₂	27.5 ± 19.1	—
2b	3-NO ₂		NH ₂	20.9 ± 16.9	—
2c	3-NO ₂	4-OMe	NH ₂	21.6 ± 5.1	—
2d	3-OMe	4-OMe	NH ₂	—2.4 ± 7.2	—
2e	3-OH	3-OH	NH ₂	8.6 ± 1.8	—
4	3-NO ₂	4-NHMe		80.6 ± 6.5	23.42 ± 3.17
5a	3-NO ₂	4-NHMe		64.3 ± 11.0	4.40 ± 1.09
5b	3-NO ₂	4-NHMe		92.8 ± 2.1	1.33 ± 0.16
5c	3-NO ₂	4-NHMe		69.7 ± 2.7	21.56 ± 2.61
5d	3-NO ₂			36.4 ± 8.4	—
5e	3-NO ₂	4-OMe		38.6 ± 39.4	—
NSC-87877	—	—	—	—	34.76 ± 4.46

L, exhibiting better potency and selectivity than NSC-87877.

2.2.3 Characterization of **5b** on enzyme kinetics

A kinetic study was performed in order to identify the inhibitory mechanism of compound **5b** (Figure 3), using the reported enzyme kinetics assays.^[28] As shown in Figure 3A, **5b** demonstrated a fast-binding inhibition of SHP1. The

fast-binding inhibition of **5b** toward SHP1 may also exclude that **5b** is a nonspecific inhibitor, because nonspecific inhibitors always show time-dependent behavior and steep inhibition curve.^[29] We further determined the inhibition modality of **5b** which inhibited SHP1 with the characteristics typical of a competitive inhibitor, as indicated by increased K_m values and unchanged V_{max} values when the

Table 2 IC_{50}^a ($\mu\text{mol}\cdot\text{L}^{-1}$) values of compound **5b** against PTPs

Compd.	SHP1	SHP2	SHP2/SHP1	PTP1B	TCPTP
5b	1.33 ± 0.16	2.99 ± 0.36	2.25	NA ^b	NA ^b
NSC-87877	34.76 ± 4.46	33.71 ± 0.72		50.04 ± 11.91	40.70 ± 6.50

^a PTP1B (Protein tyrosine phosphatase 1B); TCPTP (T-cell protein tyrosine phosphatase). The DiFMUP substrates were utilized in PTP1B/TCPTP assay, and SHP1/SHP2 assay, respectively; IC_{50} values were determined by regression analyses and expressed as means ± SD of three replications; ^b NA: No activity (compound inhibitory ratio lower than 50% at the dose of 50 $\mu\text{mol/L}$).

inhibitor concentration was increased (Figure 3C). Meanwhile, the result of the Lineweaver-Burk plot confirmed **5b** as a competitive inhibitor of SHP1 for intersecting at y-axis of a nest of lines with increased inhibitor concentration (Figure 3B). The results indicated that **5b** binds the catalytic pocket of SHP1 and behaves as a competitor to the substrate.

2.3 Predicted binding mode of **5b**

To clarify the good activity and selectivity of compound **5b** against SHP1, the molecular character and molecular docking was performed using DFT method in Gaussian 09 software and the Lamarckian genetic algorithm local search method in the AutoDock 4.2 program, respectively.^[30-31] As shown in Figure 4A, the molecular electrostatic properties of **5b** were obtained based on the stable geometry (cartesian coordinates listed in Supporting Information). It can be found that carbonyl and nitro groups were electron-withdrawing groups, and 1,3,4-thiadiazole and benzene ring were filled with π -electrons, which may benefit for molecular activity and selectivity for SHP1. In the process of molecular docking with AutoDock 4.2 software, semi empirical free energy calculation method was used to evaluate the

energy matching between protein and small molecule **5b**, and then the binding ability between protein and **5b** was scored. Additionally, the charge was added by Kollman in AutoDock 4.2, all the rotatable torsional bonds of compound **5b** were set free, the size of grid box was set to 12.6 nm × 12.6 nm × 12.6 nm points with a 0.038 nm spacing and box centre (19.281, 1.872, 8.208), and the other parameters were maintained at their default settings. Here, the X-ray crystallographic structures of SHP1 (PDB Entry: 1FPR) and SHP2 (PDB Entry: 3ZM1) were selected for molecular docking.^[32] The lowest energy conformation in the distinctly distributed cluster was selected as the representation of the docking pose of **5b** with SHP1 with −37.55 kJ/mol of binding free energy (Figure 4B). We found four residues closer to the compound **5b** as shown in Figure 4B, indicating that these residues are likely to have a crucial role in higher activity and selectivity of **5b** for SHP1. Obviously, there are stronger stabilizing interactions between these residues and **5b**, and the residues K259 and V258 have strong hydrogen-bonding interactions with acceptor nitro group of **5b** (Figure 4C), respectively. At the same time, the partially negative charge of carbonyl group in Q256 and

Figure 3 Characterization of **5b** to SHP1

(A) Time-independent inhibition of SHP1 by **5b**; (B) typical competitive inhibition of **5b** shown by Lineweaver-Burk plot; (C) the initial velocity determined with various concentrations of 6,8-difluoro-4-methylumbelliferyl phosphate (DiFMUP) at various fixed concentrations of **5b**

Figure 4 Calculated results of molecular properties for **5b** and docking simulation

(A) Molecular electrostatic properties of mapped on the 0.001 a.u. contour of the electronic density at the DFT//B3LYP/6-31G(d) level; (B) docking pose of **5b** with SHP1 (PDB Entry: 1FPR): relative position of **5b** and WPD_loop (orange), Q_loop (purple), P_loop (Yellow) and PTyr_loop (green) in SHP1; (C) strong interactions between **5b** and the nearest residues

Q487 could stabilize the π -electrons of **5b** (Figure 4C) according to π - π stacking interaction.^[33] At the other hand, the interaction mode of **5b** and SHP2 was obtained. Compared to the binding mode of **5b** and SHP1, there are less H-bond and none π - π stacking interaction between **5b** and SHP2, which might to some extent account for the good selectivity of **5b** against SHP1. Lastly, the predicted ADME properties of **5b** were displayed in Figure S15 by applying SwissADME software.^[34] We hope that the predicted results could help for more drug molecules design in the future. The calculated results might to some extent account for the good activity and selectivity of **5b** for SHP1.

3 Conclusions

In conclusion, this work offered a completely new type of SHP1 inhibitors with 5-phenyl-1,3,4-thiadiazole scaffold. The representative compound **5b** exhibited significantly better potency and selectivity for SHP1 compared to that of the control compound NSC-87877. These preliminary results could provide a possible opportunity for the development of novel SHP1 inhibitors with promising cell permeability, bioavailability, and improved pharmacological properties.

4 Experimental section

4.1 Material

All chemicals were reagent grade and used as purchased. ^1H NMR (400 MHz) spectra were recorded on a Bruker

AVIII 400 MHz spectrometer. The chemical shifts were reported in using the δ 7.26 signal of CDCl_3 (^1H NMR) and the δ 2.50 signal of $\text{DMSO}-d_6$ (^1H NMR) as internal standards and the 40.00 signal of $\text{DMSO}-d_6$ (^{13}C NMR) as internal standards. ESI mass spectra (MS) was obtained on a Shimadzu LCMS-2020. High Resolution Mass Spectrometry (HRMS) was obtained on an ESI-QTOF maxis impact. High Performance Liquid Chromatography (HPLC) was obtained on a Shimadzu2030c 3D plus.

4.2 General procedure for preparation of compounds **2a**~**2e**, **4** and **5a**~**5e**

4.2.1 General procedure for preparation of compounds **2a**~**2e**

Carboxylic acid (5.6 mmol), thiosemicarbazide (0.51 g, 5.6 mmol) and phosphorus oxychloride (10 mL) were added successively to a 50 mL round-bottom flask. The mixture was stirred at 85 $^\circ\text{C}$ for 2.5 h. The reaction solution was cooled to room temperature, then the solution was concentrated under reduced pressure. The residue was washed with saturated sodium bicarbonate solution, the mixture was filtered and the filter cake was dried *in vacuo* to give crude product. The crude product was adsorbed onto silica (2 g, 100~200 mesh) to form a dry sample, purified by silica gel column chromatography (20 g, 3 cm $\times$ 30 cm, 100~200 mesh, petroleum ether/EtOAc, $V:V=91:9\sim50:50$) to give compound **2a**~**2e**.

5-(4-(Methylamino)-3-nitrophenyl)-1,3,4-thiadiazol-2-amine (**2a**): Red solid, yield 78%. m.p. 266.7~268.3 $^\circ\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.43~8.41 (m, 1H), 8.29

(d, $J=2.0$ Hz, 1H), 7.93 (dd, $J=9.2, 2.4$ Hz, 1H), 7.36 (s, 2H), 7.09 (d, $J=9.2$ Hz, 1H), 3.01 (d, $J=4.8$ Hz, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 167.8, 155.0, 146.2, 133.8, 130.5, 123.6, 117.9, 115.4, 29.9; MS (ESI) m/z : 252 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_9\text{H}_{10}\text{N}_3\text{O}_2\text{S}$ [$\text{M} + \text{H}$] $^+$ 252.0555, found 252.0543.

5-(4-(Isopropylamino)-3-nitrophenyl)-1,3,4-thiadiazol-2-amine (**2b**): Red solid, yield 74%. m.p. 225.3~227.6 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.29 (d, $J=2.0$ Hz, 1H), 8.07 (d, $J=8.0$ Hz, 1H), 7.92 (dd, $J=9.2, 2.0$ Hz, 1H), 7.36 (s, 2H), 7.20 (d, $J=9.2$ Hz, 1H), 4.02~3.97 (m, 1H), 1.29 (d, $J=6.4$ Hz, 6H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 167.9, 154.9, 144.7, 133.9, 130.4, 123.7, 118.2, 115.8, 43.9, 22.2; MS (ESI) m/z : 280.1 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_{11}\text{H}_{13}\text{N}_5\text{NaO}_2\text{S}$ [$\text{M} + \text{Na}$] $^+$ 302.0688, found 302.0676.

Compounds **2c**~**2e** have been reported.^[35-37]

5-(4-Methoxy-3-nitrophenyl)-1,3,4-thiadiazol-2-amine (**2c**): Yellow solid, yield 48%. Purity 95.0%, m.p. 186.8~187.4 $^{\circ}\text{C}$ (lit.^[35] 210 $^{\circ}\text{C}$); ^1H NMR (400 MHz, DMSO- d_6) δ : 7.97 (d, $J=8.4$ Hz, 1H), 7.68 (s, 2H), 7.66 (d, $J=1.6$ Hz, 1H), 7.43 (dd, $J=8.4, 1.6$ Hz, 1H), 4.01 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 169.9, 154.2, 152.5, 139.1, 136.4, 126.2, 118.5, 110.7, 56.8; MS (ESI) m/z : 253 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_9\text{H}_8\text{N}_4\text{NaO}_3\text{S}$ [$\text{M} + \text{Na}$] $^+$ 275.0215, found 275.0203.

5-(3,4-Dimethoxyphenyl)-1,3,4-thiadiazol-2-amine (**2d**): White solid, yield 84%. Purity 95.6%, m.p. 122.7~123.6 $^{\circ}\text{C}$ (lit.^[36] 195~197 $^{\circ}\text{C}$); ^1H NMR (400 MHz, DMSO- d_6) δ : 7.36 (d, $J=2.0$ Hz, 1H), 7.30 (s, 2H), 7.20 (dd, $J=8.4, 2.0$ Hz, 1H), 7.01 (d, $J=8.4$ Hz, 1H), 3.82 (s, 3H), 3.80 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.0, 156.5, 150.1, 149.0, 123.8, 119.9, 111.9, 108.7, 55.6, 55.5; MS (ESI) m/z : 238 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_{10}\text{H}_{11}\text{N}_3\text{NaO}_2\text{S}$ [$\text{M} + \text{Na}$] $^+$ 260.0470, found 260.0472.

4-(5-Amino-1,3,4-thiadiazol-2-yl)benzene-1,2-diol (**2e**): White solid, yield 90%. Purity 97.9%, m.p. 253.9~255.4 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.22 (d, $J=2.0$ Hz, 1H), 7.04 (dd, $J=8.0, 2.0$ Hz, 1H), 6.85 (d, $J=8.4$ Hz, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 167.4, 156.9, 147.2, 145.6, 122.4, 118.4, 115.9, 113.2; MS (ESI) m/z : 210 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_8\text{H}_8\text{N}_3\text{O}_2\text{S}$ [$\text{M} + \text{H}$] $^+$ 210.0337, found 210.0327.

4.2.2 General procedure for preparation of compound **4**

To the solution of 3-((*tert*-butoxycarbonyl)amino)benzoic acid (201 mg, 0.85 mmol), HATU (195 mg, 0.51 mmol) and DIPEA (372 μL , 2.14 mmol) in DMF (4 mL) add compound **2a** (100 mg, 0.40 mmol), the mixture was stirred at room temperature overnight. Then add water (5 mL) to the reaction solution, the aqueous layer was extracted with EtOAc (20 mL $\times$ 3). The combined organic layers were washed with water and dried over Na_2SO_4 . The crude product was adsorbed onto silica (0.5 g, 100~200 mesh) to form a dry sample, purified by silica gel column chromatography (5 g, 1 cm diameter $\times$ 20 cm height, 100~200 mesh, petroleum ether/EtOAc, $V:V=83:17\sim 50:50$)

to obtain compound **3** (75 mg, 40%): Yellow solid, m.p. > 300 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.58 (s, 1H), 8.57 (d, $J=2.0$ Hz, 1H), 8.48~8.47 (m, 1H), 8.22 (s, 1H), 8.13~8.10 (m, 1H), 7.75 (d, $J=8.0$ Hz, 1H), 7.66~7.64 (m, 1H), 7.45 (t, $J=8.0$ Hz, 1H), 7.16 (d, $J=9.2$ Hz, 1H), 3.04 (d, $J=4.8$ Hz, 3H), 1.50 (s, 9H).

Compound **3** (70 mg, 0.15 mmol) was dissolved in HCl-dioxane (4 mol/L, 2 mL), stirred overnight at room temperature. After removal of dioxane, the solid was washed with saturated sodium bicarbonate solution, filtered and dried to obtain compound **4** (50 mg, 90%): Yellow solid, m.p. > 300 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 12.86 (s, 1H), 8.58 (d, $J=2.0$ Hz, 1H), 8.48 (d, $J=5.2$ Hz, 1H), 8.12 (dd, $J=9.2, 2.4$ Hz, 1H), 7.31 (d, $J=8.0$ Hz, 1H), 7.27 (s, 1H), 7.23~7.15 (m, 2H), 6.84 (dd, $J=8.0, 2.4$ Hz, 1H), 3.04 (d, $J=4.8$ Hz, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 158.5, 146.7, 144.6, 144.5, 136.9, 134.2, 132.2, 130.8, 129.2, 124.4, 116.9, 116.1, 115.6, 29.9; MS (ESI) m/z : 371 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_{16}\text{H}_{14}\text{N}_6\text{NaO}_3\text{S}$ [$\text{M} + \text{Na}$] $^+$ 393.0746, found 393.0746.

4.2.3 General procedure for preparation of compounds **5a**~**5e**

To the solution of amine (0.14 mmol mmol), HATU (76 mg, 0.20 mmol) and DIPEA (122 μL , 0.70 mmol) in DMF (2 mL) was added carboxylic acid (0.27 mmol), the mixture was stirred at room temperature overnight. Then water (5 mL) was added to the reaction solution, and the aqueous layer was extracted with EtOAc (20 mL) 3 times. The combined organic layers were washed with water and dried over Na_2SO_4 . The crude product was adsorbed onto silica (0.5 g, 100~200 mesh) to form a dry sample, purified by silica gel column chromatography (5 g, 1 cm $\times$ 20 cm, 100~200 mesh, DCM/MeOH, $V:V=99:1\sim 95:5$) to obtain compounds **5a**~**5e**.

N-(3-((5-(4-(Methylamino)-3-nitrophenyl)-1,3,4-thiadiazol-2-yl)carbamoyl)phenyl)furan-2-carboxamide (**5a**): Yellow solid, yield 64%. Purity 95.3%, m.p. 272.7~273.6 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 13.17 (s, 1H), 10.43 (s, 1H), 8.59 (d, $J=2.4$ Hz, 1H), 8.51~8.49 (m, 2H), 8.13 (dd, $J=9.6, 2.4$ Hz, 1H), 8.03 (d, $J=8.0$ Hz, 1H), 7.97 (s, 1H), 7.90 (d, $J=8.0$ Hz, 1H), 7.54 (t, $J=8.0$ Hz, 1H), 7.40 (d, $J=3.2$ Hz, 1H), 7.16 (d, $J=8.8$ Hz, 1H), 6.73 (dd, $J=3.6, 2.0$ Hz, 1H), 3.04 (d, $J=4.8$ Hz, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 160.5, 156.4, 147.3, 146.7, 146.0, 138.9, 134.2, 132.5, 132.4, 130.8, 129.0, 124.5, 124.4, 123.4, 120.5, 117.0, 115.5, 115.1, 112.2, 29.9; MS (ESI) m/z : 465 [$\text{M} + \text{H}$] $^+$; HRMS calcd for $\text{C}_{21}\text{H}_{16}\text{N}_6\text{NaO}_5\text{S}$ [$\text{M} + \text{Na}$] $^+$ 487.0801, found 487.0796.

5-Methyl-*N*-(3-((5-(4-(methylamino)-3-nitrophenyl)-1,3,4-thiadiazol-2-yl)carbamoyl)phenyl)furan-2-carboxamide (**5b**): Yellow solid, yield 68%. Purity 99.2%, m.p. 273.2~274.3 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 13.12 (s, 1H), 10.24 (s, 1H), 8.60 (d, $J=2.4$ Hz, 1H), 8.49 (d, $J=5.2$ Hz, 1H), 8.46 (s, 1H), 8.13 (dd, $J=8.8, 2.0$ Hz, 1H), 8.03 (d, $J=8.0$ Hz, 1H), 7.89 (d, $J=8.0$ Hz, 1H), 7.53 (t, $J=8.0$ Hz, 1H), 7.29 (d, $J=3.2$ Hz, 1H), 7.17 (d, $J=9.2$

H₂, 1H), 6.35 (dd, $J=3.2, 0.8$ Hz, 1H), 3.04 (d, $J=4.8$ Hz, 3H), 2.40 (s, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 156.4, 155.5, 146.7, 145.7, 139.1, 134.3, 130.8, 129.7, 128.9, 124.4, 123.2, 120.4, 117.0, 116.3, 115.6, 108.7, 29.9, 13.6; MS (ESI) m/z : 479 [M+H]⁺; HRMS calcd for C₂₂H₁₈N₆-NaO₅S [M+Na]⁺ 501.0957, found 501.0961.

N-(3-((5-(4-(Methylamino)-3-nitrophenyl)-1,3,4-thiadiazol-2-yl)carbamoyl)phenyl)tetrahydrofuran-2-carboxamide (**5c**): Yellow solid, yield 65%. Purity 95.5%, m.p. 219.9~221.1 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 13.10 (s, 1H), 9.90 (s, 1H), 8.59 (d, $J=2.0$ Hz, 1H), 8.49 (d, $J=5.2$ Hz, 1H), 8.43 (s, 1H), 8.13 (dd, $J=8.8, 2.0$ Hz, 1H), 8.14~8.11 (m, 1H), 7.95 (d, $J=8.0$ Hz, 1H), 7.86 (d, $J=8.0$ Hz, 1H), 7.50 (t, $J=8.0$ Hz, 1H), 7.17 (d, $J=9.2$ Hz, 1H), 4.43 (dd, $J=8.0, 5.6$ Hz, 1H), 4.06~3.99 (m, 1H), 3.88~3.83 (m, 1H), 3.04 (d, $J=4.8$ Hz, 3H), 2.27~2.18 (m, 1H), 2.06~1.97 (m, 1H), 1.92~1.85 (m, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 171.9, 146.7, 138.8, 134.2, 130.8, 128.9, 124.4, 123.3, 120.2, 116.8, 115.5, 77.9, 68.9, 30.0, 27.3, 25.1; MS (ESI) m/z : 469 [M+H]⁺; HRMS calcd for C₂₁H₂₀N₆NaO₅S [M+Na]⁺ 491.1114, found 491.1108.

N-(3-((5-(4-(Isopropylamino)-3-nitrophenyl)-1,3,4-thiadiazol-2-yl)carbamoyl)phenyl)furan-2-carboxamide (**5d**): Yellow solid, yield 49%. m.p. > 300 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 13.17 (s, 1H), 10.41 (s, 1H), 8.59 (d, $J=2.0$ Hz, 1H), 8.49~8.48 (m, 1H), 8.13~8.10 (m, 2H), 8.03~8.01 (m, 1H), 7.97 (s, 1H), 7.91 (d, $J=7.6$ Hz, 1H), 7.53 (t, $J=8.0$ Hz, 1H), 7.40 (d, $J=3.6$ Hz, 1H), 7.28 (d, $J=9.2$ Hz, 1H), 6.73 (dd, $J=3.6, 1.6$ Hz, 1H), 4.05~4.03 (m, 1H), 1.31 (d, $J=6.4$ Hz, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 165.4, 160.9, 156.4, 155.2, 153.9, 147.2, 146.0, 139.0, 132.2, 129.0, 126.7, 125.0, 124.7, 123.4, 121.8, 120.5, 116.1, 115.1, 113.6, 112.3, 48.0, 18.9; MS (ESI) m/z : 493.1 [M+H]⁺; HRMS calcd for C₂₃H₂₀N₆-NaO₅S [M+Na]⁺ 515.1114, found 515.1107.

N-(3-((5-(4-Methoxy-3-nitrophenyl)-1,3,4-thiadiazol-2-yl)carbamoyl)phenyl)furan-2-carboxamide (**5e**): White solid, yield 26%. m.p. 280.2~281.1 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 13.24 (s, 1H), 10.44 (s, 1H), 8.50 (s, 1H), 8.48 (d, $J=2.4$ Hz, 1H), 8.27 (dd, $J=8.8, 2.0$ Hz, 1H), 8.03 (d, $J=8.0$ Hz, 1H), 7.98 (d, $J=1.6$ Hz, 1H), 7.91 (d, $J=7.6$ Hz, 1H), 7.51 (t, $J=7.6$ Hz, 2H), 7.40 (d, $J=3.6$ Hz, 1H), 6.73 (dd, $J=3.6, 2.0$ Hz, 1H), 4.02 (s, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 165.3, 156.4, 152.5, 147.2, 146.0, 140.0, 139.0, 135.5, 131.9, 129.1, 126.2, 124.8, 123.4, 120.6, 119.1, 115.1, 112.3, 112.0, 57.0; MS (ESI) m/z : 466 [M+H]⁺; HRMS calcd for C₂₁H₁₅N₅NaO₆S [M+Na]⁺ 488.0641, found 488.0624.

4.3 SHP1 and related PTPs biological assay

A colorimetric assay to measure inhibition against SHP2 and SHP1 was performed in 384-well plates. Briefly, the tested compounds were solubilized in DMSO and serially diluted into concentrations for the inhibitory test and the enzymatic activities of the SHP2 and SHP1 were determined at room temperature by monitoring the dephosphorylation of 6,8-difluoro-4-methylumbelliferyl phosphate

(DiFMUP), products were then detected at a 355 nm excitation wavelength and 460 nm emission wavelength by the EnVision multilabel plate reader (Perkin-Elmer Life Sciences, Boston, MA, USA). The assays were carried out in a final volume of 50 μ L containing 60 mmol/L HEPES, pH 7.2, 75 mmol/L NaCl, 75 mmol/L KCl, 1 mmol/L ethylene diamine-*N,N,N',N'*-tetraacetate (EDTA), 0.05% Tween-20, 5 mmol/L dithiothreitol (DTT), 2 nmol/L enzyme, and indicating compound concentration. The substrate was then added to a final concentration of their respective K_m (Michaelis constant) concentrations. The initial rate of the dephosphorylation was presented by the early linear region of the enzymatic reaction kinetic curve, the inhibitory activity of the compound was continuously monitored. The inhibitor dose-response curves were analyzed using normalized IC₅₀ regression curve fitting with control based normalization.

4.4 Characterization of compound **5b** on SHP1

In the time-independent inhibition experiment, 2 nmol/L SHP1 was incubated by 1.5 μ mol/L of compound **5b** (2% DMSO as blank) for different times, and then 30 μ L of mixture of enzyme and compounds was added to a final 50 μ L of assay system to initiate the enzyme reaction. The dissociation of the enzyme-inhibitor complex was monitored by substrate hydrolysis by measuring the fluorescence using Envision microplate reader using excitation and emission wavelengths of 355 and 460 nm, respectively.

In the enzymatic kinetic experiment, the assay was carried out in a 50 μ L system containing 60 mmol/L HEPES, pH 7.2, 75 mmol/L NaCl, 75 mmol/L KCl, 1 mmol/L EDTA, 0.05% Tween-20, 5 mmol/L DTT, 2 nmol/L SHP1, substrate DiFMUP in 2-fold dilution from 400 μ mol/L to 3.125 μ mol/L, and different concentrations of the inhibitors. In the presence of the competitive inhibitor, the Michaelis-Menten equation is described as:

$$1/v = (K_m/[V_{\max}[S]])(1 + [I]/K_i) + 1/V_{\max}$$

where K_m is the Michaelis constant, v is the initial rate, $V_{\max}$ is the maximum rate, and $[S]$ is the substrate concentration. $[I]$ is concentration of free inhibitor, K_i is inhibition constant.

Supporting Information ¹H NMR, ¹³C NMR, LC-MS, HRMS and HPLC spectra for new compounds and computational details of **5b**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Tonks, N. K. *FEBS J.* **2013**, *280*, 346.
- [2] Hendriks, W.; Elson, A.; Harroch, S.; Pulido, R.; Stoker, A.; Hertog, J. D. *FEBS J.* **2013**, *280*, 708.
- [3] Frankson, R.; Yu, Z. H.; Bai, Y.; Li, Q.; Zhang, R. Y.; Zhang, Z. Y. *Cancer Res.* **2017**, *77*, 5701.
- [4] Varone, A.; Spano, D.; Corda, D. *Front. Oncol.* **2020**, *10*, 935.
- [5] Sandur, S. K.; Pandey, M. K.; Sung, B.; Aggarwal, B. B. *Mol. Cancer Res.* **2010**, *8*, 107.
- [6] Sharma, Y.; Bashir, S.; Bhardwaj, P.; Ahmad, A.; Khan, F. *Immunol.*

- Res.* **2016**, *64*, 804.
- [7] Yuan, X.; Bu, H.; Zhou, J.; Yang, C. Y.; Zhang, H. *J. Med. Chem.* **2020**, *63*, 11368.
- [8] Dempke, W. M.; Peter, U.; Klaus, F.; Timothy, C. *Oncology* **2018**, *95*, 1.
- [9] Song, M.; Park, J. E.; Park, S. G.; Lee, D. H.; Choi, H. K.; Park, B. *C. Biochem. Biophys. Res. Commun.* **2009**, *381*, 491.
- [10] Kundu, S.; Fan, K.; Cao, M.; Lindner, D. J.; Zhao, Z. J.; Borden, E.; Yi, T. L. *J. Immunol.* **2010**, *184*, 6529.
- [11] Arabaci, G.; Yi, T.; Fu, H.; Porter, M. E.; Beebe, K. D.; Pei, D. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3047.
- [12] Yi, T.; Elson, P.; Mitsuhashi, M.; Jacobs, B.; Hollovar, E.; Budd, G. T.; Spiro, T.; Triozzi, E.; Borden, E. C. *Oncotarget* **2011**, *2*, 1155.
- [13] Lu, L.; Wang, S.; Zhu, M.; Liu, Z.; Guo, M.; Xing, S.; Fu, X. *Bio-metals* **2010**, *23*, 1139.
- [14] Wang, Q.; Zhu, M.; Lu, L.; Yuan, C.; Fu, X. *Dalton. Trans.* **2011**, *40*, 12926.
- [15] Akiba, H.; Sumaoka, J.; Hamakubo, T.; Komiyama, M. *Anal. Bioanal. Chem.* **2014**, *406*, 2957.
- [16] Frija, L.; Pombeiro, A.; Kopylovich, M. N. *Eur. J. Org. Chem.* **2017**, *19*, 2670.
- [17] Yang, H.; Li, C. Y.; Wang, X. M.; Yang, Y. H.; Zhu, H. L. *Chem. Rev.* **2014**, *45*, 5572.
- [18] Wang, W. L.; Yang, D. L.; Gao, L. X.; Tang, C. L.; Ma, W. P.; Ye, H. H.; Zhang, S. Q.; Zhao, Y. N.; Xu, H. J.; Hu, Z.; Chen, X.; Fan, W. H.; Chen, H. J.; Li, J. Y.; Nan, F. J.; Li, J.; Feng, B. N. *Molecules* **2013**, *19*, 102.
- [19] Wang, W. L.; Chen, X.; Gao, L. X.; Sheng, L.; Li, J. Y.; Li, J.; Feng, B. N. *Chem. Biol. Drug Des.* **2015**, *86*, 1161.
- [20] Satheeshkumar, R.; Zhu, R.; Feng, B.; Huang, C.; Gao, Y.; Gao, L. X.; Shen, C.; Hou, T. J.; Xu, L.; Li, J.; Zhu, Y. L.; Zhou, Y. B.; Wang, W. L. *Bioorg. Med. Chem. Lett.* **2020**, *30*, 127170.
- [21] Wang, W. L.; Chen, X. Y.; Gao, Y.; Gao, L. X.; Sheng, L.; Zhu, J. Y.; Xu, L.; Ding, Z. Z.; Zhang, C.; Li, J. Y.; Li, J.; Zhou, Y. B. *Bioorg. Med. Chem. Lett.* **2017**, *27*, 5154.
- [22] Wang, W. L.; Huang, C.; Gao, L. X.; Tang, C. L.; Wang, J. Q.; Wu, M. C.; Sheng, L.; Chen, H. J.; Nan, F. J.; Li, J. Y.; Feng, B. N. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 1889.
- [23] Wang, W. L.; Luo, H.; Gao, Y.; Gao, L. X.; Sheng, L.; Zhou, Y. B.; Li, J. Y.; Li, J.; Feng, B. N. *Chin. J. Org. Chem.* **2016**, *36*, 2142 (in Chinese).
(王文龙, 骆欢, 高雅, 高立信, 盛丽, 周宇波, 李佳, 李静雅, 冯柏年, 有机化学, **2016**, *36*, 2142.)
- [24] Pac, A.; As, A.; Fal, A.; Mb, B.; Mfe, C. *J. Mol. Struct.* **2019**, *1179*, 11.
- [25] Chen, Y. N.; LaMarche, M. J.; Chan, H. M.; Fekkes, P.; Garcia, F. J.; Acker, M. G.; Antonakos, B.; Chen, C. H.; Chen, Z.; Cooke, V. G.; Dobson, J. R.; Deng, Z.; Fei, F.; Firestone, B.; Fodor, M.; Fridrich, C.; Gao, H.; Grunenfelder, D.; Hao, H. X.; Jacob, J.; Ho, S.; Hsiao, K.; Kang, Z. B.; Karki, R.; Kato, M.; Larrow, J.; La-Bonte, L. R.; Lenoir, F.; Liu, G.; Liu, S.; Majumdar, D.; Meyer, M. J.; Palermo, M.; Perez, L.; Pu, M.; Price, E.; Quinn, C.; Shaky, S.; Shultz, M. D.; Slisz, J.; Venkatesan, K.; Wang, P.; Warmuth, M.; Williams, S.; Yang, G.; Yuan, J.; Zhang, J. H.; Zhu, P.; Ramsey, T.; Keen, N. J.; Sellers, W. R.; Stams, T.; Fortin, P. D. *Nature* **2016**, *535*, 148.
- [26] Welte, S.; Baringhaus, K. H.; Schmider, W.; Müller, G.; Petry, S.; Tennagels, N. *Anal. Biochem.* **2005**, *338*, 32.
- [27] Krishnan, N.; Koveal, D.; Miller, D. H.; Xue, B.; Akshinthala, S. D.; Kragelj, J.; Jensen, M. R.; Gauss, C. M.; Page, R.; Blackledge, M.; Muthuswamy, S. K.; Peti, W.; Tonks, N. K. *Nat. Chem. Biol.* **2014**, *10*, 558.
- [28] Zhang, W.; Hong, D.; Zhou, Y.; Zhang, Y.; Shen, Q.; Li, J. Y.; Hu, L. H.; Li, J. *Biochim. Biophys. Acta* **2006**, *1760*, 1505.
- [29] McGovern, S. L.; Helfand, B. T.; Feng, B.; Shoichet, B. K. *J. Med. Chem.* **2003**, *46*, 4265.
- [30] Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Keith, T.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, O.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09*, Revision D.01, Gaussian, Inc., Wallingford, CT, **2013**.
- [31] Morris, G. M.; Huey, R.; Lindstrom, W.; Sanner, M. F.; Belew, R. K.; Goodsell, D. S.; Olson, A. J. *J. Comput. Chem.* **2009**, *30*, 2785.
- [32] Yang, J.; Cheng, Z. L.; Niu, T. Q.; Liang, X. S.; Zhao, Z. Z. J.; Zhou, G. W. *J. Biol. Chem.* **2000**, *275*, 4066.
- [33] Zhang, B.; Wang, K. B.; Wang, W.; Wang, X.; Liu, F.; Zhu, J. P.; Shi, J.; Li, L. Y.; Han, H.; Xu, K.; Qiao, H. Y.; Zhang, X.; Jiao, R. H.; Hou, K. N.; Liang, Y.; Tan, R. X.; Ge, H. M. *Nature* **2019**, *568*, 122.
- [34] Daina, A.; Michielin, O.; Zoete, V. *Sci. Rep.* **2017**, *7*, 1.
- [35] Vachala, S. D.; Bhargavi, B. *J. Chem. Pharm. Res.* **2014**, *6*, 377.
- [36] Guan, P.; Wan, Y. C.; Hou, X. B.; Wang, L.; Xu, W. F.; Tang, W. P.; Fang, H. *Bioorg. Med. Chem.* **2014**, *22*, 5766.
- [37] Jakovljević, K.; Matić, I. Z.; Stanojković, T.; Krivokuća, A.; Marković, V.; Joksović, M. D.; Mihailović, N.; Nićiforović, M.; Joksović, L. *Bioorg. Med. Chem. Lett.* **2017**, *27*, 3709.

(Lu, Y.)