以芳基碘为碘化试剂的苯酚衍生物的位点选择性 C—H 键碘化反应

张 涛^a 李尚达^{*,b} 周春林^b 王新超^b 张 孟^b
高泽众^b 李 纲^{*,b}

(^a 中北大学理学院 山西太原 030051)

(^b 中国科学院福建物质结构研究所 煤制乙二醇及相关技术重点实验室 福州 350002)

摘要 富电子苯酚类化合物的位点选择性 C—H 键碘化是一个挑战性的反应。报道了一类钯催化的、利用 4-碘基-3-硝基苯甲醚为温和的碘化试剂, 实现无保护 2-芳基苯酚和 2-苯氧乙酸类化合物的 C—H 键碘化反应。该反应适用于一系列富电子的苯酚类衍生物, 且能够获得优秀的位点选择性和良好的官能团耐受性。结果表明, 对于具有挑战性底物的 C—H 键碘化, 经由形式复分解的 C—H 键碘化反应是一个潜在的有用方法。

关键词 C—H 键碘化; 形式复分解; 位点选择性; 苯酚; 芳基碘化物; 钯催化

Site-Selective C—H Iodination of Phenol Derivatives Using Aryl Iodide as Iodinating Reagent

Zhang, Tao^a Li, Shangda^{*,b} Zhou, Chunlin^b Wang, Xinchao^b Zhang, Meng^b
Gao, Zezhong^b Li, Gang^{*,b}

(^a School of Science, North University of China, Taiyuan, Shanxi 030051)

(^b Key Laboratory of Coal to Ethylene Glycol and Its Related Technology, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou 350002)

Abstract Site-selective C—H iodination of electron-rich phenols is a challenging reaction. A Pd(II)-catalyzed C—H iodination of free 2-aryl phenols and 2-phenoxyacetic acids using 4-iodo-3-nitroanisole as the mild iodinating reagent was reported. Excellent site-selectivity and good functional group tolerance were obtained with a range of electron rich phenol derivatives. These results suggest that C—H iodination via formal metathesis is a potentially useful method for C—H iodination of challenging substrates.

Keywords C—H iodination; formal metathesis; site-selectivity; phenol; aryl iodide; palladium catalysis

1 Introduction

Phenol motifs are found in numerous synthetic intermediates, natural products and bioactive molecules.^[1] Therefore, site-selective direct C—H iodination of phenols has received wide attention. However, direct iodination of phenols utilizing electrophilic aromatic substitution (S_EAr) reaction generally leads to mixture of *ortho* and/or *para*-iodinated products (Scheme 1a). In addition, although directed *ortho*-lithiation of phenol carbamates could be used to prepare *ortho*-iodinated phenols, this method requires

harsh reaction conditions using rather strong base such as *sec*-butyllithium.^[2] In the past decades, transition-metal-catalyzed C—H iodination has emerged as a valuable method to access aryl iodides.^[3-4] Notably, a single report was disclosed on *ortho*-C—H iodination of phenol carbamates using costly cyclic hypervalent iodine(III) oxidant in the presence of a strong acid, *i.e.* trifluoromethanesulfonic acid.^[5] However, major *para*-iodinated phenols were generated instead if commonly-used electrophilic *N*-iodo-succinide (NIS) or I₂/PhI(OAc)₂ was used as the iodinating reagent with these phenol carbamates.^[5] Importantly, tran-

* Corresponding authors. E-mail: sdli@fjirsm.ac.cn; gangli@fjirsm.ac.cn.

Received June 6, 2021; revised June 7, 2021; published online June 7, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 22022111, 22071248), the Natural Science Foundation of Fujian Province (Nos. 2020J02008, 2020J01108), and the Youth Innovation Promotion Association of the Chinese Academy of Sciences (No. 2020306).

国家自然科学基金(Nos. Nos. 22022111, 22071248)、福建省自然科学基金(Nos. 2020J02008, 2020J01108)和中国科学院青年创新促进会(No. 2020306)资助项目。

sition-metal-catalyzed C—H iodination of free phenol derivatives has not been realized to date.^[6-7] Therefore, the development of a mild, non-electrophilic, and readily available reagent for C—H iodination of phenol derivatives is highly desirable.

a) possible products of iodination of phenols

b) C—H iodination of hydrocinnamic acids with ArI (our previous work)

c) C—H iodination of phenol derivatives with ArI (this work)

Scheme 1 Site-selective C—H iodination of phenol derivatives via formal metathesis

In recent years, notable advances have been made in functional group metathesis reactions.^[8a-8c] However, C—H functionalization via a metathesis reaction is extremely rare.^[8a,8d-8g] Recently, we discovered that a set of 2-nitrophenyl iodides such as 4-iodo-3-nitroanisole could enable *ortho*- and *meta*-C—H iodination of hydrocinnamic acids via a rare formal metathesis reaction (Scheme 1b).^[9] Since the 2-nitrophenyl iodides are mild iodinating reagents that are readily available and non-electrophilic, we envisioned that they could be possibly applied to the challenging site-selective C—H iodination of electron-rich phenol derivatives. Herein, we report the development of a site-selective C—H iodination of free 2-aryl phenols as well as 2-phenoxyacetic acids using 4-iodo-3-nitroanisole as the mild iodinating reagent (Scheme 1c).

2 Results and discussion

To start our investigation, commercially available 2-phenyl phenol (**1a**) was selected as the model substrate to react with 4-iodo-3-nitroanisole (Table 1), since the phenolic hydroxy group has been proved to be a good chelating group in C—H activation reactions.^[10] In our previous study of C—H iodination via formal metathesis, 39% iodinated product **2a** has been detected during the mechanistic study.^[9] However, 11% undesired arylation^[10h] product was

also produced together with 11% dibenzofuran **3a** side product that was probably generated from a competing C—O reductive elimination^[11] with 4-iodo-3-nitroanisole acting as an oxidant.^[12] Therefore, an extensive re-evaluation of reaction conditions was carried out. As shown in Table 1, product **2a** could be obtained in 23% yield with Pd(OAc)₂ as the catalyst and *N*-Formyl-Gly-OH (**L1**) as the ligand, using 0.5 equiv. of NaOAc as the base after heating at 90 °C in hexafluoroisopropanol (HFIP) for 24 h (Entry 1). Notably, no detectable arylation product was obtained. The yield of **2a** was increased remarkably when pyridine-type^[13] ligand **L2** or **L3** was employed as the ligand (Entries 2 and 3). Base was then evaluated and one equivalent of K₂CO₃ proved to be best with ligand **L3** (Entries 4~8). Subsequently, silver salt additives such as AgCl, AgOAc, and Ag₂CO₃ were tested, and it was found that 0.1 equiv. of AgCl was the most effective, leading to a slight increase of yield (Entries 9~15), which also suggested that the addition of AgCl was not crucial. However, the exact role of AgCl in the reaction is not clear at this stage. Moreover, the reaction temperature could be reduced to 60 °C nearly without change of the yield (Entries 16 and 17). Nitrogen atmosphere was proved to be not necessary (Entry 18), and other Pd catalysts such as Pd(OTFA)₂ and PdCl₂ were not superior (Entries 19 and 20). In addition, it was beneficial to shorten the reaction time to 12 h but further reducing the reaction time was not helpful (Entries 21 and 22). Notably, arylation side product was not detectable under the above reaction conditions, though it was hard to completely eliminate the generation of side product **3a**.

With the optimized reaction conditions in hand, the scope of C—H iodination of 2-aryl phenols was explored (Table 2). Good isolated yield of **2a** was received, and the yield decreased only slightly on a larger scale. Substitutions with both electron donating and withdrawing groups at the 3'- and 4'-position of the 2-aryl group were tolerated, leading to generally good yields of products (**2b~2j**). However, substrates like 1-phenylnaphthalen-2-ol afforded very low yield of desired product, possibly because they were hard to achieve a coplanar orientation that was important for C—H bond cleavage. Various functional groups including halides were also tolerated on the phenol ring of the 2-aryl phenols, affording high yields of desired products (**2k~2p**). Finally, substitutions on both aryl rings of 2-aryl phenols were also compatible with this method (**2q** and **2r**).

Based on the success in C—H iodination of 2-aryl phenols using 4-iodo-3-nitroanisole, we moved on to test this method with 2-phenoxyacetic acids that were designed for *ortho*-C—H iodination of phenols with a carboxyl directing group (Table 3). After re-evaluation of the reaction conditions, good combined yield of desired products was achieved with substrate 2-phenoxyacetic acid (**4a**) using *N*-Formyl-Gly-OH (**L1**) as the ligand without the addition of a silver salt. The products were converted to methyl esters that were isolated more easily than corresponding acids (**5a**). Pleasingly, a series of electron donating as well

Table 1 Optimization of reaction conditions^a

1a + **(2 equiv.)** $\xrightarrow[\text{base, Ag salt, HFIP (0.1 mol/L), T, 24 h}]{\text{Pd(OAc)}_2 \text{ (10 mol\%)}, \text{ligand (20 mol\%)}}$ **2a** + **3a**

L1

L2

L3

Entry	Ligand	Base (equiv.)	Ag salt (equiv.)	T/°C	Yield/% 2a/3a	Entry	Ligand	Base (equiv.)	Ag salt (equiv.)	T/°C	Yield/% 2a/3a
1	L1	NaOAc (0.5)	—	90	23/10	12	L3	K ₂ CO ₃ (1.0)	AgNO ₃ (0.1)	90	50/7
2	L2	NaOAc (0.5)	—	90	47/8	13	L3	K ₂ CO ₃ (1.0)	AgBr (0.1)	90	65/6
3	L3	NaOAc (0.5)	—	90	52/10	14	L3	K ₂ CO ₃ (1.0)	AgCl (0.2)	90	80/8
4	L3	NaOAc (1.0)	—	90	57/9	15	L3	K ₂ CO ₃ (1.0)	AgCl (0.5)	90	66/8
5	L3	Na ₂ CO ₃ (1.0)	—	90	73/10	16	L3	K ₂ CO ₃ (1.0)	AgCl (0.1)	80	82/8
6	L3	K ₂ CO ₃ (1.0)	—	90	77/7	17	L3	K ₂ CO ₃ (1.0)	AgCl (0.1)	60	81/8
7	L3	K ₂ CO ₃ (2.0)	—	90	72/10	18 ^b	L3	K ₂ CO ₃ (1.0)	AgCl (0.1)	60	80/8
8	L3	KOH (1.0)	—	90	69/6	19 ^c	L3	K ₂ CO ₃ (1.0)	AgCl (0.1)	60	82/7
9	L3	K ₂ CO ₃ (1.0)	AgCl (0.1)	90	82/8	20 ^d	L2	K ₂ CO ₃ (1.0)	AgCl (0.1)	60	25/1
10	L3	K ₂ CO ₃ (1.0)	AgOAc (0.1)	90	50/5	21^e	L3	K₂CO₃ (1.0)	AgCl (0.1)	60	85/7^f
11	L2	K ₂ CO ₃ (1.0)	Ag ₂ CO ₃ (0.1)	90	63/7	22 ^g	L3	K ₂ CO ₃ (1.0)	AgCl (0.1)	60	65/5

^a Reaction conditions: **1a** (0.2 mmol, 1.0 equiv.), 4-iodo-3-nitroanisole (0.4 mmol, 2.0 equiv.), Pd(OAc)₂ (0.02 mmol, 10 mol%), ligand (0.04 mmol, 20 mol%), base, Ag salt, HFIP (2 mL, 0.1 mol·L⁻¹), 24 h. The yields were determined by ¹H NMR using CH₂Br₂ as the internal standard. ^b Under N₂ atmosphere. ^c Pd(OTFA)₂ used instead of Pd(OAc)₂. ^d PdCl₂ used instead of Pd(OAc)₂. ^e 12 h. ^f 1-Methoxy-3-nitrobenzene (0.184 mmol) detected by ¹H NMR. ^g 6 h.

Table 2 Substrate scope of 2-aryl phenols^a

2a , 75%, 70% ^b	2b , 54%	2c , 64%	2d , 68%	2e , 71% ^c	2f , 65%	2g , 68%	2h , 63%	2i , 64%	2j , 75% ^c	2k , 75%
2l , 75%	2m , 60%	2n , 81%	2o , 80%	2p , 80%	2q , 57%	2r , 77%				

^a Reaction conditions: **1** (0.2 mmol), 4-iodo-3-nitroanisole (0.4 mmol), Pd(OAc)₂ (0.02 mmol, 10 mol%), **L3** (0.04 mmol, 20 mol%), AgCl (0.02 mmol, 10 mol%), K₂CO₃ (0.2 mmol, 1.0 equiv.), HFIP (2 mL), 12 h, 60 °C. Isolated yields. Generally less than 10% of dibenzofuran side products were detected. ^b 1 mmol of scale. ^c 90 °C, 24 h.

Table 3 Substrate scope of 2-phenoxyacetic acids^a

Reaction scheme showing the synthesis of various 2-iodo-3-substituted anisole derivatives (5a-5q) from 2-phenylphenol (4) and 4-iodo-3-nitroanisole (2.0 equiv.) under the following conditions:

Reagents: $\text{Pd}(\text{OAc})_2$ (10 mol%), **L1** (20 mol%), NaOAc (0.5 equiv.), HFIP, 90 °C, 48 h; MeI , K_2CO_3 , acetone, 60 °C, 2 h.

Products and yields (mono/di ratio):

- 5a**: 75% (mono/di = 1.78/1)^b, 71% (mono/di = 1.84/1)^{b,c}
- 5b**: 73%^b
- 5c**: 62%
- 5d**: 67%
- 5e**: 67%
- 5f**: 65%
- 5g**: 89%^b
- 5h**: 53%
- 5i**: 66%^d
- 5j**: 64%^d
- 5k**: 72% (mono/di = 1.67/1)^b
- 5l**: 73% (mono/di = 2.32/1)
- 5m**: 53% (mono/di = 7.83/1)
- 5n**: 62% (mono/di = 2.1/1)
- 5o**: 54% (mono/di = 6.71/1)
- 5p**: 60% (mono/di = 5.67/1)
- 5q**: 51% (mono/di = 1.83/1)

^a Reaction conditions: **4** (0.2 mmol, 1.0 equiv.), 4-iodo-3-nitroanisole (0.4 mmol, 2.0 equiv.), Pd(OAc)₂ (0.02 mmol, 10 mol%), **L1** (0.04 mmol, 20 mol%), NaOAc (0.1 mmol, 50 mol%), HFIP (2 mL, 0.1 mol·L⁻¹), 48 h, 90 °C. Isolated yields. ^b 24 h. ^c 1 mmol scale. ^d *N*-Ac-Leu-OH (0.04 mmol) used instead of **L1**.

as withdrawing groups at either the *ortho*- (**5b**~**5f**), *meta*- (**5g**~**5j**) or *para*-position (**5k**~**5q**) were well tolerated, leading to generally good yields of desired products. Finally, the directing group could be removed in a one-pot sequence after *ortho*-C—H iodination (Eq. 1).

Based on our results and previous works, the potential catalytic cycle for above C—H iodination is proposed (Scheme 2). First, ligand exchange would generate active Pd catalyst **A**, which coordinates with the deprotonated 2-phenyl phenol to afford complex **B**. A six-membered palladacycle **C** is then produced after C—H bond cleavage possibly through an electrophilic metalation deprotonation process. Oxidative addition with 4-iodo-3-nitroanisole gives the Pd(IV) intermediate **D**, which prefers to undergo C—I reductive elimination to afford product **2a** and complex **E**.^[14] The less preferred C—O reductive elimination leads to side product **3a**. Finally, protonolysis of complex **E** will regenerate active Pd(II) catalyst **A** and 3-nitroanisole. Notably, although the generation of 3-nitroanisole is consistent with the metathesis reaction,^[8] the proton in protonolysis is more likely from the acidic solvent HFIP rather than substrate **1a**. Therefore, the overall reaction could be considered as a formal metathesis reaction.

3 Conclusions

In summary, we have developed the Pd(II)-catalyzed

Scheme 2 Proposed catalytic cycle

C—H iodination of free 2-aryl phenols and 2-phenoxyacetic acids using 4-iodo-3-nitroanisole as the mild iodinating reagent. Excellent site-selectivity and good functional group tolerance were obtained with a range of phenol derivatives, demonstrating that this formal metathesis C—H iodination method is potentially applicable for iodinating other electron rich arenes. Further study on developing

challenging C—H iodination reactions using this method is underway in our laboratory.

4 Experimental section

4.1 General experimental information

Unless otherwise noted, commercially available reagents were purchased from commercial suppliers (such as Adamas, Strem, J&K Chemical Co., Energy Chemical, etc.), and used as received. Solvents were generally dried over 4 Å molecular sieves. Hexafluoroisopropanol (HFIP) was distilled over 3 Å molecular sieves before use. The reaction vessels used for C—H functionalization were 38 mL sealed tube or 50 mL Schlenk tube (Synthware). ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra were recorded on a Bruker AVANCE III spectrometer (400, 101 and 376 MHz, respectively) or a JEOL ECZ600S spectrometer (600 and 151 MHz, respectively). Chemical shifts are reported referenced to CDCl_3 (δ 7.26), tetramethylsilane (TMS, δ 0.00), acetone- d_6 (δ 2.05) for ^1H NMR, or CDCl_3 (δ 77.16) for ^{13}C NMR. High-resolution mass spectra (HRMS) were obtained on an Impact II UHR-TOF mass spectrometry equipped with an ESI source from a Bruker at Fujian Institute of Research on the Structure of Matter. The gas chromatography-mass spectrometry (GC-MS) spectra were measured using a TRACE1300-ISQD300 (Thermo) with a flame ionization detector. 2-Aryl phenols^[10] and 2-phenoxyacetic acids^[15] were known compounds and prepared according to the literature procedure.

4.2 General procedure for C—H iodination of 2-aryl phenols

An oven-dried 38 mL sealed tube (with a Teflon cap) equipped with a magnetic stir bar was charged with compound **1** (0.2 mmol, 1.0 equiv.), 4-iodo-3-nitroanisole (112 mg, 0.4 mmol, 2.0 equiv.), $\text{Pd}(\text{OAc})_2$ (4.5 mg, 0.02 mmol, 10 mol%), ligand (5-bromopyridin-2-ol, 6.9 mg, 0.04 mmol, 20 mol%), AgCl (2.86 mg, 0.02 mmol, 10 mol%) and K_2CO_3 (27.64 mg, 0.2 mmol, 1.0 equiv.) sequentially. HFIP (2.0 mL) was added to the mixture along the inside wall of the tube. The tube was then capped and placed into a preheated oil bath (60 °C) or hotplate (60 °C) (Note: good stirring was important for reproducibility, and splashes on the wall of the tube might lead to lower yields). The reaction was stirred for 12 h and cooled to room temperature. Then the reaction solution was treated with concentrated HCl (50 μL) and stirred for another 10 min. The crude reaction mixture was diluted with ethyl acetate (EA, 5 mL) and filtered through a short pad of Celite. The sealed tube and Celite pad were washed with an additional 20 mL of EA. The filtrate was concentrated *in vacuo*, and the resulting residue was purified by flash silica gel chromatography or preparative thin layer chromatography using petroleum ether/ethyl acetate (PE/EA, $V/V=30/1$) as the eluent.

2'-Iodo-[1,1'-biphenyl]-2-ol (**2a**): Yellow oil, 44.5 mg, 75% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, $J=7.8$ Hz, 1H), 7.49~7.42 (m, 1H), 7.36~7.31 (m, $J=7.5$, 4.7,

1.6 Hz, 2H), 7.15~7.07 (m, 2H), 7.01 (t, $J=8.5$ Hz, 2H), 4.78 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 141.9, 139.8, 131.1, 131.0, 130.5, 130.0, 128.9, 120.6, 116.0, 100.8. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_8\text{IO}$ [$\text{M}-\text{H}^+$] 294.9625, found 294.9626.

2'-Iodo-5'-methyl-[1,1'-biphenyl]-2-ol (**2b**): Yellow oil, 33.5 mg, 54% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (d, $J=8.0$ Hz, 1H), 7.32 (td, $J=7.8$, 1.8 Hz, 1H), 7.16 (d, $J=2.4$ Hz, 1H), 7.09 (dd, $J=8.0$, 1.6 Hz, 1H), 7.00 (t, $J=7.6$ Hz, 2H), 6.94 (dd, $J=8.1$, 2.2 Hz, 1H), 4.76 (s, 1H), 2.34 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.2, 141.6, 139.6, 139.0, 132.0, 131.06, 131.02, 130.4, 129.9, 120.6, 115.9, 96.5, 21.1. HRMS (ESI-TOF) calcd for $\text{C}_{13}\text{H}_{10}\text{IO}$ [$\text{M}-\text{H}^+$] 308.9782, found 308.9784.

2'-Iodo-5'-methoxy-[1,1'-biphenyl]-2-ol (**2c**): Yellow solid, 41.7 mg, 64% yield. m.p. 82.9~84.3 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.84 (d, $J=8.7$ Hz, 1H), 7.33 (td, $J=7.8$, 1.8 Hz, 1H), 7.10 (dd, $J=7.5$, 1.8 Hz, 1H), 7.01 (t, $J=8.1$ Hz, 3H), 6.90 (d, $J=3.0$ Hz, 1H), 4.81 (s, 1H), 3.80 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 160.3, 152.1, 142.7, 140.4, 130.9, 130.3, 130.0, 120.6, 116.62, 116.59, 116.0, 88.9, 55.6. HRMS (ESI-TOF) calcd for $\text{C}_{13}\text{H}_{10}\text{IO}_2$ [$\text{M}-\text{H}^+$] 324.9731, found 324.9732.

5'-Fluoro-2'-iodo-[1,1'-biphenyl]-2-ol (**2d**): Yellow oil, 42.7 mg, 68% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.92 (dd, $J=8.7$, 5.6 Hz, 1H), 7.38~7.31 (m, 1H), 7.08 (dd, $J=9.4$, 2.5 Hz, 2H), 7.02 (t, $J=7.5$ Hz, 1H), 6.98 (d, $J=8.2$ Hz, 1H), 6.88 (td, $J=8.4$, 3.0 Hz, 1H), 4.78 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 163.1 (d, $J_{\text{C-F}}=249.6$ Hz), 152.0, 144.2 (d, $J_{\text{C-F}}=7.7$ Hz), 140.9 (d, $J_{\text{C-F}}=8.0$ Hz), 130.4, 130.3, 130.2 (d, $J_{\text{C-F}}=1.4$ Hz), 120.8, 118.5 (d, $J_{\text{C-F}}=22.0$ Hz), 117.4 (d, $J_{\text{C-F}}=21.6$ Hz), 116.2, 93.6 (d, $J_{\text{C-F}}=3.4$ Hz); ^{19}F NMR (376 MHz, CDCl_3) δ : -113.35. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{FIO}$ [$\text{M}-\text{H}^+$] 312.9531, found 312.9531.

5'-Chloro-2'-iodo-[1,1'-biphenyl]-2-ol (**2e**): Reaction time 24 h; Yellow solid, 47.1 mg, 71% yield. m.p. 107.9~109.2 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.89 (d, $J=8.5$ Hz, 1H), 7.36~7.31 (m, 2H), 7.13~7.05 (m, 2H), 7.01 (t, $J=7.4$ Hz, 1H), 6.97 (d, $J=8.2$ Hz, 1H), 4.77 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.0, 144.0, 140.7, 135.0, 131.1, 130.41, 130.35, 130.1, 130.0, 120.9, 116.2, 97.8. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{ClIO}$ [$\text{M}-\text{H}^+$] 328.9236, found 328.9236.

2'-Iodo-4'-methyl-[1,1'-biphenyl]-2-ol (**2f**): Yellow oil, 40.3 mg, 65% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (s, 1H), 7.33 (td, $J=7.8$, 1.8 Hz, 1H), 7.27 (d, $J=7.6$ Hz, 1H), 7.21 (d, $J=7.7$ Hz, 1H), 7.09 (dd, $J=7.5$, 1.5 Hz, 1H), 7.00 (t, $J=7.5$ Hz, 2H), 4.77 (s, 1H), 2.39 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.3, 140.30, 140.25, 138.7, 130.9, 130.70, 130.65, 129.9, 129.7, 120.5, 115.8, 100.7, 20.74. HRMS (ESI-TOF) calcd for $\text{C}_{13}\text{H}_{10}\text{IO}$ [$\text{M}-\text{H}^+$] 308.9782, found 308.9783.

4'-Fluoro-2'-iodo-[1,1'-biphenyl]-2-ol (**2g**): Yellow oil, 42.7 mg, 68% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.72 (dd, $J=8.1$, 2.6 Hz, 1H), 7.36~7.27 (m, 2H), 7.18 (td, $J=8.3$, 2.6 Hz, 1H), 7.07 (dd, $J=7.5$, 1.5 Hz, 1H), 7.04~6.95

(m, 2H), 4.72 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 161.8 (d, $J_{\text{C-F}}=253.0$ Hz), 152.3, 138.2 (d, $J_{\text{C-F}}=3.6$ Hz), 131.8 (d, $J_{\text{C-F}}=8.2$ Hz), 130.8, 130.2, 130.1, 126.7 (d, $J_{\text{C-F}}=23.5$ Hz), 120.7, 116.0 (d, $J_{\text{C-F}}=21.0$ Hz), 116.0, 100.3 (d, $J_{\text{C-F}}=8.1$ Hz); ^{19}F NMR (376 MHz, CDCl_3) δ : -112.22. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{FIO}$ [$\text{M}-\text{H}^+$] 312.9531, found 312.9531.

4'-Chloro-2'-iodo-[1,1'-biphenyl]-2-ol (**2h**): Yellow oil, 41.4 mg, 63% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.99 (d, $J=2.1$ Hz, 1H), 7.43 (dd, $J=8.2, 2.2$ Hz, 1H), 7.33 (td, $J=8.2, 1.8$ Hz, 1H), 7.25 (d, $J=8.1$ Hz, 1H), 7.07 (dd, $J=7.6, 1.8$ Hz, 1H), 7.03~6.94 (m, 2H), 4.72 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.1, 140.7, 139.1, 134.8, 131.6, 130.6, 130.3, 130.0, 129.0, 120.8, 116.1, 100.8. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{ClIO}$ [$\text{M}-\text{H}^+$] 328.9236, found 328.9236.

4'-(*tert*-Butyl)-2'-iodo-[1,1'-biphenyl]-2-ol (**2i**): Yellow oil, 45.1 mg, 64% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.01 (d, $J=2.0$ Hz, 1H), 7.49 (dd, $J=8.0, 2.0$ Hz, 1H), 7.34 (td, $J=7.7, 1.8$ Hz, 1H), 7.27 (d, $J=7.9$ Hz, 1H), 7.11 (dd, $J=7.8, 1.8$ Hz, 1H), 7.04~6.98 (m, 2H), 4.79 (s, 1H), 1.38 (s, 9H); ^{13}C NMR (101 MHz, CDCl_3) δ : 153.5, 152.4, 138.7, 137.0, 130.9, 130.64, 130.62, 129.9, 126.2, 120.5, 115.8, 100.9, 34.7, 31.3. HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{16}\text{IO}$ [$\text{M}-\text{H}^+$] 351.0251, found 351.0253.

2'-Iodo-[1,1':4',1''-terphenyl]-2-ol (**2j**): Reaction time 24 h; Yellow solid, 55.6 mg, 75% yield. m.p. 119.2~120.7 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 8.23 (d, $J=1.8$ Hz, 1H), 7.67 (dd, $J=7.9, 1.9$ Hz, 1H), 7.61 (d, $J=7.3$ Hz, 2H), 7.48 (t, $J=7.5$ Hz, 2H), 7.41 (td, $J=8.2, 2.0$ Hz, 2H), 7.35 (td, $J=8.0, 1.6$ Hz, 1H), 7.15 (dd, $J=7.5, 1.7$ Hz, 1H), 7.06~7.00 (m, 2H), 4.79 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.3, 143.1, 140.6, 139.0, 138.4, 131.2, 130.7, 130.6, 130.1, 129.1, 128.2, 127.6, 127.3, 120.7, 116.0, 101.2. HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{12}\text{IO}$ [$\text{M}-\text{H}^+$] 370.9938, found 370.9939.

5-Fluoro-2'-iodo-[1,1'-biphenyl]-2-ol (**2k**): Yellow oil, 46.8 mg, 75% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, $J=8.0$ Hz, 1H), 7.46 (t, $J=7.5$ Hz, 1H), 7.31 (dd, $J=7.6, 1.4$ Hz, 1H), 7.13 (td, $J=7.9, 1.6$ Hz, 1H), 7.02 (td, $J=8.5, 3.0$ Hz, 1H), 6.93 (dd, $J=8.9, 4.7$ Hz, 1H), 6.82 (dd, $J=8.6, 3.0$ Hz, 1H), 4.63 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 156.7 (d, $J_{\text{C-F}}=239.1$ Hz), 148.3 (d, $J_{\text{C-F}}=2.2$ Hz), 140.9 (d, $J_{\text{C-F}}=1.5$ Hz), 140.0, 131.7 (d, $J_{\text{C-F}}=8.0$ Hz), 131.0, 130.3, 129.0, 117.0 (d, $J_{\text{C-F}}=8.2$ Hz), 116.8 (d, $J_{\text{C-F}}=23.5$ Hz), 116.5 (d, $J_{\text{C-F}}=22.9$ Hz), 100.2; ^{19}F NMR (376 MHz, CDCl_3) δ : -123.90. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{FIO}$ [$\text{M}-\text{H}^+$] 312.9531, found 312.9531.

5-Chloro-2'-iodo-[1,1'-biphenyl]-2-ol (**2l**): Yellow oil, 49.4 mg, 75% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, $J=7.9$ Hz, 1H), 7.46 (td, $J=7.5, 1.2$ Hz, 1H), 7.32~7.25 (m, 2H), 7.13 (td, $J=7.7, 1.7$ Hz, 1H), 7.08 (d, $J=2.6$ Hz, 1H), 6.92 (d, $J=8.7$ Hz, 1H), 4.74 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.0, 140.6, 140.0, 132.3, 131.1, 130.4, 130.1, 129.8, 129.0, 125.3, 117.4, 100.2. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{ClIO}$ [$\text{M}-\text{H}^+$] 328.9236, found

328.9236.

2'-Iodo-4-methyl-[1,1'-biphenyl]-2-ol (**2m**): Yellow oil, 37.2 mg, 60% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.99 (d, $J=7.9$ Hz, 1H), 7.44 (t, $J=7.5$ Hz, 1H), 7.32 (dd, $J=7.6, 1.6$ Hz, 1H), 7.10 (td, $J=7.7, 1.6$ Hz, 1H), 6.98 (d, $J=8.0$ Hz, 1H), 6.83 (d, $J=7.2$ Hz, 2H), 4.64 (s, 1H), 2.38 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 151.9, 142.0, 140.3, 139.8, 131.3, 130.2, 129.9, 128.8, 128.2, 121.5, 116.5, 101.1, 21.5. HRMS (ESI-TOF) calcd for $\text{C}_{13}\text{H}_{10}\text{IO}$ [$\text{M}-\text{H}^+$] 308.9782, found 308.9781.

2'-Iodo-4-(trifluoromethyl)-[1,1'-biphenyl]-2-ol (**2n**): Yellow oil, 59.1 mg, 81% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.03 (d, $J=8.0$ Hz, 1H), 7.49 (t, $J=7.1$ Hz, 1H), 7.32~7.25 (m, 3H), 7.22 (d, $J=7.7$ Hz, 1H), 7.16 (td, $J=7.8, 1.6$ Hz, 1H), 4.99 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.5, 140.5, 140.1, 134.2, 132.2 (q, $J_{\text{C-F}}=32.8$ Hz), 131.1, 130.9, 130.6, 129.1, 123.9 (q, $J_{\text{C-F}}=272.3$ Hz), 117.4 (q, $J_{\text{C-F}}=3.7$ Hz), 113.3 (q, $J_{\text{C-F}}=3.9$ Hz), 99.9; ^{19}F NMR (376 MHz, CDCl_3) δ : -62.69. HRMS (ESI-TOF) calcd for $\text{C}_{13}\text{H}_7\text{F}_3\text{IO}$ [$\text{M}-\text{H}^+$] 362.9499, found 362.9499.

4-Fluoro-2'-iodo-[1,1'-biphenyl]-2-ol (**2o**): Yellow oil, 50.5 mg, 80% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, $J=8.0$ Hz, 1H), 7.46 (td, $J=7.5, 1.2$ Hz, 1H), 7.31 (dd, $J=7.6, 1.6$ Hz, 1H), 7.12 (td, $J=7.8, 1.6$ Hz, 1H), 7.04 (dd, $J=9.2, 6.5$ Hz, 1H), 6.75~6.70 (m, 2H), 4.88 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 163.6 (d, $J_{\text{C-F}}=246.4$ Hz), 153.5 (d, $J_{\text{C-F}}=12.4$ Hz), 140.9, 140.0, 131.5, 131.4, 130.3, 129.0, 127.1 (d, $J_{\text{C-F}}=3.2$ Hz), 107.8 (d, $J_{\text{C-F}}=21.7$ Hz), 103.5 (d, $J_{\text{C-F}}=25.0$ Hz), 101.2; ^{19}F NMR (376 MHz, CDCl_3) δ : -111.37. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{FIO}$ [$\text{M}-\text{H}^+$] 312.9531, found 312.9531.

4-Chloro-2'-iodo-[1,1'-biphenyl]-2-ol (**2p**): Yellow oil, 53.2 mg, 80% yield. ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, $J=8.0$ Hz, 1H), 7.46 (td, $J=7.5, 1.3$ Hz, 1H), 7.29 (dd, $J=7.6, 1.4$ Hz, 1H), 7.12 (td, $J=7.9, 1.6$ Hz, 1H), 7.05~6.97 (m, 3H), 4.82 (s, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.9, 140.7, 140.0, 135.1, 131.3, 131.1, 130.4, 129.5, 129.0, 121.0, 116.4, 100.7. HRMS (ESI-TOF) calcd for $\text{C}_{12}\text{H}_7\text{ClIO}$ [$\text{M}-\text{H}^+$] 328.9236, found 328.9235.

2'-Iodo-5'-methoxy-5-methyl-[1,1'-biphenyl]-2-ol (**2q**): Yellow oil, 39 mg, 57% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.82 (d, $J=8.8$ Hz, 1H), 7.13 (d, $J=8.2$ Hz, 1H), 6.89 (d, $J=8.5$ Hz, 3H), 6.70 (dd, $J=8.7, 2.9$ Hz, 1H), 4.64 (s, 1H), 3.80 (s, 3H), 2.32 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 160.3, 149.8, 143.0, 140.3, 130.7, 130.6, 130.5, 129.8, 116.6, 116.5, 115.8, 88.9, 55.6, 20.7. HRMS (ESI-TOF) calcd for $\text{C}_{14}\text{H}_{12}\text{IO}_2$ [$\text{M}-\text{H}^+$] 338.9888, found 338.9890.

2'-Iodo-4'-methyl-4-(trifluoromethyl)-[1,1'-biphenyl]-2-ol (**2r**): Yellow oil, 58.5 mg, 77% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (s, 1H), 7.29 (d, $J=7.7$ Hz, 1H), 7.25 (d, $J=7.0$ Hz, 2H), 7.22~7.15 (m, 2H), 4.96 (s, 1H), 2.39 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.7, 141.0, 140.6, 137.3, 134.1, 132.0 (q, $J_{\text{C-F}}=32.6$ Hz), 131.3, 130.5, 130.0, 123.9 (q, $J_{\text{C-F}}=272.3$ Hz), 117.3 (q, $J_{\text{C-F}}=3.9$ Hz), 113.1 (q, $J_{\text{C-F}}=3.8$ Hz), 99.8, 20.8; ^{19}F NMR (376 MHz, CDCl_3) δ : -62.66. HRMS (ESI-TOF) calcd for $\text{C}_{14}\text{H}_9\text{F}_3\text{IO}$ [$\text{M}-\text{H}^+$] 362.9499, found 362.9499.

$F_3IO [M-H]^+$ 376.9656, found 376.9656.

4.3 General procedure for C—H iodination of 2-phenoxyacetic acids

An oven-dried 38 mL sealed tube (with a Teflon cap) equipped with a magnetic stir bar was charged with compound **4** (0.2 mmol, 1.0 equiv.), 4-iodo-3-nitroanisole (112 mg, 0.4 mmol, 2.0 equiv.), $Pd(OAc)_2$ (4.5 mg, 0.02 mmol, 10 mol%), *N*-Formyl-Gly-OH (4.1 mg, 0.04 mmol, 20 mol%), and NaOAc (8.2 mg, 0.1 mmol, 0.5 equiv.) sequentially. HFIP (2.0 mL) was added to the mixture along the inside wall of the tube. The tube was then capped and placed into a preheated oil bath (90 °C) or hotplate (90 °C) (Note: good stirring was important for reproducibility, splashes on the wall of the tube might lead to lower yields). The reaction was stirred for 48 h and cooled to room temperature. Volatile matter was removed under reduced pressure and the residue was re-dissolved in acetone (2 mL). To the solution was added K_2CO_3 (82 mg, 0.6 mmol) and MeI (24 μ L, 0.4 mmol). The reaction was stirred at 60 °C for 2 h and cooled to room temperature. The crude reaction mixture was diluted with EA (5 mL) and filtered through a short pad of Celite. The sealed tube and Celite pad were washed with an additional 20 mL of EA. The filtrate was concentrated *in vacuo* to give the crude product, which was purified by medium pressure liquid chromatography (MPLC) with a silica gel column to afford pure products, using PE/EA (*V/V*=9/1) as the eluent.

Methyl 2-(2-iodophenoxy)acetate (**5a_{mono}**): Reaction time 24 h; colourless oil, 28.0 mg, 48% yield. 1H NMR (400 MHz, $CDCl_3$) δ : 7.80 (dd, $J=7.8, 1.6$ Hz, 1H), 7.31~7.27 (m, 1H), 6.80~6.69 (m, 2H), 4.70 (s, 2H), 3.81 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 169.0, 156.8, 140.0, 129.6, 123.8, 112.5, 86.6, 66.4, 52.5. HRMS (ESI-TOF) calcd for $C_9H_9IO_3Na^+$ [$M+Na^+$] 314.9488, found 314.9489.

Methyl 2-(2,6-diiodophenoxy)acetate (**5a_{di}**): Reaction time 24 h; white solid, 22.6 mg, 27% yield. m.p. 86.2~87.9 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.76 (d, $J=7.8$ Hz, 2H), 6.59 (t, $J=7.9$ Hz, 1H), 4.61 (s, 2H), 3.88 (s, 3H). ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.2, 156.7, 140.2, 128.6, 90.6, 68.9, 52.5. HRMS (ESI-TOF) calcd for $C_9H_8I_2O_3Na^+$ [$M+Na^+$] 440.8455, found 440.8456.

Methyl 2-(2-iodo-6-methylphenoxy)acetate (**5b**): Reaction time 24 h; colorless oil, 44.7 mg, 73% yield. 1H NMR (400 MHz, $CDCl_3$) δ : 7.60 (d, $J=7.8$ Hz, 1H), 7.14 (d, $J=7.5$ Hz, 1H), 6.78 (t, $J=7.7$ Hz, 1H), 4.51 (s, 2H), 3.85 (s, 3H), 2.34 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.9, 156.1, 137.5, 132.6, 131.8, 126.7, 91.4, 69.1, 52.4, 17.1. HRMS (ESI-TOF) calcd for $C_{10}H_{11}IO_3Na^+$ [$M+Na^+$] 328.9645, found 328.9643.

Methyl 2-(2-iodo-6-methoxyphenoxy)acetate (**5c**): Yellow oil, 39.9 mg, 62% yield. 1H NMR (400 MHz, $CDCl_3$) δ : 7.33 (d, $J=7.9$ Hz, 1H), 6.87 (d, $J=8.1$ Hz, 1H), 6.80 (t, $J=8.0$ Hz, 1H), 4.60 (s, 2H), 3.82 (s, 3H), 3.81 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 169.0, 151.9, 146.7, 130.5,

126.1, 112.6, 91.7, 68.8, 55.8, 51.9. HRMS (ESI-TOF) calcd for $C_{10}H_{11}IO_4Na^+$ [$M+Na^+$] 344.9594, found 344.9594.

Methyl 2-(2-fluoro-6-iodophenoxy)acetate (**5d**): Yellow oil, 41.5 mg, 67% yield. 1H NMR (400 MHz, $CDCl_3$) δ : 7.54 (d, $J=8.0$ Hz, 1H), 7.13~6.99 (m, 1H), 6.80 (td, $J=8.1, 5.2$ Hz, 1H), 4.72 (s, 2H), 3.82 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.8, 154.1 (d, $J_{C-F}=250.8$ Hz), 145.8 (d, $J_{C-F}=11.6$ Hz), 134.8 (d, $J_{C-F}=3.4$ Hz), 126.1 (d, $J_{C-F}=7.8$ Hz), 117.5 (d, $J_{C-F}=19.8$ Hz), 91.5, 69.7 (d, $J_{C-F}=6.3$ Hz), 52.4; ^{19}F NMR (376 MHz, $CDCl_3$) δ : -126.58. HRMS (ESI-TOF) calcd for $C_9H_8FIO_3Na^+$ [$M+Na^+$] 332.9394, found 332.9395.

Methyl 2-(2-chloro-6-iodophenoxy)acetate (**5e**): Colorless oil, 43.8 mg, 67% yield. 1H NMR (400 MHz, $CDCl_3$) δ : 7.68 (d, $J=7.9$ Hz, 1H), 7.36 (d, $J=8.0$ Hz, 1H), 6.82 (t, $J=8.0$ Hz, 1H), 4.61 (s, 2H), 3.86 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.3, 153.6, 138.2, 131.0, 127.8, 127.4, 92.4, 69.0, 52.5. HRMS (ESI-TOF) calcd for $C_9H_8ClIO_3Na^+$ [$M+Na^+$] 348.9099, found 348.9099.

Methyl 2-(2-bromo-6-iodophenoxy)acetate (**5f**): Colorless oil, 48.3 mg, 65% yield. 1H NMR (400 MHz, $CDCl_3$) δ : 7.72 (d, $J=7.8$ Hz, 1H), 7.53 (d, $J=7.9$ Hz, 1H), 6.75 (t, $J=7.9$ Hz, 1H), 4.61 (s, 2H), 3.86 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.3, 154.4, 139.0, 134.1, 127.9, 116.7, 92.2, 68.9, 52.5. HRMS (ESI-TOF) calcd for $C_9H_8BrIO_3Na^+$ [$M+Na^+$] 392.8594, found 392.8594.

Methyl 2-(2-iodo-5-methylphenoxy)acetate (**5g**): Reaction time 24 h; white solid, 54.5 mg, 89% yield. m.p. 62.1~63.5 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.63 (d, $J=7.9$ Hz, 1H), 6.59 (d, $J=8.0$ Hz, 1H), 6.54 (s, 1H), 4.68 (s, 2H), 3.81 (s, 3H), 2.29 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 169.1, 156.7, 140.0, 139.5, 124.8, 113.7, 82.5, 66.4, 52.4, 21.5. HRMS (ESI-TOF) calcd for $C_{10}H_{11}IO_3Na^+$ [$M+Na^+$] 328.9645, found 328.9648.

Methyl 2-(2-iodo-5-(trifluoromethyl)phenoxy)acetate (**5h**): White solid, 38.2 mg, 53% yield. m.p. 42.1~43.6 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.92 (d, $J=8.0$ Hz, 1H), 7.02 (d, $J=8.1$ Hz, 1H), 6.89 (s, 1H), 4.75 (s, 2H), 3.82 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.3, 157.1, 140.6, 132.1 (q, $J_{C-F}=33.0$ Hz), 123.7 (q, $J_{C-F}=272.5$ Hz), 120.2 (q, $J_{C-F}=3.8$ Hz), 108.8 (q, $J_{C-F}=3.8$ Hz), 91.3, 66.3, 52.7; ^{19}F NMR (376 MHz, $CDCl_3$) δ : -62.82. HRMS (ESI-TOF) calcd for $C_{10}H_8F_3IO_3Na^+$ [$M+Na^+$] 382.9362, found 382.9362.

Methyl 2-(5-fluoro-2-iodophenoxy)acetate (**5i**): White solid, 41.0 mg, 66% yield. m.p. 60.6~62.3 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.72 (dd, $J=8.6, 6.5$ Hz, 1H), 6.55 (td, $J=8.3, 2.6$ Hz, 1H), 6.47 (dd, $J=10.3, 2.7$ Hz, 1H), 4.68 (s, 2H), 3.81 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 168.4, 163.8 (d, $J=247.6$ Hz), 157.8 (d, $J=9.9$ Hz), 140.1 (d, $J=9.4$ Hz), 110.7 (d, $J=21.9$ Hz), 101.1 (d, $J=26.4$ Hz), 79.4 (d, $J=3.7$ Hz), 66.4, 52.6; ^{19}F NMR (376 MHz, $CDCl_3$) δ : -110.72. HRMS (ESI-TOF) calcd for $C_9H_8FIO_3Na^+$ [$M+Na^+$] 332.9394, found 332.9397.

Methyl 2-(5-chloro-2-iodophenoxy)acetate (**5j**): Yellow solid, 41.8 mg, 64% yield. m.p. 70.0~71.7 °C; 1H NMR

(600 MHz, CDCl_3) δ : 7.69 (d, $J=8.4$ Hz, 1H), 6.76 (dd, $J=8.4, 2.1$ Hz, 1H), 6.69 (d, $J=2.1$ Hz, 1H), 4.69 (s, 2H), 3.81 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.4, 157.4, 140.3, 135.2, 123.9, 113.1, 83.9, 66.3, 52.6. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_8\text{ClIO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 348.9099, found 348.9097.

Methyl 2-(2-iodo-4-methylphenoxy)acetate (**5k_{mono}**): Reaction time 24 h; Yellow oil, 27.6 mg, 45% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.62 (s, 1H), 7.06 (d, $J=8.2$ Hz, 1H), 6.63 (d, $J=8.3$ Hz, 1H), 4.66 (s, 2H), 3.79 (s, 3H), 2.25 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.2, 154.8, 140.3, 133.5, 130.0, 112.6, 86.5, 66.7, 52.4, 20.1. HRMS (ESI-TOF) calcd for $\text{C}_{10}\text{H}_{11}\text{IO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 328.9645, found 328.9646.

Methyl 2-(2,6-diiodo-4-methylphenoxy)acetate (**5k_{di}**): White solid, 23.4 mg, 27% yield. m.p. 122.3~123.4 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.57 (s, 2H), 4.57 (s, 2H), 3.87 (s, 3H), 2.24 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.3, 154.5, 140.5, 138.7, 90.0, 69.0, 52.5, 19.8. HRMS (ESI-TOF) calcd for $\text{C}_{10}\text{H}_{10}\text{I}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 454.8612, found 454.8609.

Methyl 2-(2-iodo-4-methoxyphenoxy)acetate (**5l_{mono}**): White solid, 32.9 mg, 51% yield. m.p. 50.1~51.4 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.33 (d, $J=2.9$ Hz, 1H), 6.82 (dd, $J=8.9, 2.9$ Hz, 1H), 6.73 (d, $J=9.0$ Hz, 1H), 4.62 (s, 2H), 3.79 (s, 3H), 3.74 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.3, 155.3, 151.5, 125.0, 114.9, 114.3, 87.2, 67.6, 56.0, 52.4. HRMS (ESI-TOF) calcd for $\text{C}_{10}\text{H}_{11}\text{IO}_4\text{Na}$ [$\text{M}+\text{Na}^+$] 344.9594, found 344.9594.

Methyl 2-(2,6-diiodo-4-methoxyphenoxy)acetate (**5l_{di}**): White solid, 19.8 mg, 22% yield. m.p. 97.2~98.8 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.29 (s, 2H), 4.55 (s, 2H), 3.87 (s, 3H), 3.74 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.4, 157.4, 150.7, 125.3, 89.6, 69.2, 56.1, 52.5. HRMS (ESI-TOF) calcd for $\text{C}_{10}\text{H}_{10}\text{I}_2\text{O}_4\text{Na}$ [$\text{M}+\text{Na}^+$] 470.8561, found 470.8560.

Methyl 2-(2-iodo-4-(trifluoromethyl)phenoxy)acetate (**5m_{mono}**): White solid, 33.9 mg, 47% yield. m.p. 65.1~66.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.04 (s, 1H), 7.55 (d, $J=7.7$ Hz, 1H), 6.73 (d, $J=8.6$ Hz, 1H), 4.76 (s, 2H), 3.81 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.3, 159.2, 137.1 (q, $J_{\text{C-F}}=3.8$ Hz), 127.0 (q, $J_{\text{C-F}}=3.8$ Hz), 125.7 (q, $J_{\text{C-F}}=33.2$ Hz), 123.2 (q, $J_{\text{C-F}}=271.9$ Hz), 111.5, 86.2, 66.2, 52.7; ^{19}F NMR (376 MHz, CDCl_3) δ : -61.80. HRMS (ESI-TOF) calcd for $\text{C}_{10}\text{H}_8\text{F}_3\text{IO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 382.9362, found 382.9363.

Methyl 2-(2,6-diiodo-4-(trifluoromethyl)phenoxy)acetate (**5m_{di}**): White solid, 5.9 mg, 6% yield. m.p. 67.3~68.8 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.02 (s, 2H), 4.64 (s, 2H), 3.89 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 167.7, 159.7, 137.3 (q, $J_{\text{C-F}}=3.7$ Hz), 130.4 (q, $J_{\text{C-F}}=33.6$ Hz), 121.7 (q, $J_{\text{C-F}}=273.2$ Hz), 90.4, 68.8, 52.7; ^{19}F NMR (376 MHz, CDCl_3) δ : -62.29. HRMS (ESI-TOF) calcd for $\text{C}_{10}\text{H}_7\text{F}_3\text{I}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 508.8329, found 508.8329.

Methyl 2-(4-fluoro-2-iodophenoxy)acetate (**5n_{mono}**): White solid, 26.1 mg, 42% yield. m.p. 54.3~56.6 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.51 (dd, $J=7.5, 2.9$ Hz, 1H),

7.00 (td, $J=9.0, 3.0$ Hz, 1H), 6.70 (dd, $J=9.0, 4.5$ Hz, 1H), 4.65 (s, 2H), 3.80 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.9, 157.6 (d, $J_{\text{C-F}}=245.3$ Hz), 153.6 (d, $J_{\text{C-F}}=2.7$ Hz), 126.6 (d, $J_{\text{C-F}}=25.1$ Hz), 115.8 (d, $J_{\text{C-F}}=22.9$ Hz), 113.3 (d, $J_{\text{C-F}}=8.4$ Hz), 86.4 (d, $J_{\text{C-F}}=8.5$ Hz), 67.2, 52.5; ^{19}F NMR (376 MHz, CDCl_3) δ : -120.45. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_8\text{FIO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 332.9394, found 332.9395.

Methyl 2-(4-fluoro-2,6-diiodophenoxy)acetate (**5n_{di}**): White solid, 17.5 mg, 20% yield. m.p. 62.1~64.6 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.51 (d, $J=7.2$ Hz, 2H), 4.57 (s, 2H), 3.87 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.0, 158.7 (d, $J_{\text{C-F}}=254.2$ Hz), 153.6 (d, $J_{\text{C-F}}=3.5$ Hz), 126.8 (d, $J_{\text{C-F}}=24.6$ Hz), 89.2 (d, $J_{\text{C-F}}=8.7$ Hz), 69.0, 52.5; ^{19}F NMR (376 MHz, CDCl_3) δ : -114.71. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_7\text{FI}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 458.8361, found 458.8361.

Methyl 2-(4-chloro-2-iodophenoxy)acetate (**5o_{mono}**): Colorless oil, 30.7 mg, 47% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (s, 1H), 7.24 (dd, $J=8.8, 2.3$ Hz, 1H), 6.64 (d, $J=8.8$ Hz, 1H), 4.67 (s, 2H), 3.80 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.6, 155.8, 139.1, 129.3, 127.8, 113.0, 86.9, 66.6, 52.6. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_8\text{ClIO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 348.9099, found 348.9098.

Methyl 2-(4-chloro-2,6-diiodophenoxy)acetate (**5o_{di}**): White solid, 6.4 mg, 7% yield. m.p. 83.2~85.6 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.76 (s, 2H), 4.58 (s, 2H), 3.88 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 167.9, 155.8, 139.3, 132.0, 90.1, 68.9, 52.6. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_7\text{ClI}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 474.8065, found 474.8062.

Methyl 2-(4-bromo-2-iodophenoxy)acetate (**5p_{mono}**): White solid, 37.9 mg, 51% yield. m.p. 53.2~54.3 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.91 (d, $J=2.3$ Hz, 1H), 7.38 (dd, $J=8.7, 2.4$ Hz, 1H), 6.59 (d, $J=8.7$ Hz, 1H), 4.68 (s, 2H), 3.80 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.6, 156.2, 141.8, 132.3, 114.9, 113.6, 87.4, 66.5, 52.6. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_8\text{BrIO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 392.8594, found 392.8598.

Methyl 2-(4-bromo-2,6-diiodophenoxy)acetate (**5p_{di}**): White solid, 9.0 mg, 9% yield. m.p. 103.2~105.7 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.89 (s, 2H), 4.58 (s, 2H), 3.87 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 167.9, 156.3, 142.0, 119.4, 90.8, 68.9, 52.6. HRMS (ESI-TOF) calcd for $\text{C}_9\text{H}_7\text{BrI}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 518.7560, found 518.7562.

Methyl 2-((3-iodo-[1,1'-biphenyl]-4-yl)oxy)acetate (**5q_{mono}**): Yellow solid, 24.3 mg, 33% yield. m.p. 86.3~88.6 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.04 (d, $J=2.3$ Hz, 1H), 7.50 (ddd, $J=8.5, 5.4, 1.9$ Hz, 3H), 7.42 (t, $J=7.6$ Hz, 2H), 7.33 (t, $J=7.3$ Hz, 1H), 6.78 (d, $J=8.5$ Hz, 1H), 4.74 (s, 2H), 3.83 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 169.0, 156.3, 139.2, 138.5, 137.1, 129.0, 128.2, 127.5, 127.0, 112.5, 87.0, 66.5, 52.5. HRMS (ESI-TOF) calcd for $\text{C}_{15}\text{H}_{13}\text{IO}_3\text{Na}$ [$\text{M}+\text{Na}^+$] 390.9802, found 390.9801.

Methyl 2-((3,5-diiodo-[1,1'-biphenyl]-4-yl)oxy)acetate (**5q_{di}**): White solid, 17.8 mg, 18% yield. m.p. 78.2~81.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.97 (s, 2H), 7.48

(d, $J=7.4$ Hz, 2H), 7.43 (t, $J=7.4$ Hz, 2H), 7.37 (t, $J=7.1$ Hz, 1H), 4.64 (s, 2H), 3.89 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 168.2, 155.9, 141.9, 138.7, 137.5, 129.1, 128.4, 127.2, 90.6, 69.0, 52.5. HRMS (ESI-TOF) calcd for $\text{C}_{15}\text{H}_{12}\text{O}_3\text{Na} [\text{M}+\text{Na}]^+$ 516.8768, found 516.8769.

4.4 Removal of directing group of iodination product of **4g**

An oven-dried 38 mL sealed tube (with a Teflon cap) equipped with a magnetic stir bar was charged with compound **4g** (0.2 mmol, 1.0 equiv.), 4-iodo-3-nitroanisole (112 mg, 0.4 mmol, 2.0 equiv.), $\text{Pd}(\text{OAc})_2$ (4.5 mg, 0.02 mmol, 10 mol%), *N*-Formyl-Gly-OH (4.1 mg, 0.04 mmol, 20 mol%), and NaOAc (8.2 mg, 0.1 mmol, 0.5 equiv.) sequentially. HFIP (2.0 mL) was added to the mixture along the inside wall of the tube. The tube was then capped and placed into a preheated oil bath (90 °C). The reaction was stirred for 24 h and cooled to room temperature. Volatile matter was removed under reduced pressure and the residue was re-dissolved in toluene (2 mL). To the solution were added DPPA (48 μL , 0.22 mmol) and Et_3N (28 μL , 0.202 mmol). The reaction was stirred at 100 °C for 3 h and cooled to room temperature. To the solution were added glycerol (0.63 mL), EtOH (1.25 mL) and KOH solution ($m/m=1/1$, 0.5 mL). The reaction was stirred at 100 °C for 2 h and cooled to room temperature. The crude reaction mixture was adjusted to pH=5 with 1 mol·L⁻¹ HCl while stirring and then extracted with EA (10 mL×3). The organic phases were combined and dried over Na_2SO_4 . The solvent was removed under reduced pressure and the residue was purified by medium pressure liquid chromatography (MPLC) with a silica gel column to afford compound **6**, using PE/EA ($V/V=9/1$) as the eluent. 2-Iodo-5-methylphenol (**6**):^[16] Colorless oil, 35.0 mg, 75% yield. ^1H NMR (400 MHz, CDCl_3) δ : 7.50 (d, $J=8.1$ Hz, 1H), 6.83 (d, $J=2.0$ Hz, 1H), 6.51 (dd, $J=8.1$, 2.0 Hz, 1H), 5.23 (s, 1H), 2.28 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 154.7, 140.9, 137.9, 123.7, 116.0, 81.7, 21.2.

Supporting Information The Supporting Information containing the NMR spectra of new compounds reported herein is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Bringmann, G.; Gulder, T.; Gulder, T. A.; Breuning, M. *Chem. Rev.* **2011**, *111*, 563.
(b) Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, *102*, 1359.
(c) Qin, L.; Ren, L.; Wan, S.; Liu, G.; Luo, X.; Liu, Z.; Li, F.; Yu, Y.; Liu, J.; Wei, Y. *J. Med. Chem.* **2017**, *60*, 3606.
- [2] Reed, M. A.; Chang, M. T.; Snieckus, V. *Org. Lett.* **2004**, *6*, 2297.
- [3] (a) Liao, G.; Shi, B. *Acta Chim. Sinica* **2015**, *73*, 1283 (in Chinese). (廖港, 史炳峰, 化学学报, **2015**, *73*, 1283.)
(b) Petrone, D. A.; Ye, J.; Lautens, M. *Chem. Rev.* **2016**, *116*, 8003.
(c) Luo, J.; Xu, X.; Zhao, Y.; Liang, H. *Chin. J. Org. Chem.* **2017**, *37*, 2873 (in Chinese).
(d) 骆钧飞, 徐星, 赵延超, 梁洪泽, 有机化学, **2017**, *37*, 2873.)
(e) Das, R.; Kapur, M. *Asian J. Org. Chem.* **2018**, *7*, 1524.
- [4] (a) Giri, R.; Chen, X.; Yu, J. Q. *Angew. Chem., Int. Ed.* **2005**, *44*, 2112.
(b) Wan, X.; Ma, Z.; Li, B.; Zhang, K.; Cao, S.; Zhang, S.; Shi, Z. *J. Am. Chem. Soc.* **2006**, *128*, 7416.
(c) Kalyani, D.; Dick, A. R.; Anani, W. Q.; Sanford, M. S. *Org. Lett.* **2006**, *8*, 2523.
(d) Mei, T.-S.; Giri, R.; Maugel, N.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2008**, *47*, 5215.
(e) Schröder, N.; Wencel-Delord, J.; Glorius, F. *J. Am. Chem. Soc.* **2012**, *134*, 8298.
(f) Wang, X.-C.; Hu, Y.; Bonacorsi, S.; Hong, Y.; Burrell, R.; Yu, J.-Q. *J. Am. Chem. Soc.* **2013**, *135*, 10326.
(g) Urones, B.; Martínez, Á. M.; Rodríguez, N.; Arrayás, R. G.; Carretero, J. C. *Chem. Commun.* **2013**, *49*, 11044.
(h) Chu, L.; Xiao, K.-J.; Yu, J.-Q. *Science* **2014**, *346*, 451.
(i) Gao, D.-W.; Gu, Q.; You, S.-L. *ACS Catal.* **2014**, *4*, 2741.
(j) Lu, C.; Zhang, S.-Y.; He, G.; Nack, W. A.; Chen, G. *Tetrahedron* **2014**, *70*, 4197.
(k) Aihara, Y.; Chatani, N. *ACS Catal.* **2016**, *6*, 4323.
(l) Zhan, B.-B.; Liu, Y.-H.; Hu, F.; Shi, B.-F. *Chem. Commun.* **2016**, *52*, 4934.
(m) Fan, X.-M.; Guo, Y.; Li, Y.-D.; Yu, K.-K.; Liu, H.-W.; Liao, D.-H.; Ji, Y.-F. *Asian J. Org. Chem.* **2016**, *5*, 499.
(n) Li, J.; Cong, W.; Gao, Z.; Zhang, J.; Yang, H.; Jiang, G. *Org. Biomol. Chem.* **2018**, *16*, 3479.
(o) Schreiber, B. S.; Carreira, E. M. *J. Am. Chem. Soc.* **2019**, *141*, 8758.
- [5] Sun, X.; Yao, X.; Zhang, C.; Rao, Y. *Chem. Commun.* **2015**, *51*, 10014.
- [6] (a) Youn, S. W.; Cho, C. G. *Org. Biomol. Chem.* **2021**, DOI: 10.1039/d1ob00506e.
(b) Xu, X.; Luo, J. *ChemSusChem* **2019**, *12*, 4601.
- [7] (a) Bedford, R. B.; Engelhart, J. U.; Haddow, M. F.; Mitchell, C. J.; Webster, R. L. *Dalton Trans.* **2010**, *39*, 10464.
(b) John, A.; Nicholas, K. M. J. *Org. Chem.* **2012**, *77*, 5600.
(c) Sun, X.; Sun, Y.; Zhang, C.; Rao, Y. *Chem. Commun.* **2014**, *50*, 1262.
- [8] (a) Bhawal, B. N.; Morandi, B. *Angew. Chem., Int. Ed.* **2019**, *58*, 10074.
(b) Yu, B.; Zou, S.; Liu, H.; Huang, H. *J. Am. Chem. Soc.* **2020**, *142*, 18341.
(c) Yu, B.; Zou, S.; Huang, H. *J. Org. Chem.* **2021**, *86*, 7849.
(d) Rochette, E.; Desrosiers, V.; Soltani, Y.; Fontaine, F. G. *J. Am. Chem. Soc.* **2019**, *141*, 12305.
(e) Baba, K.; Tobisu, M.; Chatani, N. *Angew. Chem., Int. Ed.* **2013**, *52*, 11892.
(f) Shao, Y.; Zhang, F.; Zhang, J.; Zhou, X. *Angew. Chem., Int. Ed.* **2016**, *55*, 11485.
(g) Fan, C.; Lv, X.-Y.; Xiao, L.-J.; Xie, J.-H.; Zhou, Q.-L. *J. Am. Chem. Soc.* **2019**, *141*, 2889.
- [9] Li, S.; Zhang, C.; Fu, L.; Wang, H.; Cai, L.; Chen, X.; Wang, X.; Li, G. *CCS Chem.* **2021**, DOI: 10.31635/ccschem.021.202101156.
- [10] (a) Satoh, T.; Kawamura, Y.; Miura, M.; Nomura, M. *Angew. Chem., Int. Ed.* **1997**, *36*, 1740.
(b) Xiao, B.; Gong, T.-J.; Liu, Z.-J.; Liu, J.-H.; Luo, D.-F.; Xu, J.; Liu, L. *J. Am. Chem. Soc.* **2011**, *133*, 9250.
(c) Luo, S.; Luo, F.-X.; Zhang, X.-S.; Shi, Z.-J. *Angew. Chem., Int. Ed.* **2013**, *52*, 10598.
(d) Inamoto, K.; Kadokawa, J.; Kondo, Y. *Org. Lett.* **2013**, *15*, 3962.
(e) Zhang, C.; Ji, J.; Sun, P. *J. Org. Chem.* **2014**, *79*, 3200.
(f) Duan, S.; Xu, Y.; Zhang, X.; Fan, X. *Chem. Commun.* **2016**, *52*, 10529.
(g) Fu, L.; Li, S.; Cai, Z.; Ding, Y.; Guo, X.-Q.; Zhou, L.-P.; Yuan, D.; Sun, Q.-F.; Li, G. *Nat. Catal.* **2018**, *1*, 469.
(h) Kumar, D. R.; Gopi Krishna Reddy, A.; Satyanarayana, G. *Eur. J. Org. Chem.* **2019**, *2019*, 2472.
- [11] (a) Wei, Y.; Yoshikai, N. *Org. Lett.* **2011**, *13*, 5504.

- (b) Schmidt, B.; Riemer, M. *J. Heterocycl. Chem.* **2017**, *54*, 1287.
- [12] (a) Sun, W.-W.; Cao, P.; Mei, R.-Q.; Li, Y.; Ma, Y.-L.; Wu, B. *Org. Lett.* **2014**, *16*, 480.
(b) Tong, H.-R.; Zheng, W.; Lv, X.; He, G.; Liu, P.; Chen, G. *ACS Catal.* **2020**, *10*, 114.
(c) Zhou, T.; Jiang, M.-X.; Yang, X.; Yue, Q.; Han, Y.-Q.; Ding, Y.; Shi, B.-F. *Chin. J. Chem.* **2020**, *38*, 242.
- [13] Wang, P.; Verma, P.; Xia, G.; Shi, J.; Qiao, J. X.; Tao, S.; Cheng, P. T. W.; Poss, M. A.; Farmer, M. E.; Yeung, K.-S.; Yu, J.-Q. *Nature* **2017**, *551*, 489.
- [14] (a) Whitfield, S. R.; Sanford, M. S. *J. Am. Chem. Soc.* **2007**, *129*, 15142.
(b) Xu, L.-M.; Li, B.-J.; Yang, Z.; Shi, Z.-J. *Chem. Soc. Rev.* **2010**, *39*, 712.
(c) Hickman, A. J.; Sanford, M. S. *Nature* **2012**, *484*, 177.
(d) Powers, D. C.; Ritter, T. *Acc. Chem. Res.* **2012**, *45*, 840.
- [15] (a) Xiao, Y.; Xu, Y.; Cheon, H.; Chae, J. *J. Org. Chem.* **2013**, *78*, 5804.
(b) He, X.; Yu, Z.; Jiang, S.; Zhang, P.; Shang, Z.; Lou, Y.; Wu, J. *Bioorg. Med. Chem.* **2015**, *25*, 5601.
(c) Okaecwe, T.; Swanepoel, A.; Petzer, A.; Bergh, J.; Petzer, J. *Bioorg. Med. Chem.* **2012**, *20*, 4336.
- [16] Dudnik, A. S.; Chernyak, N.; Huang, C.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2010**, *49*, 8729.

(Fan, Y.)