三乙胺促进的邻亚甲基苯醌与 β -羰基硫代酰胺的[4+2]环加成反应 构建功能化 4*H*-色烯衍生物

南光明^a 詹靖波^a 原春鸣^a 文丽荣^{*,b} 李明^{*,a,b}

(^a 伊犁师范大学 新疆天然产物化学与应用高等学校重点实验室 新疆伊宁 835000)

(^b 青岛科技大学化学与分子工程学院 生态化工国家重点实验室培育基地 山东青岛 266042)

摘要 建立了一种通过 NEt₃ 促进的 β -羰基硫代酰胺(KTAs)和邻亚甲基苯醌(*o*-QMs)环合来合成高度取代的 4*H*-色烯衍生物的新方法. 该方法涉及一个串联序列, 该序列包括 1,6-共轭加成/*O*-环化/开环/*O*-环化. 该过程表现出良好的官能团耐受性及可扩展性, 且操作简单.

关键词 邻亚甲基苯醌; β -羰基硫代酰胺; 4*H*-色烯衍生物; [4+2]环加成

NEt₃-Promoted Construction of Functionalized 4*H*-Chromenes via [4+2] Cycloaddition Reaction of *ortho*-Quinone Methides with β -Ketothioamides

Nan, Guangming^a Zhan, Jingbo^a Yuan, Chunming^a Wen, Lirong^{*,b} Li, Ming^{*,a,b}

(^a University and College Key Lab of Natural Product Chemistry and Application in Xinjiang, Yili Normal University, Yining, Xinjiang 835000)

(^b State Key Laboratory Base of Eco-Chemical Engineering, College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, Qingdao, Shandong 266042)

Abstract A novel and efficient method for the synthesis of highly substituted 4*H*-chromene derivatives by NEt₃-promoted annulation of β -ketothioamides (KTAs) and *ortho*-quinone methides (*o*-QMs) has been developed. This protocol involves a tandem sequence including 1,6-conjugate addition/*O*-cyclization/ring-opening/*O*-cyclization. This process exhibits good functional group tolerance, operationally-simple conditions and scalability.

Keywords *ortho*-quinone methide (*o*-QMs); β -ketothioamide; 4*H*-chromene derivative; [4+2] cycloaddition

1 Introduction

Functionalized 4*H*-chromene framework, as important privileged heterocyclic scaffold, appears in a plethora of natural products and a variety of known inhibitors for a broad range of receptors.^[1] For instance, Uvafzelin is an antimicrobial agent against gram-positive and acid-fast bacteria.^[2] Troglitazone is considered to be an antidiabetic and anti-inflammatory drug.^[3] HA14-1 has been shown to synergize with various anticancer therapies.^[4] Miroestrol was considered to be a rejuvenating drug.^[5] Some are endowed with a broad spectrum of biological properties such as anticancer,^[6] antimicrobial,^[7] antimalarial,^[8] anticoagu-

lant,^[9] antifungal,^[10] and antioxidant activities.^[11] Although various approaches for the preparation of 4*H*-chromenes have been established,^[12] efficient and practical synthetic methods to access 4*H*-chromenes are still extremely desirable.

Recently β -ketothioamides (KTAs)^[13] have received much attention as multifunctional building blocks in organic synthesis and have been extensively used in various carbon-carbon and carbon-heteroatom bond-forming reactions for the construction of heterocycles (Scheme 1, modes A~D^[14-17]). Although significant progress has been made in this field, the discovery of novel reaction should be an interesting project.

* Corresponding authors. E-mail: wenlirong@qust.edu.cn; liming928@qust.edu.cn

Received April 2, 2021; revised May 26, 2021; published online June 15, 2021.

Project supported by the National Natural Science Foundation of China (No. 21572110) and the Natural Science Foundation of Shandong Province (No. ZR2019MB010).

国家自然科学基金(No. 21572110)及山东省自然科学基金(No. ZR2019MB010)资助项目.

ortho-Quinone methides (*o*-QMs),^[18] as a four-atom partner, could participate in diverse $[4+n]$ cycloadditions with various coupling partners to generate five-, six-, or seven-membered rings,^[19-20] especially oxa- $[4+2]$ cyclizations.^[21] In the construction of various oxygen-containing heterocyclic compounds, most of the $[4+2]$ -cycloadditions of *o*-QMs were confined to alkenes,^[22] alkynes^[23] and others.^[24] Recently, Li *et al.*^[25] described the organocatalytic domino reactions of *o*-QMs with β -functionalized ketones to give hydroxyl-containing chroman compounds, but the reaction of *o*-QMs with KTAs has not been developed in the synthesis for 4*H*-chromenes scaffold.

Based on our ongoing interest in heterocycle synthesis^[26] and in order to explore this new reactivity of KTAs, a novel reaction of *o*-QMs and KTAs for accessing 4*H*-chromenes was reported. It is worth noting that the carbonyl group of KTAs is involved as a “pseudoreactive site”, and the actual reaction center is involved in the thio-carbonyl group.

2 Results and discussion

During the investigation of the reaction of 3-oxo-*N*,3-diphenylpropanethioamide (**1a**) with *o*-QM **2a** promoted by NEt₃, a new product **3aa** was observed. This inspired our interest in further exploration of this transformation. The optimal reaction conditions to afford **3aa** efficiently were investigated. The results are summarized in Table 1. Initially, when the mixture of **1a** and **2a** was stirred in EtOH with 1.0 equiv. of NEt₃ at 70 °C for 10 h, **3aa** was obtained in 66% yield (Table 1, Entry 1). Only trace amount of **3aa** was afforded without any additives (Table 1, Entry 2). Considering the effect of a base on the reaction, other bases such as *N,N*-diisopropylethylamine (DIPEA), triethylenediamine (DABCO), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 4-dimethylaminopyridine (DMAP), pyridine, PPh₃ and K₂CO₃ were screened, and the results showed that these bases could not improve the yield of product **3aa** (Table 1, Entries 3–9). Next, the amount of NEt₃ was tested. The results showed that the

Table 1 Optimization of reaction conditions^{a,b}

Entry	<i>n</i> (1a) : <i>n</i> (2a)	Base (equiv.)	Solvent	<i>T</i> /°C	Yield/%
1	1 : 1	NEt ₃ (1)	EtOH	70	66
2	1 : 1	—	EtOH	70	Trace
3	1 : 1	DIPEA (1)	EtOH	70	60
4	1 : 1	DABCO (1)	EtOH	70	57
5	1 : 1	DBU (1)	EtOH	70	Trace
6	1 : 1	DMAP (1)	EtOH	70	65
7	1 : 1	Pyridine (1)	EtOH	70	35
8	1 : 1	PPh ₃ (1)	EtOH	70	32
9	1 : 1	K ₂ CO ₃ (1)	EtOH	70	Trace
10	1 : 1	NEt ₃ (0.5)	EtOH	70	77
11	1 : 1	NEt ₃ (0.2)	EtOH	70	72
12	1 : 1	NEt ₃ (0.5)	—	90	40
13	1 : 1	NEt ₃ (0.5)	HFIP	70	72
14	1 : 1	NEt ₃ (0.5)	CH ₃ OH	70	63
15	1 : 1	NEt ₃ (0.5)	TFE	70	65
16	1 : 1	NEt ₃ (0.5)	CH ₃ CN	70	65
17	1 : 1	NEt ₃ (0.5)	DCE	70	63
18	1 : 1	NEt ₃ (0.5)	THF	70	55
19	1 : 1	NEt ₃ (0.5)	Toluene	70	50
20	1 : 1	NEt ₃ (0.5)	EtOH	80	75
21	1 : 1	NEt ₃ (0.5)	EtOH	60	69
22	1 : 1.2	NEt ₃ (0.5)	EtOH	70	85
23	1.2 : 1	NEt ₃ (0.5)	EtOH	70	75
24	1 : 1.5	NEt ₃ (0.5)	EtOH	70	84
25 ^c	1 : 1.2	NEt ₃ (0.5)	EtOH	70	72

^a Reaction conditions unless otherwise noted: **1a** (0.2 mmol), **2a** (0.2 mmol), solvent (2 mL), 10 h. ^b Isolated yields. ^c EtOH (1 mL).

Scheme 1 Some reaction types of KTAs

yield of **3aa** was increased from 66% to 77% when the amount of NEt_3 was reduced to 0.5 equiv. (Table 1, Entry 10), whereas further reducing the amounts of NEt_3 did not improve the yield of **3aa** (Entry 11). A solventless reaction of **1a** and **2a** was also attempted with 0.5 equiv. of NEt_3 at 90 °C for 10 h, and **3aa** was obtained in 40% yield (Table 1, Entry 12). Different solvents such as 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP), CH_3OH , TFE, CH_3CN , DCE, THF, and toluene were examined, but they did not give satisfactory results (Table 1, Entries 13~19). The effect of temperature on the reaction was evaluated, and the results suggested that 70 °C favored this reaction (Table 1, Entries 20, 21). Finally, the mole ratio of **1a** and **2a** was tested, and the results indicated the excess of **2a** can cause an obvious improvement in the yield of **3aa** (Table 1, Entries 22~24). However, increasing the reaction system concentration was not conducive to the reaction (Table 1, Entry 25). Consequently, the optimal reaction conditions were established by employing **1a** and **2a** in a mole ratio of 1 : 1.2 with 0.5 equiv. of NEt_3 in EtOH at 70 °C for 10 h.

Under the optimized reaction conditions, the process could easily be applied to a variety of KTAs **1**, and the results are summarized in Table 2. As one can see from

Table 2 Synthesis of 4*H*-chromenes **3aa**~**3ya**^{a,b}

^a Reaction conditions: **1** (0.3 mmol), **2a** (0.36 mmol), NEt_3 (0.15 mmol), EtOH (3 mL). ^b Isolated yields.

Table 2, when R of KTAs **1** was an aromatic group or a heterocyclic ring, and R^1 was phenyl, the reactions proceeded efficiently to provide the desired products **3aa**~**3oa** in 52%~85% yields. When R was phenyl and R^1 of the KTAs **1** was an aromatic group, the reactions showed similar reactivity to provide the desired products **3qa**~**3va** in 50%~77% yields. Unfortunately, however, when R^1 was 4- $\text{NO}_2\text{C}_6\text{H}_4$, a complex mixture was observed (**3pa**). Additionally, when R^1 was an aliphatic group such as methyl, the reaction gave the desired product **3wa** in 66% yield, but when two methyl groups or a morpholine group were involved, different reactivity was observed and the target products **3xa** and **3ya** were not obtained. Since X-ray analysis of **3aa** revealed that there existed an intramolecular hydrogen bond between amino and carbonyl group, helping to further stabilize the structures of the product, so we inferred that N—H bond possibly played an indispensable role to promote this reaction.

Next, the scope of *o*-QMs **2** was explored (Table 3), and the results revealed that various *o*-QMs **2a**~**2g** could be tolerated well, and the reactions proceeded smoothly to afford the corresponding 4*H*-chromene derivatives **3ab**~**3ag** in moderate to good yields. It is noteworthy that when R^2 was an electron-donating group (**3ae**~**3ag**), the yields relative to halogen substitution (**3ab**~**3ad**) were higher.

Table 3 Synthesis of 4*H*-chromenes **3ab**~**3ag**^{a,b}

Compd.	R^2	Yield/%
3ab	6-F	67
3ac	6-Cl	65
3ad	6-Br	68
3ae	6- CH_3	73
3af	6-OMe	76
3ag	8-OMe	78

^a Reaction conditions: **1a** (0.3 mmol), **2** (0.36 mmol), NEt_3 (0.15 mmol), EtOH (3 mL). ^b Isolated yields.

The structures of all newly synthesized 4*H*-chromenes were identified by their IR, ^1H NMR, ^{13}C NMR, and HRMS spectra and unequivocally confirmed by X-ray diffraction analysis of the monocrystal of **3aa** (CCDC 1583563).

Next, our attention was shifted to elucidate a reasonable mechanism of this transformation. Notably, during the investigation of this transformation reaction under the standard conditions, we noticed a trace intermediate by thin-layer chromatography (TLC), which disappeared gradually with time expanding. In order to obtain this intermediate,

the reaction of **1a** and **2a** was conducted at 25 °C for 1 h. To our delight, a chroman intermediate **4** (Its structure was confirmed by X-ray diffraction analysis. No. CCDC 1814241) was obtained in 65% yield (Scheme 2, a). When the reaction of **1a** and **2a** was performed without NEt₃ at 25 °C for 1 h, only 15% of **4** was provided (Scheme 2, b). Next, to indicate the effect of NEt₃ on the reaction, two comparative experiments were carried out with intermediate **4** as the substrate at 70 °C for 10 h. It was found that NEt₃ is essential in the conversion of intermediate **4** to product **3aa** (Scheme 2, c). Additionally, a wet lead acetate paper was placed in the reaction bottle, and the paper changed from white to black as the reaction proceeded, verifying hydrogen sulfide gas was generated during the reaction.

On the basis of the experimental results above, a possi-

ble mechanism is proposed (Scheme 3). First, KTAs **1** are mediated by Et₃N and deprotonate to form the anions of β -(haloaryl)thioacetanilides, which trigger 1,6-conjugate addition involving Michael addition with *o*-QMs **2** to generate the intermediate [A] according to type C with chemo- and regioselectivity, followed by an intramolecular *O*-cyclization, forming the kinetically favored intermediate **4**. Subsequently, in the presence of triethylamine the C—O bond would collapse to regenerate the ring-open intermediate [A—H][−], which underwent intramolecular *O*-cyclization to form the more stable conjugated thioenol intermediate [B]. Finally, the thermodynamically more stable 4*H*-chromene **3** is afforded irreversibly with elimination of H₂S. Obviously, the present reaction involved novel processes via combined modes C and D of KTAs.

Scheme 2 Control experiments

Scheme 3 Proposed possible mechanism

To demonstrate the practical application of the present protocol, a gram-scale experiment was carried out for the reaction of **1a** (4 mmol, 1.02 g) with **2a** (4.8 mmol, 1.49 g) under the optimal reaction conditions, and the coupling product **3aa** was obtained in 68% yield (1.45 g) (Scheme 4).

Scheme 4 Gram scale experiment

3 Conclusions

In summary, an efficient method for the synthesis of highly substituted 4*H*-chromenes by NEt_3 -promoted annulation of KTAs and *o*-QMs was developed. Mechanistic study of the reaction intermediate and control experiments indicated that the annulation reaction underwent a cascade pathway including 1,6-conjugate addition/*O*-cyclization/ring-opening/*O*-cyclization. Unlike previous reports, the present reaction involved novel relay processes and exhibited high bond-forming efficiency. This process exhibits good functional group tolerance, operationally-simple conditions and scalability. This reaction not only provides an efficient method for the construction of biologically important 4*H*-chromenes but also greatly enriches the chemistry of *o*-QMs and KTAs.

4 Experimental section

4.1 General Information

Unless noted, all commercial reagents and solvents were used without further purification. Melting points were recorded on a RY-1 microscopic melting apparatus and uncorrected. ^1H NMR spectra were recorded on 500 MHz and ^{13}C NMR spectra were recorded on 125 MHz by using a Bruker Avance 500 spectrometer. Chemical shifts were reported relative to tetramethylsilane (TMS). High-resolution mass spectra were obtained on a Waters Xevo G2-XS QToF spectrometer with an ESI source. The X-ray single-crystal diffraction was performed on Saturn 724+ instrument. The IR spectra were recorded on an iS10 FT-IR spectrometer, and only major peaks are reported in cm^{-1} . Silica gel (200~300 mesh) for column chromatography and silica GF254 for TLC were produced by Qingdao Marine Chemical Company (China).

4.2 General procedure

KTAs **1** (0.3 mmol, 1.0 equiv.), *o*-QMs **2** (0.36 mmol, 1.2 equiv.) and NEt_3 (0.15 mmol, 0.5 equiv.) were dissolved in EtOH (3.0 mL). The reaction mixture was stirred at 70 °C. After the reaction was completed (detected by

TLC, petroleum ether/ethyl acetate, $V:V=50:1$), the solvent was directly removed under reduced pressure and the crude mixture was purified by silica gel flash column chromatography (petroleum ether/ethyl acetate, $V:V=50:1$) to afford the pure products **3**.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3aa**): White solid, 135.6 mg, 85% yield. m.p. 169~171 °C; ^1H NMR ($\text{DMSO}-d_6$, 500 MHz) δ : 12.73 (s, 1H), 7.43~7.42 (m, 5H), 7.36 (t, $J=7.4$ Hz, 2H), 7.28~7.25 (m, 3H), 7.22~7.17 (m, 3H), 7.15 (t, $J=7.7$ Hz, 1H), 6.74 (s, 1H), 6.57 (s, 2H), 4.95 (s, 1H), 1.22 (s, 18H); ^{13}C NMR ($\text{DMSO}-d_6$, 125 MHz) δ : 194.2, 159.5, 152.3, 148.6, 141.3, 139.5, 138.1, 137.4, 129.9, 129.7, 129.4, 128.4, 128.3, 127.7, 126.9, 125.9, 124.8, 123.0, 121.9, 116.4, 91.6, 41.7, 34.7, 30.6; IR (KBr) ν : 3630, 3060, 2956, 2916, 2869, 1633, 1597, 1432, 1325, 1235, 1152, 1116, 754, 699 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{38}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 532.2852, found 532.2851.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(4-fluorophenyl)methanone (**3ba**): Pale yellow solid, 115.4 mg, 70% yield. m.p. 156~158 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.25 (s, 1H), 7.49 (d, $J=7.9$ Hz, 2H), 7.42 (t, $J=7.8$ Hz, 2H), 7.22~7.18 (m, 4H), 7.15~7.07 (m, 3H), 7.02 (t, $J=8.6$ Hz, 2H), 6.60 (s, 2H), 4.99 (s, 1H), 4.90 (s, 1H), 1.30 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 193.2, 162.9 ($^1J_{\text{C-F}}=248.3$ Hz) 160.2, 152.1, 148.5, 137.7, 137.4, 137.2, 135.7, 129.1, 128.6 ($^3J_{\text{C-F}}=8.3$ Hz), 128.5, 127.4, 127.2, 125.2, 124.6, 123.4, 122.5, 116.2, 114.9 ($^2J_{\text{C-F}}=21.7$ Hz), 90.8, 42.1, 34.1, 30.1; IR (KBr) ν : 3631, 3060, 2957, 2921, 2861, 1635, 1599, 1585, 1558, 1464, 1432, 1329, 1316, 1217, 1151, 1018, 908, 847, 768, 754, 689 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{37}\text{NO}_3\text{F}$ $[\text{M}+\text{H}]^+$ 550.2757, found 550.2756.

(4-Chlorophenyl)(4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)methanone (**3ca**): White solid, 110 mg, 65% yield. m.p. 155~157 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.26 (s, 1H), 7.48 (d, $J=8.0$ Hz, 2H), 7.42 (t, $J=7.9$ Hz, 2H), 7.30~7.28 (m, 2H), 7.22~7.17 (m, 2H), 7.13~7.06 (m, 5H), 6.58 (s, 2H), 4.99 (s, 1H), 4.86 (s, 1H), 1.30 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 193.0, 160.2, 152.1, 148.3, 140.0, 137.4, 137.2, 135.8, 134.7, 129.2, 129.0, 128.2, 127.9, 127.5, 127.1, 125.2, 124.7, 123.5, 122.6, 116.2, 90.7, 42.0, 34.1, 30.1; IR (KBr) ν : 3636, 3074, 3001, 2961, 2921, 2866, 1627, 1599, 1571, 1532, 1457, 1434, 1327, 1237, 1209, 1151, 1027, 907, 830, 765, 758, 695 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{37}\text{N}-\text{O}_3\text{Cl}$ $[\text{M}+\text{H}]^+$ 566.2462, found 566.2468.

(4-Bromophenyl)(4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)methanone (**3da**): White solid, 115.4 mg, 63% yield. m.p. 170~172 °C; ^1H NMR ($\text{DMSO}-d_6$, 500 MHz) δ : 12.69 (s, 1H), 7.51 (d, $J=8.2$ Hz, 2H), 7.43~7.42 (m, 4H), 7.27~7.23 (m, 3H), 7.21~7.18 (m, 1H), 7.15~7.11 (m, 3H), 6.74 (s, 1H), 6.57 (s, 2H), 4.93 (s, 1H), 1.23 (s, 18H); ^{13}C NMR ($\text{DMSO}-d_6$, 125 MHz) δ : 192.8, 159.6, 152.5, 148.3, 140.5, 139.5, 138.2, 137.5, 131.3, 129.8, 129.6, 129.0, 128.2, 127.5, 125.8, 124.9, 123.2, 123.0, 122.1, 116.4, 91.4, 41.5, 34.7, 30.6; IR

(KBr) ν : 3636, 3097, 3001, 2960, 2880, 1625, 1600, 1586, 1568, 1531, 1462, 1433, 1328, 1237, 1209, 1185, 1151, 1010, 906, 829, 766, 758, 695 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{37}\text{NO}_3\text{Br}$ $[\text{M} + \text{H}]^+$ 610.1957, found 610.1954.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(4-(trifluoromethyl)phenyl)methanone (**3ea**): White solid, 93.6 mg, 52% yield. m.p. 210 ~ 212 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.35 (s, 1H), 7.58 (d, $J=8.1$ Hz, 2H), 7.50 (d, $J=7.8$ Hz, 2H), 7.43 (t, $J=7.9$ Hz, 2H), 7.25 ~ 7.17 (m, 4H), 7.13 ~ 7.10 (m, 2H), 7.07 (t, $J=7.3$ Hz, 1H), 6.56 (s, 2H), 5.00 (s, 1H), 4.78 (s, 1H), 1.29 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 192.6, 160.3, 152.2, 148.2, 145.0, 137.4, 137.0, 135.9, 130.6 ($^2J_{\text{C-F}}=32.5$ Hz), 129.2, 129.0, 127.5, 126.9, 126.7, 125.3, 125.0, 124.9, 123.4, 122.7, 116.2, 90.6, 41.9, 34.1, 30.1; IR (KBr) ν : 3607, 3074, 2964, 2907, 2875, 1634, 1611, 1597, 1545, 1498, 1457, 1432, 1320, 1238, 1180, 1167, 1125, 1107, 1064, 938, 910, 843, 770, 756 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{37}\text{NO}_3\text{F}_3$ $[\text{M} + \text{H}]^+$ 600.2726, found 600.2728.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(2-fluorophenyl)methanone (**3fa**): White solid, 112.1 mg, 68% yield. m.p. 178 ~ 180 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.29 (s, 1H), 7.51 (d, $J=7.8$ Hz, 2H), 7.42 (t, $J=7.9$ Hz, 2H), 7.34 ~ 7.25 (m, 1H), 7.22 ~ 7.16 (m, 2H), 7.11 ~ 7.10 (m, 2H), 7.07 ~ 6.97 (m, 4H), 6.55 (s, 2H), 4.96 (s, 1H), 4.75 (s, 1H), 1.28 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 189.6, 159.8, 152.0, 148.0, 137.6, 137.1, 135.5, 130.2, 129.5 ($^2J_{\text{C-F}}=18.1$ Hz), 129.3, 129.1, 127.4, 126.9, 125.2, 124.7, 123.9, 123.5, 122.7, 116.1, 115.6 ($^2J_{\text{C-F}}=18.1$ Hz), 91.9, 41.7, 34.1, 30.1; IR (KBr) ν : 3616, 3074, 2959, 2912, 2870, 1636, 1614, 1596, 1561, 1464, 1431, 1330, 1219, 1180, 1157, 754 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{37}\text{NO}_3\text{F}$ $[\text{M} + \text{H}]^+$ 550.2757, found 550.2760.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(3-fluorophenyl)methanone (**3ga**): White solid, 108.8 mg, 66% yield. m.p. 160 ~ 162 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.28 (s, 1H), 7.52 (d, $J=7.6$ Hz, 2H), 7.45 (t, $J=7.8$ Hz, 2H), 7.36 ~ 7.31 (m, 1H), 7.25 ~ 7.20 (m, 2H), 7.17 ~ 7.08 (m, 4H), 7.01 (d, $J=7.5$ Hz, 1H), 6.85 (d, $J=9.0$ Hz, 1H), 6.69 (s, 2H), 5.03 (s, 1H), 4.88 (s, 1H), 1.33 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 192.5, 162.2 ($^1J_{\text{C-F}}=247.2$ Hz), 160.3, 152.2, 148.4, 143.5, 137.5, 137.1, 135.8, 129.7 ($^3J_{\text{C-F}}=7.9$ Hz), 129.2, 129.0, 127.4, 127.2, 125.3, 124.7, 123.4, 122.6, 122.1, 116.2, 115.7 ($^2J_{\text{C-F}}=21.2$ Hz), 113.7 ($^2J_{\text{C-F}}=22.3$ Hz), 90.7, 41.9, 34.1, 30.1; IR (KBr) ν : 3622, 3074, 2958, 2912, 2870, 1638, 1588, 1562, 1465, 1436, 1330, 1217, 1177, 1145, 963, 850, 788, 754 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{36}\text{NO}_3\text{NaF}$ $[\text{M} + \text{Na}]^+$ 572.2577, found 572.2576.

(3-Chlorophenyl)(4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)methanone (**3ha**): White solid, 101.9 mg, 60% yield. m.p. 165 ~ 167 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.28 (s, 1H), 7.50 (d, $J=7.6$ Hz, 2H), 7.43 (t, $J=7.9$ Hz, 2H), 7.35 ~ 7.33 (m, 1H),

7.28 ~ 7.25 (m, 1H), 7.23 ~ 7.18 (m, 2H), 7.07 ~ 7.15 (m, 5H), 6.63 (s, 2H), 5.00 (s, 1H), 4.86 (s, 1H), 1.32 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 192.3, 160.3, 152.2, 148.3, 143.3, 137.4, 137.1, 135.9, 134.0, 129.3, 129.1, 129.0, 128.8, 127.4, 127.1, 126.7, 125.2, 124.7, 124.4, 123.4, 122.6, 116.2, 90.6, 42.0, 34.2, 30.1; IR (KBr) ν : 3630, 3063, 2959, 2912, 2870, 1630, 1610, 1596, 1563, 1540, 1457, 1433, 1326, 1237, 1180, 1103, 941, 760, 742, 692 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{37}\text{NO}_3\text{Cl}$ $[\text{M} + \text{H}]^+$ 566.2462, found 566.2468.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(*p*-tolyl)methanone (**3ia**): White solid, 126.1 mg, 77% yield. m.p. 150 ~ 152 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.25 (s, 1H), 7.48 (d, $J=7.9$ Hz, 2H), 7.40 (t, $J=7.9$ Hz, 2H), 7.21 ~ 7.17 (m, 2H), 7.16 ~ 7.11 (m, 6H), 7.09 ~ 7.06 (m, 1H), 6.59 (s, 2H), 4.99 (s, 1H), 4.96 (s, 1H), 2.39 (s, 3H), 1.28 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 194.5, 159.9, 152.0, 148.8, 138.8, 138.8, 137.5, 135.5, 129.1, 128.7, 127.5, 127.3, 126.6, 125.1, 124.3, 123.5, 122.4, 116.1, 91.1, 42.0, 34.1, 30.1, 21.4; IR (KBr) ν : 3636, 3002, 2959, 2917, 2870, 1627, 1599, 1534, 1457, 1434, 1327, 1238, 1210, 1178, 1150, 907, 828, 765, 695 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{39}\text{NO}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 568.2828, found 568.2830.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(4-ethylphenyl)methanone (**3ja**): White solid, 125.9 mg, 75% yield. m.p. 171 ~ 173 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.24 (s, 1H), 7.48 (d, $J=8.0$ Hz, 2H), 7.40 (t, $J=7.9$ Hz, 2H), 7.22 ~ 7.18 (m, 8H), 7.10 ~ 7.07 (m, 1H), 6.56 (s, 2H), 5.01 (s, 1H), 4.97 (s, 1H), 2.69 (q, $J=7.6$ Hz, 2H), 1.30 ~ 1.26 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 125 MHz) δ : 194.5, 160.0, 152.0, 148.8, 145.2, 139.0, 137.5, 135.6, 129.1, 127.5, 127.3, 126.8, 125.1, 124.3, 123.5, 122.4, 116.1, 91.1, 42.0, 34.1, 30.1, 28.8, 15.4; IR (KBr) ν : 3634, 3000, 2958, 2866, 1635, 1612, 1599, 1586, 1545, 1465, 1456, 1434, 1353, 1324, 1240, 1200, 1178, 1158, 1120, 1020, 936, 908, 834, 755, 690 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{38}\text{H}_{41}\text{NO}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 582.2984, found 582.2985.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(4-methoxyphenyl)methanone (**3ka**): White solid, 123.0 mg, 73% yield. m.p. 186 ~ 188 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.23 (s, 1H), 7.50 (d, $J=8.0$ Hz, 2H), 7.43 (t, $J=7.6$ Hz, 2H), 7.28 (d, $J=8.0$ Hz, 2H), 7.25 ~ 7.11 (m, 5H), 6.90 (d, $J=8.3$ Hz, 2H), 6.63 (s, 2H), 5.07 (s, 1H), 5.00 (s, 1H), 3.87 (s, 3H), 1.31 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 193.9, 160.3, 160.0, 152.0, 148.9, 137.5, 137.4, 135.6, 134.1, 129.1, 129.0, 128.5, 127.5, 127.4, 125.1, 124.3, 123.4, 122.3, 116.1, 113.3, 91.1, 55.2, 42.1, 34.1, 30.2; IR (KBr) ν : 3625, 3060, 2952, 2917, 2849, 1634, 1598, 1576, 1460, 1433, 1303, 1241, 1168, 1149, 1030, 928, 908, 843, 772, 754, 689 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{39}\text{NO}_4\text{Na}$ $[\text{M} + \text{Na}]^+$ 584.2777, found 584.2766.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(4-(methylthio)phenyl)methanone (**3la**): Yellow solid, 130.0 mg, 75% yield. m.p. 158 ~ 160 °C; ^1H

NMR (CDCl₃, 500 MHz) δ : 13.24 (s, 1H), 7.47 (d, J =7.9 Hz, 2H), 7.41 (t, J =7.8 Hz, 2H), 7.21~7.07 (m, 9H), 6.59 (s, 2H), 4.98 (s, 1H), 4.97 (s, 1H), 2.51 (s, 3H), 1.29 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 193.6, 160.1, 152.1, 148.7, 139.9, 138.1, 137.4, 135.6, 129.1, 129.0, 127.4, 127.2, 125.5, 125.2, 124.5, 123.5, 122.4, 116.2, 91.0, 42.0, 34.1, 30.1, 15.3; IR (KBr) ν : 3627, 3056, 2951, 2926, 2870, 1635, 1592, 1563, 1459, 1433, 1332, 1315, 1238, 1209, 1176, 1150, 1102, 1087, 928, 907, 832, 771, 754, 686 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₇H₄₀NO₃S [M+H]⁺ 578.2733, found 578.2729.

Benzo[d][1,3]dioxol-5-yl(4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)methanone (**3ma**): White solid, 110.5 mg, 64% yield. m.p. 170~171 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.18 (s, 1H), 7.47 (d, J =8.2 Hz, 2H), 7.40 (t, J =7.9 Hz, 2H), 7.22~7.16 (m, 3H), 7.13~7.08 (m, 2H), 6.82~6.77 (m, 2H), 6.73 (d, J =1.1 Hz, 1H), 6.63 (s, 2H), 5.99 (dd, J =4.8, 1.3 Hz, 2H), 5.02 (s, 1H), 4.99 (s, 1H), 1.30 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 193.2, 160.1, 152.1, 148.8, 148.2, 147.2, 137.5, 137.4, 135.6, 129.1, 129.0, 127.4, 125.2, 124.4, 123.4, 122.4, 121.0, 116.1, 107.9, 107.7, 101.1, 90.9, 42.1, 34.2, 30.1; IR (KBr) ν : 3551, 3056, 2959, 2907, 2875, 1629, 1610, 1599, 1586, 1542, 1488, 1502, 1461, 1435, 1356, 1308, 1240, 1181, 1168, 1106, 1077, 1039, 938, 898, 860, 808, 783, 761, 652 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₇H₃₈NO₅ [M+H]⁺ 576.2750, found 576.2751.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(naphthalen-2-yl)methanone (**3na**): Light yellow solid, 123.9 mg, 71% yield. m.p. 212~214 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.39 (s, 1H), 7.85 (d, J =7.9 Hz, 1H), 7.81 (d, J =8.4 Hz, 1H), 7.74 (d, J =7.8 Hz, 1H), 7.66 (s, 1H), 7.55~7.46 (m, 4H), 7.43 (t, J =7.9 Hz, 2H), 7.36~7.34 (m, 1H), 7.22~7.18 (m, 2H), 7.14~7.05 (m, 3H), 6.55 (s, 2H), 5.01 (s, 1H), 4.96 (s, 1H), 1.20 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 194.2, 160.2, 152.0, 148.6, 138.9, 137.5, 137.3, 135.6, 133.4, 132.6, 129.2, 129.0, 128.5, 127.7, 127.7, 127.4, 126.6, 126.2, 126.0, 125.2, 124.5, 124.3, 123.4, 122.5, 116.2, 91.1, 42.1, 34.0, 30.0; IR (KBr) ν : 3634, 3014, 2959, 2907, 2866, 1632, 1611, 1597, 1585, 1549, 1498, 1456, 1424, 1325, 1320, 1181, 1148, 1113, 1096, 932, 900, 860, 805, 758, 694, 654 cm⁻¹; HRMS (ESI-TOF) calcd for C₄₀H₄₀NO₃ [M+H]⁺ 582.3008, found 582.3013.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(thiophen-2-yl)methanone (**3oa**): Yellow solid, 96.8 mg, 60% yield. m.p. 160~162 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.53 (s, 1H), 7.47 (d, J =8.0 Hz, 2H), 7.45 (s, 1H), 7.40 (t, J =7.9 Hz, 2H), 7.35 (d, J =3.6 Hz, 1H), 7.34~7.31 (m, 1H), 7.23~7.17 (m, 2H), 7.16~7.08 (m, 2H), 7.06~7.01 (m, 1H), 6.89 (s, 2H), 5.44 (s, 1H), 5.02 (s, 1H), 1.32 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 183.7, 161.3, 152.2, 149.0, 144.8, 137.2, 136.4, 135.8, 129.1, 129.0, 128.6, 128.4, 128.0, 127.5, 126.9, 125.2, 124.6, 123.3, 122.5, 116.2, 91.0, 41.5, 34.3, 30.2; IR (KBr) ν : 3614, 3086, 2958, 1633, 1582, 1431, 1320, 1235, 1182, 801, 760, 743, 691 cm⁻¹; HRMS (ESI-TOF) calcd

for C₃₄H₃₆NO₃S [M+H]⁺ 538.2416, found 538.2415.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-((4-fluorophenyl)amino)-4*H*-chromen-3-yl)(phenyl)methanone (**3qa**): White solid, 113.8 mg, 69% yield. m.p. 168~170 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.24 (s, 1H), 7.44~7.42 (m, 2H), 7.39 (d, J =7.2 Hz, 1H), 7.34 (t, J =7.3 Hz, 2H), 7.22~7.18 (m, 3H), 7.15~7.07 (m, 5H), 6.57 (s, 2H), 4.99 (s, 1H), 4.94 (s, 1H), 1.29 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 194.4, 160.1, 159.7 (¹ J_{C-F} =244.5 Hz), 152.0, 148.6, 141.4, 137.4, 135.6, 133.3, 129.0 (³ J_{C-F} =9.9 Hz), 128.1, 127.4, 126.5, 125.2, 124.4, 124.3, 123.4, 116.1, 115.9 (² J_{C-F} =22.6 Hz), 90.9, 41.9, 34.1, 30.1; IR (KBr) ν : 3635, 3296, 3059, 3009, 2958, 2912, 2866, 1636, 1614, 1599, 1556, 1510, 1444, 1407, 1326, 1230, 1195, 1160, 1118, 1034, 938, 906, 832, 756, 700, 676 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₆H₃₇NO₃F [M+H]⁺ 550.2757, found 550.2760.

(2-((4-Chlorophenyl)amino)-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-4*H*-chromen-3-yl)(phenyl)methanone (**3ra**): White solid, 129.1 mg, 76% yield. m.p. 168~170 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.27 (s, 1H), 7.49~7.29 (m, 7H), 7.24~7.17 (m, 3H), 7.16~7.05 (m, 3H), 6.55 (s, 2H), 4.98 (s, 1H), 4.94 (s, 1H), 1.28 (s, 18H); ¹³C{¹H} NMR (CDCl₃, 125 MHz) δ : 194.6, 159.8, 152.1, 148.6, 141.3, 137.3, 136.0, 135.6, 129.7, 129.2, 129.1, 128.1, 127.5, 127.3, 126.5, 125.3, 123.6, 123.4, 116.1, 91.3, 41.9, 34.1, 30.1; IR (KBr) ν : 3634, 3320, 3060, 3005, 2957, 2921, 2870, 1635, 1614, 1587, 1550, 1490, 1461, 1443, 1402, 1325, 1236, 1181, 1158, 1118, 1094, 938, 905, 838, 825, 757, 699, 651 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₆H₃₇NO₃Cl [M+H]⁺ 566.2462, found 566.2465.

(2-((4-Bromophenyl)amino)-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-4*H*-chromen-3-yl)(phenyl)methanone (**3sa**): White solid, 135.6 mg, 74% yield. m.p. 175~177 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.26 (s, 1H), 7.51 (d, J =8.6 Hz, 2H), 7.42~7.29 (m, 5H), 7.21~7.19 (m, 3H), 7.14~7.07 (m, 3H), 6.55 (s, 2H), 4.97 (s, 1H), 4.94 (s, 1H), 1.28 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 194.7, 159.8, 152.1, 148.6, 141.3, 137.3, 136.6, 135.6, 132.1, 129.1, 128.1, 127.5, 127.3, 126.5, 125.3, 123.9, 123.4, 117.3, 116.1, 91.4, 41.9, 34.1, 30.1; IR (KBr) ν : 3634, 3334, 3056, 3000, 2956, 2917, 2870, 1635, 1613, 1587, 1548, 1489, 1460, 1442, 1398, 1326, 1223, 1182, 1157, 1118, 1075, 937, 905, 836, 821, 757, 699, 650 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₆H₃₇NO₃Br [M+H]⁺ 610.1957, found 610.1956.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-((4-(trifluoromethyl)phenyl)amino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ta**): White solid, 89.9 mg, 50% yield. m.p. 210~212 °C; ¹H NMR (CDCl₃, 500 MHz) δ : 13.35 (s, 1H), 7.66 (d, J =8.4 Hz, 2H), 7.59 (d, J =8.3 Hz, 2H), 7.41 (t, J =7.2 Hz, 1H), 7.36 (t, J =7.3 Hz, 2H), 7.23~7.21 (m, 3H), 7.17~7.10 (m, 3H), 6.55 (s, 2H), 4.99 (s, 1H), 4.98 (s, 1H), 1.28 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ : 195.3, 159.5, 152.1, 148.5, 141.1, 140.8, 137.1, 135.7, 129.3, 129.2, 128.1, 127.6, 127.1, 126.5, 126.4, 125.8 (² J_{C-F} =32.9 Hz), 125.4, 124.1 (¹ J_{C-F} =271.6 Hz), 123.5, 121.6,

116.1, 91.9, 41.9, 34.1, 30.1; IR (KBr) ν : 3630, 3070, 2957, 2912, 2870, 1633, 1610, 1577, 1555, 1443, 1410, 1325, 1235, 1188, 1167, 1114, 1069, 937, 905, 840, 761, 700 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{37}\text{NO}_3\text{F}_3$ $[\text{M} + \text{H}]^+$ 600.2726, found 600.2728.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(*p*-tolylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ua**): White solid, 126.1 mg, 77% yield. m.p. 135~137 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.23 (s, 1H), 7.37 (d, $J=7.9$ Hz, 3H), 7.33 (t, $J=7.2$ Hz, 2H), 7.22~7.17 (m, 5H), 7.14~7.05 (m, 3H), 6.57 (s, 2H), 4.96 (s, 1H), 4.92 (s, 1H), 2.38 (s, 3H), 1.28 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 194.0, 160.2, 152.0, 148.7, 141.7, 137.6, 135.6, 134.7, 134.3, 129.7, 129.0, 128.8, 128.0, 127.6, 127.3, 126.5, 125.1, 123.5, 122.6, 116.1, 90.8, 42.0, 34.1, 30.1, 20.9; IR (KBr) ν : 3633, 3051, 2957, 2907, 2866, 1635, 1608, 1588, 1550, 1441, 1329, 1236, 1182, 1155, 938, 905, 814, 756, 701 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{39}\text{NO}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 568.2828, found 568.2832.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-((4-methoxyphenyl)amino)-4*H*-chromen-3-yl)(phenyl)methanone (**3va**): White solid, 124.7 mg, 74% yield. m.p. 168~170 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.22 (s, 1H), 7.40 (d, $J=8.9$ Hz, 2H), 7.38 (d, $J=7.2$ Hz, 1H), 7.34 (t, $J=7.2$ Hz, 2H), 7.21 (d, $J=6.9$ Hz, 2H), 7.19~7.12 (m, 2H), 7.09~7.06 (m, 2H), 6.96 (d, $J=8.9$ Hz, 2H), 6.58 (s, 2H), 4.98 (s, 1H), 4.92 (s, 1H), 3.86 (s, 3H), 1.29 (s, 18H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 125 MHz) δ : 193.8, 160.3, 156.8, 152.0, 148.7, 141.7, 137.6, 135.5, 130.1, 129.0, 128.8, 128.0, 127.6, 127.3, 126.5, 125.1, 124.3, 123.4, 116.1, 114.3, 90.5, 55.5, 42.0, 34.1, 30.1; IR (KBr) ν : 3619, 3008, 2956, 1636, 1612, 1558, 1513, 1441, 1327, 1242, 1183, 1150, 1028, 936, 905, 835, 753, 702, 677 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{39}\text{NO}_4\text{Na}$ $[\text{M} + \text{Na}]^+$ 584.2777, found 584.2775.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-(methylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3wa**): White solid, 93.0 mg, 66% yield. m.p. 190~192 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 11.32~11.14 (m, 1H), 7.34~7.27 (m, 3H), 7.18 (t, $J=7.1$ Hz, 1H), 7.12~7.09 (m, 4H), 7.04 (t, $J=7.3$ Hz, 1H), 6.52 (s, 2H), 4.94 (s, 1H), 4.81 (s, 1H), 3.19 (d, $J=5.0$ Hz, 3H), 1.28 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 192.9, 163.6, 151.8, 148.8, 142.2, 138.0, 135.4, 129.0, 128.4, 127.9, 127.8, 127.2, 126.4, 124.8, 123.4, 116.0, 89.4, 41.9, 34.1, 30.1, 27.2; IR (KBr) ν : 3648, 3065, 3003, 2953, 2912, 2875, 1637, 1613, 1534, 1447, 1414, 1339, 1239, 1196, 1182, 1109, 1024, 906, 766, 703 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{31}\text{H}_{36}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 470.2695, found 470.2697.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-6-fluoro-2-(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ab**): Yellow solid, 110.5 mg, 67% yield. m.p. 208~210 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.24 (s, 1H), 7.48 (d, $J=7.8$ Hz), 7.45~7.37 (m, 3H), 7.34 (t, $J=7.3$ Hz, 2H), 7.21 (d, $J=7.2$ Hz, 1H), 7.18 (d, $J=7.3$ Hz, 2H), 7.09 (dd, $J=8.8$, 4.4 Hz, 1H), 6.89 (td, $J=8.6$, 2.5 Hz, 1H), 6.85~6.79 (m, 1H), 6.56 (s, 2H), 5.02 (s, 1H), 4.89 (s, 1H), 1.30 (s, 18H);

^{13}C NMR (CDCl_3 , 125 MHz) δ : 194.5, 159.9, 159.4 ($^1J_{\text{C-F}}=243.7$ Hz), 152.2, 144.6, 141.4, 137.2, 137.0, 135.8, 129.2, 129.0, 128.1, 126.4, 124.6, 123.4, 122.5, 117.4, 115.3 ($^2J_{\text{C-F}}=23.7$ Hz), 114.3 ($^2J_{\text{C-F}}=24.5$ Hz), 90.1, 42.3, 34.2, 30.1; IR (KBr) ν : 3624, 3065, 3003, 2959, 2912, 2875, 1638, 1622, 1598, 1559, 1495, 1460, 1432, 1328, 1284, 1229, 1205, 1187, 1156, 1028, 936, 895, 870, 820, 754, 696, 655 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{36}\text{NO}_3\text{FNa}$ $[\text{M} + \text{Na}]^+$ 572.2577, found 572.2578.

(6-Chloro-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ac**): White solid, 110.4 mg, 65% yield. m.p. 118~120 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.20 (s, 1H), 7.47~7.39 (m, 5H), 7.35 (t, $J=7.3$ Hz, 2H), 7.22~7.14 (m, 4H), 7.10 (s, 1H), 7.07 (d, $J=8.6$ Hz, 1H), 6.54 (s, 2H), 5.02 (s, 1H), 4.87 (s, 1H), 1.30 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 194.5, 159.7, 152.3, 147.2, 141.3, 137.1, 137.0, 135.8, 130.0, 129.2, 129.1, 128.8, 128.1, 127.5, 126.4, 124.7, 123.4, 122.6, 117.5, 90.4, 42.1, 34.2, 30.1; IR (KBr) ν : 3622, 3065, 2958, 2917, 2866, 1634, 1597, 1558, 1499, 1484, 1459, 1430, 1323, 1286, 1240, 1177, 1154, 1119, 1028, 935, 818, 752, 696, 652 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{37}\text{NO}_3\text{Cl}$ $[\text{M} + \text{H}]^+$ 566.2462, found 566.2464.

(6-Bromo-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-2-(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ad**): White solid, 124.6 mg, 68% yield. m.p. 213~215 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.20 (s, 1H), 7.46 (d, $J=7.8$ Hz, 2H), 7.44~7.38 (m, 3H), 7.35 (t, $J=7.3$ Hz, 2H), 7.30 (dd, $J=8.6$, 1.9 Hz, 1H), 7.25 (s, 1H), 7.22~7.17 (m, 3H), 7.01 (d, $J=8.6$ Hz, 1H), 6.54 (s, 2H), 5.02 (s, 1H), 4.88 (s, 1H), 1.30 (s, 18H); ^{13}C NMR (CDCl_3 , 125 MHz) δ : 194.5, 159.6, 152.3, 147.7, 141.3, 137.1, 136.9, 135.8, 131.7, 130.4, 129.7, 129.2, 129.1, 128.1, 126.4, 124.7, 123.4, 122.6, 117.9, 117.6, 90.5, 42.0, 34.2, 30.1; IR (KBr) ν : 3625, 3060, 2948, 2907, 2866, 1633, 1597, 1577, 1545, 1500, 1483, 1458, 1433, 1343, 1243, 1214, 1175, 1154, 1120, 1028, 939, 915, 819, 769, 757, 702, 652 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{36}\text{H}_{36}\text{NO}_3\text{BrNa}$ $[\text{M} + \text{Na}]^+$ 632.1776, found 632.1774.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-6-methyl-2-(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ae**): White solid, 119.5 mg, 73% yield. m.p. 193~195 °C; ^1H NMR (CDCl_3 , 500 MHz) δ : 13.26 (s, 1H), 7.48 (d, $J=7.8$ Hz, 2H), 7.42~7.37 (m, 3H), 7.34 (t, $J=7.2$ Hz, 2H), 7.21~7.16 (m, 3H), 7.02 (d, $J=8.2$ Hz, 1H), 6.98 (d, $J=8.5$ Hz, 1H), 6.92 (s, 1H), 6.57 (s, 2H), 4.98 (s, 1H), 4.87 (s, 1H), 2.25 (s, 3H), 1.29 (s, 18H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 125 MHz) δ : 194.3, 160.3, 152.0, 146.7, 141.6, 137.7, 137.4, 135.6, 134.7, 129.3, 129.1, 128.9, 128.1, 128.0, 127.1, 126.5, 124.4, 123.5, 122.4, 115.8, 91.3, 42.1, 34.2, 30.1, 20.8; IR (KBr) ν : 3634, 3065, 3005, 2945, 2912, 2870, 1628, 1584, 1538, 1496, 1459, 1431, 1346, 1253, 1202, 1152, 1117, 1027, 937, 814, 759, 703 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{37}\text{H}_{39}\text{NO}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 568.2828, found 568.2825.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-6-methoxy-2-

(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3af**): White solid, 128.1 mg, 76% yield. m.p. 170~172 °C; ¹H NMR (CDCl₃, 500 MHz) δ: 13.27 (s, 1H), 7.48 (d, *J*=7.8 Hz, 2H), 7.44~7.37 (m, 3H), 7.34 (t, *J*=7.2 Hz, 2H), 7.23~7.15 (m, 3H), 7.06 (d, *J*=8.9 Hz, 1H), 6.77~6.69 (m, 1H), 6.66~6.60 (m, 1H), 6.58 (s, 2H), 4.99 (s, 1H), 4.87 (s, 1H), 3.73 (s, 3H), 1.29 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ: 194.2, 160.4, 156.6, 152.1, 142.8, 141.6, 137.4, 135.6, 129.1, 128.9, 128.4, 128.0, 126.5, 124.4, 123.4, 122.4, 116.9, 113.3, 113.0, 90.8, 55.6, 42.5, 34.2, 30.1; IR (KBr) ν: 3640, 3060, 2998, 2951, 2911, 2860, 1632, 1597, 1544, 1496, 1458, 1433, 1332, 1237, 1200, 1157, 1120, 1041, 944, 823, 761, 704 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₇H₃₉NO₄Na [M+Na]⁺ 584.2777, found 584.2776.

(4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-7-methoxy-2-(phenylamino)-4*H*-chromen-3-yl)(phenyl)methanone (**3ag**): Light yellow solid, 131.4 mg, 78% yield. m.p. 158~160 °C; ¹H NMR (CDCl₃, 500 MHz) δ: 13.28 (s, 1H), 7.48~7.47 (m, 2H), 7.42 (t, *J*=7.9 Hz, 2H), 7.39~7.36 (m, 1H), 7.34~7.31 (m, 2H), 7.21~7.17 (m, 3H), 7.02~7.00 (m, 1H), 6.66~6.64 (m, 2H), 6.54 (s, 2H), 4.96 (s, 1H), 4.87 (s, 1H), 3.78 (s, 3H), 1.28 (s, 18H); ¹³C NMR (CDCl₃, 125 MHz) δ: 194.5, 159.9, 158.9, 151.9, 149.1, 141.6, 137.9, 137.4, 135.5, 129.5, 129.1, 128.9, 128.0, 126.5, 124.5, 123.4, 122.5, 119.4, 111.5, 101.4, 91.2, 55.6, 41.4, 34.1, 30.1; IR (KBr) ν: 3000, 2951, 2903, 2866, 1640, 1620, 1577, 1548, 1465, 1433, 1418, 1345, 1287, 1193, 1161, 1122, 1030, 976, 832, 752, 699 cm⁻¹; HRMS (ESI-TOF) calcd for C₃₇H₄₀NO₄ [M+H]⁺ 562.2957, found 562.2967.

4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-2-hydroxy-*N*,2-diphenylchromane-3-carboxamide (**4**): White solid, 110.2 mg, 65% yield. m.p. 170~172 °C; ¹H NMR (DMSO-*d*₆, 500 MHz) δ: 11.65 (s, 1H), 9.34 (s, 1H), 8.11 (d, *J*=7.6 Hz, 2H), 7.64~7.59 (m, 2H), 7.51 (t, *J*=7.6 Hz, 2H), 7.27 (t, *J*=7.8 Hz, 2H), 7.20~7.15 (m, 5H), 6.92 (t, *J*=7.6 Hz, 1H), 6.75~6.70 (m, 2H), 6.61 (s, 1H), 6.26 (d, *J*=11.4 Hz, 1H), 5.49 (d, *J*=9.7 Hz, 1H), 1.26 (s, 18H); ¹³C NMR (DMSO-*d*₆, 125 MHz) δ: 197.1, 194.0, 155.3, 152.0, 139.5, 138.9, 137.5, 134.8, 133.5, 129.4, 129.0, 128.8, 128.7, 127.1, 126.7, 124.4, 118.9, 115.8, 65.5, 60.2, 34.9, 30.9; HRMS (ESI-TOF) calcd for C₃₆H₃₉NO₃SNa [M+Na]⁺ 588.2548, found 588.2550.

Supporting Information X-Ray crystallographic data of **3aa** and **4**, mechanistic studies, copies of ¹H NMR, ¹³C NMR of the products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Pratap, R.; Ram, V. *J. Chem. Rev.* **2014**, *114*, 10476.
(b) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893.
(c) Lu, Z.-Y.; Lin, Z.-J.; Wang, W.-L.; Du, L.; Zhu, T.-J.; Fang, Y.-C.; Gu, Q.-Q.; Zhu, W.-M. *J. Nat. Prod.* **2008**, *71*, 543.
- (d) Costa, M.; Dias, T. A.; Brito, A.; Proença, F. *Eur. J. Med. Chem.* **2016**, *123*, 487.
(e) Das, S. G.; Srinivasan, B.; Hermanson, D. L.; Bleeker, N. P.; Doshi, J. M.; Tang, R.; Beck, W. T.; Xing, C. *J. Med. Chem.* **2011**, *54*, 5937.
(f) Nicolaou, K. C.; Roecker, A. J.; Barluenga, S.; Pfeifferkorn, J. A.; Cao, G.-Q. *ChemBioChem* **2001**, *2*, 460.
- [2] (a) Zhang, Z.; ElSohly, H. N.; Li, X.-C.; Khan, S. I.; Broedel, S. E.; Rauli, R. E.; Cihlar, R. L.; Burandt, C.; Walker, L. A. *J. Nat. Prod.* **2003**, *66*, 548.
(b) Hufford, C. D.; Oguntimein, B. O.; Engen, D. V.; Muthard, D.; Clardy, J. *J. Am. Chem. Soc.* **1980**, *102*, 7365.
- [3] (a) Azziz, R.; Ehrmann, D.; Legro, R. S.; Whitcomb, R. W.; Hanley, R.; Fereshetian, A. G.; O'Keefe, M.; Ghaziz, M. N. *J. Clin. Endocrinol. Metab.* **2001**, *86*, 1626.
(b) Yaghoub, P.; Fatemeh, Z.; Khalil, E.; Mahshid, T.; Laleh, M.; Mahnaz, A.; Arash, K.; Reza, S.; Ehsan, F.-M.; Ali, A. *J. Nanosci. Nanotechnol.* **2020**, *20*, 3206.
- [4] Das, S. G.; Doshi, J. M.; Tian, D.; Addo, S. N.; Srinivasan, B.; Hermanson, D. L.; Xing, C. *J. Med. Chem.* **2009**, *52*, 5937.
- [5] Chansakaow, S.; Ishikawa, T.; Seki, H.; Sekine, K.; Okada, M.; Chaichantipyuth, C. *J. Nat. Prod.* **2000**, *63*, 173.
- [6] (a) Tian, D.; Das, S. G.; Doshi, J. M.; Peng, J.; Lin, J.; Xing, C. *Cancer Lett.* **2008**, *259*, 198.
(b) Gourdeau, H.; Leblond, L.; Hamelin, B.; Desputeau, C.; Dong, K.; Kianicka, I.; Custeau, D.; Boudreau, C.; Geerts, L.; Cai, S.-X.; Drewe, J.; Labrecque, D.; Kasibhatla, S.; Tseng, B. *Mol. Cancer Ther.* **2004**, *3*, 1375.
- [7] (a) Groweiss, A.; Cardellina, J. H.; Boyd, M. R. *J. Nat. Prod.* **2000**, *63*, 1537.
(b) Sabry, N. M.; Mohamed, H. M.; Khattab, E. S. A.; Motlaq, S. S.; El-Agrody, A. M. *Eur. J. Med. Chem.* **2011**, *46*, 765.
(c) Creagh, T.; Ruckle, J. L.; Tolbert, D. T.; Giltner, J.; Eiznhamer, D. A.; Dutta, B.; Flavin, M. T.; Xu, Z.-Q. *Antimicrob. Agents Chemother.* **2001**, *45*, 1379.
- [8] (a) Bonsignore, L.; Loy, G.; Secci, D.; Calignano, A. *Eur. J. Med. Chem.* **1993**, *28*, 517.
(b) Alvey, L.; Prado, S.; Saint-Joanis, B.; Michel, S.; Koch, M.; Cole, S. T.; Tillequin, F.; Janin, Y. L. *Eur. J. Med. Chem.* **2009**, *44*, 2497.
- [9] Santhisudha, S.; Sreekanth, T.; Murali, S.; Kumar, B. V.; Devi, M. A.; Reddy, C. S. *Cardiovasc. Hematol. Agents Med. Chem.* **2016**, *14*, 167.
- [10] Mori, K.; Audran, G.; Monti, H. *Synlett* **1998**, 259.
- [11] Pietta, P.-G. *J. Nat. Prod.* **2000**, *63*, 1035.
- [12] (a) Shi, Y.-L.; Shi, M. *Org. Biomol. Chem.* **2007**, *5*, 1499.
(b) Mei, G.-J.; Xu, S.-L.; Zheng, W.-Q.; Bian, C.-Y.; Shi, F. *J. Org. Chem.* **2018**, *83*, 1414.
(c) Wang, M.; Tang, B.-C.; Xiang, J.-C.; Cheng, Y.; Wang, Z.-X.; Ma, J.-T.; Wu, Y.-D.; Wu, A.-X. *J. Org. Chem.* **2018**, *83*, 3409.
(d) Xu, W.; Li, Q.; Wang, W.; Zheng, H.; Zhang, F.; Hu, Y. *RSC Adv.* **2015**, *5*, 56333.
(e) Bhunia, A.; Yetra, S. R.; Gonnade, R. G.; Biju, A. T. *Org. Biomol. Chem.* **2016**, *14*, 5612.
(f) Ma, C.; Zhao, Y. *Org. Biomol. Chem.* **2018**, *16*, 703.
(g) Wu, B.; Gao, X.; Yan, Z.; Chen, M.-W.; Zhou, Y.-G. *Org. Lett.* **2015**, *17*, 6134.
(h) Tahtoui, C.; Demailly, A.; Guidemann, C.; Joyeux, C.; Schneider, P. *J. Org. Chem.* **2010**, *75*, 3781.
- [13] (a) Zhang, L.; Dong, J.; Xu, X.; Liu, Q. *Chem. Rev.* **2016**, *116*, 287.
(b) Guo, W.-S.; Wen, L.-R.; Li, M. *Org. Biomol. Chem.* **2015**, *13*, 1942.
(c) Luo, X.; Ge, L.-S.; An, X.-L.; Jin, J.-H.; Wang, Y.; Sun, P.-P.; Deng, W.-P. *J. Org. Chem.* **2015**, *80*, 4611.
(d) Singh, M. S.; Nandi, G. C.; Samai, S. *Green Chem.* **2012**, *14*, 447.
(e) Huang, F.; Wu, P.; Wang, L.; Chen, J.; Sun, C.; Yu, Z. *J. Org. Chem.* **2014**, *79*, 10553.

- (f) Huang, F.; Wu, P.; Wang, L.; Chen, J.; Sun, C. *Yu, Z. Chem. Commun.* **2014**, 50, 12479.
- (g) Nandi, G. C.; Soumini, K. *J. Org. Chem.* **2016**, 81, 11909.
- (h) Nandi, G. C.; Singh, M. S. *J. Org. Chem.* **2016**, 81, 5824.
- (i) Ansari, M. A.; Yadav, D.; Soni, S.; Srivastava, A.; Singh, M. S. *J. Org. Chem.* **2019**, 84, 5404.
- [14] Wen, L.-R.; Men, L.-B.; He, T.; Ji, G.-J.; Li, M. *Chem.-Eur. J.* **2014**, 20, 5028.
- [15] Man, N.-N.; Wang, J.-Q.; Zhang, L.-M.; Wen, L.-R.; Li, M. *J. Org. Chem.* **2017**, 82, 5566.
- [16] Guo, W.-S.; Xin, X.; Zhao, K.-L.; Wen, L.-R.; Li, M. *RSC Adv.* **2015**, 5, 70429.
- [17] Li, M.; Hou, Y.-L.; Wen, L.-R.; Gong, F.-M. *J. Org. Chem.* **2010**, 75, 8522.
- [18] (a) Zhao, K.; Zhi, Y.; Shu, T.; Valkonen, A.; Rissanen, K.; Enders, D. *Angew. Chem., Int. Ed.* **2016**, 55, 12104.
(b) Wang, J.-Y.; Hao, W.-J.; Tu, S.-J.; Jiang, B. *Org. Chem. Front.* **2020**, 7, 1743.
(c) Mao, W.; Lin, S.; Zhang, S.; Lu, H.; Jia, J.; Xu, Z. *Org. Chem. Front.* **2020**, 7, 856.
(d) Zhou, S.-J.; Cheng, X.; Hu, C.-X.; Xu, G.-Y.; Xiao, W.-J.; Xuan, J. *Sci. China: Chem.* **2021**, 64, 61.
- [19] For examples on [4+1] cyclizations of *o*-hydroxyphenyl substituted *p*-QMs: (a) Xiong, Y.-J.; Shi, S.-Q.; Hao, W.-J.; Tu, S.-J.; Jiang, B. *Org. Chem. Front.* **2018**, 5, 3483.
(b) Liu, L.; Yuan, Z.; Pan, R.; Zeng, Y.; Lin, A.; Yao, H.; Huang, Y. *Org. Chem. Front.* **2018**, 5, 623.
(c) Zhi, Y.; Zhao, K.; Essen, C.; Rissanen, K.; Enders, D. *Org. Chem. Front.* **2018**, 5, 1348.
(d) Liou, Y.-C.; Karanam, P.; Jang, Y.-J.; Lin, W. *Org. Lett.* **2019**, 21, 8008.
(e) Tan, J.-P.; Yu, P.; Wu, J.-H.; Chen, Y.; Pan, J.; Jiang, C.; Ren, X.; Zhang, H.-S.; Wang, T. *Org. Lett.* **2019**, 21, 7298.
- [20] For examples on [4+3] cyclizations of *o*-hydroxyphenyl substituted *p*-QMs: (a) Chen, K.; Hao, W.-J.; Tu, S.-J.; Jiang, B. *Green Chem.* **2019**, 21, 675.
(b) Li, W.; Yuan, H.; Liu, Z.; Zhang, Z.; Cheng, Y.; Li, P. *Adv. Synth. Catal.* **2018**, 360, 2460.
(c) Liu, Q.; Li, S.; Chen, X.-Y.; Rissanen, K.; Enders, D. *Org. Lett.* **2018**, 20, 3622.
(d) Jiang, F.; Yuan, F.-R.; Jin, L.-W.; Mei, G.-J.; Shi, F. *ACS Catal.* **2018**, 8, 10234.
(e) Sun, M.; Ma, C.; Zhou, S.-J.; Lou, S.-F.; Xiao, J.; Jiao, Y.; Shi, F. *Angew. Chem., Int. Ed.* **2019**, 58, 8703.
- [21] For examples on [4+2] cyclizations of *o*-hydroxyphenyl substituted *p*-QMs: (a) Liu, S.; Lan, X.-C.; Chen, K.; Hao, W.-J.; Li, G.; Tu, S.-J.; Jiang, B. *Org. Lett.* **2017**, 19, 3831.
(b) Zhang, L.; Liu, Y.; Liu, K.; Liu, Z.; He, N.; Li, W. *Org. Biomol. Chem.* **2017**, 15, 8743.
(c) Zhang, Z.-P.; Xie, K.-X.; Yang, C.; Li, M.; Li, X. *J. Org. Chem.* **2018**, 83, 364.
(d) Mei, G.-J.; Xu, S.-L.; Zheng, W.-Q.; Bian, C.-Y.; Shi, F. *J. Org. Chem.* **2018**, 83, 1414.
(e) Jiang, X.-L.; Wu, S.-F.; Wang, J.-R.; Mei, G.-J.; Shi, F. *Adv. Synth. Catal.* **2018**, 360, 4225.
(f) Zhou, J.-Y.; Ma, C.; Zhang, Y.-Z.; Wu, Q.; Shi, F. *Org. Biomol. Chem.* **2018**, 16, 9382.
(g) Satbhaiya, S.; Khonde, N. S.; Rathod, J.; Gonnade, R.; Kumar, P. *Eur. J. Org. Chem.* **2019**, 3127.
(h) Huang, H.-M.; Wu, X.-Y.; Leng, B.-R.; Zhu, Y.-L.; Meng, X.-C.; Hong, Y.; Jiang, B.; Wang, D.-C. *Org. Chem. Front.* **2020**, 7, 414.
(i) Zhang, J.-R.; Jin, H.-S.; Wang, R.-B.; Zha, L.-M. *Adv. Synth. Catal.* **2019**, 361, 4811.
(j) Yuan, F.-R.; Jiang, F.; Chen, K.-W.; Mei, G.-J.; Wu, Q.; Shi, F. *Org. Biomol. Chem.* **2019**, 17, 2361.
(k) Tan, J.-P.; Zhang, H.; Jiang, Z.; Chen, Y.; Ren, X.; Jiang, C.; Wang, T. *Adv. Synth. Catal.* **2020**, 362, 1058.
(l) Cheng, X.; Zhou, S.-J.; Xu, G.-Y.; Wang, L.; Yang, Q.-Q.; Xuan, J. *Adv. Synth. Catal.* **2020**, 362, 523.
(m) Zhang, Z.-P.; Chen, L.; Li, X.; Cheng, J.-P. *J. Org. Chem.* **2018**, 83, 2714.
- [22] (a) Zhao, K.; Zhi, Y.; Shu, T.; Valkonen, A.; Rissanen, K.; Enders, D. *Angew. Chem., Int. Ed.* **2016**, 55, 12104.
(b) Liu, S.; Lan, X.-C.; Chen, K.; Hao, W.-J.; Li, G.; Tu, S.-J.; Jiang, B. *Org. Lett.* **2017**, 19, 3831.
(c) Jiang, X.-L.; Wu, S.-F.; Wang, J.-R.; Mei, G.-J.; Shi, F. *Adv. Synth. Catal.* **2018**, 360, 4225.
- [23] (a) Mei, G.-J.; Xu, S.-L.; Zheng, W.-Q.; Bian, C.-Y.; Shi, F. *J. Org. Chem.* **2018**, 83, 1414.
(b) Wang, C.-S.; Cheng, Y.-C.; Zhou, J.; Mei, G.-J.; Wang, S.-L.; Shi, F. *J. Org. Chem.* **2018**, 83, 13861.
- [24] (a) Satbhaiya, S.; Khonde, N. S.; Rathod, J.; Gonnade, R.; Kumar, P. *Eur. J. Org. Chem.* **2019**, 3127.
(b) Zhang, L.; Zhou, X.; Li, P.; Liu, Z.; Liu, Y.; Sun, Y.; Li, W. *RSC Adv.* **2017**, 7, 39216.
(c) Cheng, Y.-C.; Wang, C.-S.; Li, T.-Z.; Gao, F.; Jiao, Y.; Shi, F. *Org. Biomol. Chem.* **2019**, 17, 6662.
- [25] Li, W.; Shi, Z.; Zhou, X. *CN 106674178*, **2017**.
- [26] (a) Yuan, W.-K.; Cui, T.; Liu, W.; Wen, L.-R.; Li, M. *Org. Lett.* **2018**, 20, 1513.
(b) Guo, W.-S.; Gong, H.; Zhang, Y.-A.; Wen, L.-R.; Li, M. *Org. Lett.* **2018**, 20, 6394.
(c) Zhang, L.-B.; Zhu, M.-H.; Ni, S.-F.; Wen, L.-R.; Li, M. *ACS Catal.* **2019**, 9, 1680.
(d) Yuan, W.-K.; Zhu, M.-H.; Geng, R.-S.; Ren, G.-Y.; Zhang, L.-B.; Wen, L.-R.; Li, M. *Org. Lett.* **2019**, 21, 1654.

(Zhao, C.)