无催化剂一锅法构建取代磺酰吡唑

马豪杰^{a,b} 周小强^{b,c} 韩波^a 李然^a 侯雪艳^a
姬星月^a 张玉琦^{*,a} 黄国生^{*,b} 王记江^a

(^a 延安大学化学与化工学院 延安大学新能源与新功能材料重点实验室

陕西化学反应工程重点实验室 陕西延安 716000)

(^b 兰州大学功能有机分子化学国家重点实验室 兰州 730000)

(^c 渭南师范学院化学与材料学院 陕西渭南 714099)

摘要 开发了一种新型有效的在无催化剂条件下由苯磺酰肼和戊烷-2,4-二酮构建取代吡唑的方法,该方法具有官能团适用性好、步骤经济、操作简单、试剂便宜、反应条件温和及可进行按比例放大等优势。

关键词 无催化剂; 磺酰吡唑; 戊烷-2,4-二酮; 苯磺酰肼

A Catalyst-Free One-Pot Protocol for the Construction of Substituted Sulfonyl Pyrazoles

Ma, Haojie^{a,b} Zhou, Xiaoqiang^{b,c} Han, Bo^a Li, Ran^a Hou, Xueyan^a
Ji, Xingyue^a Zhang, Yuqi^{*,a} Huang, Guosheng^{*,b} Wang, Jijiang^a

(^a Key Laboratory of New Energy & New Functional Materials, Shaanxi Key Laboratory of Chemical Reaction Engineering, College of Chemistry and Chemical Engineering, Yan'an University, Yan'an, Shaanxi 716000)

(^b State Key Laboratory of Applied Organic Chemistry, Lanzhou University, Lanzhou 730000)

(^c College of Chemistry and Material, Weinan Normal University, Weinan, Shaanxi 714099)

Abstract A novel and efficient protocol for the construction of substituted pyrazoles from benzenesulfonylhydrazides and pentane-2,4-diones under catalyst-free conditions was developed. This protocol features excellent tolerance to a variety of functional groups, step economy, simple operation with inexpensive reagents, mild reaction conditions, and can be scaled-up.

Keywords catalyst-free; sulfonyl pyrazole; pentane-2,4-dione; benzenesulfonylhydrazide

1 Introduction

Pyrazoles and their derivatives, as an important class of compounds, have extensive applications in the fields of pharmaceutical industry, agrochemicals and materials science.^[1] Molecules with this valuable motif exhibit various biological and pharmaceutical activities, ranging from anticancer^[2] to antiinflammatory,^[3] antibacterial,^[4] antidiabetic,^[5] antiobesity,^[6] and antiviral.^[7] Structural cores containing pyrazole rings have been found as the core struc-

tures of many drugs,^[8] such as Penthiopyrad, Lonazolac, Rimnabant, Razaxaban, Celebrex and Deracoxib (Figure 1). Particularly, Celebrex, served as a selective COX-2 inhibitor, is top 200 brand name drug by total US prescriptions in 2012.^[9] Additionally, pyrazoles are used as ligands in coordination chemistry, directing groups for C—H activations and precursors to N-heterocyclic carbenes (NHCs). Especially, sulfonyl pyrazoles are also used as herbicidal agents^[10] and antioxidants.^[11] Therefore, it is of great value to develop a new method to introduce sulfonyl group to

* Corresponding authors. E-mail: yqzhang@iccas.ac.cn; hgs@lzu.edu.cn

Received April 16, 2021; revised April 27, 2021; published online June 21, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21663032, 31701679, 22061041), the PhD Research Startup Foundation of Yan'an University (Nos. YDBK2019-30, YDBK2015-11), the Municipal Special Fund of Yan'an for High-Level Talents (No. 2019-29), the Yan'an University Training Program of Innovation and Entrepreneurship for Undergraduates (No. D2019016), the Natural Science Basic Research Program of Shaanxi Province (No. 2019JQ-902) and the Scientific Research Program Funded by Shaanxi Provincial Education Department (No. 19JK0287).

国家自然科学基金(Nos. 21663032, 31701679, 22061041)、延安大学博士科研启动基金(Nos. YDBK2019-30, YDBK2015-11)、延安大学高层次人才专项基金(No. 2019-29)、延安大学大学生创新创业训练计划(No. D2019016)、陕西省科技厅自然科学基金基础研究基金(No. 2019JQ-902)和陕西省教育厅自然科学专项研究基金(No. 19JK0287)资助项目。

pyrazoles.

Figure 1 Structures of some commercial pyrazoles

Due to the importance of these heterocycles, a variety of synthetic approaches have been developed for the preparation of pyrazole scaffold, such as oxidative coupling of enaminones and nitriles,^[12] 1,3-dipolar cycloaddition based on alkenes or alkynes,^[13] substituted hydrazines with 1,3-dicarbonyl compounds^[14] or unsaturated aldehydes and ketones,^[15] functionalization of unsubstituted pyrazoles,^[16] metal-catalyzed oxidative carbonylations of arylhydrazines and alkynes.^[17] Among these methods, the most efficient and simplest procedure is based on the condensation-cyclization reaction of substituted hydrazines and 1,3-dicarbonyl compounds. A variety of catalysts have been employed in this transformation, such as Cu salts,^[18] Sc(OTf)₃,^[19] polystyrene supported sulfonic acid,^[20] H₂SO₄,^[21] layered zirconium sulfophenyl phosphonate.^[22] In 2018, Wang's group^[18b] obtained pyrazoles from phenylhydrazine and pentane-2,4-dione under Cu catalyst (Scheme 1, a). The same year, Yu's group^[18a] reported a novel synthesis of pyrazoles in the presence of Cu_{1.5}P-Mo₁₂O₄₀ (Scheme 1, b). In 2009, Wu's group^[19] presented a Sc-catalyzed reaction leading to the formation of pyrazoles (Scheme 1, c). However, there are few methods to acquire substituted sulfonyl pyrazoles.^[23] Additionally, in

(a) Wang's group

(b) Yu's group

(c) Wu's group

(d) This work

Scheme 1 Synthetic approaches toward pyrazoles

the presented methods, there are still some drawbacks that limit their practical applications, including requirements of strongly acid or transition-metal catalysts, harsh and sensitive reaction conditions, long reaction times and low yields. Therefore, there is a great need to develop a simple synthetic route with mild reaction conditions, environmental friendly, high yields and availability of industrial production for sulfonyl pyrazole and its derivatives. Wei's group^[24] reported effective and environmentally approaches for the synthesis of a variety of polycyclic compounds under additive-free or metal catalyst-free conditions. Organic reaction without catalyst will represent a promising option for cleaner and cheaper technologies in industry.^[25] Herein, we designed a novel route for the synthesis of substituted pyrazoles from benzenesulfonylhydrazides and pentane-2,4-diones under catalyst-free conditions.

2 Results and discussion

To initiate our study, the reaction conditions were screened for the formation of pyrazole by using 4-methylbenzenesulfonylhydrazide (**1a**) and pentane-2,4-dione (**2a**) as model substrates (Table 1). The reaction of **1a** (0.2 mmol, 1 equiv.) with **2a** (0.3 mmol, 1.5 equiv.) was conducted in the presence of CBr₄ (1 equiv.) in dimethylacetamide (DMSO) at 60 °C, and was able to isolate the desired 3,5-dimethyl-1-tosyl-1H-pyrazole (**3aa**) in 50% yield (Table 1, Entry 1). Then the reaction was carried out without CBr₄ in order to test whether CBr₄ is necessary. The results showed that the yield of **3aa** did not change (Table 1, Entry 2). Encouraged by this result, we continued

Table 1 Optimization of reaction conditions^a

Entry	Catalyst	Solvent	Temp./°C	Yield ^b /%
1	CBr ₄	DMSO	60	50
2	—	DMSO	60	50
3	—	DMA	60	^c nd
4	—	1,4-Dioxane	60	83
5	—	DMF	60	70
6	—	Toluene	60	94
7	—	CH ₃ CN	60	97
8	—	DCE	60	92
9	—	H ₂ O	60	nd
10	—	CH ₃ CN	40	88
11	—	CH ₃ CN	80	94
^c 12	—	CH ₃ CN	60	95
^d 13	—	CH ₃ CN	60	96

^a Reaction conditions: **1a** (0.2 mmol, 1 equiv.), **2a** (0.3 mmol, 1.5 equiv.), solvent (1.5 mL), 2 h. ^b Isolated yields. ^c Reaction was performed under N₂ atmosphere. ^d Reaction was performed under O₂ atmosphere. ^e nd=not detected. DMA=dimethylacetamide, DMF=N,N-dimethylformamide.

to explore the optimal reaction conditions. Furthermore, the investigation of the solvent effects on the yield of the target product was carried out. The results illustrated that toluene, CH₃CN, and 1,2-dichloroethane (DCE) were better reaction solvents in this transformation, and the highest yield of 97% was acquired in CH₃CN (Table 1, Entries 2~9). In addition, the screening of various reaction temperatures showed that the yield of **3aa** decreased when the reaction was conducted at higher or lower temperatures. The results suggested that 60 °C was favorable for the formation of the target product (Table 1, Entries 7, 10, 11). The effects of reaction atmosphere on the product yield were also studied. The results showed that air was better than N₂ and O₂ (Table 1, Entries 7, 12, 13). Finally, the optimized reaction conditions are as follows: 4-methylbenzenesulfonylhydrazide (**1a**) (0.2 mmol), pentane-2,4-dione (**2a**) (0.3 mmol), in 1.5 mL of CH₃CN at 60 °C under air (Table 1, Entry 7).

With the optimized reaction conditions in hand, the scope and generality of the reaction regard to different

Table 2 Synthesis of 3,5-dimethyl-1-(phenylsulfonyl)-1H-pyrazoles from substituted benzenesulfonylhydrazides and pentane-2,4-dione^a

^a Reaction conditions: **1** (0.2 mmol, 1 equiv.), **2a** (0.3 mmol, 1.5 equiv.), in 1.5 mL of CH₃CN at 60 °C under air for 2 h.

benzenesulfonylhydrazides were investigated. The results are summarized in Table 2. A diverse array of benzenesulfonylhydrazides, bearing electron-donating, electron-withdrawing or sterically hindered group substituted aromatic rings were examined. All these benzenesulfonylhydrazides reacted smoothly with pentane-2,4-dione (**2a**) and the desired 3,5-dimethyl-1-(phenylsulfonyl)-1H-pyrazoles were obtained efficiently in good yields (Table 2, **3aa**~**3la**), ensuring a broad range of substrates. The yields remained relatively stable with varying the nature of the substituents on the benzenesulfonylhydrazide. Concerning the steric effect of the substituents on benzenesulfonylhydrazide, the substrates bearing mono or multi methyl or fluorine groups afforded the desired products in good to excellent yields (Table 2, **3aa**, **3ca**, **3ea**, **3fa**). Additionally, the substrates containing heteroaromatic and condensed ring structures were also able to produce the products efficiently (Table 2, **3ka**~**3la**).

To further explore the applicability and compatibility of this synthetic route, the scope of diketone under the same reaction conditions was examined (Table 3). A range of diketone underwent the reaction conditions smoothly and gave the corresponding products in excellent yields, such as 2,6-dimethylheptane-3,5-dione, 1,3-di-phenylpropane-1,3-dione, etc. (Table 3, **3aa**~**3af**). Heptane-3,5-dione reacted excellently with yield up to 98%, while 3-methylpentane-2,4-dione and 3-chloropentane-2,4-dione gave moderate results (Table 3, **3ac**, **3ab**, **3af**).

Table 3 Synthesis of 3,5-dimethyl-1-tosyl-1H-pyrazoles from 4-methylbenzenesulfonylhydrazide and substituted pentane-2,4-diones^a

^a Reaction conditions: **1a** (0.2 mmol, 1 equiv.), **2** (0.3 mmol, 1.5 equiv.), in 1.5 mL of CH₃CN at 60 °C under air for 2 h.

To showcase the practicality of this method, the gram

scale reaction was further carried out. 4-Methylbenzenesulfonylhydrazide (**1a**) and pentane-2,4-dione (**2a**) was scaled up to 20 mmol under standard reaction conditions. The corresponding product 3,5-dimethyl-1-tosyl-1H-pyrazole (**3aa**) was obtained in 94% yield without a significant loss (Scheme 2).

Scheme 2 Gram scale reaction

On the basis of the above results and previous literature reports,^[26] a plausible mechanism is proposed for the formation of the substituted pyrazoles as illustrated in Scheme 3. The reaction is initiated by the reaction of 4-methylbenzenesulfonylhydrazide (**1a**) and pentane-2,4-dione (**2a**) to give the intermediate **A**, followed by the elimination of H₂O to form intermediate **B**. Then intramolecular nucleophilic addition of **B** produces the intermediate **C**, which converts to the final product **3aa** by the elimination of H₂O.

3 Conclusions

In conclusion, a novel and efficient protocol for the installation of substituted pyrazoles in one step by using benzenesulfonylhydrazides and pentane-2,4-diones as starting materials under catalyst-free conditions was successfully disclosed. The method demonstrated excellent tolerance to a variety of functional groups, giving the corresponding substituted pyrazoles in good to excellent yields under mild conditions. Comprehensively, this catalyst-free system was found to be highly effective although on a 20 mmol scale. Step economy, simple operation with inexpensive reagents and mild reaction conditions render this novel and efficient strategy to be highly useful. This approach will promote the development of the synthesis of pyrazoles in pharmaceutical industry, agrochemicals and materials science.

4 Experimental section

4.1 General experimental details

¹H NMR spectra were recorded at 400 MHz. Chemical shifts were referenced to tetramethylsilane ($\delta=0.00$) as an internal standard in CDCl₃. ¹³C NMR spectra were obtained at 100 MHz and were calibrated with CDCl₃ ($\delta=77.00$). The high resolution mass spectra (HRMS) were recorded on an FT-ICR mass spectrometer using electrospray ionization (ESI). Products were purified by flash chromatography on 200~300 mesh silica gels. Unless otherwise noted, commercially available reagents and solvents were used without further purification. All melting points were determined without correction.

4.2 Typical procedure for the preparation of 3,5-dimethyl-1-tosyl-1H-pyrazoles 3

A test tube was charged with **1** (0.2 mmol), **2** (0.3 mmol) in CH₃CN (1.5 mL). Then the reaction mixture was stirred at 60 °C (oil bath temperature) under air atmosphere. After cooling to room temperature, the solvent was extracted with ethyl acetate and washed with brine, dried with Na₂SO₄. After the solvent was evaporated *in vacuo*, the residues were purified by column chromatography, eluting with petroleum ether/ethyl acetate to afford pure **3a**.

3,5-Dimethyl-1-tosyl-1H-pyrazole (**3aa**): Yellow solid. m.p. 93~95 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 7.83 (d, $J=8.4$ Hz, 2H), 7.30 (d, $J=8.3$ Hz, 2H), 5.90 (s, 1H), 2.49 (s, 3H), 2.41 (s, 3H), 2.20 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ : 153.31, 145.08, 143.97, 135.24, 129.81, 127.45, 110.66, 21.56, 13.75, 13.01; HRMS calcd for C₁₂H₁₅N₂O₂S [M+H]⁺ 251.0849, found 251.0851.

3,5-Dimethyl-1-(phenylsulfonyl)-1H-pyrazole (**3ba**): White solid. m.p. 79~81 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 7.95 (d, $J=7.2$ Hz, 2H), 7.62 (t, $J=7.5$ Hz, 1H), 7.52 (t, $J=7.7$ Hz, 2H), 5.91 (s, 1H), 2.50 (s, 3H), 2.21 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ : 153.59, 144.19, 138.26, 133.94, 129.23, 127.47, 110.84, 13.80, 13.08; HRMS calcd for C₁₁H₁₃N₂O₂S [M+H]⁺ 237.0692, found 237.0695.

Scheme 3 Proposed mechanism

1-(Mesitylsulfonyl)-3,5-dimethyl-1*H*-pyrazole (**3ca**): Yellow solid. m.p. 106~108 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 6.96 (s, 2H), 5.88 (s, 1H), 2.56 (s, 6H), 2.45 (s, 3H), 2.30 (s, 3H), 2.15 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 151.78, 144.01, 143.58, 140.62, 132.63, 131.96, 109.37, 22.36, 21.00, 13.71, 12.83; HRMS calcd for C₁₄H₁₉N₂O₂S [M + H]⁺ 279.1162, found 279.1159.

1-((4-Methoxyphenyl)sulfonyl)-3,5-dimethyl-1*H*-pyrazole (**3da**): Yellow solid. m.p. 127~129 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.90 (d, *J*=9.0 Hz, 2H), 6.97 (d, *J*=9.0 Hz, 2H), 5.89 (s, 1H), 3.86 (s, 3H), 2.49 (s, 3H), 2.20 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 163.87, 153.20, 143.85, 129.86, 129.68, 114.40, 110.56, 55.67, 13.82, 13.10; HRMS calcd for C₁₂H₁₅N₂O₃S [M + H]⁺ 267.0798, found 267.0799.

1-((4-Fluorophenyl)sulfonyl)-3,5-dimethyl-1*H*-pyrazole (**3ea**): Yellow solid. m.p. 88~90 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.02~7.96 (m, 2H), 7.23~7.16 (m, 2H), 5.92 (s, 1H), 2.50 (s, 3H), 2.20 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 165.78 (d, *J*=256 Hz), 153.82, 144.23, 134.27 (d, *J*=3 Hz), 130.49 (d, *J*=10 Hz), 116.60 (d, *J*=23 Hz), 110.92, 13.76, 13.07; HRMS calcd for C₁₁H₁₂FN₂O₂S [M + H]⁺ 255.0598, found 255.0602.

1-((2,4-Difluorophenyl)sulfonyl)-3,5-dimethyl-1*H*-pyrazole (**3fa**): White solid. m.p. 106~108 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.16~8.09 (m, 1H), 7.10~7.03 (m, 1H), 6.94~6.87 (m, 1H), 5.94 (s, 1H), 2.57 (s, 3H), 2.17 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 166.67 (dd, *J*=259, 12 Hz), 160.02 (dd, *J*=260, 13 Hz), 154.06, 145.64, 132.94 (d, *J*=11 Hz), 122.81 (dd, *J*=13, 4 Hz), 112.28 (dd, *J*=22, 4 Hz), 110.35, 105.85 (dd, *J*=26, 24 Hz), 13.75, 13.04 (d, *J*=4 Hz); HRMS calcd for C₁₁H₁₁F₂N₂O₂S [M + H]⁺ 273.0504, found 273.0501.

1-((2-Chlorophenyl)sulfonyl)-3,5-dimethyl-1*H*-pyrazole (**3ga**): Yellow solid. m.p. 97~99 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.16 (d, *J*=8.1 Hz, 1H), 7.62~7.51 (m, 1H), 7.51~7.39 (m, 2H), 5.93 (s, 1H), 2.59 (s, 3H), 2.16 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 153.53, 146.05, 136.08, 134.93, 132.66, 131.93, 131.78, 127.21, 109.68, 13.75, 13.44; HRMS calcd for C₁₁H₁₂ClN₂O₂S [M + H]⁺ 271.0303, found 271.0305.

1-((3-Bromophenyl)sulfonyl)-3,5-dimethyl-1*H*-pyrazole (**3ha**): Yellow solid. m.p. 93~95 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.09 (s, 1H), 7.88 (d, *J*=7.7 Hz, 1H), 7.73 (d, *J*=7.2 Hz, 1H), 7.41 (t, *J*=8.0 Hz, 1H), 5.94 (s, 1H), 2.51 (s, 3H), 2.21 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 154.08, 144.36, 139.82, 136.99, 130.71, 130.22, 126.06, 123.01, 111.12, 13.78, 13.08; HRMS calcd for C₁₁H₁₂BrN₂O₂S [M + H]⁺ 314.9798, found 314.9802.

3,5-Dimethyl-1-((4-(trifluoromethyl)phenyl)sulfonyl)-1*H*-pyrazole (**3ia**): Yellow solid. m.p. 96~98 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.10 (d, *J*=8.3 Hz, 2H), 7.80 (d, *J*=8.4 Hz, 2H), 5.95 (s, 1H), 2.52 (s, 3H), 2.21 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 154.39, 144.56, 141.59, 135.42 (q, *J*=33 Hz), 128.12, 126.40 (q, *J*=4 Hz), 122.91 (q, *J*=272 Hz), 111.29, 13.77,

13.07; HRMS calcd for C₁₂H₁₂F₃N₂O₂S [M + H]⁺ 305.0566, found 305.0562.

3,5-Dimethyl-1-((4-nitrophenyl)sulfonyl)-1*H*-pyrazole (**3ja**): Yellow solid. m.p. 178~180 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.37 (d, *J*=8.9 Hz, 2H), 8.16 (d, *J*=8.9 Hz, 2H), 5.97 (s, 1H), 2.53 (s, 3H), 2.21 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 154.80, 150.63, 144.73, 143.44, 128.94, 124.43, 111.50, 13.76, 13.07; HRMS calcd for C₁₁H₁₂N₃O₄S [M + H]⁺ 282.0543, found 282.0541.

2-((3,5-Dimethyl-1*H*-pyrazol-1-yl)sulfonyl)pyridine (**3ka**): Yellow solid. m.p. 80~82 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 9.14 (d, *J*=1.9 Hz, 1H), 8.84 (dd, *J*=4.9, 1.6 Hz, 1H), 8.31~8.22 (m, 1H), 7.50 (dd, *J*=8.0, 4.9 Hz, 1H), 5.95 (s, 1H), 2.53 (s, 3H), 2.20 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 154.46, 154.32, 148.14, 144.57, 135.37, 134.98, 123.76, 111.26, 13.76, 13.06; HRMS calcd for C₁₀H₁₂N₃O₂S [M + H]⁺ 238.0645, found 238.0646.

3,5-Dimethyl-1-(naphthalen-2-ylsulfonyl)-1*H*-pyrazole (**3la**): White solid. m.p. 104~106 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 8.57 (s, 1H), 8.00~7.91 (m, 2H), 7.87 (d, *J*=8.8 Hz, 2H), 7.71~7.55 (m, 2H), 5.89 (s, 1H), 2.55 (s, 3H), 2.19 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 153.52, 144.14, 135.24, 135.07, 131.84, 129.56, 129.45, 129.28, 127.86, 127.70, 122.05, 110.78, 13.75, 13.10; HRMS calcd for C₁₅H₁₅N₂O₂S [M + H]⁺ 287.0849, found 287.0850.

3,4,5-Trimethyl-1-tosyl-1*H*-pyrazole (**3ab**): Yellow solid. m.p. 143~145 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.80 (d, *J*=8.4 Hz, 2H), 7.28 (d, *J*=8.3 Hz, 2H), 2.40 (s, 6H), 2.15 (s, 3H), 1.83 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 153.62, 144.81, 139.53, 135.51, 129.75, 127.39, 117.15, 21.56, 12.34, 11.24, 7.93; HRMS calcd for C₁₃H₁₇N₂O₂S [M + H]⁺ 265.1005, found 265.1009.

3,5-Diethyl-1-tosyl-1*H*-pyrazole (**3ac**): Yellow oil. ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.82 (d, *J*=8.4 Hz, 2H), 7.29 (d, *J*=8.7 Hz, 2H), 5.97 (s, 1H), 2.92 (q, *J*=7.5 Hz, 2H), 2.59 (q, *J*=7.6 Hz, 2H), 2.40 (s, 3H), 1.26 (t, *J*=7.5 Hz, 3H), 1.18 (t, *J*=7.6 Hz, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 158.96, 150.53, 144.94, 135.40, 129.72, 127.47, 107.16, 21.59, 21.55, 20.35, 13.04, 12.82; HRMS calcd for C₁₄H₁₉N₂O₂S [M + H]⁺ 279.1162, found 279.1160.

3,5-Di-*iso*-propyl-1-tosyl-1*H*-pyrazole (**3ad**): Yellow oil. ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.81 (d, *J*=8.1 Hz, 2H), 7.28 (d, *J*=8.2 Hz, 2H), 5.98 (s, 1H), 3.60 (p, *J*=6.8 Hz, 1H), 2.93 (p, *J*=6.9 Hz, 1H), 2.40 (s, 3H), 1.25 (d, *J*=6.8 Hz, 6H), 1.18 (d, *J*=6.9 Hz, 6H); ¹³C NMR (100 MHz, Chloroform-*d*) δ: 163.29, 156.20, 144.79, 135.59, 129.57, 127.50, 104.16, 27.98, 26.26, 23.36, 22.02, 21.56; HRMS calcd for C₁₆H₂₃N₂O₂S [M + H]⁺ 307.1475, found 307.1478.

3,5-Diphenyl-1-tosyl-1*H*-pyrazole (**3ae**): White solid. m.p. 119~121 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ: 7.86~7.82 (m, 2H), 7.62 (d, *J*=8.4 Hz, 2H), 7.49~7.35 (m, 8H), 7.19 (d, *J*=8.2 Hz, 2H), 6.60 (s, 1H), 2.35 (s,

3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 155.11, 149.39, 145.24, 134.86, 131.33, 129.96, 129.58, 129.41, 129.24, 128.65, 128.01, 127.76, 127.14, 126.41, 109.44, 21.62; HRMS calcd for $\text{C}_{22}\text{H}_{19}\text{N}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 375.1162, found 375.1163.

4-Chloro-3,5-dimethyl-1-tosyl-1*H*-pyrazole (**3af**): White solid. m.p. 133~135 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.84 (d, $J=7.8$ Hz, 2H), 7.33 (d, $J=8.0$ Hz, 2H), 2.49 (s, 3H), 2.42 (s, 3H), 2.21 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 150.81, 145.63, 139.21, 134.63, 129.99, 127.66, 113.68, 21.62, 11.79, 11.15; HRMS calcd for $\text{C}_{12}\text{H}_{14}\text{ClN}_2\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 285.0459, found 285.0456.

Supporting Information ^1H NMR and ^{13}C NMR spectra of products **3**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Brown, A. W. *Adv. Heterocycl. Chem.* **2018**, 126, 55.
(b) Elkanzi, N. A. A. *J. Chin. Chem. Soc.* **2018**, 65, 189.
(c) Pizzuti, L.; Barschak, A. G.; Stefanello, F. M.; Farias, M. D.; Lencina, C.; Roesch-Ely, M.; Cunico, W.; Moura, S.; Pereira, C. M. *P. Curr. Org. Chem.* **2014**, 18, 115.
(d) Janin, Y. L. *Chem. Rev.* **2012**, 112, 3924.
(e) Zhang, Q.; Tang, M. *Org. Lett.* **2019**, 21, 1917.
(f) Sun, J.; Qiu, J. K.; Zhu, Y. L.; Guo, C.; Hao, W. J.; Jiang, B.; Tu, S. J. *J. Org. Chem.* **2015**, 80, 8217.
(g) Li, G. L.; Huang, H. F.; Yang, J.; Duan, H. Z. *Chin. J. Org. Chem.* **2021**, 41, 1466 (in Chinese).
(李光磊, 黄海丰, 杨军, 段红珍, 有机化学, **2021**, 41, 1466.)
(h) Chen, Y.; Zhu, J. N.; Zhao, S. Y. *Chin. J. Org. Chem.* **2019**, 39, 1923 (in Chinese).
(陈樱, 祝家楠, 赵圣印, 有机化学, **2019**, 39, 1923.)
(i) Chen, D. X.; Xiong, H. L.; Yang, H. W.; Tang, J.; Cheng, G. B. *FirePhysChem* **2021**, 1, 71.
(j) Daling, I. L.; Shkineva, T. K.; Vatsadze, I. A.; Kormanov, A. V.; Kozev, A. M.; Suponitsky, K. Y.; Pivkina, A. N.; Sheremetev, A. B. *FirePhysChem* **2021**, 1, 83.
(k) Zhu, X.; Hu, S.; Jiang, L.; Yang, W.; Yang, G. *Sci. Sin.: Chim.* **2016**, 46, 1188.
- [2] (a) Abdel-Magid, A. F. *ACS Med. Chem. Lett.* **2014**, 5, 730.
(b) Wang, S. H.; Zhang, B. H.; Chen, J.; Zheng, Y. Y.; Feng, N.; Ma, A. J.; Xu, X. T.; Abdullah, M. A. *Chin. J. Org. Chem.* **2020**, 40, 15 (in Chinese).
(王少华, 张帮红, 陈洁, 郑莹莹, 冯娜, 马爱军, 徐学涛, Abdullah M. Asiri, 有机化学, **2020**, 40, 15.)
- [3] (a) Habeeb, A. G.; Rao, P. N. P.; Knaus, E. E. *J. Med. Chem.* **2001**, 44, 3039.
(b) Li, Y.; Wan, J. P. *Chin. J. Org. Chem.* **2020**, 40, 3889 (in Chinese).
(李毅, 万结平, 有机化学, **2020**, 40, 3889.)
(c) Wang, Y. C.; Wang, H. S.; Huang, G. B.; Huang, F. P.; Hu, K.; Pan, Y. M. *Tetrahedron* **2014**, 70, 1621.
- [4] (a) Sridhar, R.; Perumal, P. T.; Etti, S.; Shanmugam, G.; Ponnuswamy, M. N.; Prabavathy, V. R.; Mathivanan, N. *Bioorg. Med. Chem. Lett.* **2004**, 14, 6035.
(b) Liu, J. J.; Sun, J.; Fang, Y. B.; Yang, Y. A.; Jiao, R. H.; Zhu, H. L. *Org. Biomol. Chem.* **2014**, 12, 998.
(c) Zeng, J. L.; Xu, Z. H.; Ma, J. A. *Chin. J. Org. Chem.* **2020**, 40, 1105 (in Chinese).
(曾俊良, 许志红, 马军安, 有机化学, **2020**, 40, 1105.)
- [5] Beberitz, G. R.; Argentieri, G.; Battle, B.; Brennan, C.; Balkan, B.; Burkey, B. F.; Eckhardt, M.; Gao, J.; Kapa, P.; Strohschein, R. J.; Schuster, H. F.; Wilson, M.; Xu, D. D. *J. Med. Chem.* **2001**, 44, 2601.
- [6] Chang, C. P.; Wu, C. H.; Song, J. S.; Chou, M. C.; Wong, Y. C.; Lin, Y. C.; Yeh, T. K.; Sadani, A. A.; Ou, M. H.; Chen, K. H.; Chen, P. H.; Kuo, P. C.; Tseng, C. T.; Chang, K. H.; Tseng, S. L.; Chao, Y. S.; Hung, M. S.; Shia, K. S. *J. Med. Chem.* **2013**, 56, 9920.
- [7] (a) Ouyang, G.; Cai, X. J.; Chen, Z.; Song, B. A.; Bhadury, P. S.; Yang, S.; Jin, L. H.; Xue, W.; Hu, D. Y.; Zeng, S. J. *Agric. Food Chem.* **2008**, 56, 10160.
(b) Wu, Z.; Li, X.; Zhou, X.; Yang, W.; Ye, Y.; Song, B.; Yang, S. *Sci. Sin.: Chim.* **2016**, 46, 1195.
- [8] (a) Xu, Z.; Gao, C.; Ren, Q. C.; Song, X. F.; Feng, L. S.; Lv, Z. S. *Eur. J. Med. Chem.* **2017**, 139, 429.
(b) Ansari, A.; Ali, A.; Asif, M.; Shamsuzzaman. *New J. Chem.* **2017**, 41, 16.
- [9] (a) McGrath, N. A.; Brichacek, M.; Njardarson, J. T. *J. Chem. Educ.* **2010**, 87, 1348.
(b) Ilardi, E. A.; Vitaku, E.; Njardarson, J. T. *J. Chem. Educ.* **2013**, 90, 1403.
(c) Zhang, J.; Shao, Y.; Wang, H. X.; Luo, Q.; Chen, J. J.; Xu, D. M.; Wan, X. B. *Org. Lett.* **2014**, 16, 3312.
- [10] (a) Katsuhiko, K.; Masahiro, S.; Takuo, K. *Plant Physiol.* **1979**, 64, 774.
(b) Su, S. M.; Zhu, M.; Zhang, D. Q.; Yuan, D. K. *Chin. J. Org. Chem.* **2019**, 39, 2026 (in Chinese).
(苏诗淼, 朱墨, 张大强, 袁德凯, 有机化学, **2019**, 39, 2026.)
- [11] Kazutoshi, W.; Yasuhiro, M.; Katsuhiko, I.; Toshiaki, W.; Satoshi, Y.; Hiroyoshi, N. *Redox Rep.* **2003**, 8, 151.
- [12] Neumann, J. J.; Suri, M.; Glorius, F. *Angew. Chem., Int. Ed.* **2010**, 49, 7790.
- [13] Shao, N.; Chen, T.; Zhang, T.; Zhu, H.; Zheng, Q.; Zou, H. *Tetrahedron* **2014**, 70, 795.
- [14] (a) Knorr, L. *Ber. Dtsch. Chem. Ges.* **1883**, 16, 2597.
(b) Wang, Z. X.; Qin, H. L. *Green Chem.* **2004**, 6, 90.
(c) Katritzky, A. R. *Handbook of Heterocyclic Chemistry*, Pergamon, Oxford, **1985**, p. 416.
(d) Hayter, M. J.; Bray, D. J.; Clegg, J. K.; Lindoy, L. F. *Synth. Commun.* **2006**, 36, 707.
(e) Sanchez-Carmona, M. A.; ContrerasCruz, D. A.; Miranda, L. D. *Org. Biomol. Chem.* **2011**, 9, 6506.
(f) Vaddula, B. R.; Varma, R. S.; Leazer, J. *Tetrahedron Lett.* **2013**, 54, 1538.
(g) Zakerinasab, B.; Nasser, M. A.; Hassani, H.; Samiadel, M. M. *Res. Chem. Intermed.* **2016**, 42, 3169.
(h) Polshettiwar, V.; Varma, R. S. *Tetrahedron Lett.* **2008**, 49, 397.
- [15] (a) Liu, H. L.; Jiang, H. F.; Zhang, M.; Yao, W. J.; Zhu, Q. H.; Tang, Z. *Tetrahedron Lett.* **2008**, 49, 3805.
(b) Grotjahn, D. B.; Van, S.; Combs, D.; Lev, D. A.; Schneider, C.; Rideout, M.; Meyer, C.; Hernandez, G.; Mejorado, L. *J. Org. Chem.* **2002**, 67, 9200.
(c) Bagley, M. C.; Lubinu, M. C.; Mason, C. *Synlett* **2007**, 704.
(d) Sar, D.; Bag, R.; Yashmeen, A.; Bag, S. S.; Punniyamurthy, T. *Org. Lett.* **2015**, 17, 5308.
(e) Zhang, H.; Wei, Q.; Zhu, G.; Qu, J.; Wang, B. *Tetrahedron Lett.* **2016**, 57, 2633.
- [16] (a) McLaughlin, M.; Marcantonio, K.; Chen, C. Y.; Davies, I. W. *J. Org. Chem.* **2008**, 73, 4309.
(b) Dishington, A.; Feron, J. L.; Gill, K.; Graham, M. A.; Hollingsworth, I.; Pink, J. H.; Roberts, A.; Simpson, I.; Tatton, M. *Org. Lett.* **2014**, 16, 6120.
- [17] Hu, J.; Chen, S.; Sun, Y.; Yang, J.; Rao, Y. *Org. Lett.* **2012**, 14, 5030.
- [18] (a) Yang, G. P.; He, X.; Yu, B.; Hu, C. W. *Appl. Organomet. Chem.* **2018**, 32, 4532.
(b) Wang, H. F.; Sun, X. L.; Zhang, S. L.; Liu, G. L.; Wang, C. J.; Zhu, L. L.; Zhang, H. *Synlett* **2018**, 29, 2689.
(c) Venkateswarlu, V.; Kour, J.; Aravinda Kumar, K. A.; Verma, P. K.; Lakshma Reddy, G.; Hussain, Y.; Tabassum, A.; Balgotra, S.;

- Gupta, S.; Hudwekar, A. D.; Vishwakarma, R. A.; Sawant, S. D. *RSC Adv.* **2018**, 8, 26523.
- [19] Xiong, W.; Chen, J. X.; Liu, M. C.; Ding, J. C.; Wu, H. Y.; Su, W. K. *J. Braz. Chem. Soc.* **2009**, 20, 367.
- [20] Polshettiwar, V.; Varma, R. S. *Tetrahedron Lett.* **2008**, 49, 397.
- [21] Wang, Z.; Qin, H. *Green Chem.* **2004**, 6, 90.
- [22] Curini, M.; Rosati, O.; Campagna, V.; Montanari, F.; Cravotto, G.; Boccalini, M. *Synlett* **2005**, 2927.
- [23] Zhao, X.; Zhang, L.; Li, T.; Liu, G.; Wang, H.; Lu, K. *Chem. Commun.* **2014**, 50, 13121.
- [24] (a) Liu, Y.; Meng, Y. N.; Huang, X. J.; Qin, F. H.; Wu, D. P.; Shao, Q. J.; Guo, Zh. Y.; Li, Q.; Wei, W. T. *Green Chem.* **2020**, 22, 4593.
(b) Huang, X. J.; Qin, F. H.; Liu, Y.; Wu, Sh. P.; Li, Q.; Wei, W. T. *Green Chem.* **2020**, 22, 3952.
(c) Kang, Q. Q.; Wu, W. F.; Li, Q.; Wei, W. T. *Green Chem.* **2020**, 22, 3060.
- [25] (a) Qin, F. H.; Huang, X. J.; Liu, Y.; Liang, H.; Li, Q.; Cao, Zh.; Wei, W. T.; He, W. M. *Chin. Chem. Lett.* **2020**, 31, 3267.
(b) Wang, G. D.; Guo, Y. H.; Wan, J. P. *Chin. J. Org. Chem.* **2020**, 40, 645 (in Chinese).
(王国防, 郭艳辉, 万结平, 有机化学, **2020**, 40, 645.)
(c) Dong, D. Q.; Li, G. H.; Chen, D. M.; Sun, Y. Y.; Han, Q. Q.; Wang, Z. L.; Xu, X. M.; Yu, X. Y. *Chin. J. Org. Chem.* **2020**, 40, 1766 (in Chinese).
(董道青, 李光辉, 陈德茂, 孙媛媛, 韩晴晴, 王祖利, 徐鑫明, 于贤勇, 有机化学, **2020**, 40, 1766.)
- [26] Fustero, S.; Sanchez-Rosello, M.; Barrio, P.; Simon-Fuentes, A. *Chem. Rev.* **2011**, 111, 6984.

(Zhao, C.)