碱催化 α -氰基- β -甲基烯基(杂)芳基酮苯增环反应合成多取代苯

安逸 张放 蔡志华 杜广芬*

(石河子大学化工学院 新疆兵团化工绿色过程重点实验室 新疆石河子 832000)

摘要 发展了一种非金属催化下高效构筑多取代苯的新方法. 以 10 mol% 的 Cs_2CO_3 为催化剂, 一系列 α -氰基- β -甲基烯基(杂)芳基酮可在极其温和的反应条件下与丁炔二酸酯发生[4+2]环加成/脱水芳构化苯增环反应, 以 62%~94% 的收率生成 1,2-二酯基-3-(杂)芳基-4-氰基苯衍生物.

关键词 α -氰基- β -甲基烯基(杂)芳基酮; 丁炔二酸酯; [4+2]环加成反应; 1,2-二酯基-3-(杂)芳基-4-氰基苯衍生物

Direct Assembly of Polysubstituted Benzenes via Base-Catalyzed Benzannulation Reaction of α -Cyano- β -methylalkenyl-(hetero)aryl Ketones

An, Yi Zhang, Fang Cai, Zhihua Du, Guangfen*

(Key Laboratory for Green Processing of Chemical Engineering of Xinjiang Bingtuan, School of Chemistry and Chemical Engineering, Shihezi University, Shihezi, Xinjiang 832000)

Abstract A mild and transition-metal free method for rapid construction of benzene frameworks has been developed. Under the catalysis of 10 mol% Cs_2CO_3 , a variety of α -cyano- β -methylalkenyl(hetero)aryl ketones undergo [4+2] annulation with different dialkyl butynedioates efficiently to produce 1,2-diesteryl-3-(hetero)aryl-4-cyanobenzene derivatives in 62%~94% yields.

Keywords α -cyano- β -methylalkenyl(hetero)aryl ketones; dialkyl butynedioates; [4+2] annulation; 1,2-diesteryl-3-(hetero)-aryl-4-cyanobenzene derivatives

1 Introduction

Polysubstituted benzenes are fundamental structural motifs found in various natural products, biologically active compounds, pharmaceuticals and functional materials.^[1] As a result, different methods have been developed for the construction of these important compounds. Traditionally, electrophilic or nucleophile substitution reactions^[2] of benzenes and transition-metal-catalyzed cross coupling reactions^[3] of halogenated benzenes provide powerful strategies for the synthesis of polysubstituted benzene derivatives. Alternatively, transition-metal-catalyzed [2+2+2] cyclotrimizations^[4] and [4+2] annulations^[5] have been proven to be facile methods for the preparation of benzene frameworks from acyclic substrates. Despite remarkable progress made in this area, these aforementioned methods often suffer from harsh reaction conditions and heavy met-

al wastes. Therefore, the development of mild and efficient methods for direct assembly of polysubstituted benzenes under transition-metal-free^[6] conditions is highly significant.

As important C4 synthon partners, α -cyano- β -methyl-enones have been successfully used in the synthesis of different aromatic compounds. In 2016, Ye and Wang^[7] independently pioneered an oxidative N-heterocyclic carbene (NHC)-catalyzed [4+2] annulation between α -cyano- β -methyl-enones and enals, which provides a novel method for the synthesis of polysubstituted benzenes. In the same year, Ye and coworkers^[8] reported another NHC-catalyzed annulation of α -cyano- β -methyl-enones and α -bromo-enals, which offers an efficient protocol for facile access to hetero-1,3,5-trisubstituted benzenes. Later, the same group^[9] further developed a more convenient method for direct assembly of 1,3,5-trisubstituted benzenes via 1,8-diazabi-

* Corresponding author. E-mail: duguangfen@shzu.edu.cn

Received April 27, 2021; revised June 7, 2021; published online June 28, 2021.

Project supported by the National Natural Science Foundation of China (No. 21662029), the Excellent Young Teachers Plan of Bingtuan (Nos. 2017CB001, CZ027203) and the International Cooperation Project of Shihezi University (No. GJHZ201801).

国家自然科学基金(No. 21662029)、兵团优秀青年教师(Nos. 2017CB001, CZ027203)和石河子大学国际合作(No. GJHZ201801)资助项目.

cyclo[5.4.0]undec-7-ene (DBU)-mediated annulation of α -cyano- β -methyleneones and α , β -unsaturated carboxylic acids. In 2018, Shi and co-workers^[10] documented a base promoted tandem reaction of α -cyano- β -methyleneones and α , β -unsaturated enones, which provides an efficient method for the construction of polysubstituted benzenes. We^[11] have been interested in the synthesis of aromatic derivatives through nontransition-metal catalysed reactions. Recently, we^[12] found that α -cyano- β -methyleneones can undergo [4+2] benzannulation reaction with the active carbon-carbon triple bond of benzyne to produce polysubstituted naphthalenes. Based on this previous work, we envisioned that α -cyano- β -methylalkenyl(hetero)aryl ketones could undergo the similar annulation reaction with the active carbon-carbon triple bond of dialkyl butynediates to construct polysubstituted benzenes.

2 Results and discussion

With this idea in mind, the study was initiated by using the readily available enone **1a** and dimethyl butynedioate (DMAD) **2a** as model substrates to optimize the reaction conditions. In the presence of 20 mol% Cs₂CO₃, enone **1a** reacted with dimethyl butynedioate (DMAD) **2a** smoothly in acetonitrile at room temperature to produce polysubstituted benzene **3a** in 84% yield within 3 h (Table 1, Entry 1). Encouraged by this success, several other common bases were subsequently tested for the reaction. Inorganic bases, such as Na₂CO₃, K₂CO₃, NaOH and KOH, can promote the annulation efficiently to afford **3a** in good yields (Table 1, Entries 2~5). Organic bases triethylamine, the Hünig's base and DBU catalyzed the reaction to give **3a** in high yields (Table 1, Entries 6~8). Sodium acetate can also catalyze the annulation in 84% yield (Table 1, Entry 9), but prolonged reaction time is required. Increasing the amount of Cs₂CO₃ to 50 mol% did not improve the reaction yield (Table 1, Entry 10). Interestingly, reducing the base loading to 10 mol% resulted in higher reaction yield (Table 1, Entries 11~13). However, further lowering the base loading to 5 mol% led to dramatic decrease of the reaction yield (Table 1, Entry 14). A brief evaluation of the reaction media revealed that acetonitrile was the best choice in terms of reaction yield (Table 1, Entries 15~18). Increasing the amount of DMAD to 1.5 equiv. cannot improve the yield of **3a** (Table 1, Entry 19). Finally, the control experiment showed that no desired product was formed without the addition of a base (Table 1, Entry 20).

With the optimal reaction conditions in hand, the substrate scope of the annulation reaction was then evaluated and the results were summarized in Table 2. A variety of α -cyano- β -methylalkenylaryl ketones bearing electron-neutral, electron-donating and electron-withdrawing substituents on the β -aryl ring participated in the reaction efficiently, giving the corresponding 1,2-diesteryl-3-aryl-4-cyanobenzene derivatives in high yields (Table 2, Entries 1~5). In addition, different positions of the substituents

Table 1 Optimization of the reaction conditions^a

Entry	Base	Time/h	Solvent	Yield ^b /%
1	Cs ₂ CO ₃ (20 mol%)	3	CH ₃ CN	84
2	Na ₂ CO ₃ (20 mol%)	5	CH ₃ CN	73
3	K ₂ CO ₃ (20 mol%)	5	CH ₃ CN	80
4	NaOH (20 mol%)	5	CH ₃ CN	75
5	KOH (20 mol%)	5	CH ₃ CN	83
6	Et ₃ N (20 mol%)	4	CH ₃ CN	72
7	DIPEA (20 mol%)	12	CH ₃ CN	83
8	DBU (20 mol%)	5	CH ₃ CN	90
9	AcONa (20 mol%)	36	CH ₃ CN	85
10	Cs ₂ CO ₃ (50 mol%)	3	CH ₃ CN	77
11	Cs ₂ CO ₃ (10 mol%)	3	CH ₃ CN	94
12	Et ₃ N (10 mol%)	5	CH ₃ CN	89
13	DBU (10 mol%)	5	CH ₃ CN	92
14	Cs ₂ CO ₃ (5 mol%)	5	CH ₃ CN	75
15	Cs ₂ CO ₃ (10 mol%)	24	Toluene	30
16	Cs ₂ CO ₃ (10 mol%)	5	CH ₂ Cl ₂	70
17	Cs ₂ CO ₃ (10 mol%)	3	ClCH ₂ CH ₂ Cl	70
18	Cs ₂ CO ₃ (10 mol%)	4	1,4-Dioxane	78
19 ^c	Cs ₂ CO ₃ (10 mol%)	3	CH ₃ CN	93
20	No base	24	CH ₃ CN	0

^a **1a** (0.10 mmol), **2a** (0.12 mmol), solvent (1.0 mL). ^b Isolated yield. ^c **1a** (0.10 mmol), **2a** (0.15 mmol).

had no apparent impact on the reaction yield (Table 2, Entries 6~10). β -Naphthyl substituted enone underwent the annulation to afford the desired product **3k** in 73% yield (Table 2, Entry 11). β -Heteroaryl substituted enone **1l** was proved to be successful candidate for the cyclization, producing substituted benzene **3l** in 81% yield (Table 2, Entry 12). Interestingly, β -Alkyl substituted enones coupled with DMAD efficiently to furnish the corresponding hexa-substituted benzenes in good yields (Table 2, Entries 13 and 14). β , β -Dimethyl substituted enone also performed very well, delivering the product **3o** in 72% yield (Table 2, Entry 15). Enones with electron-donating substituents on the α -aryl ring underwent the cyclization to give the corresponding products in high yields (Table 2, Entries 16 and 17). α -Heteroaryl substituted enone **1r** also reacted smoothly to form the desired product **3r** in 81% yield (Table 2, Entry 18). Diethyl acetylenedicarboxylate (**2b**) and di-*tert*-butyl acetylenedicarboxylate (**2c**) were all suitable for the annulation reaction, giving the corresponding products in excellent yields (Table 2, Entries 19 and 20). Owing to the dramatic decrease of the electrophilicity, when ethyl propiolate **2d** was used instead of DMAD, no desired product was formed (Table 2, Entry 21). Enonates **1s**, **1t** and enone **1u** cannot undergo this annulation, and the starting materials were recovered in high yields (Table

Table 2 Evaluation of substrate scope^a

Entry	Ketone	Product	3, Yield ^b / % of 3
1	 1a	 3a	94 (92) ^c
2	 1b	 3b	75
3	 1c	 3c	91
4	 1d	 3d	89
5	 1e	 3e	90
6	 1f	 3f	84
7	 1g	 3g	90
8	 1h	 3h	75

Continued

Entry	Ketone	Product	3, Yield ^b % of 3
9	1i		3i , 89
10	1j		3j , 82
11	1k		3k , 73
12	1l		3l , 81
13	1m		3m , 80
14	1n		3n , 75
15	1o		3o , 72
16	1p		3p , 87
17	1q		3q , 80

Continued

Entry	Ketone	Product	3, Yield ^b / % of 3
18	1r	3r , 81	
19	1a	3s , 92	
20	1a	3t , 90	
21 ^d	1a	—	ND
22	1s	—	ND
23	1t	—	ND
24	1u	—	ND

^a Reaction conditions: **1a** (0.10 mmol), **2** (0.12 mmol), Cs₂CO₃ (10 mol%), CH₃CN (1.0 mL), 3 h, room temperature. ^b Isolated yield. ^c **1a** (5 mmol), **2** (6 mmol), Cs₂CO₃ (10 mol%), CH₃CN (30 mL), 6 h, room temperature. ^d Ethyl propiolate was used instead of **2a**.

2, Entries 22–24). We assumed that the acidity of the β -methyl protons of these compounds was weaker than that of enone **1a**. Therefore, the dienolate intermediate **I** (Scheme 1) might be difficult to be generated and the following annulation reaction cannot be initiated. Notably, the benzannulation reaction can be conducted on a gram-scale, and a high yield can be maintained (Table 2, Entry 1).

The product **3a** was crystallized from dichloromethane (DCM) and *n*-hexane, and its structure was clearly confirmed by single-crystal X-ray analysis (Figure 1).^[13]

Based on previous studies on [4+2] annulation reactions of α -cyano- β -methyleneones,^[7–10] a possible mechanism was proposed and depicted in Scheme 1. In the presence of Cs₂CO₃, enone **1a** undergoes deprotonation to generate dienolate intermediate **I**, which might undergo

Figure 1 Crystal structure of **3a**

Scheme 1 Proposed mechanism

cyclization with diethyl butynedioate to produce species **II**. The following protonation of **II** will lead to the formation of intermediate **III**. And the final product could be produced by the dehydrative aromatization of **III**.

3 Conclusions

In summary, an annulation reaction of α -cyano- β -methylalkenyl(hetero)aryl ketones and dialkyl butynedioates has been demonstrated. The mild and transition-metal-free conditions, simple procedure and generally high yields provide a convenient protocol for rapid construction of 1,2-diesterly-3-(hetero)aryl-4-cyanobenzene derivatives. Further studies on the synthesis of other aromatic derivatives through reactions of α -cyano- β -methyleneones are underway in our lab.

4 Experimental section

4.1 General methods

Unless otherwise indicated, all reactions were conducted under air with magnetic stirring bar. Column chromatography was performed with silica gel (200~300 mesh) and analytical thin-layer chromatography (TLC) on silica gel 60-F₂₅₄. ¹H NMR (400 MHz, CDCl₃) and ¹³C NMR (100 MHz, CDCl₃) were recorded on a Bruker-DMX 400 spectrometer in CDCl₃ with tetramethylsilane as an internal standard. α -Cyano- β -methyleneones and enonates were prepared according to literature procedure. Diethyl butynedioate **2** was purchased from Aladdin. Cs₂CO₃ was purchased from Bide Pharmatech. All other chemicals were obtained from commercial supplies and used as received without any further purification. Anhydrous toluene, DCM and CH₃CN were obtained from commercial suppliers. Petroleum ether has a boiling point range of 60~90 °C.

4.2 General procedure

10% Cs₂CO₃ (1.38 mg, 0.01 mmol) was added to a stirred solution of enone **1a** (24.71 mg, 0.10 mmol) and diethyl butynedioate **2a** (17.05 mg, 0.12 mmol, 1.2 equiv.) in dry CH₃CN (1 mL) under air. The resulting mixture was stirred at room temperature for 3 h. The solvent was then removed under reduced pressure and the crude material was purified by flash column chromatography (silica gel,

PE/EtOAc (*V*: *V*=5:1) to give the desired product dimethyl 2'-cyano[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3a**): White solid, 35.0 mg, 94% yield. m.p. 171.1~171.6 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 8.12 (s, 1H), 7.63~7.57 (m, 2H), 7.55~7.49 (m, 3H), 7.49~7.45 (m, 3H), 7.44~7.38 (m, 2H), 3.94 (s, 3H), 3.60 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ : 167.3, 164.8, 147.0, 145.3, 136.9, 135.3, 134.8, 130.8, 130.3, 129.4, 129.3, 129.1, 128.9, 128.8, 128.4, 116.1, 116.0, 53.1, 52.5; HR-MS (APCI) calcd for C₂₃H₁₈NO₄ [M+H]⁺ 372.1230, found 372.1225.

Dimethyl 2'-cyano-4-methoxy-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3b**): White solid, 31 mg, 75% yield. m.p. 195.7~196.2 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 8.10 (s, 1H), 7.61~7.57 (m, 2H), 7.56~7.46 (m, 3H), 7.37~7.30 (m, 2H), 7.01~6.96 (m, 2H), 3.94 (s, 3H), 3.86 (s, 3H), 3.65 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ : 167.3, 164.9, 160.6, 146.7, 145.3, 135.4, 134.2, 130.8, 130.1, 129.2, 129.1, 128.3, 116.4, 115.7, 114.3, 55.4, 53.1, 52.5; HRMS (APCI) calcd for C₂₄H₂₀NO₅ [M+H]⁺ 402.1335, found 402.1329.

Dimethyl 4-chloro-2'-cyano-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3c**): White solid, 36.5 mg, 91% yield. m.p. 228.8~229.3 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 8.09 (s, 1H), 7.60~7.35 (m, 7H), 7.43~7.34 (m, 2H), 3.95 (s, 3H), 3.60 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ : 167.1, 164.6, 145.7, 145.5, 135.9, 135.3, 135.2, 135.1, 131.0, 130.2, 130.0, 129.4, 129.3, 129.2, 129.1, 128.4, 115.9, 99.9, 53.2, 52.6; HRMS (APCI) calcd for C₂₃H₁₆ClNO₄ 405.0773, found 405.0775.

Dimethyl 4-bromo-2'-cyano-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3d**): White solid, 40 mg, 89% yield. m.p. 193.1~193.6 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 8.09 (s, 1H), 7.68~7.64 (m, 2H), 7.50~7.45 (m, 5H), 7.42~7.37 (m, 2H), 3.94 (s, 3H), 3.60 (s, 3H); ¹³C NMR (101 MHz, Chloroform-*d*) δ : 167.1, 164.6, 145.7, 145.5, 135.8, 135.2, 135.1, 132.1, 131.0, 130.4, 130.0, 129.4, 129.0, 128.4, 124.1, 116.0, 115.9, 53.2, 52.6; HRMS (APCI) calcd for C₂₃H₁₇BrNO₄ [M+H]⁺ 450.0335, found 450.0328.

Dimethyl 2'-cyano-4-fluoro-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3e**): White solid, 35 mg, 90% yield. m.p. 203.8~204.3 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 8.14 (s, 1H), 7.62~7.69 (m, 2H), 7.55~7.50 (m, 3H), 7.42~7.35 (m, 2H), 7.20~7.14 (m, 2H), 3.94 (s, 3H), 3.64 (s, 3H); ¹³C NMR (100 MHz, Chloroform-*d*) δ : 167.2, 164.6, 163.25 (*d*, *J*=248 Hz), 147.1, 144.2, 136.8, 134.9, 131.23 (*d*, *J*=4 Hz), 130.9, 130.5, 129.5, 128.8 (*d*, *J*=22 Hz), 116.2, 116.1, 115.6 (*d*, *J*=22 Hz), 53.2, 52.6; ¹⁹F NMR (376 MHz, Chloroform-*d*) δ : -111.47; HRMS (APCI) calcd for C₂₃H₁₇FNO₄ [M+H]⁺ 390.1136, found 390.1129.

Dimethyl 2'-cyano-2-methyl-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3f**): White solid, 32 mg, 84% yield. m.p. 175.1~175.6 °C; ¹H NMR (400 MHz, Chloroform-*d*) δ : 8.00 (s, 1H), 7.49~7.44 (m, 3H), 7.43~7.39 (m, 2H), 7.38~7.28 (m, 3H), 7.26~7.22 (m, 1H), 3.93 (s, 3H),

3.63 (s, 3H), 2.25 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.4, 164.8, 147.4, 144.6, 136.8, 135.6, 135.1, 134.9, 130.6, 130.5, 129.4, 129.3, 129.2, 129.0, 128.4, 126.1, 117.6, 115.6, 53.1, 52.6, 19.9; HRMS (APCI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 386.1386, found 386.1379.

Dimethyl 2'-cyano-3-methyl-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3g**): White solid, 34.6 mg, 90% yield. m.p. 238.1~238.6 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.11 (s, 1H), 7.50~7.44 (m, 3H), 7.43~7.37 (m, 5H), 7.32~7.26 (m, 1H), 3.94 (s, 3H), 3.60 (s, 3H), 2.45 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.3, 164.8, 147.2, 145.3, 138.6, 136.9, 135.3, 134.7, 130.7, 130.2, 130.1, 129.5, 129.2, 129.1, 128.7, 128.4, 125.9, 116.2, 116.0, 53.1, 52.5, 21.4; HRMS (APCI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 386.1386, found 386.1380.

Dimethyl 2'-cyano-3-methoxy-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3h**): White solid, 30 mg, 75% yield. m.p. 210.3~210.8 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.06 (s, 1H), 7.49~7.38 (m, 6H), 7.30 (dd, $J=7.6$, 1.6 Hz, 1H), 7.12~7.00 (m, 2H), 3.92 (s, 3H), 3.84 (s, 3H), 3.61 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.5, 164.9, 156.4, 144.3, 135.4, 134.6, 131.1, 131.0, 130.6, 130.5, 129.2, 129.1, 128.3, 126.1, 120.8, 118.2, 116.1, 111.3, 55.5, 53.0, 52.5; HRMS (APCI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 402.1335, found 402.1329.

Dimethyl 3-chloro-2'-cyano-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3i**): White solid, 35.5 mg, 89% yield. m.p. 178.5~179.0 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.10 (d, $J=1.2$ Hz, 1H), 7.59~7.55 (m, 1H), 7.52~7.44 (m, 6H), 7.43~7.37 (m, 2H), 3.95 (s, 3H), 3.61 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.0, 164.5, 145.5, 145.4, 138.6, 135.4, 135.0, 134.8, 131.0, 130.2, 130.1, 129.5, 129.4, 129.0, 128.9, 128.4, 127.1, 116.0, 115.8, 53.2, 52.6; HRMS (APCI) calcd for $\text{C}_{23}\text{H}_{17}\text{ClNO}_4$ $[\text{M}+\text{H}]^+$ 406.0840, found 406.0834.

Dimethyl 2'-cyano-3,5-difluoro-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3j**): White solid, 33.5 mg, 82% yield. m.p. 216.6~217.1 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.09 (s, 1H), 7.51~7.47 (m, 3H), 7.41~7.37 (m, 2H), 7.16~7.10 (m, 2H), 7.00~6.90 (m, 1H), 3.95 (s, 3H), 3.61 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 166.9, 164.4, 164.0 (dd, $J=250$, 13 Hz), 145.6, 144.43, 144.41, 144.3, 139.9, 139.8 (t, $J=10$ Hz), 135.8, 134.9, 131.1, 130.0, 129.5, 129.0, 128.5, 116.0, 115.5, 112.2 (dd, $J=19$, 8 Hz), 105.0 (t, $J=25$ Hz), 53.2, 52.6; ^{19}F NMR (376 MHz, Chloroform-*d*) δ : -107.80 (s, 2F); HRMS (APCI) calcd for $\text{C}_{23}\text{H}_{16}\text{F}_2\text{NO}_4$ $[\text{M}+\text{H}]^+$ 408.1041, found 408.1037.

Dimethyl 6-cyano-5-(naphthalen-2-yl)-[1,1'-biphenyl]-2,3-dicarboxylate (**3k**): White solid, 30.7 mg, 73% yield. m.p. 255.2~260.0 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.23 (s, 1H), 8.10~8.07 (m, 1H), 8.00 (s, 1H), 7.92 (m, 2H), 7.70 (dd, $J=8.4$, 2.0 Hz, 1H), 7.59~7.54 (m, 2H), 7.49~7.41 (m, 5H), 3.95 (s, 3H), 3.62 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.3, 164.8, 147.0, 145.4, 135.3, 134.8, 134.2, 133.3, 133.0, 130.9, 130.5, 129.3, 129.1, 128.8, 128.7, 128.5, 128.4, 127.7,

127.2, 126.8, 125.9, 116.3, 116.2, 53.1, 52.5; HRMS (APCI) calcd for $\text{C}_{27}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 422.1386, found 422.1381.

Dimethyl 6-cyano-5-(thiophen-2-yl)-[1,1'-biphenyl]-2,3-dicarboxylate (**3l**): White solid, 30.4 mg, 81% yield. m.p. 163.1~163.6 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.22 (s, 1H), 7.71 (dd, $J=3.6$, 0.8 Hz, 1H), 7.52 (dd, $J=5.2$, 1.2 Hz, 1H), 7.50~7.45 (m, 3H), 7.41~7.37 (m, 2H), 7.19 (dd, $J=5.2$, 4 Hz, 1H), 3.95 (s, 3H), 3.58 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.1, 164.6, 146.0, 139.1, 137.8, 135.2, 134.6, 131.0, 129.9, 129.3, 129.0, 128.9, 128.6, 128.5, 128.4, 116.4, 114.5, 53.2, 52.5; HRMS (APCI) calcd for $\text{C}_{21}\text{H}_{16}\text{NO}_4\text{S}$ $[\text{M}+\text{H}]^+$ 378.0794, found 378.0787.

Dimethyl 4-chloro-2'-cyano-6'-methyl-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3m**): White solid, 33 mg, 80% yield. m.p. 255.2~255.7 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.52~7.47 (m, 2H), 7.46~7.40 (m, 3H), 7.39~7.30 (m, 2H), 7.26~7.22 (m, 2H), 3.92 (s, 3H), 3.51 (s, 3H), 2.18 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.5, 166.9, 147.4, 142.6, 136.5, 136.0, 135.3, 135.2, 134.8, 132.5, 130.1, 129.3, 129.0, 128.7, 128.4, 115.9, 53.0, 52.6, 18.03; HRMS (APCI) calcd for $\text{C}_{24}\text{H}_{19}\text{ClNO}_4$ $[\text{M}+\text{H}]^+$ 420.0997, found 420.0990.

Dimethyl 2'-cyano-6'-propyl-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3n**): White solid, 31 mg, 75% yield. m.p. 200.0~200.5 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.52~7.41 (m, 6H), 7.38~7.28 (m, 4H), 3.91 (s, 3H), 3.51 (s, 3H), 2.60~2.50 (m, 2H), 1.44~1.33 (m, 2H), 0.72 (t, $J=8$ Hz, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.7, 167.2, 148.7, 142.2, 139.7, 136.7, 136.2, 136.0, 132.6, 129.0, 128.9, 128.8, 128.76, 128.71, 128.4, 116.3, 115.9, 52.9, 52.5, 32.8, 24.3, 14.3; HRMS (APCI) calcd for $\text{C}_{26}\text{H}_{24}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 414.1699, found 414.1694.

Dimethyl 6-cyano-5-methyl-[1,1'-biphenyl]-2,3-dicarboxylate (**3o**): White solid, 22.1 mg, 71.5% yield. m.p. 208.2~208.7 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.95 (s, 1H), 7.48~7.42 (m, 3H), 7.37~7.32 (m, 2H), 3.93 (s, 3H), 3.57 (s, 3H), 2.67 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 67.4, 164.9, 144.4, 143.7, 135.3, 133.7, 130.7, 130.1, 129.2, 128.9, 128.4, 117.7, 115.8, 53.0, 52.4, 21.0; HRMS (APCI) calcd for $\text{C}_{18}\text{H}_{16}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 310.1073, found 310.1067.

Dimethyl 2'-cyano-4'-methyl-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3p**): White solid, 33 mg, 87% yield. m.p. 197.3~197.8 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.10 (s, 1H), 7.60~7.57 (m, 2H), 7.54~7.46 (m, 3H), 7.34~7.27 (m, 4H), 3.94 (s, 3H), 3.63 (s, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.4, 164.8, 147.0, 145.5, 139.2, 137.0, 134.8, 132.3, 130.7, 130.1, 129.4, 129.1, 128.9, 128.8, 116.3, 116.1, 53.1, 52.5, 21.4; HRMS (APCI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 386.1386, found 386.1380.

Dimethyl 2'-cyano-4'-methoxy-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3q**): White solid, 32 mg, 80% yield. m.p. 200.6~201.1 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.09 (s, 1H), 7.58~7.52 (m, 2H), 7.50~7.45 (m,

3H), 7.45~7.2 (m, 2H), 7.06~7.00 (m, 2H), 3.93 (s, 3H), 3.86 (s, 3H), 3.59 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.4, 164.8, 160.2, 147.0, 145.1, 137.0, 135.0, 134.2, 130.8, 130.5, 129.4, 129.0, 128.8, 127.4, 116.4, 116.3, 113.9, 55.2, 53.1, 52.6; HRMS (APCI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 402.1335, found 402.1329.

Dimethyl 6-cyano-5-(thiophen-2-yl)-[1,1'-biphenyl]-3,4-dicarboxylate (**3r**): White solid, 31 mg, 81% yield. m.p. 199.7~200.2 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.14 (s, 1H), 7.61~7.57 (m, 2H), 7.55~7.49 (m, 4H), 7.24 (dd, $J=3.6, 1.2$ Hz, 1H), 7.14 (dd, $J=5.2, 3.6$ Hz, 1H), 3.94 (s, 3H), 3.73 (s, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 167.0, 164.5, 147.2, 138.2, 136.7, 136.0, 134.3, 131.0, 130.7, 130.2, 129.5, 128.9, 128.8, 128.5, 127.2, 117.2, 115.9, 53.2, 52.7; HRMS (APCI) calcd for $\text{C}_{21}\text{H}_{16}\text{NO}_4\text{S}$ $[\text{M}+\text{H}]^+$ 378.0794, found 378.0788.

Diethyl 2'-cyano-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3s**): White solid, 37.1 mg, 92% yield. m.p. 171.2~171.7 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 8.11 (s, 1H), 7.62~7.40 (m, 10H), 4.39 (q, $J=7.2$ Hz, 2H), 4.05 (q, $J=7.2$ Hz, 2H), 1.37 (t, $J=7.2$ Hz, 3H), 0.97 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 166.7, 164.4, 146.9, 145.2, 137.0, 135.4, 134.9, 131.3, 130.2, 129.3, 129.2, 128.8, 128.3, 116.2, 115.8, 62.3, 61.7, 14.0, 13.5; HRMS (APCI) calcd for $\text{C}_{25}\text{H}_{22}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 400.1543, found 400.1536.

Di-*tert*-butyl 2'-cyano-[1,1':3',1''-terphenyl]-4',5'-dicarboxylate (**3t**): White solid, 41 mg, 90% yield. m.p. 262.9~263.4 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.89 (s, 1H), 7.59~7.38 (m, 10H), 1.59 (d, $J=1.6$ Hz, 9H), 1.23 (d, $J=1.6$ Hz, 9H); ^{13}C NMR (100 MHz, Chloroform-*d*) δ : 165.4, 164.0, 146.2, 145.1, 137.4, 135.9, 135.4, 133.7, 129.5, 129.4, 129.2, 129.0, 128.9, 128.8, 128.2, 116.3, 115.1, 83.1, 82.9, 28.0, 27.5; HRMS (APCI) calcd for $\text{C}_{29}\text{H}_{29}\text{NO}_4$ $[\text{M}+\text{e}]^-$ 455.2096, found 455.2103.

Supporting Information Single crystal data of compound **3a**. ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra of compounds **3a**~**3t**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Lvan Otterlo, W. A. L.; de Koning, C. B. *Chem. Rev.* **2009**, *109*, 3743.
(b) Fuhr, L.; Rousseau, M.; Plauth, A.; Schroeder, F. C.; Sauer, S. J. *Nat. Prod.* **2015**, *78*, 1160.
(c) Boland, S.; Bourin, A.; Alen, J.; Geraets, J.; Schroeders, P.; Castermans, K.; Kindt, N.; Boumans, N.; Panitti, L.; Fransen, S.; Vanormel ingen, J.; Stassen, J. M.; Leysen, D.; Defert, O. *J. Med. Bioorg. Chem. Lett.* **2015**, *58*, 4309.
(d) Shirota, Y. *J. Mater. Chem.* **2005**, *15*, 75.
- [2] (a) Kirkham, J. D.; Butlin, R. J.; Harrity, J. P. A. *Angew. Chem., Int. Ed.* **2012**, *51*, 6402.
(b) Fillion, E.; Dumas, A. M. *J. Org. Chem.* **2008**, *73*, 2920.
- [3] (a) Ritleng, V.; Sirlin, C.; Pfeffer, M. *Chem. Rev.* **2002**, *102*, 1731.
(b) Ciana, C. L.; Phipps, R. J.; Brandt, J. R.; Meyer, F. M.; Gaunt, M. J. *Angew. Chem., Int. Ed.* **2011**, *50*, 458.
- (c) Gustafson, J. L.; Lim, D.; Barrett, K. T.; Miller, S. J. *Angew. Chem., Int. Ed.* **2011**, *50*, 5125.
- (d) Zhao, F.; Zhang, Y.-F.; Wen, J.; Yu, D.-G.; Wei, J.-B.; Xi, Z.; Shi, Z.-J. *Org. Lett.* **2013**, *15*, 3230.
- [4] (a) Roglans, A.; Pla-Quintana, A.; Solà, M. *Chem. Rev.* **2021**, *121*, 1894.
(b) Chien, C.-W.; Teng Gary, Y.-H.; Honda, T.; Ojima, I. *J. Org. Chem.* **2018**, *83*, 11623.
(c) Okamoto, S.; Yamada, T.; Tanabe, Y. K.; Sakai, M. *Organometallics* **2018**, *37*, 4431.
(d) More, A. A.; Ramana, C. V. *J. Org. Chem.* **2016**, *81*, 3400.
(e) García-Lacuna, J.; Dominguez, G.; Blanco-Urgoiti, J.; Perez-Castells, J. *Org. Lett.* **2018**, *20*, 5219.
(f) Méndez-Gálvez, C.; Böhme, M.; Leino, R.; SavelaEur, R. *J. Org. Chem.* **2020**, *11*, 1708.
- [5] (a) Hein, S. J.; Lehnher, D.; Arslan, H.; Uribe, F. J.; Dichtel, W. R. *Acc. Chem. Res.* **2017**, *50*, 2776.
(b) Poudel, T. N.; Tamargo, R. J. I.; Cai, H.; Lee, Y. R. *Asian J. Org. Chem.* **2018**, *7*, 985.
(c) Takikawa, H.; Nishii, A.; Takiguchi, H.; Yagishita, H.; Tanaka, M.; Hirano, K.; Uchiyama, M.; Ohmori, K.; Suzuki, K. *Angew. Chem., Int. Ed.* **2020**, *59*, 12440.
(d) Song, C. L.; Dong, X.; Wang, Z. J.; Liu, K.; Chiang, C. W.; Lei, A. W. *Angew. Chem., Int. Ed.* **2019**, *58*, 12206.
- [6] (a) Jia, Q. F.; Lin, Y. Q.; Li, Y. H. *Chin. J. Org. Chem.* **2020**, *40*, 1502 (in Chinese).
(贾乾发, 李娅琼, 林银河, 有机化学, **2020**, *40*, 1502.)
(b) Wang, D.; Lin, J. H.; Zhu, Y. N.; Huang, Y. *Adv. Synth. Catal.* **2021**, *363*, 1873.
(c) Wu, J. C.; Mou, C.-L.; Chi, Y. G. *Chin. J. Chem.* **2018**, *36*, 333.
(d) Chang, M. J.; Huang, Y. *Chem. Asian J.* **2019**, *14*, 2588.
(e) Shu, W. M.; Zheng, K. L.; Ma, J. R.; Wu, A. X. *Org. Lett.* **2015**, *17*, 5216.
(f) Jia, F. C.; Xu, C.; Zhou, Z. W.; Cai, Q.; Wu, Y. D.; Wu, A. X. *Org. Lett.* **2016**, *18*, 5232.
(g) Zhou, Q.-F.; Xin, B.-T.; Lu, T.; Xue, S. *Chin. J. Org. Chem.* **2009**, *29*, 1462 (in Chinese).
(周庆发, 辛波涛, 陆涛, 薛松, 有机化学, **2009**, *29*, 1462.)
- [7] (a) Zhang, C. L.; Ye, S. *Org. Lett.* **2016**, *18*, 6408.
(b) Jia, Q.; Wang, J. *Org. Lett.* **2016**, *18*, 2212.
- [8] Zhang, C. L.; Gao, Z. H.; Liang, Z. Q.; Ye, S. *Adv. Synth. Catal.* **2016**, *358*, 2862.
- [9] Zhang, C. L.; Zhang, Z.-F.; Xia, Z. H.; Han, Y. F.; Ye, S. *J. Org. Chem.* **2018**, *83*, 12507.
- [10] Zhu, C. Z.; Wei, Y.; Shi, M. *Adv. Synth. Catal.* **2018**, *360*, 808.
- [11] (a) Wang, W. H.; Wan, H. W.; Du, G. F.; Dai, B.; He, L. *Org. Lett.* **2019**, *21*, 3496.
(b) Pian, J. X.; He, L.; Du, G. F.; Guo, H.; Dai, B. *J. Org. Chem.* **2014**, *79*, 5820.
(c) Xie, P.; Yang, S. S.; Guo, Y. Y.; Cai, Z. H.; Dai, B.; He, L. *J. Org. Chem.* **2020**, *85*, 8872.
(d) Li, Z. J.; Wang, W. H.; Jian, H.; Li, W. J.; Dai, B.; He, L. *Chin. Chem. Lett.* **2019**, *30*, 386.
(e) Feng, Z. N.; Luo, J. Y.; Zhang, Y.; Du, G.-F.; He, L. *Org. Biomol. Chem.* **2019**, *17*, 4700.
- [12] Wang, Q.; An, Y.; Du, G. F.; Cai, Z. H.; Dai, B.; He, L. *J. Org. Chem.* **2020**, *85*, 14210.
- [13] For more X-ray crystal structure information of **3a** (CCDC: 2077708) see Supporting Information. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

(Cheng, F.)