光催化下二芳基碘鎓盐参与的自由基芳基化反应研究进展

孙媛媛^a 宋敬城^a 秦启学^a 张恩选^b 韩晴晴^a
 杨少慧^a 王祖利^{*,a} 岳 珊^a 董道青^a

(^a 青岛农业大学化学与药学院 山东青岛 266109.)

(^b 凯莱英生命科学技术(天津)有限公司 天津 300457)

摘要 二芳基高价碘盐在有机合成中被广泛应用。总结了二芳基高价碘试剂在可见光诱导的自由基芳基化反应中的研究进展。根据反应类型,分为碳碳键的形成、碳硫键的形成和碳磷键的形成三部分进行综述。

关键词 二芳基高价碘; 可见光; 自由基反应

Recent Progress in Radical Arylation Reaction with Diaryliodonium Salts under Photocatalysis

Sun, Yuanyuan^a Song, Jingcheng^a Qin, Qixue^a Zhang, Enxuan^b Han, Qingqing^a
 Yang, Shaohui^a Wang, Zuli^{*,a} Yue, Shan^a Dong, Daoqing^a

(^a College of Chemistry and Pharmaceutical Sciences, Qingdao Agricultural University, Qingdao, Shandong 266109)

(^b Asymchem Life Science (Tianjin) Co. Ltd., Tianjin 300457)

Abstract Diaryliodonium salts have been widely applied in organic transformations. According to the reaction type, the use of diaryliodonium salts in visible light induced arylation reactions is summarized from the following three parts: C—C bond formation, C—S bond formation and C—P bond formation.

Keywords diaryliodonium salt; visible light; radical reaction

Because of its environmental friendliness, stability and high selectivity, diaryliodonium salt as a kind of highly electrophilic arylation reagent, has been widely used in organic synthetic chemistry.^[1] Generally, diaryliodonium salts have high electrophilicity, which can be regarded as the equivalent of aryl cation. It can react with nucleophiles to form monosubstituted arylates. The reaction of diaryliodonium salts with various nucleophiles such as P(O)—OH,^[2] oxygen^[3] nitrogen,^[4] carbon^[5] nucleophiles and so on^[6] has been successfully realized. Additionally, cyclization reactions induced by electrophilic diaryliodonium salts were also established.^[7] Pyridines,^[8] quinolines,^[9] oxindole derivatives,^[10] polycyclic compounds^[11] and other cyclic compounds^[12] have been efficiently obtained via cyclic reaction using diaryliodonium salts.

Visible light induced chemical reactions have received great attention in recent years.^[13] At present, visible light

induced coupling reaction,^[14] cyclization reaction,^[15] oxidation reaction^[16] and other reactions^[14b,17] have all been successfully realized efficiently. With the assistance of visible light, diaryliodonium salts could be converted to aryl radical, which could participate in a series of reactions. In this review, we would like to describe the recent advances in visible light induced arylation reaction using diaryliodonium salts via radical mechanism. According to the reaction type, this review is divided into the following three parts: C—C bond formation, C—S bond formation and C—P bond formation (Scheme 1).

1 C—C Bond formation

As early as 2013, Xiao^[18] and Chatani^[19] group realized arylation of arenes and heteroarenes with diaryliodonium salts by photoredox catalysis independently (Scheme 2). Photoinduced single-electron-transfer from excited photo-

* Corresponding author. E-mail: wangzulichem@163.com

Received June 3, 2021; revised July 3, 2021; published online July 23, 2021

Project supported by the National Natural Science Foundation of China (Nos. 21772107, 21901010) and the Key Research and Development Plan of Shandong Province (No. 2019GSF108017).

国家自然科学基金(Nos. 21772107, 21901010)及山东省重点研发计划(No. 2019GSF108017)资助项目。

Scheme 1 Reactions triggered by diaryliodonium salts

catalyst to Ar_2I^+ , generating aryl radicals, oxidation reaction by higher valent photocatalyst were involved in the proposed mechanism.

Scheme 2 Arylation of arenes and heteroarenes

Quinoline and pyridine *N*-oxides are highly reactive substrates and lots of related reactions involving quinoline and pyridine *N*-oxides are presented in recent years.^[20] The visible-light-promoted C(2) selective arylation of quinoline and pyridine *N*-oxides, with diaryliodonium tetrafluoroborate as an arylation reagent was realized by Song group (Scheme 3).^[21] This transformation provided an efficient way for the synthesis of 2-aryl-substituted quinoline and

Scheme 3 Arylation of quinoline and pyridine *N*-oxides

pyridine *N*-oxides. In the reaction mechanism, oxidative quenching of excited eosin Y^* with diaryliodonium salt gave aryl radical, which followed by addition reaction with quinoline or pyridine *N*-oxide, oxidation by eosin Y^+ and deprotonation occurred in sequence to generate the desired products (**3D**).

The palladium-catalyzed arylation of vinyl arenes with diaryliodonium salts in the presence of visible light was reported by Gu group.^[22] When the reaction was irradiated by visible light, the kinetic isotope effect ($k_{\text{H}}/k_{\text{D}}$) was changed from 3.6 (under darkness) to 1.1, indicating that the C—H functionalization step was the rate determining step under darkness and significantly accelerated by the irradiation of visible light. The catalytic cycle is proposed in Scheme 4. Two radical species **4A** and **4B** were formed from the irradiation of Ph_2IOTf . Radical addition of **4B** to vinyl arenes would afford **4C**, which followed by association with Pd^{II} quickly to provide palladium species **4D**. Then the olefin products were isolated via oxidation of **4D** by **4A** through single-electron-transfer and base-mediated elimination (Scheme 4).

Glorius group^[23] realized a dual gold and photoredox catalytic system for the synthesis of α -arylated ether products via intermolecular three-component oxyarylation reactions of simple alkenes (Scheme 5). Using aryldiazonium salts as the aryl coupling partner, the iridium complex $[\text{Ir}(\text{ppy})_2(\text{dtbbpy})]\text{PF}_6$ as photocatalyst, visible light from blue LEDs allows the formation of products efficiently. A mechanistic hypothesis is depicted in Scheme 5. The alkyl gold(I) complex **5D** was generated from nucleophilic attack of the methanol solvent to double bond assisted by cationic gold(I). Aryl radical which generated from the oxidative quenching of excited iridium catalyst $[\text{Ir}(\text{ppy})_2(\text{dtbbpy})]^+$ by diaryliodonium salts via single electron transfer (SET) reacted with alkyl gold(I) complex **5D** to deliver the gold(II) species **5A**. Then oxidation of the gold(II) species by oxidized Ir catalyst to gold(III) species **5C** and reductive elimination occurred in sequence to give the desired products.

In 2014, Yu group^[24] realized the first example of isoquinoline synthesis via visible light-promoted vinyl isocyanide insertion with diaryliodonium salts at room temperature (Scheme 6). Products with various functional groups can be synthesized from readily available vinyl isocyanides and diaryliodonium salts. Aryl radical **7A** which was generated from the oxidative quenching of excited state $\text{fac-Ir}(\text{ppy})_3^*$ with diaryliodonium salts reacted with vinyl isocyanide to produce the imidoyl radical intermediate **7B**. Then homolytic aromatic substitution (HAS), oxidation and deprotonation took place to deliver the product isoquinoline (Scheme 7).

Zhu group^[25] established a novel photocatalytic arylation of unactivated alkenes using diaryliodonium salts as the arylation reagent (Scheme 8). A variety of desired products with promising yields were isolated at room temperature. Control experiments implied that the phenyl

Scheme 6 Synthesis of isoquinolines via visible light-promoted insertion of vinyl isocyanides

Scheme 7 Proposed mechanism for the synthesis of isoquinolines

radical may be involved in the catalytic cycle. The oxidative quenching of photoexcited Ir(III)* species with diaryliodonium salts produces the aryl radical **7A**, which was trapped by the C—C double bond to form alkyl radical **7B**. Then the products were generated via intramolecular addition, 1,2-hydrogen atom transfer (1,2-HAT), SET oxidation and deprotonation reaction.

In 2020, the consecutive domino arylation of *ortho*-hydroxyaryl enaminones with *in situ* photogenerated aryl radicals from diaryliodonium salts for the synthesis of iso-flavones was described by Mkrtchyan group (Scheme 9).^[26] The desired products were obtained with good to excellent yields. Additionally, diazonium salts were also suitable for this reaction instead of diaryliodonium salts. The reaction of diaryliodonium salts and the excited photocatalyst Ru(II) complexes occurred to produce aryl radical **9A** via single electron transfer (SET). Then aryl radical was attacked by enaminone to afford the radical intermediate **9B**, which was followed by oxidation to give carbocation **9C**. Subsequently, pyrone ring **9D** was formed via intramolecular electrophilic cyclization. After the elimination of dimethylamine, the corresponding products were formed.

Scheme 8 Photocatalytic acylation of unactivated alkenes

2 C—S Bond formation

1,4-Diazoniabicyclo[2.2.2]octane-1,4-disulfinate (DAB-SO) was an efficient and readily available reagent for sulfonylation reaction.^[27] Manolikakes *et al.*^[28] reported a photoredox-catalyzed three-component synthesis of *N*-aminosulfonamides starting from diaryliodonium salts, hydrazines and sulfur dioxide (Scheme 10). *N*-Aminosulfonamides were obtained with moderate to good yields.

Scheme 9 Consecutive domino arylation of *ortho*-hydroxyaryl enaminones

Scheme 10 Three-component synthesis of *N*-aminosulfonamides with sulfur dioxide surrogates

A highly selective transfer of one aryl moiety was observed when unsymmetrical diaryliodonium salts was applied to this reaction. Reductive quenching of PC* with the hydrazine-sulfur dioxide complex **10A** and oxidative quenching of PC* with diaryliodonium salts were proposed in the reaction mechanism.

A simple and efficient visible-light mediated three-component reaction of thiols, diaryliodonium salts and DABSO for the synthesis of thiosulfonates was developed by Volla group (Scheme 11).^[29] A radical-radical coupling of sulfonyl radicals and arylsulfonyl radicals produced by photocatalysis, which were formed from DABSO and diaryliodonium salts, was involved in the reaction mechanism. The unsymmetrical thiosulfonates were isolated with good to excellent yields. Besides diaryliodonium salts, aryldiazonium tetrafluoroborates could also give the products with good to excellent yields.

Scheme 11 Three-component coupling of thiols, iodonium salts and SO₂

In 2020, the visible-light photoredox catalyzed sulfonylation of imidazoheterocycles was developed by Piguel group (Scheme 12).^[30] Various of C-3 sulfonylated imidazoheterocycles were obtained from imidazopyridines, diaryliodonium salts and DABCO with moderate to good yields. When the radical scavengers 2,2,6,6-tetra-methyl-piperidine-1-oxyl (TEMPO) or 2,6-di-*tert*-butyl-4-methyl-phenol (BHT) was added to the reaction system, no product was isolated. These results indicated that a radical pathway was possible. Stern-Volmer fluorescence quenching experiments indicated that the catalytic cycle would start with an oxidative quenching of the excited EosinY* with diaryliodonium salts, giving the radical cation EosinY^{•+} and the aryl radical **12A**. Aryl radical can be trapped by SO₂ to give sulfonyl radical **12B**. Then the alkyl radical intermediate **12C** was formed from the reaction of sulfonyl radical and 2-phenylimidazo[1,2-*a*]pyridine, which followed by oxidation and deprotonation to deliver the final products.

Scheme 12 Sulfonylation of various imidazopyridines under visible-light

In 2017, a selective construction of organic sulfurs in different valence states from common thiosulfate salts was developed by Jiang group (Scheme 13).^[31] The sulfoxidation and sulfenylation reaction can be successfully tuned via a facile variation of the atmosphere under photocatalyzed conditions. Late-stage sulfoxidation for pharmaceuticals and sugar derivatives was proved to be highly compatible. From the proposed reaction mechanism, we can

see that electron transfer between diaryliodonium salt and EY*, energy-transfer between $^3\text{O}_2$ and EY* were involved.

In 2018, a visible-light-induced three-component reaction of *N*-propargyl aromatic amines, diaryliodonium salts and sulfur dioxide for the preparation of 3-arylsulfonylquinoline derivatives was discovered by Zhang group (Scheme 14).^[32] Simple operation and cheap photocatalyst make this protocol very practical. Proposed reaction mechanism was described. Aryl radical **14A** was generated from the excited Eosin Y* and diaryliodonium salts via oxidative quenching reaction. Subsequently, sulfonyl radical **14B** which was formed from the reaction of aryl radical and DABSO regioselectively attack *N*-propargyl aromatic to provide alkenyl radical **14C**, which followed by intramolecular cyclization, deprotonation, oxidation by Eosin Y⁺⁺ and aromatization to deliver the final product. In the same year, Manolikakes group^[33] found that *N*-aryl-acrylamides (Scheme 15, a) or arylpropynoates (Scheme 15, b)^[34] were also good substrates that can react with diaryliodonium salts and DABSO via cyclization. It should be mentioned that no additional photocatalyst was needed for these transformations.

Spirocyclohexadienones as privileged scaffolds like natural products, drugs and functional materials have recently gained significant attention.^[35] In 2020, metal free sulfonylative spirocyclization of alkynyl amides via insertion of sulfur dioxide was disclosed by Volla group (Scheme 16).^[36] These reactions proceed under metal free conditions and *in situ* generation of aryl sulfonyl radicals from DABSO was proposed for this transformation. Unfortunately, *N*-benzylacrylamide did not efficiently react with diaryliodonium salts to give the desired products azaspiro[4,5]-decanes. Besides diaryliodonium salts, aryl diazonium salts were also good substrates for this sulfonylative spirocyclization reaction.

Scheme 13 Sulfoxidation and sulfenylation with organic thiosulfate salts

Scheme 14 Three-component synthesis of 3-arylsulfonylquinoline derivatives

Scheme 15 Visible-light mediated cyclization reported by Manolikakes group

Scheme 16 Visible light mediated access to azaspiro[4,5]trienones with diaryliodonium salts

3 C—P bond formation

The visible light induced approach to aryl phosphonates by the simple combination of diaryliodonium salts with phosphites was realized by Lakhdar group (Scheme 17)^[37] electron paramagnetic resonance (EPR), NMR, and density functional theory (DFT) calculations were conducted to

understand the postulated reaction mechanism. The electron donor-acceptor (EDA) complex **17A** was formed between diaryliodonium salts and phosphites. The weak halogen bond between phosphorus lone pair and the I—C bond was proposed. Aryl radical **17A** (detected by EPR spectroscopy) and the phosphorus radical cation **17C** were generated from the decomposition of EDA complex. The final products were formed via nucleophilic displacement of phosphonium salt **17D** with the base. It was worth mentioning that radical-chain process was also possible for this system.

Scheme 17 Reaction of diaryliodonium salts with phosphites

The visible-light-induced arylation of tertiary phosphites

phines with aryl(mesityl)iodonium triflates to produce the quaternary phosphonium salts was described by Karchava group (Scheme 18).^[38] No photocatalyst was needed. Additionally, sterically congested phosphines, which are challenging under metal-based catalytic methods reacted smoothly in this reaction. Photo-excited EDA complexes **18A** between diaryliodonium salts and phosphines perhaps facilitate this transformation, which was followed by single-electron transfer from the phosphorous atom to the iodonium center to give aryl radical **18B** and the phosphorous radical cation **18C**. Then the desired phosphonium salts was isolated via combination the above radical species.

Scheme 18 Arylation of tertiary phosphines with aryl(mesityl)iodonium triflates.

4 Conclusion

In the past decades, diaryliodonium salts have been proved to be a powerful reagent for visible light induced arylation reaction via radical mechanism. Various reagents have successfully reacted with diaryliodonium salts to construct C—X (X=C, S, P) bonds. In most cases, oxidative quenching of the excited photocatalyst by diaryliodonium salts was proposed in the reaction mechanism. Despite these achievements, some challenges still exist to be solved. For example, more carbon-hetero bonds types (C—O, C—N, etc.) need to be explored. Additionally, asymmetric arylation reaction using diaryliodonium salts is still difficult. Visible light induced arylation reaction which can be conducted under catalyst free conditions also deserve our further attentions. We anticipate that solution of these problems, and further development of visible light induced arylation reaction using diaryliodonium salts will be realized in future.

References

- [1] (a) Chen, J.; Qu, H. M.; Peng, J.; Chen, C. *Chin. J. Org. Chem.* **2015**, 35, 937 (in Chinese). (陈静, 曲红梅, 彭静, 陈超, 有机化学, **2015**, 35, 937.) (b) Novák, Z.; Aradi, K.; Tóth, B.; Tolnai, G. *Synlett* **2016**, 27, 1456. (c) Ghosh, M. K.; Rout, N. *ChemistrySelect* **2020**, 5, 13644. (d) Chen, X.-J.; Gui, Q.-W.; Yi, R.; Yu, X.; Wu, Z.-L.; Huang, Y.; Cao, Z.; He, W.-M. *Org. Biomol. Chem.* **2020**, 18, 5234. (e) Takenaga, N.; Kumar, R.; Dohi, T. *Front Chem.* **2020**, 8, 599026.
- [2] Xiong, B. Q.; Feng, X. F.; Zhu, L. Z.; Chen, T. Q.; Zhou, Y. B.; Au, C. T.; Yin, S. F. *ACS Catal.* **2015**, 5, 537.
- [3] (a) Jalalian, N.; Petersen, T. B.; Olofsson, B. *Chem.-Eur. J.* **2012**, 18, 14140. (b) Shi, W. M.; Li, X. H.; Liang, C.; Mo, D. L. *Adv. Synth. Catal.* **2017**, 359, 4129.
- [4] (a) Li, J.; Liu, L. *RSC Adv.* **2012**, 2, 10485. (b) Carroll, M. A.; Wood, R. A. *Tetrahedron* **2007**, 63, 11349. (c) Duan, W. *Chin. J. Org. Chem.* **2020**, 40, 4387 (in Chinese). (段伟良, 有机化学, **2020**, 40, 4387.)
- [5] (a) Dohi, T.; Ito, M.; Yamaoka, N.; Morimoto, K.; Fujioka, H.; Kita, Y. *Angew. Chem., Int. Ed.* **2010**, 49, 3334. (b) Li, H.; Yang, P.; Xu, Z.; Du, Z.; Fu, Y. *Chin. J. Org. Chem.* **2020**, 40, 2476 (in Chinese). (李怀贵, 杨澎, 许峙, 杜正银, 傅颖, 有机化学, **2020**, 40, 2476.)
- [6] (a) Umierski, N.; Manolikakes, G. *Org. Lett.* **2013**, 15, 188. (b) Zhu, K.; Song, Z.; Wang, Y.; Zhang, F. *Org. Lett.* **2020**, 22, 9356.
- [7] (a) Wen, L. R.; Dou, Q.; Wang, Y. C.; Zhang, J. W.; Guo, W. S.; Li, M. *J. Org. Chem.* **2017**, 82, 1428. (b) Guo, W.; Li, S.; Tang, L.; Li, M.; Wen, L.; Chen, C. *Org. Lett.* **2015**, 17, 1232. (c) Wu, X.; Yang, Y.; Han, J.; Wang, L. *Org. Lett.* **2015**, 17, 5654. (d) Chen, Z.; Wang, W.-F.; Yang, H.; Wu, X.-F. *Org. Lett.* **2020**, 22, 1980.
- [8] Sheng, J.; Wang, Y.; Su, X.; He, R.; Chen, C. *Angew. Chem., Int. Ed.* **2017**, 56, 4824.
- [9] Wang, Y.; Chen, C.; Peng, J.; Li, M. *Angew. Chem., Int. Ed.* **2013**, 52, 5323.
- [10] Shi, L.; Wang, Y.; Yang, H.; Fu, H. *Org. Biomol. Chem.* **2014**, 12, 4070.
- [11] Wang, G.; Chen, C.; Peng, J. *Chem. Commun.* **2016**, 52, 10277.
- [12] (a) Tang, G.; Li, X.; Xu, J.; Zhang, P.; Gao, Y.; Wu, J.; Zhao, Y. *Synlett* **2014**, 25, 2009. (b) Wang, L.; Han, J.; Yang, Y.; Wu, X.; Mao, S.; Yu, J. *Synlett* **2014**, 25, 1419. (c) Pan, C.; Wang, L.; Han, J. *Org. Lett.* **2020**, 22, 4776.
- [13] (a) Ye, S. Q.; Li, X. F.; Xie, W. L.; Wu, J. *Eur. J. Org. Chem.* **2020**, 2020, 1274. (b) Guo, Y. Q.; Wang, R. G.; Song, H. J.; Liu, Y. X.; Wang, Q. M. *Org. Lett.* **2020**, 22, 709. (c) Krylov, I. B.; Lopat'eva, E. R.; Subbotina, I. R.; Nikishin, G. I.; Yu, B.; Terent'ev, A. O. *Chin. J. Catal.* **2021**, 42, 1700. (d) Dong, D.; Sun, Y.; Li, G.; Yang, H.; Wang, Z.; Xu, X. *Chin. J. Org. Chem.* **2020**, 40, 4071 (in Chinese). (董道青, 孙媛媛, 李光辉, 杨欢, 王祖利, 徐鑫明, 有机化学, **2020**, 40, 4071.) (e) Dong, D.-Q.; Yang, H.; Shi, J.-L.; Si, W.-J.; Wang, Z.-L.; Xu, X.-M. *Org. Chem. Front.* **2020**, 7, 2538.
- [14] (a) Zhong, J. J.; Meng, Q. Y.; Chen, B.; Tung, C. H.; Wu, L. Z. *Acta*

- Chim. Sinica* **2017**, 75, 34 (in Chinese).
(钟建基, 孟庆元, 陈彬, 佟振合, 吴骊珠, 化学学报, **2017**, 75, 34.)
- (b) Li, Z. L.; Jin, J.; Huang, S. H. *Chin. J. Org. Chem.* **2020**, 40, 563 (in Chinese).
(李祯龙, 金健, 黄莎华, 有机化学, **2020**, 40, 563.)
- (c) Kong, Y. L.; Xu, W. X.; Ye, F. X.; Weng, J. Q. *Chin. J. Org. Chem.* **2019**, 39, 3065 (in Chinese).
(孔瑶蕾, 徐雯秀, 叶飞霞, 翁建全, 有机化学, **2019**, 39, 3065.)
- (d) Guan, B. C.; Xu, X. L.; Wang, H.; Li, X. N. *Chin. J. Org. Chem.* **2016**, 36, 1564 (in Chinese).
(关保川, 许孝良, 王红, 李小年, 有机化学, **2016**, 36, 1564.)
- (e) Bagdi, A. K.; Rahman, M.; Bhattacharjee, D.; Zyryanov, G. V.; Ghosh, S.; Chupakhin, O. N.; Hajra, A. *Green Chem.* **2020**, 22, 6632.
- (f) Zhang, G. T.; Bian, C. L.; Lei, A. W. *Chin. J. Catal.* **2015**, 36, 1428.
- (g) Xie, L.-Y.; Liu, Y.-S.; Ding, H.-R.; Gong, S.-F.; Tan, J.-X.; He, J.-Y.; Cao, Z.; He, W.-M. *Chin. J. Catal.* **2020**, 41, 1168.
- [15] (a) Meng, X. X.; Kang, Q. Q.; Zhang, J. Y.; Li, Q.; Wei, W. T.; He, W. M. *Green Chem.* **2020**, 22, 1388.
(b) Liu, Y.; Chen, X. L.; Sun, K.; Li, X. Y.; Zeng, F. L.; Liu, X. C.; Qu, L. B.; Zhao, Y. F.; Yu, B. *Org. Lett.* **2019**, 21, 4019.
(c) Davies, J.; Booth, S. G.; Essafi, S.; Dryfe, R. A. W.; Leonori, D. *Angew. Chem., Int. Ed.* **2015**, 54, 14017.
(d) Zhang, Y.; Sun, K.; Lv, Q.; Chen, X.; Qu, L.; Yu, B. *Chin. Chem. Lett.* **2019**, 30, 1361.
(e) He, W.-B.; Gao, L.-Q.; Chen, X.-J.; Wu, Z.-L.; Huang, Y.; Cao, Z.; Xu, X.-H.; He, W.-M. *Chin. Chem. Lett.* **2020**, 31, 1895.
(f) Yang, W.-C.; Chen, C.-Y.; Li, J.-F.; Wang, Z.-L. *Chin. J. Catal.* **2021**, 42, 1865.
- [16] (a) Zhang, X.; Rakesh, K. P.; Ravindar, L.; Qin, H. L. *Green Chem.* **2018**, 20, 4790.
(b) Lin, J. Q.; Han, Q.; Ding, Y. *Chem. Rec.* **2018**, 18, 1531.
(c) He, S.; Chen, X.; Zeng, F.; Lu, P.; Peng, Y.; Qu, L.; Yu, B. *Chin. Chem. Lett.* **2020**, 31, 1863.
- [17] (a) Zhou, W.; Jiang, Y.; Chen, L.; Liu, K.; Yu, D. *Chin. J. Org. Chem.* **2020**, 40, 3697 (in Chinese).
(周文俊, 蒋元旭, 陈亮, 刘开兴, 余达刚, 有机化学, **2020**, 40, 3697.)
(b) Zhou, M. D.; Qin, P. T.; Jing, L. K.; Sun, J.; Du, H. W. *Chin. J. Org. Chem.* **2020**, 40, 598 (in Chinese).
(周明东, 覃丕涛, 经理珂, 孙京, 杜海武, 有机化学, **2020**, 40, 598.)
(c) Peng, S.; Lin, Y. W.; He, W. M. *Chin. J. Org. Chem.* **2020**, 40, 541 (in Chinese).
(彭莎, 林英武, 何卫民, 有机化学, **2020**, 40, 541.)
(d) Du, J. B.; Chen, Y. G.; Zuo, Z. W. *Chin. J. Org. Chem.* **2020**, 40, 3646 (in Chinese).
(杜建波, 陈跃刚, 左智伟, 有机化学, **2020**, 40, 3646.)
(e) Chen, Y. F.; Zhao, H.; Cheng, D. P.; Li, X. N.; Xu, X. L. *Chin. J. Org. Chem.* **2020**, 40, 1297 (in Chinese).
(陈跃峰, 赵赫, 程冬萍, 李小年, 许孝良, 有机化学, **2020**, 40, 1297.)
(f) Li, G.-H.; Han, Q.-Q.; Sun, Y.-Y.; Chen, D.-M.; Wang, Z.-L.; Xu, X.-M.; Yu, X.-Y. *Chin. Chem. Lett.* **2020**, 31, 3255.
(g) Chen, D.; Sun, Y.; Dong, D.; Han, Q.; Wang, Z. *Chin. J. Org. Chem.* **2020**, 40, 4267 (in Chinese).
(陈德茂, 孙媛媛, 董道青, 韩晴晴, 王祖利, 有机化学, **2020**, 40, 4267.)
(h) Meng, N.; Lv, Y.; Liu, Q.; Liu, R.; Zhao, X.; Wei, W. *Chin. Chem. Lett.* **2021**, 32, 258.
- (i) Gan, Z. Y.; Li, G. Q.; Yang, X. B.; Yan, Q. L.; Xu, G. Y.; Li, G. Y.; Jiang, Y. Y.; Yang, D. S. *Sci. China: Chem.* **2020**, 63, 1652.
- (j) Gui, Q.-W.; Teng, F.; Li, Z.-C.; Xiong, Z.-Y.; Jin, X.-F.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Chin. Chem. Lett.* **2021**, 32, 1907.
- (k) Jiang, T. G.; Wang, K.; Guo, T.; Wu, X. Y.; Zhang, G. K. *Chin. J. Catal.* **2020**, 41, 161.
- (l) Li, X. B.; Xiong, J.; Xu, Y.; Feng, Z. J.; Huang, J. T. *Chin. J. Catal.* **2019**, 40, 424.
- (m) Ren, D. D.; Shen, R. C.; Jiang, Z. M.; Lu, X. Y.; Li, X. *Chin. J. Catal.* **2020**, 41, 31.
- (n) Ren, Z. X.; Liu, X. J.; Zhuge, Z. H.; Gong, Y. Y.; Sun, C. Q. *Chin. J. Catal.* **2020**, 41, 180.
- [18] Liu, Y.-X.; Xue, D.; Wang, J.-D.; Zhao, C.-J.; Zou, Q.-Z.; Wang, C.; Xiao, J. *Synlett* **2013**, 24, 507.
- [19] Tobisu, M.; Furukawa, T.; Chatani, N. *Chem. Lett.* **2013**, 42, 1203.
- [20] (a) Hu, H.; Hu, X.; Liu, Y. *Angew. Chem., Int. Ed.* **2020**, 59, 18975.
(b) Yuan, Y.; Jiang, M.; Wang, T.; Xiong, Y.; Li, J.; Guo, H.; Lei, A. *Chem. Commun.* **2019**, 55, 11091.
(c) Xie, L.-Y.; Peng, S.; Liu, F.; Liu, Y.-F.; Sun, M.; Tang, Z.-L.; Jiang, S.; Cao, Z.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, 7, 7193.
(d) Peng, S.; Song, Y.-X.; He, J.-Y.; Tang, S.-S.; Tan, J.-X.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, 30, 2287.
(e) Du, S.; Pi, C.; Wan, T.; Wu, Y.; Cui, X. *Adv. Synth. Catal.* **2019**, 361, 1766.
(f) Dong, D. Q.; Li, L. X.; Li, G. H.; Deng, Q.; Wang, Z. L.; Long, S. *Chin. J. Catal.* **2019**, 40, 1494.
(g) Yu, H.; Pi, C.; Wang, Y.; Cui, X.; Wu, Y. *Chin. J. Org. Chem.* **2018**, 38, 124 (in Chinese).
(余海洋, 皮超, 王勇, 崔秀灵, 吴养洁, 有机化学, **2018**, 38, 124.)
(h) Xie, L.-Y.; Qu, J.; Peng, S.; Liu, K.-J.; Wang, Z.; Ding, M.-H.; Wang, Y.; Cao, Z.; He, W.-M. *Green Chem.* **2018**, 20, 760.
(i) Yuan, J.-W.; Qu, L.-B. *Chin. Chem. Lett.* **2017**, 28, 981.
- [21] Li, D. Z.; Liang, C.; Jiang, Z. X.; Zhang, J. Z.; Zhuo, W. T.; Zou, F. Y.; Wang, W. P.; Gao, G. L.; Song, J. Z. *J. Org. Chem.* **2020**, 85, 2733.
- [22] Feng, J.; Li, B.; Jiang, J.; Zhang, M.; Ouyang, W.; Li, C.; Fu, Y.; Gu, Z. *Chin. J. Chem.* **2018**, 36, 11.
- [23] Hopkinson, M. N.; Sahoo, B.; Glorius, F. *Adv. Synth. Catal.* **2014**, 356, 2794.
- [24] Jiang, H.; Cheng, Y. Z.; Wang, R. Z.; Zhang, Y.; Yu, S. Y. *Chem. Commun.* **2014**, 50, 6164.
- [25] Chen, Y.; Shu, C.; Luo, F.; Xiao, X.; Zhu, G. *Chem. Commun.* **2018**, 54, 5373.
- [26] Mkrtchyan, S.; Iaroshenko, V. O. *Chem. Commun.* **2020**, 56, 2606.
- [27] (a) Qiu, G.; Zhou, K.; Gao, L.; Wu, J. *Org. Chem. Front.* **2018**, 5, 691.
(b) Qiu, G.; Lai, L.; Cheng, J.; Wu, J. *Chem. Commun.* **2018**, 54, 10405.
(c) Li, G. H.; Dong, D. Q.; Deng, Q.; Yan, S. Q.; Wang, Z. L. *Synthesis* **2019**, 51, 3313.
(d) Ni, B.; Zhang, B.; Han, J.; Peng, B.; Shan, Y.; Niu, T. *Org. Lett.* **2020**, 22, 670.
- [28] Liu, N. W.; Liang, S.; Manolikakes, G. *Adv. Synth. Catal.* **2017**, 359, 1308.
- [29] Nair, A. M.; Kumar, S.; Halder, I.; Volla, C. M. R. *Org. Biomol. Chem.* **2019**, 17, 5897.
- [30] Breton-Patient, C.; Naud-Martin, D.; Mahuteau-Betzer, F.; Piguel, S. *Eur. J. Org. Chem.* **2020**, 2020, 6653.
- [31] Li, Y.; Wang, M.; Jiang, X. *ACS Catal.* **2017**, 7, 7587.
- [32] Sun, D. L.; Yin, K.; Zhang, R. H. *Chem. Commun.* **2018**, 54, 1335.

- [33] Chen, Z.; Liu, N.-W.; Bolte, M.; Ren, H.; Manolikakes, G. *Green Chem.* **2018**, *20*, 3059.
- [34] Liu, N.-W.; Chen, Z.; Herbert, A.; Ren, H.; Manolikakes, G. *Eur. J. Org. Chem.* **2018**, 2018, 5725.
- [35] (a) Gravel, E.; Poupon, E. *Nat. Prod. Rep.* **2010**, *27*, 32.
(b) Han, Y.; Zheng, H.; Zhang, Y. Y.; Yan, C. G. *Chin. Chem. Lett.* **2020**, *31*, 1337.
(c) Ye, Y. S.; Jiang, N. N.; Yang, X. W.; Xu, G. *Chin. Chem. Lett.* **2020**, *31*, 2433.
(d) Ma, X.; Luan, X. *Chin. J. Org. Chem.* **2020**, *40*, 1785 (in Chinese).
(马小军, 栾新军, 有机化学, **2020**, *40*, 1785.)
(e) Nie, Q. Y.; Ji, Z. Y.; Tang, G. L. *Chin. J. Org. Chem.* **2020**, *40*, 3467 (in Chinese).
(聂秋玥, 季珍瑜, 唐功利, 有机化学, **2020**, *40*, 3467.)
(f) Yang, M.; Cao, S. X.; He, Z. J. *Chin. J. Org. Chem.* **2019**, *39*, 2235 (in Chinese).
(杨梅, 曹仕轩, 贺峥杰, 有机化学, **2019**, *39*, 2235.)
(g) Yue, Y. S.; Xie, Z. F.; Chu, Y. C.; Xue, S. S. *Chin. J. Org. Chem.* **2020**, *40*, 501 (in Chinese).
(岳永双, 解正峰, 褚义成, 薛松松, 有机化学, **2020**, *40*, 501.)
- [36] Nair, A. M.; Halder, I.; Khan, S.; Volla, C. M. R. *Adv. Synth. Catal.* **2020**, *362*, 224.
- [37] Lecroq, W.; Bazille, P.; Morlet-Savary, F.; Breugst, M.; Lalevée, J.; Gaumont, A.-C.; Lakhdar, S. *Org. Lett.* **2018**, *20*, 4164.
- [38] Bugaenko, D. I.; Volkov, A. A.; Livantsov, M. V.; Yurovskaya, M. A.; Karchava, A. V. *Chem.-Eur. J.* **2019**, *25*, 12502.

(Zhao, C.)