利用 aza-Michael/Michael 串联环化反应合成螺巴比妥哌啶酮衍生物

刘雪华^{†,a} 王川川^{†,b} 王新璐^a 马志伟^b 孟 蕾^c
丁德刚^b 刘俊桃^b 陈亚静^{*,a}

(^a 郑州大学药学院 郑州 450001)

(^b 河南牧业经济学院理学部 郑州 450046)

(^c 河南省兽药饲料监察所 郑州 450008)

摘要 发展了巴比妥衍生烯烃化合物与 *N*-烷氧基丙烯酰胺之间的串联 aza-Michael/分子内 Michael 加成反应。该反应条件温和, 适用于克级制备, 为具有生物活性的螺巴比妥分子的高效合成提供了一种实用的方法。

关键词 螺巴比妥; 哌啶酮; 串联反应; Michael 加成

Synthesis of Spirobarbiturate Piperidin-2-one Derivatives via Cascade Aza-Michael/Michael Cyclization Reaction

Liu, Xuehua^{†,a} Wang, Chuanchuan^{†,b} Wang, Xinlu^a Ma, Zhiwei^b Meng, Lei^c
Ding, Degang^b Liu, Juntao^b Chen, Yajing^{*,a}

(^a School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001)

(^b Faculty of Science, Henan University of Animal Husbandry and Economy, Zhengzhou 450046)

(^c Henan Institute of Veterinary Drug and Feed Control, Zhengzhou 450008)

Abstract A highly efficient cascade aza-Michael/intramolecular-Michael reaction employing barbiturate-derived alkenes and *N*-alkoxy acrylamides has been developed. The reaction proceeds smoothly under mild conditions and is amenable to gram scale synthesis, providing a practical tool in the synthesis of bioactive spirobarbiturate piperidin-2-ones molecules in moderate to good yields.

Keywords spirobarbiturate; piperidin-2-one; cascade reaction; Michael addition

1 Introduction

Spirobarbiturates constitute a class of structurally diverse and remarkable nitrogen-containing spiroheterocyclic frameworks and possess importantly biological and pharmacological properties, such as anticonvulsant, antibacterial, α -glucosidase inhibitor, TNF- α converting enzyme (TACE) inhibitor and anticancer activities (Figure 1).^[1] Over the past decades, intrigued by the biological significances of these spiro architectures, many efforts have been invested on the development of powerful synthetic strategies for the construction of various spirobarbi-

turates.^[2] On the other hand, piperidin-2-one motif is also an ubiquitous structural moiety in a large variety of bioactive natural and synthetic compounds.^[3] Based on the principle of the hybridization of dominant skeletons,^[4] it is of great value to study the synthesis and activity of spirobarbiturate piperidin-2-ones. Actually, spirobarbiturate piperidin-2-one derivatives, in which the piperidin-2-one moieties are fused at the 5-position of the barbituric nucleus, have been proven to be potent inhibitors of TACE, indicating a potential application in rheumatoid arthritis and diabetes mellitus.^[5] However, traditional methods for constructing spirobarbiturate piperidin-2-one scaffolds rely on

* Corresponding author. E-mail: chenyj@zzu.edu.cn

Received May 26, 2021; revised July 2, 2021; published online July 23, 2021.

Project supported by the Key Scientific and Technological Project of Henan Province (Nos. 202102110067, 212102110439), the Natural Science Foundation of Henan Province (Nos. 202300410188, 212300410152) and the Henan University of Animal Husbandry and Economy (Nos. 2019HNUAHEDF011, XKYCXXJ2020006).

河南省重点研发与推广专项(科技攻关)(Nos. 202102110067, 212102110439)、河南省自然科学基金(Nos. 202300410188, 212300410152)和河南牧业经济学院(Nos. 2019HNUAHEDF011, XKYCXXJ2020006)资助项目。

[†] 共同第一作者(These authors contributed equally to this work).

time-consuming multistep synthesis.^[6] Hence, there is a great demand for developing concise and powerful protocols to construct functionalized spirobarbiturate piperidin-2-ones.

Figure 1 Bioactive molecules with spirobarbiturate scaffolds

On the other hand, spirobarbiturates were always prepared by employing barbituric acid building blocks as the starting materials.^[7] Particularly, barbiturate-derived alkenes, robust and easily accessible building blocks, participated $[2+m]$ annulations provide various simple and feasible synthetic strategies for the construction of structurally diverse and complex spirobarbiturates.^[8] Recently, we developed a practical strategy for the synthesis of spirobarbiturate-pyrrolidinones via a cascade $[3+2]$ cycloaddition reaction between barbiturate-derived alkenes and *N*-alkoxy α -haloamides under mild conditions (Scheme 1, A).^[9] However, we failed to synthesize pharmaceutically useful spirobarbiturate piperidin-2-one using barbiturate-derived alkene and *N*-OBn β -bromopropanamide as the substrates. Actually, the cascade $[4+2]$ annulations are powerful synthetic strategies for the construction of six-membered rings.^[10] Particularly, annulations involving *N*-alkoxyl acrylamides have been successfully applied in the synthesis of piperidin-2-ones. Li and co-workers^[11] reported the cascade cycloaddition between 3-methyleneindolinones and *in-situ* formed *N*-alkoxyl acrylamides from α -bromoamides via base promoted elimination process, providing racemic spirooxindole piperidin-2-ones. Then, Comesse and co-workers^[12] realized the cascade aza-Michael/Michael reaction employing *N*-alkoxyl acrylamides as key partners for the preparation of piperidin-2-ones. Following these success, Sun group^[13] achieved the first enantioselective

synthesis of spirooxindole piperidin-2-ones via squaramide catalyzed aza-Michael/Michael cyclization of 3-methyleneindolinones with *N*-alkoxyl acrylamides. These elegant studies prove that *N*-alkoxyl acrylamides could be recognized as a class of powerful four-atom synthons for the synthesis of piperidin-2-ones through cascade aza-Michael/Michael cyclization with activated alkenes. Inspired by our previous work,^[14] we conceived that the cascade aza-Michael/ Michael $[4+2]$ annulation between barbiturate-derived alkenes and *N*-alkoxyacrylamides could deliver spirobarbiturate piperidin-2-one scaffolds (Scheme 1, B).

Scheme 1 Our work on the synthesis of spirobarbiturate scaffolds

2 Results and discussion

With this mindset, barbiturate-derived alkene (**1a**) and *N*-benzyloxy acrylamide (**2a**) were selected as model substrates to optimize the reaction conditions (Table 1). Based on the previous study, the reaction was firstly conducted at 25 °C for 48 h with the use of K_2CO_3 and 4Å MS in CH_2Cl_2 under air atmosphere and the desired product **3aa** could be generated in 58% yield (Entry 1). While replacing K_2CO_3 with other bases (Na_2CO_3 , Cs_2CO_3 and K_2HPO_4), the target spirobarbiturate **3aa** was generated in reduced yields (Entries 2~4). Next, a variety of solvents such as CH_3CN , $CHCl_3$, toluene, dichloroethane (DCE) and tetrahydrofuran (THF) were screened and CH_3CN displayed the best result, giving **3aa** in 69% yield (Entry 5). To further optimize the cascade transformation, stoichiometry of the substrates and base were also examined (Entries 10~12). It was found that a 1.0 : 1.5 : 1.5 molar ratio of **1a** : **2a** : K_2CO_3 provided superior result (Entry 12). Controlled experiment conducted in the absence of 4Å MS led to a lower yield (Entry 13). Furthermore, higher reaction temperature (40 °C) has no effect on the yield (Entry 14).

Considering the step-economy, the direct one-pot cyclization of 1,3-dimethylbarbituric acid, benzaldehyde and *N*-benzyloxy acrylamide **2a** was further studied. However, no desired spirobarbiturate product **3aa** was detected.

Table 1 Optimization of the reaction conditions^a

Entry	Base	1a/2a/Base	Solvent	Temp./°C	Yield ^b /%
1	K ₂ CO ₃	1.0/1.2/1.2	CH ₂ Cl ₂	25	58
2	Na ₂ CO ₃	1.0/1.2/1.2	CH ₂ Cl ₂	25	54
3	CS ₂ CO ₃	1.0/1.2/1.2	CH ₂ Cl ₂	25	13
4	K ₂ HPO ₄	1.0/1.2/1.2	CH ₂ Cl ₂	25	Trace
5	K ₂ CO ₃	1.0/1.2/1.2	CH ₃ CN	25	69
6	K ₂ CO ₃	1.0/1.2/1.2	CHCl ₃	25	67
7	K ₂ CO ₃	1.0/1.2/1.2	Toluene	25	21
8	K ₂ CO ₃	1.0/1.2/1.2	DCE	25	29
9	K ₂ CO ₃	1.0/1.2/1.2	THF	25	33
10	K ₂ CO ₃	1.0/1.5/1.2	CH ₃ CN	25	74
11	K ₂ CO ₃	1.0/2.0/1.2	CH ₃ CN	25	74
12	K ₂ CO ₃	1.0/1.5/1.5	CH ₃ CN	25	83
13 ^c	K ₂ CO ₃	1.0/1.5/1.5	CH ₃ CN	25	52
14	K ₂ CO ₃	1.0/1.5/1.5	CH ₃ CN	40	83

^a Unless otherwise noted, all reactions were carried out with **1a** (0.2 mmol), **2a**, base and 4Å MS (100 mg) in 2 mL of solvent under air atmosphere.

^b Isolated yields. ^c Without 4Å MS.

Moreover, the step-by-step operation was tried. Barbiturate-derived alkene **1a** could be smoothly synthesized from 1,3-dimethylbarbituric acid and benzaldehyde in CH₃CN via Knoevenagel condensation reaction. After the complete consumption of 1,3-dimethylbarbituric acid, *N*-benzyloxy acrylamide (**2a**), K₂CO₃ and 4Å MS were added into the reaction mixture without purification. Nevertheless, the desired product **3aa** remained undetected. The probable reason might be that the complex reaction system affected the cyclization reaction. Moreover, the cyclization of demethylated unsubstituted *N*-H barbiturate-derived alkene and *N*-benzyloxy acrylamide (**2a**) under the optimized reaction conditions was further studied. Unfortunately, no desired spiro-product had been detected.

With the optimized conditions in hand, the substrate scope of the cascade aza-Michael/Michael cyclization for the synthesis of spirobarbiturate piperidin-2-ones under the optimized conditions was next examined (Table 2). Firstly, a wide range of aryl substituted barbiturate-derived alkenes **1** were evaluated with *N*-benzyloxy acrylamide **2a**. In general, the electronic effect and steric effect of the aryl substituents seemed to have no remarkable influence on the reaction outcome. And the reactions of aryl substituted barbiturate-derived alkenes **1** with either electron-donating groups (such as OMe and Me) or electron-withdrawing groups (such as F, Cl, Br, CN, CF₃ and NO₂) on the different position of benzene ring all could afford the desired spirobarbiturates **3** in moderate to excellent yields (**3aa**~**3ra**). In addition, for the barbiturate-derived alkenes with

the use of ring-condensed naphthyl (**1s**~**1t**) and piperonyl (**1u**) as the R¹ group, good yields (71%~87%) were also obtained (**3sa**~**3ua**). Moreover, the heteroaromatic (3-thienyl) substituted alkene **1v** also was an applicable substrate for the cyclization, offering the corresponding product **3va** in 35% yield. However, no desired spirobarbiturate product **3wa** was observed when the R¹ group was an aliphatic isopropyl substituent. Considering that the barbiturate-derived alkene substrates bearing large substituents, such as **1q** (2-CF₃C₆H₄), **1r** (2-NO₂C₆H₄) and **1t** (1-naphthyl), all could provide the desired products (**3qa**, **3ra** and **3ta**) with excellent yields, we think that the main reason for the failure of **1w** is not the steric effect and the conjugate effect of the aryl substituent may play an important role for the successful cyclization.

Next, the scope and limitations of *N*-protected acrylamides were also examined. *N*-Methoxy acrylamides could provide the desired product **3ab** in 77% yield. Encouraged by the feasibility of this cascade sequence, we further explored more challenging acrylamides with substituent at α or β position. α-Substituted methacrylamide **2c** could afford the desired product *trans*-**3ac** in 26% yield and only one diastereoisomer was observed on the ¹H NMR of the crude product. To our delight, α-phenyl substituted acrylamide **2d** could afford the desired product **3ad** in 74% yield. In sharp contrast, the steric hindrance at β positions of acrylamides showed significant impact on the reaction outcome. The crotonamide and cinnamamide derivatives **2e** and **2f** failed to provide the desired products **3ae** and **3af** under the optimized reaction conditions or even at 80 °C. Moreover, the cyclization of *N*-butyl acrylamide (**2g**) and unsubstituted *N*-H acrylamide **2h** was also studied. No desired products **3ag** and **3ah** were detected probably owing to the weaker acidity of such substrates compared to *N*-alkoxy acrylamides.

To further illustrate the preparative utility of this cascade cyclization, a gram-scale reaction was carried out under the optimized conditions (Scheme 2). The spirobarbiturate product **3pa** was obtained in a comparable yield (88%).

Scheme 2 Gram scale synthesis of **3pa**

On the basis of previous report, we speculate that the reaction might occur through a cascade aza-Michael/Michael pathway. As depicted in Scheme 3, the sequence is initiated by the aza-Michael addition between *N*-alkoxyacrylamide **2** and barbiturate-derived alkene **1** under the

Table 2 Scope of the reaction^a

K_2CO_3 , 4Å MS
 CH_3CN , r.t., 48 h

Variation of 1

3aa, 83% yield	3ba, 72% yield	3ca, 79% yield	3da, 83% yield	3ea, 78% yield	3fa, 77% yield
3ga, 62% yield	3ha, 63% yield	3ia, 70% yield	3ja, 45% yield	3ka, 72% yield	3la, 66% yield
3ma, 73% yield	3na, 89% yield	3oa, 77% yield	3pa, 93% yield	3qa, 90% yield	3ra, 83% yield
					Variation of 2 3ab, 77% yield
3sa, 87% yield	3ta, 74% yield	3ua, 71% yield	3va, 35% yield	3wa, NP	
3ac, 26% yield ^b	3ad, 74% yield (<i>dr</i> = 9:1) ^c	3ae, NP	3af, NP	3ag, NP	3ah, NP

^a Reaction conditions: **1** (0.2 mmol), **2** (0.3 mmol), K₂CO₃ (0.3 mmol), 4Å MS (100 mg), MeCN (2.0 mL), 25 °C, air atmosphere. The listed yield is isolated yield.

^b Only one diastereoisomer was observed and the *trans*-configuration is determined by NOESY spectroscopy. ^c Total yield and the *dr* value is determined by ¹H NMR.

Scheme 3 Putative reaction mechanism

promotion of base. The resulting aza-Michael adduct **A** then undergoes an intramolecular Michael addition to generate the desired spirobarbiturate piperidin-2-one **3** via intermediate **B**.

3 Conclusions

In conclusion, barbiturate-derived alkenes and *N*-alkoxyacrylamides can undergo a tandem aza-Michael/Michael cyclization under mild conditions, affording spirobarbiturate piperidin-2-ones in good to excellent yields. This cascade formal [4+2] cyclization was successfully performed under air atmosphere with good tolerance of a wide range of functional groups. This strategy not only provides a practical method for the synthesis of pharmaceutically useful spirobarbiturates, but also further expands the synthetic utility of *N*-alkoxyacrylamides on the synthesis of aza-heterocycles. Further expansion of new tandem reactions based on *N*-alkoxyacrylamides together with the enantioselective synthesis of aza-heterocycles, as well as investigation of the biological activity of the produced spirobarbiturate piperidin-2-ones, are currently underway in our laboratory.

4 Experimental section

4.1 Experimental details

All reactions were performed in oven-dried glassware with magnetic stirring. Unless otherwise stated, all reagents were purchased from commercial suppliers and used without further purification. All solvents were purified and dried according to standard methods prior to use. Organic solutions were concentrated under reduced pressure on a rotary evaporator or an oil pump. ¹H NMR, ¹³C NMR and ¹⁹F NMR spectra were recorded at 400 or 600, 100 or 151, and 565 MHz on a Bruker instrument, respectively. Chemical shifts are reported referenced to an internal SiMe₄ standard for ¹H NMR and chloroform-*d* (δ 77.16) for ¹³C NMR. High-resolution mass spectra (HRMS) were recorded on a Thermo Q-Exactive Spectrometer (ESI source). Reactions were monitored through thin layer chromatography (TLC) on silica gel-precoated glass plates. Flash column chromatography was performed using silica gel (300~400 mesh).

4.2 General procedures for the cascade reaction

To a 10 mL test tube equipped with a magneton were sequentially added barbiturate derived alkene **1**^[9] (0.2 mmol, 1.0 equiv.), acrylamide **2**^[13] (0.3 mmol, 1.5 equiv.), K₂CO₃ (0.3 mmol, 1.5 equiv.), 4 Å MS (100 mg) and MeCN (2.0 mL) under air atmosphere. Then the tube was sealed and stirred at room temperature for 48 h. The mixture was concentrated under reduced pressure and the residue was purified via flash chromatograph on silica gel to give the desired product **3**.

8-(Benzyloxy)-2,4-dimethyl-7-phenyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3aa**): White powder, 70 mg, 83% yield (eluent: ethyl acetate/hexane, *V*: *V*=1:1).

1). m.p. 159~161 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.37~7.20 (m, 6H), 7.09~7.07 (m, 2H), 6.99~6.97 (m, 2H), 5.09~5.06 (m, 2H), 4.72 (d, *J*=10.0 Hz, 1H), 3.36~3.27 (m, 1H), 3.14 (s, 3H), 3.03 (s, 3H), 2.84~2.77 (m, 1H), 2.64~2.56 (m, 1H), 2.19~2.13 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.1, 168.9, 167.3, 149.9, 135.0, 133.5, 129.6, 129.4, 128.8, 128.6, 128.4, 127.6, 76.5, 71.3, 56.1, 30.3, 29.2, 28.6, 27.6; HRMS (ESI) calcd for C₂₃H₂₄N₃O₅ [M+H]⁺ 422.1710, found 422.1707.

8-(Benzyloxy)-7-(4-methoxyphenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ba**): White powder, 65 mg, 72% yield (eluent: ethyl acetate/hexane, *V*: *V*=1:1). m.p. 129~130 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.30~7.21 (m, 3H), 7.04~6.99 (m, 4H), 6.82~6.80 (m, 2H), 5.06~5.03 (m, 2H), 4.71 (d, *J*=10.0 Hz, 1H), 3.80 (s, 3H), 3.32~3.22 (m, 1H), 3.15 (s, 3H), 3.07 (s, 3H), 2.84~2.77 (m, 1H), 2.60~2.52 (m, 1H), 2.18~2.12 (m, 1H); ¹³C NMR (151 MHz, CDCl₃) δ: 169.2, 168.6, 167.4, 160.3, 150.0, 135.1, 129.4, 128.9, 128.7, 128.4, 125.2, 113.9, 76.4, 70.7, 56.3, 55.4, 30.2, 29.1, 28.6, 27.5; HRMS (ESI) calcd for C₂₄H₂₆N₃O₆ [M+H]⁺ 452.1816, found 452.1812.

8-(Benzyloxy)-2,4-dimethyl-7-(*p*-tolyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ca**): White powder, 69 mg, 79% yield (eluent: ethyl acetate/hexane, *V*: *V*=1:1). m.p. 134~136 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.30~7.21 (m, 3H), 7.10 (d, *J*=8.0 Hz, 2H), 7.02~7.00 (m, 2H), 6.95 (d, *J*=8.0 Hz, 2H), 5.06 (d, *J*=10.0 Hz, 1H), 5.03 (s, 1H), 4.71 (d, *J*=10.0 Hz, 1H), 3.32~3.23 (m, 1H), 3.14 (s, 3H), 3.05 (s, 3H), 2.85~2.78 (m, 1H), 2.60~2.51 (m, 1H), 2.33 (s, 3H), 2.19~2.13 (m, 1H); ¹³C NMR (151 MHz, CDCl₃) δ: 169.2, 168.7, 167.4, 150.0, 139.5, 135.1, 130.4, 129.4, 129.2, 128.7, 128.4, 127.5, 76.5, 71.1, 56.2, 30.2, 29.1, 28.6, 27.3, 21.3; HRMS (ESI) calcd for C₂₄H₂₆N₃O₅ [M+H]⁺ 436.1867, found 436.1863.

8-(Benzyloxy)-7-(4-fluorophenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3da**): Light yellow powder, 73 mg, 83% yield (eluent: ethyl acetate/hexane, *V*: *V*=1:1). m.p. 124~126 °C; ¹H NMR (600 MHz, CDCl₃) δ: 7.31~7.24 (m, 3H), 7.08~7.06 (m, 2H), 7.03~6.98 (m, 4H), 5.11 (s, 1H), 5.06 (d, *J*=9.6 Hz, 1H), 4.71 (d, *J*=10.2 Hz, 1H), 3.25~3.19 (m, 1H), 3.16 (s, 3H), 3.06 (s, 3H), 2.80~2.76 (m, 1H), 2.59~2.53 (m, 1H), 2.16~2.13 (m, 1H); ¹³C NMR (151 MHz, CDCl₃) δ: 169.0, 168.4, 167.1, 163.1 (d, ¹*J*_{C-F}=249.2 Hz), 149.9, 135.0, 129.7 (d, ³*J*_{C-F}=9.1 Hz), 129.5 (d, ⁴*J*_{C-F}=3.0 Hz), 129.4, 128.9, 128.5, 115.6 (d, ²*J*_{C-F}=21.1 Hz), 76.5, 70.1, 56.3, 30.1, 29.2, 28.6, 28.2; ¹⁹F NMR (565 MHz, CDCl₃) δ: -111.3; HRMS (ESI) calcd for C₂₃H₂₃FN₃O₅ [M+H]⁺ 440.1616, found 440.1614.

8-(Benzyloxy)-7-(4-chlorophenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ea**): White powder, 71 mg, 78% yield (eluent: ethyl acetate/hexane, *V*: *V*=1:1). m.p. 89~90 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.33~7.25 (m, 5H), 7.06~7.01 (m, 4H), 5.10 (s, 1H), 5.07 (d, *J*=10.4 Hz, 1H), 4.73 (d, *J*=10.0 Hz, 1H), 3.24~3.14 (m, 4H), 3.07 (s, 3H), 2.81~2.75 (m,

1H), 2.57~2.49 (m, 1H), 2.17~2.11 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.9, 168.1, 166.9, 149.8, 135.3, 134.9, 132.2, 129.4, 129.3, 128.9, 128.8, 128.5, 76.4, 70.0, 56.2, 30.0, 29.3, 28.6, 28.3; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{ClN}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 456.1321, found 456.1317.

8-(Benzyloxy)-7-(4-bromophenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3fa**): White powder, 77 mg, 77% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 90~91 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.43 (d, $J=8.4$ Hz, 2H), 7.33~7.25 (m, 3H), 7.06~7.04 (m, 2H), 6.96 (d, $J=8.4$ Hz, 2H), 5.09~5.06 (m, 2H), 4.73 (d, $J=10.0$ Hz, 1H), 3.22~3.13 (m, 4H), 3.07 (s, 3H), 2.81~2.75 (m, 1H), 2.57~2.49 (m, 1H), 2.17~2.11 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.9, 168.1, 166.9, 149.8, 134.9, 132.8, 131.7, 129.6, 129.4, 128.9, 128.5, 123.5, 76.4, 70.0, 56.1, 30.0, 29.3, 28.6, 28.4; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}^{79}\text{BrN}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 500.0816, found 500.0811.

4-(8-(Benzyloxy)-2,4-dimethyl-1,3,5,9-tetraoxo-2,4,8-triazaspiro[5.5]undecan-7-yl)benzonitrile (**3ga**): White powder, 55 mg, 62% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 119~121 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.59~7.57 (m, 2H), 7.36~7.27 (m, 3H), 7.21~7.19 (m, 2H), 7.07~7.05 (m, 2H), 5.21 (s, 1H), 5.08 (d, $J=10.0$ Hz, 1H), 4.76 (d, $J=10.4$ Hz, 1H), 3.20 (s, 3H), 3.15~3.06 (m, 4H), 2.81~2.74 (m, 1H), 2.55~2.47 (m, 1H), 2.17~2.11 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.6, 167.6, 166.5, 149.7, 139.2, 134.7, 132.2, 129.5, 129.1, 128.9, 128.7, 118.1, 113.2, 76.4, 69.5, 56.3, 29.9, 29.4, 29.2, 28.7; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{23}\text{N}_4\text{O}_5$ $[\text{M}+\text{H}]^+$ 447.1663, found 447.1659.

8-(Benzyloxy)-2,4-dimethyl-7-(4-(trifluoromethyl)phenyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ha**): White powder, 62 mg, 63% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 147~148 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.55 (d, $J=6.6$ Hz, 2H), 7.31~7.21 (m, 5H), 7.02~7.01 (m, 2H), 5.21 (s, 1H), 5.09 (d, $J=9.6$ Hz, 1H), 4.74 (d, $J=10.2$ Hz, 1H), 3.19~3.12 (m, 4H), 3.05 (s, 3H), 2.80~2.77 (m, 1H), 2.56~2.53 (m, 1H), 2.16~2.15 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.8, 167.9, 166.7, 149.8, 138.0, 134.9, 131.5 (q, $^2J_{\text{C-F}}=33.2$ Hz), 129.4, 129.0, 128.57, 128.55, 125.5 (q, $^3J_{\text{C-F}}=3.0$ Hz), 123.8 (q, $^1J_{\text{C-F}}=271.8$ Hz), 76.5, 69.8, 56.3, 30.0, 29.4, 28.9, 28.6; ^{19}F NMR (565 MHz, CDCl_3) δ : -62.8; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{23}\text{F}_3\text{N}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 490.1584, found 490.1581.

8-(Benzyloxy)-2,4-dimethyl-7-(4-nitrophenyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ia**): White powder, 65 mg, 70% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 152~154 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.18~8.16 (m, 1H), 7.87~7.83 (m, 1H), 7.51~7.46 (m, 2H), 7.35~7.25 (m, 3H), 7.07~7.06 (m, 2H), 5.23 (s, 1H), 5.10 (d, $J=10.4$ Hz, 1H), 4.77 (d, $J=10.4$ Hz, 1H), 3.19 (s, 3H), 3.17~3.10 (m, 1H), 3.06 (s, 3H), 2.85~2.78 (m, 1H), 2.57~2.49 (m, 1H), 2.19~2.13 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.6, 167.4, 166.4, 149.7, 148.3, 141.2, 134.8, 129.5, 129.3, 129.2,

128.7, 123.5, 76.5, 69.1, 56.4, 29.8, 29.5 (2C), 28.7; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_7$ $[\text{M}+\text{H}]^+$ 467.1561, found 467.1558.

8-(Benzyloxy)-7-(3-methoxyphenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ja**): White powder, 41 mg, 45% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 180~181 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.30~7.20 (m, 4H), 7.04~7.03 (m, 2H), 6.88~6.86 (m, 1H), 6.65~6.63 (m, 2H), 5.10 (d, $J=10.0$ Hz, 1H), 5.03 (s, 1H), 4.75 (d, $J=10.0$ Hz, 1H), 3.75 (s, 3H), 3.35~3.26 (m, 1H), 3.15 (s, 3H), 3.05 (s, 3H), 2.84~2.77 (m, 1H), 2.62~2.54 (m, 1H), 2.19~2.13 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.1, 168.8, 167.3, 159.8, 150.0, 135.0 (2C), 129.6, 129.4, 128.7, 128.4, 119.7, 115.0, 113.3, 76.6, 71.1, 56.1, 55.4, 30.3, 29.2, 28.6, 27.6; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{N}_3\text{O}_6$ $[\text{M}+\text{H}]^+$ 452.1816, found 452.1814.

8-(Benzyloxy)-7-(3-fluorophenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ka**): White powder, 63 mg, 72% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 169~170 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.31~7.24 (m, 4H), 7.05~7.02 (m, 3H), 6.86~6.82 (m, 2H), 5.10~5.08 (m, 2H), 4.73 (d, $J=10.2$ Hz, 1H), 3.27~3.21 (m, 1H), 3.17 (s, 3H), 3.07 (s, 3H), 2.81~2.76 (m, 1H), 2.58~2.53 (m, 1H), 2.17~2.13 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.9, 168.5, 167.0, 162.7 (d, $^1J_{\text{C-F}}=249.2$ Hz), 149.9, 136.3 (d, $^3J_{\text{C-F}}=7.6$ Hz), 135.0, 130.2 (d, $^3J_{\text{C-F}}=7.6$ Hz), 129.4, 128.9, 128.5, 123.5 (d, $^4J_{\text{C-F}}=3.0$ Hz), 116.5 (d, $^2J_{\text{C-F}}=21.1$ Hz), 114.9 (d, $^2J_{\text{C-F}}=22.7$ Hz), 76.6, 70.4, 56.1, 30.2, 29.3, 28.6, 28.2; ^{19}F NMR (565 MHz, CDCl_3) δ : -111.5; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{FN}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 440.1616, found 440.1610.

8-(Benzyloxy)-7-(3-chlorophenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3la**): White powder, 60 mg, 66% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 188~189 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.33~7.22 (m, 5H), 7.07~7.05 (m, 3H), 6.95 (d, $J=7.6$ Hz, 1H), 5.10 (d, $J=10.4$ Hz, 1H), 5.02 (s, 1H), 4.73 (d, $J=10.4$ Hz, 1H), 3.30~3.21 (m, 1H), 3.16 (s, 3H), 3.07 (s, 3H), 2.83~2.76 (m, 1H), 2.60~2.52 (m, 1H), 2.18~2.12 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.8, 168.5, 167.0, 149.8, 135.7, 134.9, 134.7, 129.8, 129.7, 129.5, 128.9, 128.5, 127.9, 125.9, 76.7, 70.5, 56.1, 30.2, 29.2, 28.6, 28.0; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{ClN}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 456.1321, found 456.1317.

8-(Benzyloxy)-2,4-dimethyl-7-(3-nitrophenyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ma**): White powder, 68 mg, 73% yield (eluent: ethyl acetate/hexane, $V:V=1:1$). m.p. 175~176 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.14~8.12 (m, 2H), 7.39~7.25 (m, 5H), 7.09~7.07 (m, 2H), 5.27 (s, 1H), 5.10 (d, $J=10.4$ Hz, 1H), 4.78 (d, $J=10.4$ Hz, 1H), 3.21 (s, 3H), 3.14~3.05 (m, 4H), 2.81~2.75 (m, 1H), 2.55~2.47 (m, 1H), 2.18~2.12 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 168.6, 167.6, 166.6, 149.7, 148.1, 136.1, 134.8, 134.3, 129.5, 129.4, 129.1, 128.6, 124.2, 122.8, 76.6, 69.5, 56.3, 29.9, 29.4, 28.8, 28.7; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_7$ $[\text{M}+$

$\text{H}]^+$ 467.1561, found 467.1556.

8-(Benzyloxy)-7-(2-methoxyphenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3na**): light yellow powder, 80 mg, 89% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 184~185 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.32~7.25 (m, 4H), 7.19~7.17 (m, 3H), 7.00~6.96 (m, 1H), 6.82 (d, $J=8.4$ Hz, 1H), 5.55 (s, 1H), 4.94 (d, $J=10.4$ Hz, 1H), 4.84 (d, $J=10.8$ Hz, 1H), 3.70 (s, 3H), 3.21~3.10 (m, 4H), 3.03~2.88 (m, 4H), 2.42~2.30 (m, 2H); ^{13}C NMR (151 MHz, CDCl_3) δ : 170.0, 168.4, 167.5, 156.4, 150.7, 135.3, 130.2, 129.3, 128.6 (2C), 128.4, 122.8, 121.0, 110.4, 76.4, 63.3, 55.7, 55.0, 30.0, 29.3, 28.6, 26.1; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{N}_3\text{O}_6$ $[\text{M}+\text{H}]^+$ 452.1816, Found 452.1811.

8-(Benzyloxy)-2,4-dimethyl-7-(*o*-tolyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3oa**): White powder, 67 mg, 77% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 132~133 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.30~7.14 (m, 7H), 6.86 (d, $J=7.2$ Hz, 2H), 5.43 (s, 1H), 5.03 (d, $J=9.6$ Hz, 1H), 4.50 (d, $J=9.6$ Hz, 1H), 3.55~3.46 (m, 1H), 3.12 (s, 3H), 3.08 (s, 3H), 2.85~2.69 (m, 2H), 2.25~2.19 (m, 1H), 2.10 (s, 3H); ^{13}C NMR (151 MHz, CDCl_3) δ : 170.2, 168.8, 168.1, 150.0, 136.6, 135.0, 131.5, 131.3, 129.4, 129.3, 128.6, 128.4, 127.4, 125.8, 76.9, 66.6, 55.0, 30.6, 29.3, 28.6, 27.8, 19.0; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{N}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 436.1867, found 436.1863.

8-(Benzyloxy)-7-(2-bromophenyl)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3pa**): White powder, 93 mg, 93% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 176~178 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.54~7.52 (m, 1H), 7.37~7.33 (m, 1H), 7.31~7.28 (m, 3H), 7.27~7.19 (m, 4H), 5.58 (s, 1H), 4.96 (d, $J=10.4$ Hz, 1H), 4.78 (d, $J=10.4$ Hz, 1H), 3.25~3.16 (m, 4H), 3.10~3.02 (m, 1H), 2.99 (s, 3H), 2.48~2.36 (m, 2H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.0, 168.0, 167.0, 150.3, 134.9, 133.7, 133.4, 130.8, 129.5 (3C), 128.8, 128.6, 127.8, 123.4, 76.4, 68.3, 54.3, 30.0, 29.7, 29.0, 25.5; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}^{79}\text{BrN}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 500.0816, found 500.0811.

8-(Benzyloxy)-2,4-dimethyl-7-(2-(trifluoromethyl)phenyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3qa**): White powder, 88 mg, 90% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 183~184 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.66~7.59 (m, 2H), 7.53~7.46 (m, 2H), 7.26~7.24 (m, 3H), 7.17~7.14 (m, 2H), 5.55 (s, 1H), 4.93 (d, $J=10.0$ Hz, 1H), 4.76 (d, $J=10.0$ Hz, 1H), 3.26 (s, 3H), 3.22~3.14 (m, 1H), 3.01 (s, 3H), 3.00~2.94 (m, 1H), 2.59~2.52 (m, 1H), 2.42~2.34 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.2, 167.5, 167.2, 150.2, 134.5, 133.9, 132.0, 129.9, 129.5, 129.4, 128.8, 128.43, 128.36 (q, $^2J_{\text{C-F}}=30.2$ Hz), 126.6 (q, $^3J_{\text{C-F}}=6.0$ Hz), 124.2 (q, $^1J_{\text{C-F}}=274.8$ Hz), 76.2, 64.5, 54.3, 29.8, 29.4, 28.9, 27.5; ^{19}F NMR (565 MHz, CDCl_3) δ : -57.0; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{23}\text{F}_3\text{N}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 490.1584, found 490.1578.

8-(Benzyloxy)-2,4-dimethyl-7-(2-nitrophenyl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ra**): White

powder, 77 mg, 83% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 169~170 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.01~7.99 (m, 1H), 7.73~7.69 (m, 1H), 7.53~7.49 (m, 2H), 7.32~7.21 (m, 5H), 5.69 (s, 1H), 5.02 (d, $J=10.8$ Hz, 1H), 4.94 (d, $J=10.8$ Hz, 1H), 3.37 (s, 3H), 3.02 (s, 3H), 3.00~2.93 (m, 1H), 2.90~2.82 (m, 1H), 2.49~2.42 (m, 1H), 2.26~2.19 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.9, 167.5, 167.4, 150.3, 148.2, 135.0, 134.0, 132.7, 130.0, 129.6, 129.2, 128.9, 128.6, 125.4, 76.5, 64.5, 54.3, 29.5, 29.3, 29.04, 28.97; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_7$ $[\text{M}+\text{H}]^+$ 467.1561, found 467.1558.

8-(Benzyloxy)-2,4-dimethyl-7-(naphthalen-2-yl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3sa**): Yellow powder, 82 mg, 87% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 86~87 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.84~7.74 (m, 3H), 7.54~7.48 (m, 3H), 7.26~7.14 (m, 4H), 6.97~6.96 (m, 2H), 5.25 (s, 1H), 5.09 (d, $J=10.0$ Hz, 1H), 4.75 (d, $J=10.0$ Hz, 1H), 3.35~3.26 (m, 1H), 3.11 (s, 3H), 2.97 (s, 3H), 2.88~2.82 (m, 1H), 2.65~2.57 (m, 1H), 2.23~2.17 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.2, 168.5, 167.3, 149.8, 135.0, 133.6, 132.9, 131.0, 129.5, 128.8, 128.4, 128.3, 128.0, 127.9, 127.5, 127.1, 126.9, 124.6, 76.5, 71.3, 56.3, 30.2, 29.2, 28.6, 27.8; HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{26}\text{N}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 472.1867, found 472.1863.

8-(Benzyloxy)-2,4-dimethyl-7-(naphthalen-1-yl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ta**): Yellow powder, 70 mg, 74% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 121~122 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.88~7.84 (m, 2H), 7.50~7.46 (m, 4H), 7.42~7.41 (m, 1H), 7.28~7.26 (m, 1H), 7.20~7.17 (m, 2H), 6.98 (d, $J=7.2$ Hz, 2H), 5.83 (s, 1H), 5.09 (d, $J=10.8$ Hz, 1H), 4.71 (d, $J=10.8$ Hz, 1H), 3.49~3.43 (m, 1H), 2.98~2.93 (m, 1H), 2.76 (s, 3H), 2.72 (s, 3H), 2.68~2.63 (m, 1H), 2.30~2.26 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.1, 169.0, 167.5, 149.6, 135.2, 133.8, 130.6, 130.0, 129.5 (3C), 128.9, 128.8, 128.5, 126.8, 126.1, 126.0, 124.7, 121.4, 76.6, 65.6, 55.3, 30.3, 29.1, 28.4, 26.5; HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{26}\text{N}_3\text{O}_5$ $[\text{M}+\text{H}]^+$ 472.1867, Found 472.1863.

7-(Benzo[d][1,3]dioxol-5-yl)-8-(benzyloxy)-2,4-dimethyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ua**): Yellow powder, 66 mg, 71% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 166~167 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.31~7.25 (m, 3H), 7.09 (d, $J=7.2$ Hz, 2H), 6.70 (d, $J=7.8$ Hz, 1H), 6.60 (s, 1H), 6.49 (d, $J=7.8$ Hz, 1H), 5.96 (s, 2H), 5.06 (d, $J=10.2$ Hz, 1H), 4.99 (s, 1H), 4.73 (d, $J=10.2$ Hz, 1H), 3.28~3.20 (m, 1H), 3.16 (s, 3H), 3.10 (s, 3H), 2.82~2.78 (m, 1H), 2.55~2.49 (m, 1H), 2.18~2.14 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.1, 168.6, 167.3, 150.1, 148.4, 148.1, 135.1, 129.4, 128.8, 128.5, 127.2, 121.5, 108.1, 107.8, 101.6, 76.6, 70.8, 56.3, 30.2, 29.2, 28.7, 27.5; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{24}\text{N}_3\text{O}_7$ $[\text{M}+\text{H}]^+$ 466.1609, found 466.1604.

8-(Benzyloxy)-2,4-dimethyl-7-(thiophen-3-yl)-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3va**): White

powder, 30 mg, 35% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 154~155 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.30~7.22 (m, 4H), 7.09 (s, 1H), 7.00 (d, $J=7.2$ Hz, 2H), 6.92 (d, $J=4.8$ Hz, 1H), 5.34 (s, 1H), 5.07 (d, $J=9.6$ Hz, 1H), 4.67 (d, $J=9.6$ Hz, 1H), 3.22~3.17 (m, 4H), 3.14 (s, 3H), 2.77~2.72 (m, 1H), 2.60~2.54 (m, 1H), 2.16~2.13 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.1, 168.4, 167.5, 150.1, 135.0, 134.7, 129.4, 128.7, 128.4, 126.9, 126.6, 125.0, 76.8, 66.5, 56.2, 30.2, 29.2, 28.7, 28.5; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{N}_3\text{O}_5\text{S} [\text{M}+\text{H}]^+$ 428.1275, found 428.1270.

8-Methoxy-2,4-dimethyl-7-phenyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ab**): White powder, 53 mg, 77% yield (eluent: ethyl acetate/hexane, $V: V=1:1$). m.p. 162~163 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.36~7.29 (m, 3H), 7.13~7.10 (m, 2H), 5.32 (s, 1H), 3.64 (s, 3H), 3.37~3.26 (m, 1H), 3.20 (s, 3H), 3.03 (s, 3H), 2.78~2.65 (m, 2H), 2.20~2.13 (m, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 169.3, 168.9, 167.3, 149.9, 133.5, 129.6, 128.6, 127.3, 70.2, 62.1, 56.2, 30.2, 29.2, 28.5, 27.9; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{20}\text{N}_3\text{O}_5 [\text{M}+\text{H}]^+$ 346.1397, found 346.1394.

8-(Benzyloxy)-2,4,10-trimethyl-7-phenyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ac**): White powder, 23 mg, 26% yield (eluent: ethyl acetate/hexane, $V: V=1:3$). m.p. 171~172 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.41~7.33 (m, 3H), 7.30~7.26 (m, 3H), 7.16~7.15 (m, 2H), 6.97~6.96 (m, 2H), 4.82~4.76 (m, 3H), 3.51~3.41 (m, 1H), 3.09 (s, 3H), 3.04 (s, 3H), 2.36~2.30 (m, 1H), 2.17~2.11 (m, 1H), 1.46 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (151 MHz, CDCl_3) δ : 172.0, 170.0, 167.3, 150.6, 135.4, 133.4, 129.8, 129.4, 128.82, 128.76, 128.5, 127.8, 76.8, 71.8, 56.3, 34.5, 30.6, 29.2 (2C); HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{N}_3\text{O}_5 [\text{M}+\text{H}]^+$ 436.1867, found 436.1860.

8-(Benzyloxy)-2,4-dimethyl-7,10-diphenyl-2,4,8-triazaspiro[5.5]undecane-1,3,5,9-tetraone (**3ad**): White powder, 74 mg, 74% yield (eluent: ethyl acetate/hexane, $V: V=1:3$). m.p. 174~175 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.45~7.39 (m, 7H), 7.34~7.32 (m, 1H), 7.30~7.25 (m, 3H), 7.14~7.13 (m, 4H), 4.93 (s, 1H), 4.86 (d, $J=10.8$ Hz, 1H), 4.76~4.71 (m, 2H), 3.14 (s, 3H), 3.01 (s, 3H), 2.87 (dd, $J=15.0, 12.6$ Hz, 1H), 2.33 (dd, $J=15.0, 7.2$ Hz, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 170.2, 169.9, 167.2, 150.6, 138.6, 135.2, 133.4, 130.0, 129.6, 128.96, 128.94, 128.90, 128.85, 128.5, 128.0, 127.7, 76.9, 71.9, 56.0, 46.3, 31.4, 29.3 (2C); HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{28}\text{N}_3\text{O}_5 [\text{M}+\text{H}]^+$ 498.2023, found 498.2018.

Supporting Information Spectroscopic characterization of the products **3**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- (a) Barakat, A.; Islam, M. S.; Al-Majid, A. M.; Ghabbour, H. A.; Fun, H.-K.; Javed, K.; Imad, R.; Yousuf, S.; Choudhary, M. I.; Wadood, A. *Bioorg. Med. Chem.* **2015**, *23*, 6740.
(b) Shi, C.; Zhang, Y.; Wang, T.; Lu, W.; Zhang, S.; Guo, B.; Chen, Q.; Luo, C.; Zhou, X.; Yang, Y. *J. Med. Chem.* **2019**, *62*, 2950.
- (c) Ramachary, D. B.; Reddy, Y. V.; Banerjee, A.; Banerjee, S. *Org. Biomol. Chem.* **2011**, *9*, 7282.
- (d) Ziarani, G. M.; Asadi, S.; Faramarzi, S.; Amanlou, M. *Iran. J. Pharm. Res.* **2015**, *14*, 1105.
- (e) Kesharwani, S.; Sahu, N. K.; Kohli, D. V. *Pharm. Chem. J.* **2009**, *43*, 315.
- (f) Bhaskarachar, R. K.; Revanasiddappa, V. G.; Hegde, S.; Bala-krishna, J. P.; Reddy, S. Y. *Med. Chem. Res.* **2015**, *24*, 3516.
- (a) Roy, P.; Anjum, S. R.; Ramachary, D. B. *Org. Lett.* **2020**, *22*, 2897.
(b) Han, B.; Huang, W.; Ren, W.; He, G.; Wang, J.-H.; Peng, C. *Adv. Synth. Catal.* **2015**, *357*, 561.
- (c) Kotha, S.; Sreevani, G. *Synthesis* **2018**, *50*, 4883.
- (d) Nagaraju, S.; Sathish, K.; Paplal, B.; Kashinath, D. *Tetrahedron Lett.* **2017**, *58*, 2865.
- (e) Zhao, H.-W.; Tian, T.; Li, B.; Yang, Z.; Pang, H.-L.; Meng, W.; Song, X.-Q.; Chen, X.-Q. *J. Org. Chem.* **2015**, *80*, 10380.
- (a) Huang, X.; Pham, K.; Yi, W.; Zhang, X.; Clamens, C.; Hyatt, J. H.; Jasinsk, J. P.; Tayvah, U.; Zhang, W. *Adv. Synth. Catal.* **2015**, *357*, 3820.
(b) Shcherbakova, V.; Dibchak, D.; Snisarenko, M.; Skalenko, Y.; Denisenko, A. V.; Kuznetsova, A. S.; Mykhailiuk, P. K. *J. Org. Chem.* **2021**, *86*, 2200.
(c) Yang, W.; Xu, X.; Zhang, Z. *Chin. J. Org. Chem.* **2014**, *34*, 1161 (in Chinese).
(杨维芳, 徐晓丽, 张站斌, 有机化学, **2014**, *34*, 1161.)
- (a) Wang, L.-Z.; Li, X.-Q.; An, Y.-S. *Org. Biomol. Chem.* **2015**, *13*, 5497.
(b) Tu, S.; Xu, W.; Qi, F.; He, W.; Fei, B. *Chin. J. Org. Chem.* **2019**, *39*, 3269 (in Chinese).
(徐双燕, 徐武双, 齐芬, 何卫江, 费宝丽, 有机化学, **2019**, *39*, 3269.)
(c) Wei, B.; Zhou, J.; Xu, J.-J.; Cui, J.; Ping, F.-F.; Ling, J.-J.; Chen, Y.-J. *Eur. J. Med. Chem.* **2019**, *184*, e111779.
- Duan, J.; Jiang, B.; Chen, L.; Lu, Z.; Barbosa, J.; Pitts, W. *WO 2003053941*, **2003**.
- Freeman-Cook, K. D.; Reiter, L. A.; Noe, M. C.; Antipas, A. S.; Danley, D. E.; Datta, K.; Downs, J. T.; Eisenbeis, S.; Eskra, J. D.; Garmene, D. J.; Greer, E. M.; Griffiths, R. J.; Guzman, R.; Hardink, J. R.; Janat, F.; Jones, C. S.; Martinelli, G. J.; Mitchell, P. G.; Laird, E. R.; Liras, J. L.; Lopresti-Morrow, L. L.; Pandit, J.; Reilly, U. D.; Robertson, D.; Vaughn-Bowser, M. L.; Wolf-Gouvêa, L. A.; Yocum, S. A. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 6529.
- (a) Patel, D. M.; Patel, H. M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 18667.
(b) Zhu, S.; Chen, C.; Duan, K.; Sun, Y.-M.; Li, S.-S.; Liu, Q.; Xiao, J. *J. Org. Chem.* **2019**, *84*, 8440.
(c) Wang, B.; Leng, H.-J.; Yang, X.-Y.; Han, B.; Rao, C.-L.; Liu, L.; Peng, C.; Huang, W. *RSC Adv.* **2015**, *5*, 88272.
(d) Lomlim, L.; Einsiedel, J.; Heinemann, F. W.; Meyer, K.; Gmeiner, P. *J. Org. Chem.* **2008**, *73*, 3608.
(e) Yin, G.; Li, Y.; Fan, L. *Chin. J. Org. Chem.* **2021**, *41*, 1234 (in Chinese).
(殷国栋, 李源, 范玲, 有机化学, **2021**, *41*, 1234.)
- (a) Frankowski, S.; Gajda, T.; Albrecht, L. *Adv. Synth. Catal.* **2018**, *360*, 1822.
(b) Gao, Z.; Wang, C.; Zhou, L.; Yuan, C.; Xiao, Y.; Guo, H. *Org. Lett.* **2018**, *20*, 4302.
(c) Liu, H.; Liu, Y.; Yuan, C.; Wang, G.-P.; Zhu, S.-F.; Wu, Y.; Wang, B.; Sun, Z.; Xiao, Y.; Zhou, Q.-L.; Guo, H. *Org. Lett.* **2016**, *18*, 1302.
(d) Gao, X.; Li, Z.; Yang, W.; Liu, Y.; Chen, W.; Zhang, C.; Zheng, L.; Guo, H. *Org. Biomol. Chem.* **2017**, *15*, 5298.
(e) Yang, W.-J.; Sun, Q.; Sun, J.; Yan, C.-G. *Org. Chem. Front.* **2018**, *5*, 2754.
(f) Zhao, H.-W.; Tian, T.; Pang, H.-L.; Li, B.; Chen, X.-Q.; Yang, Z.; Meng, W.; Song, X.-Q.; Zhao, Y.-D.; Liu, Y.-Y. *Adv. Synth.*

- Catal.* **2016**, 358, 2619.
- (g) Liu, D.; Sun, J.; Xie, J.; Shi, H.; Yan, C.-G. *J. Org. Chem.* **2021**, 86, 1827.
- (h) Liu, Y.; Tang, Y.; Luo, Y.; Zhu, Y.; Yang, J.; Wei, X.; Liu, X.; Zhao, Y. *Chin. J. Org. Chem.* **2019**, 39, 2980 (in Chinese).
(刘钰, 汤永星, 罗月阳, 朱叶婷, 杨进, 魏欣雨, 刘雄, 赵云辉, 有机化学, **2019**, 39, 2980.)
- [9] Wang, C.-C.; Zhou, J.; Ma, Z.-W.; Chen, X.-P.; Chen, Y.-J. *Org. Biomol. Chem.* **2019**, 17, 9200.
- [10] (a) Xu, P.-W.; Yu, J.-S.; Chen, C.; Cao, Z.-Y.; Zhou, F.; Zhou, J. *ACS Catal.* **2019**, 9, 1820.
(b) Lopes, S. M. M.; Cardoso, A. L.; Lemos, A.; Pinho e Melo, T. M. V. D. *Chem. Rev.* **2018**, 118, 11324.
(c) Guranova, N.; Golubev, P.; Bakulina, O.; Dar'in, D.; Kantin, G.; Krasavin, M. *Org. Biomol. Chem.* **2021**, 19, 3829.
(d) Yang, L.; Wei, L.; Wan, J.-P. *Chem. Commun.* **2018**, 54, 7475.
(e) Luo, T.; Xu, H.; Liu, Y. *ChemistrySelect* **2019**, 4, 10621.
- (f) Jia, J.; Yu, A.; Liu, X.; Meng, X. *Chin. J. Org. Chem.* **2019**, 39, 2175 (in Chinese).
(贾继茹, 于爱敏, 刘旭光, 孟祥太, 有机化学, **2019**, 39, 2175.)
- (g) Jiang, C.; Yu, P.-P.; Zhang, Q.; Xu, H.-D.; Shen, M.-H. *Chin. Chem. Lett.* **2019**, 30, 266.
- [11] Liu, Y.; Xue, J.; Sun, Z.; Liu, D.; Xing, Y.; Li, Y. *Asian J. Org. Chem.* **2016**, 5, 43.
- [12] Champetier, P.; Castillo-Aguilera, O.; Taillier, C.; Brière, J. F.; Dalla, V.; Oudeyer, S.; Comesse, S. *Eur. J. Org. Chem.* **2019**, 7703.
- [13] Tang, Q.-G.; Cai, S.-L.; Wang, C.-C.; Lin, G.-Q.; Sun, X.-W. *Org. Lett.* **2020**, 22, 3351.
- [14] (a) Wang, C.-C.; Qu, Y.-L.; Liu, X.-H.; Ma, Z.-W.; Yang, B.; Liu, Z.-J.; Chen, X.-P.; Chen, Y.-J. *J. Org. Chem.* **2021**, 86, 3546.
(b) Wang, C.-C.; Ma, Z.-W.; Qu, Y.-L.; Liu, Z.-J.; Chen, X.-P.; Zhou, J.; Chen, Y.-J. *Chem.-Asian J.* **2020**, 15, 560.
(c) Zhou, J.; Zhang, H.; Chen, X.-L.; Qu, Y.-L.; Zhu, Q.; Feng, C.-G.; Chen, Y.-J. *J. Org. Chem.* **2019**, 84, 9179.

(Zhao, C.)