钴(II)-三联吡啶配位聚合物催化的苄位直接氧化

刘建奇 范伟伟* 熊航行 江京耘 詹红菊

(荆楚理工学院化工与药学院 湖北荆门 448000)

摘要 通过苄位 C—H 直接氧化制备了一系列芳香酮类化合物. 该反应体系采用钴(II)-三联吡啶配位聚合物作为催化剂, 叔丁基过氧化氢(TBHP)作为氧化剂, Na_2CO_3 作为助催化剂, 水作为溶剂, 以较高的收率(78%~99%)制得了 19 种芳香酮类化合物. 该方法底物适用范围广, 具有较高的化学选择性和官能团耐受性. 通过 2,2,6,6-四甲基哌啶-1-氧(TEMPO)的自由基捕获实验及对反应过程中中间体的监测, 提出了一种合理的自由基反应机理.

关键词 苄位直接氧化; 配位聚合物; 叔丁基过氧化氢; 酮类

Benzylic Oxidation Catalyzed by Cobalt(II)-Terpyridine Coordination Polymers

Liu, Jianqi Fan, Weiwei* Xiong, Hangxing Jiang, Jingyun Zhan, Hongju
(College of Chemical Engineering and Pharmacy, Jingchu University of Technology, Jingmen, Hubei 448000)

Abstract Direct benzylic C—H oxidation for the synthesis of aromatic ketones was developed. Employing cobalt(II)-terpyridine coordination polymers as catalyst, *tert*-butyl hydroperoxide (TBHP) as oxidant and Na_2CO_3 as activator, 19 aromatic ketones were prepared with good to excellent yields (78%~99%) in water. The reaction showed a broad range of substrates with good functional group tolerance and chemical selectivity. By control experiments, a trapping experiment using 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) and detection of intermediates during reaction, a reasonable radical mechanism for this type of reaction was also demonstrated.

Keywords direct benzylic oxidation; coordination polymer; *tert*-butyl hydroperoxide; ketone

1 Introduction

Direct C—H oxidation is one of the most fundamental and valuable transformations in organic synthesis.^[1-3] For decades, applying direct C—H oxidation for the synthesis of ketones has attracted extensive research interest, as aromatic ketones are important intermediates for the synthesis of pharmaceuticals, natural products, agrochemicals, etc. For example, benzoylpyridines are key intermediates for the synthesis of many pharmaceutical products.^[4-6] For metal-catalyzed C—H oxidation, plenty of transition-metal catalysts have been developed, such as Ru,^[7] Re,^[8] Au,^[9] Cu,^[10] Co,^[11] Fe,^[12-13] Mn,^[14,15] Ni,^[16] Cr,^[17] Zn,^[18] and V,^[19] as well as Pd-Cu^[20] and Mo-Cu^[21] bimetal systems. In addition, several metal-free catalytic systems have been reported, including *N*-hydroxyphthalimide/ O_2 ,^[22] *N*-heterocyclic carbenes,^[23] I_2 /pyridine/ BuOOH (TBHP),^[24] and KBr-Oxone,^[25,26] etc. Despite significant progress, there

are still many challenges to overcome for catalytic C—H oxidation. It is a growing concern that most reaction media are environmentally unfriendly, although in most known examples the reactions were carried out under mild conditions. Towards a perspective of green chemistry, the use of benign oxidizers (O_2 , H_2O_2 , TBHP, etc.) and greener solvents such as H_2O and alcohols are desired.^[7,27,28]

In metal catalysis, the replacement of noble metals with earth-abundant metals is a desirable aim. Employing nitrogen based ligands instead of expensive, air-sensitive and toxic phosphorus ligands is also highly anticipated. To address this, a group of cobalt and manganese coordination polymers based on readily available 4'-(4-pyridyl)-2,2':6',2''-terpyridine ligands have been previously developed by the Zhang group.^[29-33] These catalysts were air- and bench-stable, and exhibited excellent catalytic efficiency for carbonyl, alkene and alkyne hydroboration or

* Corresponding author. E-mail: fwwjm@foxmail.com

Received April 14, 2021; revised June 10, 2021; published online August 9, 2021.

Project supported by the Hubei Provincial Scientific and Technological Innovation Team Project (No. T2020023) and the Science and Technology Plan Project of Jingmen City (No. 2020ZDYF002).

湖北省科技创新群体项目(No. T2020023)和荆门市科技计划(No. 2020ZDYF002)资助项目.

hydrofunctionalization. We were interested in exploring whether such metal coordination polymers could be utilized in oxidation catalysis. Herein, we report the facile and mild oxidation of benzylic C—H bonds of aryl ketones with TBHP in aqueous media.

Recently, a metal-free TBHP system for benzylic oxidation has been revealed,^[34] however, excessive oxidant (12 equiv.) and high temperature were required. Although several Co(II) catalysts for benzylic C—H oxidation with TBHP, including Co(acac)₂, Co(II) porphyrin and Co(II) complex with 2,4,6-tris(2-pyridyl)-1,3,5-triazine, have also appeared^[35–37] (Scheme 1), none of them could be applied in aqueous media. Our study provides an alternative Co(II) catalyst with high catalytic activity in aqueous media, and also proposes a new model of direct C—H oxidation that employs one-dimensional (1-D) coordination polymer (CP) instead of conventional small molecules.

Scheme 1 Benzylic C—H oxidation with ^tBuOOH catalyzed by Co(II) complexes

2 Results and discussion

To evaluate the catalytic efficiency of coordination polymer **1**, the oxidation of diphenylmethane was selected as a model reaction. The influence of reaction conditions, including the catalysts, oxidants, activators and solvents was investigated (Table 1). It was pleased to observe that diphenylmethane could be efficiently oxidized by TBHP in water using **1** as a catalyst in the presence of Na₂CO₃, leading to complete conversion to benzophenone (Entry 1). Control experiments revealed that no product was observed

in the absence of catalyst (Entry 2) and a much lower yield of 35% was observed with CoCl₂·6H₂O as a catalyst (Entry 3), proving the role of **1** as an effective catalyst. It was further found that the choice of oxidizer is crucial to high efficiency. In the absence of TBHP or replacing it with other oxidizers (O₂, H₂O₂, ^tBuOO^tBu), the oxidation of diphenylmethane did not occur at all (Entries 4~7). The solvent effect was also found to be significant. The yields dramatically fall in other solvents such as MeCN, MeOH, dioxane or EtOH (Entries 8~11). Adding a base as an activator is also a key to high efficiency, which could significantly increase yield from 13% to 99% (Entries 1 and 12). But the selection of base additive was found to be less important. The reactions with K₂CO₃, NaOH, Et₃N or ^tBuOK as an activator were all effective, offering excellent yields ranging from 90% to 99% (Entries 13~16).

Table 1 Optimization of reaction conditions^a

Entry	Catalyst	Oxidant	Activator	Solvent	Time/h	Yield ^b /%
1	1	TBHP	Na ₂ CO ₃	H ₂ O	7	>99
2	—	TBHP	Na ₂ CO ₃	H ₂ O	10	—
3	CoCl ₂ ·6H ₂ O	TBHP	Na ₂ CO ₃	H ₂ O	10	35
4	1	—	Na ₂ CO ₃	H ₂ O	10	—
5	1	O ₂ ^c	Na ₂ CO ₃	H ₂ O	10	—
6	1	H ₂ O ₂	Na ₂ CO ₃	H ₂ O	10	—
7	1	^t BuOO ^t Bu	Na ₂ CO ₃	H ₂ O	10	—
8	1	TBHP	Na ₂ CO ₃	MeCN	10	62
9	1	TBHP	Na ₂ CO ₃	MeOH	10	33
10	1	TBHP	Na ₂ CO ₃	Dioxane	10	6
11	1	TBHP	Na ₂ CO ₃	EtOH	10	11
12	1	TBHP	—	H ₂ O	7	13
13	1	TBHP	K ₂ CO ₃	H ₂ O	7	>99
14	1	TBHP	NaOH	H ₂ O	7	>99
15	1	TBHP	Et ₃ N	H ₂ O	7	>99
16	1	TBHP	^t BuOK	H ₂ O	7	90

^a All reactions were performed on a 1-mmol scale in 3 mL solvent with 3 mol% catalyst, 5 equiv. oxidant and 3 mol% activator. ^b Determined by GC analysis with decane as an internal standard. ^c Oxygen balloon.

Under the optimal reaction conditions, the scope of benzylic substrates was investigated, and the results are summarized in Table 2. For diphenylmethane type substrates (Table 2, **2a**~**2c**), substituted benzophenones were observed with excellent yields (95%~99%), and functional groups such as Cl and COOH were tolerated, even though it was assumed that the introduction of COOH group would consume the alkali activator and thus inhibited the reaction. One possible explanation is that the substrate is insoluble in water and the tardy neutralization reaction could not prevent the activator from promoting the catalytic process. For fluorene type substrates (**2d**~**2g**), the reaction afforded oxidation products with high to excellent yields (80%~99%) and some electron-withdrawing groups (Br and acetyl) were tolerant. However, the intro-

duction of stronger electron-withdrawing group such as NO₂ caused an evident decline of yield (**2g**). 9,10-Dihydroanthracene was oxidized completely on its both benzyl sites to afford anthraquinone with a high yield (**2h**). 4-Benzylpyridine and xanthene were also suitable for this method, showing excellent yields (**2i**, **2j**). In addition, the catalytic method has been successfully applied to ethylbenzene and its derivatives, and good to excellent yields (78%~95%) were observed (**2k**~**2s**). Both electron-donating group (OCH₃) and electron-withdrawing groups (Cl, Br, I) were tolerated, while in the case of 2-bromoethylbenzene the yield was slightly lower, possi-

bly due to the steric effect (**2s**).

The oxidation protocol was also applied for a gram-scale synthesis. Under the same reaction condition, a 10 mmol scale of diphenylmethane (1.68 g) was oxidized completely, resulting in the production of **2a** in 97% isolated yield.

Finally, the possible mechanism was investigated to gain insights into the oxidative process. As the yield did not vary under the N₂ atmosphere, and the substrate could not be oxidized by O₂ (Table 1, Entry 5), the TBHP is the main oxidant. In a trapping experiment using 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO), the TEMPO adduct **4** was confirmed by LC-MS (Scheme 2, a). That indicated

Table 2 Substrate scope for the benzylic oxidation^a

99% ^b (99% ^c)	99% (99%)	98% ^d (95%)	99% (99%)	95% (93%)	99% (98%)

85% (80%)	99% ^d (95%)	97% (96%)	99% (97%)	97% (95%)	96% ^d (89%)

93% (90%)	99% (86%)	97% (88%)	98% (87%)	95% (89%)	96% (90%)

			84% (78%)		

^a All reactions were performed on a 1-mmol scale in 3 mL solvent. ^b Yield determined by GC analysis with decane as an internal standard. ^c Isolated yield. ^d Yield determined by ¹H NMR with 1,4-dioxane as an internal standard. ^e 8 equiv. of TBHP was added.

Scheme 2 Trapping experiment (a), incomplete oxidation of 9,10-dihydroanthracene (b) and diphenylmethane (c)

the abstraction of benzylic hydrogen through a radical mechanism during the reaction. This is also supported by the incomplete oxidation of 9,10-dihydroanthracene. When TBHP was inadequate (4 equiv. instead of 8 equiv.), the dehydrogenated product anthracene (**5**) was obtained as a byproduct (Scheme 2, b). During the oxidation of diphenylmethane, a key peroxide intermediate **6** was also observed (Scheme 2, c). It was isolated and confirmed by ^1H NMR and ^{13}C NMR spectroscopies.

Based on the above results, a plausible reaction mechanism was proposed as shown in Scheme 3. Firstly, Co(II) complex could react with TBHP to form a Co(III)-peroxy adduct, releasing $^t\text{BuOO}\cdot$ and $^t\text{BuO}\cdot$ while regenerating the Co(II) complex.^[38-39] Then the abstraction of benzylic hydrogen took place, affording radical **7** resulting from the reaction with $^t\text{BuO}\cdot$.^[7,21] The newly generated $^t\text{BuOO}\cdot$ could also abstract benzylic hydrogens, but the rate was much less than that of $^t\text{BuO}\cdot$.^[40] Therefore, it would mainly react with radical **7** to form the peroxide intermediate **6**. Finally, **6** decomposed to give the resulting ketone product.^[21,34] It was believed that the Co(II) coordination polymers might have facilitated the decomposition of peroxide intermediate **6**, as it was known that its spontaneous decomposition could only be carried out at a much higher temperature (130 °C) with longer reaction time.^[34]

Scheme 3 Possible reaction mechanism

3 Conclusions

In summary, an efficient procedure for the benzylic oxidation in water, using cobalt(II)-terpyridine coordination polymer as a catalyst and TBHP as an oxidant was developed. A broad scope of substrates have been explored and

good to excellent yields of the corresponding ketone products were obtained. Further studies on the structural modification of the current coordination polymer catalyst for improved performance in more challenging direct C—H oxidation reactions are underway.

4 Experimental section

4.1 Materials and methods

All the Chemicals and solvents were purchased from commercial suppliers and used as purchased without any further purification. Unless stated otherwise, all the reactions were carried out in a 10 mL test tube with rubber stopper under air atmosphere. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker Avance III HD 400MHz spectrometer at ambient temperature with CDCl_3 as solvent. For the calculation of conversions of some substrates, dioxane was added as the internal standard. Conversions of most substrates were determined by a TRACE 1300 gas chromatography equipped with a Phenomenex Zebron ZB-1701 capillary column. Mass spectrometry (LC-MS) was performed on a Thermo Scientific LCQ Fleet Ion Trap Mass Spectrometer.

4.2 Preparation of Co(II) coordination polymer (catalyst)

Co(II) coordination polymer was prepared according the literatures.^[32-33] 4'-(4-Pyridyl)-2,2':6',2''-terpyridine (310 mg, 1 mmol) was dissolved in 40 mL of CH_2Cl_2 and then placed in a 100 mL test tube. After 20 mL of MeOH was layered on the top of the solution, a solution of $\text{CoCl}_2\cdot 6\text{H}_2\text{O}$ (238 mg, 1 mmol) in MeOH (40 mL) was slowly added. The tube was sealed and stored at room temperature. Four weeks later, the crystals were collected and washed with EtOH, and then dried in vacuo. Finally, light brown crystals were obtained (285 mg, 52% yield) and the structure was confirmed by a X-ray single crystal diffractometer.

4.3 General oxidation procedure

Substrate (1 mmol), TBHP (0.65 g, 70% in water), Co(II) coordination polymer (13 mg, 0.03 mmol based on $\text{Co}(\text{L})\text{Cl}_2$), Na_2CO_3 (3.2 mg, 0.03 mmol) and water (3 mL) were added to a 10 mL test tube equipped with a stir bar. The reaction mixture was stirred at 70 °C for 7 h. After the reaction, the reaction mixture was cooled down to room temperature and extracted 3 times with CH_2Cl_2 (10~15 mL). The CH_2Cl_2 solution was diluted and analyzed by GC with decane as an internal standard. For substrates **2c**, **2h** and **2l**, the CH_2Cl_2 solution was evaporated and analyzed by ^1H NMR with 1,4-dioxane as an internal standard.

In a repeat experiment, the CH_2Cl_2 solution was dried with anhydrous sodium sulfate and then evaporated. The crude product was purified through silica gel column chromatography using petroleum ether and ethyl acetate as eluent.

Benzophenone (**2a**): White solid (181 mg, 99% yield). m.p. 46~48 °C (lit.^[7] 47.5~47.9 °C); ^1H NMR (400

MHz, CDCl₃) δ : 7.80 (dt, J =8.4, 1.6 Hz, 4H), 7.63~7.53 (m, 2H), 7.52~7.43 (m, 4H); ¹³C NMR (101 MHz, CDCl₃) δ : 196.82, 137.52, 132.44, 130.07, 128.27.

(4-Chlorophenyl)(phenyl)methanone (**2b**): White solid (216 mg, 99% yield). m.p. 75~78 °C (lit.^[41] 77.8~78.4 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.80~7.70 (m, 4H), 7.64~7.57 (m, 1H), 7.52~7.42 (m, 4H); ¹³C NMR (101 MHz, CDCl₃) δ : 195.43, 138.85, 137.19, 135.83, 132.63, 131.45, 129.91, 128.61, 128.39.

2-Benzoylbenzoic acid (**2c**):^[42] White solid (215 mg, 95% yield). m.p. 126~128 °C (123~125 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.07 (d, J =7.8 Hz, 1H), 7.68 (ddd, J =10.3, 8.7, 4.2 Hz, 3H), 7.56 (ddd, J =10.2, 8.4, 4.3 Hz, 2H), 7.40 (dd, J =16.4, 8.5 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 197.11, 170.71, 142.60, 136.96, 133.28, 133.26~132.94 (m), 130.92, 129.58, 129.45, 128.48, 127.85, 127.71.

9H-Fluoren-9-one (**2d**): Yellow solid (1789 mg, 99% yield). m.p. 83~84 °C (lit.^[43] 79~81 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.65~7.56 (m, 2H), 7.48~7.39 (m, 4H), 7.30~7.20 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) δ : 193.97, 144.42, 134.72, 134.12, 129.09, 124.31, 120.34.

2-Bromo-9H-fluoren-9-one (**2e**): Yellow solid (241 mg, 93% yield). m.p. 145~146 °C (lit.^[43] 143~146 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.76 (d, J =1.8 Hz, 1H), 7.66 (dt, J =7.3, 0.8 Hz, 1H), 7.61 (dd, J =7.9, 1.9 Hz, 1H), 7.53~7.49 (m, 2H), 7.40 (d, J =7.9 Hz, 1H), 7.32 (ddd, J =7.4, 5.0, 3.5 Hz, 1H); ¹³C NMR (101 MHz, CDCl₃) δ : 192.44, 143.68, 143.01, 137.11, 135.76, 135.05, 133.69, 129.44, 127.58, 124.63, 122.92, 121.74, 120.45.

2-Acetyl-9H-fluoren-9-one (**2f**): Yellow solid (218 mg, 98% yield). m.p. 156~158 °C (lit.^[43] 153~156 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.29~8.05 (m, 2H), 7.73 (d, J =7.4 Hz, 1H), 7.64 (dd, J =11.4, 4.0 Hz, 2H), 7.56 (tt, J =8.9, 4.4 Hz, 1H), 7.40 (td, J =7.4, 1.0 Hz, 1H), 2.65 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 196.64, 192.79, 148.50, 143.26, 137.78, 135.06, 134.92, 134.87, 134.31, 130.25, 124.68, 124.19, 121.29, 120.46, 26.78.

2-Nitro-9H-fluoren-9-one (**2g**): Yellow solid (180 mg, 80% yield). m.p. 220~222 °C (lit.^[44] 222~223 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.49 (d, J =2.1 Hz, 1H), 8.42 (dd, J =8.2, 2.2 Hz, 1H), 7.82~7.75 (m, 1H), 7.72~7.66 (m, 2H), 7.62 (td, J =7.5, 1.1 Hz, 1H), 7.46 (td, J =7.4, 1.0 Hz, 1H); ¹³C NMR (101 MHz, CDCl₃) δ : 190.97, 149.74, 148.73, 142.30, 135.46, 135.03, 134.98, 131.02, 129.96, 125.12, 121.80, 120.72, 119.58.

Anthracene-9,10-dione (**2h**): Yellow-green solid (198 mg, 95% yield). m.p. 279~281 °C (lit.^[7] 277.9~278.4 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.32 (dt, J =7.0, 3.5 Hz, 4H), 7.84~7.78 (m, 4H); ¹³C NMR (101 MHz, CDCl₃) δ : 183.18, 134.14, 133.52, 127.24.

Phenyl(pyridin-4-yl)methanone (**2i**): Yellow solid (176 mg, 96% yield). m.p. 69~71 °C (lit.^[4] 68~70 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.82 (dd, J =4.4, 1.6 Hz, 2H), 7.90~7.77 (m, 2H), 7.72~7.62 (m, 1H), 7.59 (dd, J =4.4, 1.6 Hz, 2H), 7.56~7.49 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) δ : 195.21, 150.40, 144.37, 135.89, 133.58, 130.17,

128.69, 122.91.

9H-Xanthen-9-one (**2j**): Beige solid (190 mg, 97% yield). m.p. 172~174 °C (lit.^[7] 172.9~173.4 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.36 (dd, J =8.0, 1.6 Hz, 2H), 7.74 (ddd, J =8.7, 7.1, 1.7 Hz, 2H), 7.51 (dd, J =8.4, 0.6 Hz, 2H), 7.39 (ddd, J =8.1, 7.2, 1.0 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ : 177.27, 156.21, 134.84, 126.77, 123.93, 121.88, 118.00.

Acetophenone (**2k**):^[7] Colorless oil (115 mg, 95% yield). ¹H NMR (400 MHz, CDCl₃) δ : 7.97 (dt, J =8.5, 1.7 Hz, 2H), 7.61~7.53 (m, 1H), 7.52~7.42 (m, 2H), 2.61 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 198.25, 137.11, 133.16, 128.60, 128.34, 26.68.

1-Naphthalen-1-yl-ethanone (**2l**):^[41] Colorless oil (151 mg, 89% yield). ¹H NMR (400 MHz, CDCl₃) δ : 8.83~8.69 (m, 1H), 7.96~7.81 (m, 3H), 7.58 (ddd, J =8.5, 6.8, 1.5 Hz, 1H), 7.52~7.41 (m, 2H), 2.70 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 201.81, 135.34, 133.93, 133.02, 130.09, 128.70, 128.38, 128.02, 126.40, 125.97, 124.30, 29.94.

4-Acetylbiphenyl (**2m**): Beige solid (176 mg, 90% yield). m.p. 120~122 °C (lit.^[7] 128.4~128.8 °C); ¹H NMR (400 MHz, CDCl₃) δ : 8.10~7.99 (m, 2H), 7.73~7.67 (m, 2H), 7.67~7.60 (m, 2H), 7.53~7.45 (m, 2H), 7.41 (d, J =7.3 Hz, 1H), 2.65 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 197.8, 145.8, 139.88, 135.8, 128.9, 128.9, 128.2, 127.2, 127.2, 26.6.

1-(4-Methoxyphenyl)ethanone (**2n**): White solid (129 mg, 86% yield). m.p. 37~38 °C (lit.^[7] 38.2~38.5 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.96~7.89 (m, 2H), 6.96~6.88 (m, 2H), 3.85 (s, 3H), 2.54 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 196.76, 163.47, 130.57, 130.27, 113.66, 55.44, 26.33.

1-(4-Chlorophenyl)ethanone (**2o**):^[21] Colorless oil (136 mg, 88% yield). ¹H NMR (400 MHz, CDCl₃) δ : 7.94~7.78 (m, 2H), 7.40 (dd, J =8.9, 2.2 Hz, 2H), 2.57 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 196.69, 139.41, 135.33, 129.69, 128.79, 26.49.

1-(4-Bromophenyl)ethanone (**2p**): White solid (173 mg, 87% yield). m.p. 50~52 °C (lit.^[7] 51.4~52.2 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.85~7.77 (m, 2H), 7.62~7.53 (m, 2H), 2.58 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 197.00, 135.77, 131.87, 129.85, 128.29, 26.57.

1-(4-Iodophenyl)ethanone (**2q**): Beige solid (219 mg, 89% yield). m.p. 82~84 °C (lit.^[7] 84.4~84.8 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.89~7.76 (m, 2H), 7.73~7.60 (m, 2H), 2.58 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 197.3, 137.9, 136.3, 129.7, 101.1, 26.4.

1-(3-Bromophenyl)ethanone (**2r**):^[21] Yellow oil (191 mg, 90% yield). ¹H NMR (400 MHz, CDCl₃) δ : 8.07 (t, J =1.8 Hz, 1H), 7.93~7.81 (m, 1H), 7.68 (ddd, J =8.0, 2.0, 1.0 Hz, 1H), 7.34 (t, J =7.9 Hz, 1H), 2.59 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 196.60, 138.76, 135.94, 131.33, 130.21, 126.87, 122.94, 26.63.

1-(2-Bromophenyl)ethanone (**2s**):^[21] Yellow oil (167 mg, 78% yield). ¹H NMR (400 MHz, CDCl₃) δ : 7.61 (dd, J =7.9, 1.0 Hz, 1H), 7.47 (dd, J =7.6, 1.7 Hz, 1H), 7.37

(td, $J=7.5$, 1.1 Hz, 1H), 7.32~7.26 (m, 1H), 2.63 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 201.42, 141.46, 133.86, 131.83, 128.94, 127.47, 118.92, 30.36.

4.4 Gram-scale reaction

Diphenylmethane (1.7 g), TBHP (6.5 g, 70% in water), Co(II) coordination polymer (100 mg), Na_2CO_3 (30 mg) and water (30 mL) were added to a 100 mL flask equipped with a stir bar. The reaction mixture was stirred at 70 °C for 7 h. After the reaction, the reaction mixture was cooled down to room temperature and extracted 3 times with CH_2Cl_2 (15 mL). The CH_2Cl_2 solution was then evaporated. The crude product was purified through silica gel column chromatography using petroleum ether and ethyl acetate as eluent.

4.5 Trapping experiment

4-Benzylpyridine (169 mg, 1 mmol), TBHP (0.65 g, 70% in water), TEMPO (780 mg, 5 mmol), Co(II) coordination polymer (13 mg, 0.03 mmol based on Co(L)Cl₂), Na_2CO_3 (3.2 mg, 0.03 mmol) and water (3 mL) were added to a 10 mL test tube equipped with a stir bar. The reaction mixture was stirred at 70 °C for 7 h. After the reaction, the reaction mixture was cooled down to room temperature and extracted 3 times with CH_2Cl_2 (10~15 mL). The CH_2Cl_2 solution was diluted and analyzed by LC-MS. The products were isolated by silica gel column chromatography using petroleum ether and ethyl acetate as eluent.

Supporting Information Preparation of catalyst, ^1H NMR and ^{13}C NMR spectra, LC-MS spectrum. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Karimov, R. R.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2018**, *57*, 4234.
- [2] Wu, J.; Zhu, J.; Li, H.; Wu, C.; Shen, R.; Yu, L. *Chin. J. Org. Chem.* **2019**, *39*, 3328 (in Chinese). (吴锦雯, 朱佳雯, 李慧, 吴春雷, 沈润溥, 余乐茂, 有机化学, **2019**, *39*, 3328.)
- [3] Sun, Q.; Sun, W. *Chin. J. Org. Chem.* **2020**, *40*, 3686 (in Chinese). (孙强盛, 孙伟, 有机化学, **2020**, *40*, 3686.)
- [4] Ren, L.; Wang, L.; Lv, Y.; Li, G.; Gao, S. *Org. Lett.* **2015**, *17*, 2078.
- [5] Ren, L.; Gao, S. *Chin. J. Org. Chem.* **2017**, *37*, 1338 (in Chinese). (任兰会, 高爽, 有机化学, **2017**, *37*, 1338.)
- [6] Pieber, B.; Kappe, C. O. *Green Chem.* **2013**, *15*, 320.
- [7] Bo, C.; Bu, Q.; Li, X.; Ma, G.; Wei, D.; Guo, C.; Dai, B.; Liu, N. *J. Org. Chem.* **2020**, *85*, 4324.
- [8] Peng, H.; Lin, A.; Zhang, Y.; Jiang, H.; Zhou, J.; Cheng, Y.; Zhu, C.; Hu, H. *ACS Catal.* **2012**, *2*, 163.
- [9] Li, H.; Li, Z.; Shi, Z. *Tetrahedron* **2009**, *65*, 1856.
- [10] Sterckx, H.; De Houwer, J.; Mensch, C.; Caretti, I.; Tehrani, K. A.; Herrebout, W. A.; Van Doorslaer, S.; Maes, B. U. W. *Chem. Sci.* **2016**, *7*, 346.
- [11] Hruszkewycz, D. P.; Miles, K. C.; Thiel, O. R.; Stahl, S. S. *Chem. Sci.* **2017**, *8*, 1282.
- [12] Miao, C.; Zhao, H.; Zhao, Q.; Xia, C.; Sun, W. *Catal. Sci. Technol.* **2016**, *6*, 1378.
- [13] Liu, P.; Liu, Y.; Wong, E. L.; Xiang, S.; Che, C. *Chem. Sci.* **2011**, *2*, 2187.
- [14] Shen, D.; Miao, C.; Wang, S.; Xia, C.; Sun, W. *Org. Lett.* **2014**, *16*, 1108.
- [15] Talsi, E. P.; Samsonenko, D. G.; Ottenbacher, R. V.; Bryliakov, K. P. *ChemCatChem* **2017**, *9*, 4580.
- [16] Urgoitia, G.; Sanmartin, R.; Herrero, M. T.; Domínguez, E. *Chem. Commun.* **2015**, *51*, 4799.
- [17] Yamazaki, S. *Org. Lett.* **1999**, *1*, 2129.
- [18] Muthupandi, P.; Sekar, G. *Tetrahedron Lett.* **2011**, *52*, 692.
- [19] Xia, J.; Cormier, K. W.; Chen, C. *Chem. Sci.* **2012**, *3*, 2240.
- [20] Jeong, Y.; Moon, Y.; Hong, S. *Org. Lett.* **2015**, *17*, 3252.
- [21] Li, P.; Wang, Y.; Wang, X.; Wang, Y.; Liu, Y.; Huang, K.; Hu, J.; Duan, L.; Hu, C.; Liu, J. *J. Org. Chem.* **2020**, *85*, 3101.
- [22] Yang, G.; Zhang, Q.; Miao, H.; Tong, X.; Xu, J. *Org. Lett.* **2005**, *7*, 263.
- [23] Alanthadka, A.; Devi, E. S.; Nagarajan, S.; Sridharan, V.; Suvitha, A.; Maheswari, C. U. *Eur. J. Org. Chem.* **2016**, *2016*, 4872.
- [24] Zhang, J.; Wang, Z.; Wang, Y.; Wan, C.; Zheng, X.; Wang, Z. *Green Chem.* **2009**, *11*, 1973.
- [25] Moriyama, K.; Takemura, M.; Togo, H. *Org. Lett.* **2012**, *14*, 2414.
- [26] Yin, L.; Wu, J.; Xiao, J.; Cao, S. *Tetrahedron Lett.* **2012**, *53*, 4418.
- [27] Sheldon, R. A. *Chem. Soc. Rev.* **2012**, *41*, 1437.
- [28] Waffel, D.; Alkan, B.; Fu, Q.; Chen, Y.; Schmidt, S.; Schulz, C.; Wiggers, H.; Muhler, M.; Peng, B. *ChemPlusChem* **2019**, *84*, 1155.
- [29] Zhang, G.; Zeng, H.; Li, S.; Johnson, J.; Mo, Z.; Neary, M. C.; Zheng, S. *Dalton Trans.* **2020**, *49*, 2610.
- [30] Zhang, G.; Li, S.; Wu, J.; Zeng, H.; Mo, Z.; Davis, K.; Zheng, S. *Org. Chem. Front.* **2019**, *6*, 3228.
- [31] Fan, W.; Li, L.; Zhang, G. *J. Org. Chem.* **2019**, *84*, 5987.
- [32] Zhang, G.; Wu, J.; Li, S.; Cass, S.; Zheng, S. *Org. Lett.* **2018**, *20*, 7893.
- [33] Wu, J.; Zeng, H.; Cheng, J.; Zheng, S.; Golen, J. A.; Manke, D. R.; Zhang, G. *J. Org. Chem.* **2018**, *83*, 9442.
- [34] Tan, J.; Zheng, T.; Yu, Y.; Xu, K. *RSC Adv.* **2017**, *7*, 15176.
- [35] Zhang, Z.; Li, J.; Yao, Y.; Sun, S. *Cryst. Growth Des.* **2015**, *15*, 5028.
- [36] Azarkamanzad, Z.; Farzaneh, F.; Maghami, M.; Simpson, J. *New J. Chem.* **2019**, *43*, 12020.
- [37] Han, X.; Zhou, Z.; Wan, C.; Xiao, Y.; Qin, Z. *Synthesis* **2013**, *45*, 615.
- [38] Zhang, J.; Biradar, A. V.; Pramanik, S.; Emge, T. J.; Asefa, T.; Li, J. *Chem. Commun.* **2012**, *48*, 6541.
- [39] Asgharpour, Z.; Farzaneh, F.; Abbasi, A. *RSC Adv.* **2016**, *6*, 95729.
- [40] Bryant, J. R.; Mayer, J. M. *J. Am. Chem. Soc.* **2003**, *125*, 10351.
- [41] Wang, T.; Jing, X.; Chen, C.; Yu, L. *J. Org. Chem.* **2017**, *82*, 9342.
- [42] Cheng, K.; Zhao, B.; Qi, C. *RSC Adv.* **2014**, *4*, 48698.
- [43] Li, L.; Liu, Z.; Tang, S.; Li, J.; Ren, X.; Yang, G.; Li, H.; Yuan, B. *Catal. Commun.* **2019**, *127*, 34.
- [44] Zhang, X.; Ji, X.; Su, R.; Weeks, B. L.; Zhang, Z.; Deng, S. *ChemPlusChem* **2013**, *78*, 703.

(Zhao, C.)