无光催化剂条件下紫外光引发芳基醛或芳基酮与炔烃环化反应 构筑茚酮化合物

肖玉娟^a 杨 阳^a 张 凡^b 冯亚栋^{*,a,b} 崔秀灵^{*,b}

(^a 厦门华厦学院环境与公共健康学院 检测技术和智能仪器研究所 厦门 361024)

(^b 华侨大学生物医学学院 分子药物教育部工程研究中心 福建省分子医学重点实验室 福建省高校精准医学与分子诊断重点实验室 厦门市海洋与基因药物重点实验室 厦门 361021)

摘要 报道了一种紫外光引发的芳基醛或芳基酮与炔烃的环化反应来构筑茚酮化合物。该反应具有效率较高、原子经济性高、原料简单易得、环境友好、底物适应性较强等优势,在紫外光照射,氧气为氧化剂,无光催化剂的条件下可以顺利进行。

关键词 紫外光; 环化反应; 无光催化剂; 茚酮

UV-Light-Initiated Construction of Indenones through Cyclization of Aryl Aldehydes or Aryl Ketones with Alkynes Avoiding Photocatalyst

Xiao, Yujuan^a Yang, Yang^a Zhang, Fan^b Feng, Yadong^{*,a,b} Cui, Xiuling^{*,b}

(^a College of Environment and Public Health, Institute of Analytical Technology and Smart Instruments, Xiamen Huaxia University, Xiamen 361024)

(^b Engineering Research Center of Molecular Medicine, Ministry of Education, Key Laboratory of Fujian Molecular Medicine, Key Laboratory of Precision Medicine and Molecular Diagnosis of Fujian Universities, Key Laboratory of Xiamen Marine and Gene Drugs, School of Biomedical Sciences, Huaqiao University, Xiamen 361021)

Abstract The first UV-light-initiated cyclization of aryl aldehydes or aryl ketones with alkynes has been developed for the preparation of indenones. This reaction could proceed smoothly under UV-light with O₂ as an oxidant and photocatalyst-free condition, which features high efficiency, high atom economy, easily available starting materials, environmental friendliness, and tolerance to broad functional groups.

Keywords UV-light; cyclization; photocatalyst-free; indenones

1 Introduction

Indenones are widely found as structural motifs in various pharmaceutical compounds and natural product family (Scheme 1a).^[1] As a result, the establishment of easy and effective approaches for indenones starting from readily available materials has been continuously desired. With the rapid development of transition metal-catalyzed C—H bond functionalization in recent decades,^[2] undoubtedly, the direct coupling cyclization of aryl aldehydes with alkynes via C(sp²)—H bond functionalization is the most straightforward and atom-economical pathway for substituted indenones. For example, Cheng *et al.*^[3]

reported the direct annulation of aryl aldehyde with alkynes using Rh as a catalyst via *in situ* directing group formation in NH₂NHAc/AcOH system. Notably, Adhikary *et al.*^[4] reported a PIDA mediated radical oxidative annulation of aromatic aldehydes with 1,2-diphenylethyne with di-*t*-butyl peroxide (DTBP) as an oxidant towards the synthesis of 2,3-diarylindenones in 2018. Recently, we^[5] have developed a ferrocene-initiated oxidative cyclization of benzaldehyde with alkyne for a direct access to substituted indenones, in which a radical process was involved (Scheme 1b).

On the other hand, much attention has recently focused on the photocatalysis in the field of C—H functionalization

* Corresponding authors. E-mail: fengyd@hxxy.edu.cn; cuixl@hqu.edu.cn

Received July 7, 2021; revised July 29, 2021; published online August 9, 2021.

Project supported by the Fujian Education and Scientific Research Project for Young and Middle-aged Teachers (No. JAT190990).

福建省中青年骨干教师教育科研(Nos. JAT190990)资助项目.

Scheme 1 Selected examples of the indenone motif in pharmaceuticals and procedure to indenones

due to its ability to promote bond formation under green and mild reaction conditions.^[6] For example, MacMillan *et al.*^[7] reported the first direct C(sp²)—H bond trifluoromethylation of (hetero)arenes using photoredox catalysts in 2011. Sanford *et al.*^[8] reported the C—H bond arylation of phenylpyridines using dual catalysis of Pd and Ru photocatalysis. Since then, some strategies for C—H bond functionalization using transition metal-based photocatalysts have been developed.^[9] Notably, in 2012, König *et al.*^[10] developed the first intermolecular C—H bond arylation of heteroarenes with aryl diazonium salts under metal-free conditions using the organic dye eosin Y as a photoredox catalyst. Then, more and more organic photocatalysts have been employed in C—H bond functionalization instead of transition metal-based photocatalysts, such as iodo-bodipys, rhodamine, acridine, porphyrins, phenothiazines, and anthraquinones.^[11] As well-known, UV-initiated radical polymerization has been demonstrated to exhibit high efficiency due to the high initiator decomposition rate and the low activation energy.^[12] Driven by developing easy and green method in C—H bond functionalization,^[13] herein, we describe the first successful implementation of UV-initiated cyclization of aryl aldehydes or aryl ketones with alkynes using O₂ as a green oxidant for the preparation of indenones without photocatalyst or photoinitiator (Scheme 1c). This protocol features high efficiency, high atom economy, easily available materials, environmental friendliness, and tolerance to broad functional groups.

2 Results and discussion

Initially, benzaldehyde (**1a**) and diphenylethyne (**2a**)

were chosen as the standard substrates for the desired coupling cyclization reaction. The results revealed that 2,3-diphenyl-1*H*-inden-1-one (**3a**) was obtained in 41% yield in CH₃CN at 120 °C under UV-light (λ_{max} = 365 nm, 250 W) in O₂ for 24 h, and benzoic acid was detected in large amount as the by-product (Table 1, Entry 1). Only trace of target product **3a** was detected by GC-MS when the reaction was carried out without UV-light or under N₂, and 23% yield of **3a** was obtained under air (Table 1, Entries 2~4). When the power of UV-light was reduced to 200 and 150 W, the yield of **3a** was decreased to 25% and 13% (Table 1, Entries 5 and 6). Moreover, CH₃CN was proved to favor this reaction better than other solvents, including CH₃OH, EtOH, 1,2-dichloroethane (DCE), *N,N*-dimethylformamide (DMF), dimethyl sulfoxide (DMSO), and toluene (Table 1, Entry 3 vs Entries 7~12). In addition, the yield of **3a** was increased to 63% when the reaction was carried out at 140 °C (Table 1, Entries 13~16). Fortunately, when the reaction time was prolonged to 48 h, the yield of **3a** could be increased to 82% (Table 1, Entries 17~21). However, when the loading of **1a** was decreased to 0.25 and 0.10 mmol, the yield of **3a** was decreased to 28% and trace, respectively, and benzoic acid was detected

Table 1 Optimization of reaction conditions^a

Entry	UV-light (λ_{max} , P)	Solvent	Temp./°C	t/h	Yield ^b /%
1	365 nm, 250 W	CH ₃ CN	120	24	41
2	—	CH ₃ CN	120	24	Trace
3 ^c	365 nm, 250 W	CH ₃ CN	120	24	23
4 ^d	365 nm, 250 W	CH ₃ CN	120	24	Trace
5	365 nm, 200 W	CH ₃ CN	120	24	25
6	365 nm, 150 W	CH ₃ CN	120	24	13
7	365 nm, 250 W	CH ₃ OH	120	24	Nd
8	365 nm, 250 W	EtOH	120	24	Nd
9	365 nm, 250 W	DCE	120	24	11
10	365 nm, 250 W	DMF	120	24	Trace
11	365 nm, 250 W	DMSO	120	24	Nd
12	365 nm, 250 W	Toluene	120	24	20
13	365 nm, 250 W	CH ₃ CN	110	24	30
14	365 nm, 250 W	CH ₃ CN	130	24	57
15	365 nm, 250 W	CH ₃ CN	140	24	63
16	365 nm, 250 W	CH ₃ CN	150	24	60
17	365 nm, 250 W	CH ₃ CN	140	32	69
18	365 nm, 250 W	CH ₃ CN	140	40	75
19	365 nm, 250 W	CH ₃ CN	140	48	82
20	365 nm, 250 W	CH ₃ CN	140	56	73
21	365 nm, 250 W	CH ₃ CN	140	72	74
22 ^e	365 nm, 250 W	CH ₃ CN	140	48	28
23 ^f	365 nm, 250 W	CH ₃ CN	140	48	Trace

^a Reaction conditions: **1a** (0.50 mmol), **2a** (0.10 mmol), solvent (1.5 mL), reflux under O₂. ^b Isolated yields. ^c Under air. ^d Under N₂. ^e **1a** (0.25 mmol). ^f **1a** (0.10 mmol). Nd = not detected.

as the main by-product (Table 1, Entries 22~23). Based on the results, under the optimal reaction conditions, the target product **3a** was obtained in 82% yield in CH₃CN at 140 °C under O₂ atmosphere and UV-light ($\lambda_{\max}$ = 365 nm, 250 W) for 48 h (Table 1, Entry 19).

With the optimized reaction conditions in hand, the scope of the substrates was examined (Table 2). 1,2-Diphenylethyne (**2a**) reacted smoothly with benzaldehyde (**1a**) and its derivatives (**1b**~**1i**) to give the substituted indenones (**3a**~**3i**) in moderate to good yields (57%~84%). Methyl group at 2-, 3-, and 4- position of benzaldehyde provided the corresponding products **3b**, **3c**, and **3d** in 57%, 63%, and 84% yields, which indicated that steric effect of groups slightly impacted on this transformation. When 3-methylbenzaldehyde (**1c**) was introduced in the reaction, only 6-methyl-2,3-diphenyl-1*H*-inden-1-one (**3e**) was obtained with 63% yield at the less steric hindrance. Halogen groups, including F, Cl, and Br at 4-position of benzaldehydes provided the corresponding products **3e**~**3g** in 70%, 68%, and 72% yields, respectively. Other substituents, such as CH₃O and CF₃, could be well tolerated and gave the corresponding products in satisfactory yields (**3h**, **3i**) (80%, 77%), which indicated that electron density on the moiety of the benzaldehydes did not significantly influence on the efficiency of the C—H functionalization cyclization. Benzaldehyde (**1a**) could also react smoothly with substituted alkynes (**2b**~**2d**) to give the desired products **3j**~**3l** in 71%~78% yields. Chloro group at 2- and 4-position of diarylacetylenes (**2b**, **2c**) provided the corresponding products **3j** and **3k** in 71% and 74% yields. Moreover, CF₃ group at 4-position of diarylacetylene could

also react with benzaldehyde (**1a**) smoothly under the standard reaction conditions to give the corresponding product **3l** in 78% yield.

To demonstrate the reaction pathway, some control experiments were carried out (Scheme 2). Firstly, when 2-oxo-2-phenylacetic acid (**4**) and benzoyl chloride (**5**), being common photochemical reagents affording acyl radical,^[14] reacted with 1,2-diphenylethyne (**2a**) under the standard conditions, only trace of target product **3a** was detected by GC-MS and **2a** was recovered in 97% and 96% yields (Schemes 2a and 2b). When 2-hydroxy 1,2-diphenylethanone (**6**) and 2,2-diethoxy-1-phenylethanone (**7**), also being common photoinitiators, were introduced into the reaction, **3a** was obtained in moderate yields (55% and 62%) (Schemes 2c and 2d). It could be shown that **6** and **7** have good ability to generate acyl radical under UV-light at 365 nm in this reaction system. On the other hand, when alkyl alkynes and unsymmetric alkynes, such as oct-4-yne (**8**) and ethynylbenzene (**9**), were introduced into the reaction, the corresponding products **3m** or **3n** (or **3n'**) were not detected, and the coupling product (*E*)-chalcone (**10**) was detected by GC-MS (Schemes 2e and 2f). In addition, when olefins were introduced into the reaction, such as (*E*)-1,2-diphenylethene (**11**) and styrene (**13**), the corresponding products **3o** and **3p** (or **3p'**) were detected in trace yields, and the coupling products (*E*)-1,2,3- triphenylprop-2-en-1-one (**12**) and (*E*)-chalcone (**10**) were detected by GC-MS, respectively (Schemes 2g and 2h).

When the reaction was carried out at room temperature or with a photoinitiator at room temperature, such as benzophenone (BP), the target product **3a** was detected in

Table 2 Scope of substrates^a

^a Reaction conditions: **1** (0.50 mmol), **2** (0.10 mmol), solvent (1.5 mL), reflux under O₂; isolated yield.

Scheme 2 Control experiments with other substrates

trace amount, and benzoic acid (**14**) was detected in large

amount as expected, which indicated that the acyl radical was generated from the dissociation of benzaldehyde with the assistance of UV-light and high temperature (Schemes 3a, 3b and Table 1, Entry 2). On the other hand, when the reaction was carried out under N₂, **3a** was detected in trace amount and the coupling product (*E*)-1,2,3-triphenylprop-2-en-1-one (**12**) was detected by GC-MS, which indicated that O₂ was crucial in the final cyclization step (Scheme 3c). When a radical scavenger, 2,2,6,6-tetramethylpiperidinoxy (TEMPO), was introduced into the reaction (Scheme 3d), only trace of **3a** was detected, and compound **15** was generated as the by-product, which was detected by HRMS. It was suggested that the acyl radical might be the key intermediate in this reaction.

Scheme 3 Control experiments with other conditions

Based on our previous work and literatures,^[15] a plausible reaction mechanism for the UV-initiated radical C—H cyclization of aryl aldehydes or aryl ketones with alkynes is proposed and shown in Scheme 4. Firstly, the acyl radical **A** could be generated from the dissociation of aryl aldehydes or aryl ketones (**1**) under UV-light. Then, the acyl radical **A** attacked at 1,2-diphenylethyne (**2a**) to give the alkenyl radical intermediate **B**. Subsequently, this radical **B** attacked at the benzene ring via *ortho*-C—H bond activation to form the cyclohexadienyl radical **C**, which was oxidized by O₂^[16] from air to yield the product **3a**.

Scheme 4 Proposed reaction mechanism

3 Conclusions

In summary, the first UV-initiated radical C—H functionalization cyclization of aryl aldehydes or aryl ketones with alkynes for the preparation of indenones to avoid photocatalyst or photoinitiator has been demonstrated. This transformation could proceed smoothly featuring high efficiency, high atom economy, cheap reactants, environmental friendliness, easy operation, and tolerance to broad functional groups. Moreover, this work might not only broaden the research in constructing indenones, but also provide a fast-track strategy for atom/step economical synthesis of useful pharmaceutical molecules.

4 Experimental section

4.1 General experimental information

All manipulations were conducted under O₂ atmosphere. All commercial materials and solvents were used directly without further purification. UV-light (250 W, $\lambda_{\max}$ = 365 nm) was used for the light irradiation. NMR spectra (¹H, ¹³C, ¹⁹F) were recorded with a Bruker spectrometer (400 MHz), using CDCl₃ as the solvent and tetramethylsilane (TMS) as the internal standard at 298 K. Chemical shifts for ¹H NMR were relative to the signal of CHCl₃ at δ 7.26. Chemical shifts for ¹³C NMR were relative to the signal of CHCl₃ at δ 77.2. Column chromatography was performed on silica gel (70~230 mesh ASTM) using the reported eluents. Thin-layer chromatography (TLC) was carried out on 15 cm × 4 plates with a layer thickness of 0.2 mm (silica gel 60 F254).

4.2 Experimental method

To a 20 mL tube were added **1** (0.50 mmol) and **2** (0.10 mmol) and followed by addition of CH₃CN (1.5 mL). The formed mixture was stirred in 140 °C oil bath under UV-light ($\lambda_{\max}$ = 365 nm, 250 W) in O₂ for 48 h. The reaction was monitored by TLC. The solution was then quenched with H₂O (10 mL), extracted with ethyl acetate (10 mL × 3). The combined organic phases were washed with saturated sodium bicarbonate solution, dried over sodium sulfate, filtered, and evaporated under vacuum. The crude product was purified by column chromatography on silica gel [eluent: *V*(petroleum ether) : *V*(ethyl acetate) = 200 : 1 to 100 : 1] to afford the desired products **3**.

4.3 Characterization of the product

2,3-Diphenyl-1*H*-inden-1-one (3a): Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red solid, yield 82% (23.0 mg). m.p. 150~154 °C (Lit.^[17] 151~153 °C); ¹H NMR (400 MHz, CDCl₃) δ : 7.59 (d, *J* = 7.0 Hz, 1H), 7.46~7.34 (m, 6H), 7.32~7.24 (m, 6H), 7.15 (d, *J* = 7.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 196.7, 155.5, 145.4, 133.6, 132.9, 132.6, 130.9, 130.9, 130.2, 129.5, 129.2, 129.0, 128.7, 128.3, 127.9, 123.2, 121.5.

7-Methyl-2,3-diphenyl-1*H*-inden-1-one (3b):^[18] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 57% (17.0 mg). ¹H NMR (400 MHz, CDCl₃) δ : 7.43~7.33 (m, 5H), 7.29~7.22 (m, 6H), 7.05 (d, *J* = 7.8 Hz, 1H), 6.96 (d, *J* = 7.2 Hz, 1H), 2.61 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 197.8, 154.5, 145.8, 138.1, 133.1, 132.8, 132.6, 132.4, 131.1, 130.2, 129.2, 128.9, 128.8, 128.1, 127.7, 127.2, 119.4, 17.5.

6-methyl-2,3-diphenyl-1*H*-inden-1-one (3c):^[18] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 63% (18.6 mg). ¹H NMR (400 MHz, CDCl₃) δ : 7.41 (ddd, *J* = 10.0, 8.1, 5.8 Hz, 4H), 7.33~7.28 (m, 2H), 7.22~7.13 (m, 6H), 7.09 (d, *J* = 7.5 Hz, 1H), 1.79 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 197.1, 158.4, 142.2, 138.0, 138.0, 136.1, 133.8, 133.2, 131.0, 130.9, 130.0, 129.0, 128.6, 128.0, 128.0, 127.7, 121.1, 19.6.

5-Methyl-2,3-diphenyl-1*H*-inden-1-one (3d): Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red solid, Yield 84% (25.0 mg). m.p. 180~184 °C (Lit.^[19] 178~180 °C); ¹³C NMR (100 MHz, CDCl₃) δ : 196.4, 155.1, 145.9, 144.6, 133.1, 133.0, 131.1, 130.2, 129.3, 129.1, 129.0, 128.7, 128.6, 128.2, 127.8, 123.2, 122.7, 22.3; ¹H NMR (400 MHz, CDCl₃) δ : 7.47 (d, *J* = 7.3 Hz, 1H), 7.44~7.34 (m, 5H), 7.24 (d, *J* = 6.1 Hz, 5H), 7.07 (d, *J* = 7.3 Hz, 1H), 6.93 (s, 1H), 2.34 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 196.4, 155.1, 145.9, 144.6, 133.0, 133.0, 131.0, 130.1, 129.3, 129.1, 129.0, 128.7, 128.5, 128.2, 127.8, 123.2, 122.7, 22.3.

5-Fluoro-2,3-diphenyl-1*H*-inden-1-one (3e):^[20] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 70% (21.0 mg). ¹H NMR (400 MHz, CDCl₃) δ : 7.57 (dd, *J* = 7.9, 5.2 Hz, 1H), 7.45~7.33 (m, 5H), 7.26 (d, *J* = 2.3 Hz, 5H), 6.93 (ddd, *J* = 8.9, 8.0, 2.2 Hz, 1H), 6.87 (dd, *J* = 8.5, 2.1 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 195.0, 166.7 (d, *J* = 254.2 Hz), 153.3 (d, *J* = 2.3 Hz), 148.8 (d, *J* = 9.4 Hz), 133.9, 132.4, 130.6, 130.2, 129.7, 129.2, 128.6, 128.3, 128.2, 126.7 (d, *J* = 3.1 Hz), 125.0 (d, *J* = 9.8 Hz), 114.6 (d, *J* = 23.1 Hz), 110.4 (d, *J* = 25.8 Hz); ¹⁹F NMR (376 MHz, CDCl₃) δ : -104.0.

5-Chloro-2,3-diphenyl-1*H*-inden-1-one (3f): Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red solid, yield 68% (21.5 mg). m.p. 192~194 °C (Lit.^[21] 190~192 °C). ¹H NMR (400 MHz, CDCl₃) δ : 7.51 (d, *J* = 7.6 Hz, 1H), 7.43 (dd, *J* = 6.5, 3.7 Hz, 3H), 7.39~7.33 (m, 2H), 7.29~7.24 (m, 6H), 7.12 (d, *J* = 1.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 195.2, 154.2, 147.4, 139.9, 133.7, 132.3, 130.5, 130.2, 129.8, 129.2, 129.1, 128.7, 128.6, 128.3, 128.3, 124.1, 122.2.

5-Bromo-2,3-diphenyl-1*H*-inden-1-one (**3g**): Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red solid, yield 72% (26.0 mg). m.p. 168~174 °C (Lit.^[18] 167~168 °C). ¹H NMR (400 MHz, CDCl₃) δ: 7.51~7.40 (m, 5H), 7.39~7.33 (m, 2H), 7.27 (d, *J* = 4.6 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 195.4, 154.4, 147.4, 133.6, 132.3, 131.8, 130.4, 130.2, 129.8, 129.5, 129.2, 128.6, 128.5, 128.3, 124.9, 124.3.

5-Methoxy-2,3-diphenyl-1*H*-inden-1-one (**3h**): Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1). Orange-red solid, yield 80% (25.0 mg). m.p. 158~162 °C (Lit.^[18] 157~159 °C). ¹H NMR (400 MHz, CDCl₃) δ: 7.55 (d, *J* = 7.8 Hz, 1H), 7.44~7.32 (m, 5H), 7.28~7.22 (m, 5H), 6.68 (dt, *J* = 7.9, 2.1 Hz, 2H), 3.84 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 195.3, 164.6, 153.4, 148.1, 134.1, 132.9, 131.1, 130.2, 129.3, 129.0, 128.7, 128.2, 127.9, 125.1, 123.7, 110.7, 110.5, 55.9.

2,3-Diphenyl-5-(trifluoromethyl)-1*H*-inden-1-one (**3i**):^[20] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 77% (27.0 mg). ¹H NMR (400 MHz, CDCl₃) δ: 7.68 (d, *J* = 7.4 Hz, 1H), 7.61 (s, 1H), 7.46 (dd, *J* = 4.1, 2.4 Hz, 3H), 7.42~7.32 (m, 3H), 7.31~7.25 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ: 195.3, 154.9, 146.3, 135.3 (d, *J* = 32.3 Hz), 133.9, 133.6, 132.1, 130.3, 130.2, 130.0, 129.3, 128.5, 128.4, 128.4, 126.7 (q, *J* = 4.2 Hz), 123.8 (d, *J* = 272.9 Hz), 123.0, 117.9 (d, *J* = 3.6 Hz); ¹⁹F NMR (376 MHz, CDCl₃) δ: -63.1.

2,3-Bis(2-chlorophenyl)-1*H*-inden-1-one (**3j**):^[18] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 71% (25.0 mg). ¹H NMR (400 MHz, CDCl₃) δ: 7.60 (d, *J* = 6.6 Hz, 1H), 7.44 (d, *J* = 8.0 Hz, 1H), 7.34 (ddd, *J* = 26.2, 11.7, 4.5 Hz, 4H), 7.25~7.05 (m, 5H), 6.93 (d, *J* = 5.3 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 194.8, 156.4, 144.7, 134.4, 133.8, 133.8, 132.1, 131.7, 130.5, 130.5, 130.2, 130.1, 130.1, 129.7, 129.4, 129.4, 127.2, 126.6, 123.4, 123.4, 122.2.

2,3-Bis(4-chlorophenyl)-1*H*-inden-1-one (**3k**):^[18] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 74% (26.0 mg). ¹H NMR (400 MHz, CDCl₃) δ: 7.62~7.57 (m, 1H), 7.45~7.37 (m, 3H), 7.35~7.29 (m, 3H), 7.25 (dd, *J* = 4.4, 2.3 Hz, 2H), 7.22~7.16 (m, 2H), 7.12 (d, *J* = 7.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 196.0, 154.5, 144.8, 135.7, 134.2, 133.9, 133.9, 131.7, 131.4, 131.0, 130.7, 130.0, 129.6, 129.1, 128.8, 123.5, 121.4.

2,3-Bis(4-(trifluoromethyl)phenyl)-1*H*-inden-1-one (**3l**):^[18] Eluent: petroleum ether/ethyl acetate (*V* : *V* = 200 : 1), red oil, yield 78% (32.6 mg). ¹H NMR (400 MHz, CDCl₃) δ: 7.72 (d, *J* = 8.1 Hz, 2H), 7.64 (d, *J* = 6.6 Hz, 1H), 7.52 (dd, *J* = 17.0, 8.1 Hz, 4H), 7.43 (td, *J* = 7.7, 1.2 Hz, 1H), 7.39~7.33 (m, 3H), 7.12 (d, *J* = 7.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 195.5, 155.2, 144.5, 136.2, 134.1, 134.0, 131.7 (q, *J* = 32.0 Hz), 130.4, 130.4, 130.0, 129.0, 126.3 (q, *J* = 3.7 Hz), 125.4 (q, *J* = 3.8 Hz), 124.2 (q, *J* = 269.0 Hz), 123.9 (q, *J* = 271.0 Hz), 123.8, 121.7; ¹⁹F NMR (376 MHz, CDCl₃) δ: -62.8, -62.8.

Supporting Information ¹H NMR and ¹³C NMR spectra of compounds **3a**~**3l**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Huebner, C. F.; Donoghue, E. M.; Plummer, A. J.; Furness, D. A. *J. Med. Chem.* **1966**, *9*, 830.
(b) Jeffrey, J. L.; Sarpong, R. *Org. Lett.* **2009**, *11*, 5450.
(c) Barbera, J.; Rakitin, O. A.; Ros, M. B.; Torroba, T. *Angew. Chem., Int. Ed.* **1998**, *37*, 296.
(d) Morinaka, K.; Ubukata, T.; Yokoyama, Y. *Org. Lett.* **2009**, *11*, 3890.
(e) Anstead, G. M.; Wilson, S. R.; Katzenellenbogen, J. A. *J. Med. Chem.* **1989**, *32*, 2163.
(f) Froimowitz, M.; Wu, K. M.; Moussa, A.; Haidar, R. M.; Jurayj, J.; George, C.; Gardner, E. L. *J. Med. Chem.* **2000**, *43*, 4981.
(g) Gao, K.; Yoshikai, N. *Chem. Commun.* **2012**, *48*, 4305.
(h) Kunitobu, Y.; Matsuki, T.; Takai, K. *Org. Lett.* **2012**, *12*, 2948.
(i) Vasilyev, A. V.; Walspurger, S.; Pale, P.; Sommer, J. *Tetrahedron Lett.* **2004**, *45*, 3379.
(j) Sheng, Y.; You, Y.; Cao, Z.; Liu, L.; Wu, H. C. *Analyst* **2018**, *143*, 2411.
- [2] (a) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174.
(b) Lyons, T. W.; Sanford, M. S. *Chem. Rev.* **2010**, *110*, 1147.
(c) Giri, R.; Shi, B.; Engle, K. M.; Maugel, N.; Yu, J. *Chem. Soc. Rev.* **2009**, *38*, 3242.
(d) McMurray, L.; O'Hara, F.; Gaunt, M. J. *Chem. Soc. Rev.* **2011**, *40*, 1885.
(e) Ylijoki, K. E. O.; Stryker, J. M. *Chem. Rev.* **2013**, *113*, 2244.
(f) Wang, Z.; Xie, P.; Xia, X. *Chin. Chem. Lett.* **2018**, *29*, 47.
(g) Zhang, Y.; Sun, K.; Lv, K.; Chen, X.; Qu, L.; Yu, B. *Chin. Chem. Lett.* **2019**, *30*, 1361.
(h) Suo, J.; Du, J.; Jiang, Y.; Chen, D.; Ding, C.; Hou, X. *Chin. Chem. Lett.* **2019**, *30*, 1512.
(i) Pan, Y.; Chen, G.; Shen, C.; He, W.; Ye, L. *Org. Chem. Front.* **2016**, *3*, 491.
(j) Li, W.; Yin, G.; Huang, L.; Xiao, Y.; Fu, Z.; Xin, L.; Liu, F.; Li, Z.; He, W. *Green Chem.* **2016**, *18*, 4879.
(k) Gao, Q.; Hao, W.; Liu, F.; Tu, S.; Wang, S.; Li, G.; Jiang, B. *Chem. Commun.* **2016**, *52*, 900.
(l) Zhou, P.; Wang, J.; Zhang, T.; Li, G.; Hao, W.; Tu, S.; Jiang, B. *Chem. Commun.* **2018**, *54*, 164.
(m) Zhu, S.; Zhou, J.; Wu, Q.; Hao, W.; Tu, S.; Jiang, B. *Org. Chem. Front.* **2020**, *7*, 2975.
(n) Tian, S.; Luo, T.; Zhu, Y.; Wang, J. *Chin. Chem. Lett.* **2020**, *31*, 3072.
(o) Butenschön, H. *Angew. Chem., Int. Ed.* **2008**, *47*, 5287.
(p) Harmata, M. *Chem. Commun.* **2010**, *46*, 8886.
(q) Lv, N.; Chen, Z.; Liu, Y.; Liu, Z.; Zhang, Y. *Org. Lett.* **2017**, *19*, 2588.
(r) Yu, W.; Zhang, W.; Liu, Z.; Zhang, Y. *Chem. Commun.* **2016**, *52*, 6837.
- [3] Chen, S.; Yu, J.; Jiang, Y.; Chen, F.; Cheng, J. *Org. Lett.* **2013**, *15*, 4754.
- [4] Banerji, B.; Majumder, L.; Adhikary, S. *ChemistrySelect* **2018**, *3*, 1381.
- [5] Feng, Y.; Zhang, H.; Yu, Y.; Yang, L.; Cui, X. *Eur. J. Org. Chem.* **2019**, *16*, 2740.
- [6] (a) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, *113*, 5322.
(b) Romero, N. A.; Nicewicz, D. A. *Chem. Rev.* **2016**, *116*, 10075.
(c) Ravelli, D.; Fagnoni, M.; Albin, A. *Chem. Soc. Rev.* **2013**, *42*, 97.
- [7] Nagib, D. A.; MacMillan, D. W. C. *Nature* **2011**, *480*, 224.

- [8] Kalyani, D.; McMurtrey, K. B.; Neufeldt, S. R.; Sanford, M. *J. Am. Chem. Soc.* **2011**, *133*, 18566.
- [9] (a) Tobisu, M.; Furukawa, T.; Chatani, N. *Chem. Lett.* **2013**, *42*, 1203.
(b) Gomes, F.; Narbonne, V.; Blanchard, F.; Maestri, G. *Org. Chem. Front.* **2015**, *2*, 464.
(c) Liang, L.; Xie, M.; Wang, H.; Niu, H.; Qu, G.; Guo, H. *J. Org. Chem.* **2017**, *82*, 966.
(d) Xue, D.; Jia, Z.; Zhao, C.; Zhang, Y.; Wang, C.; Xiao, J. *Chem.-Eur. J.* **2014**, *20*, 2960.
(e) Zhang, J.; Chen, J.; Zhang, X.; Lei, X. *J. Org. Chem.* **2014**, *79*, 10682.
(f) Xu, H.; Jiang, Z. *Chin. J. Org. Chem.* **2020**, *40*, 3483 (in Chinese).
(许荷欢, 江智勇, 有机化学, **2020**, *40*, 3483.)
(g) Jin, J.-K.; Xia, H.-M.; Zhang, F.-L.; Wang, Y.-F. *Chin. J. Org. Chem.* **2020**, *40*, 2185 (in Chinese).
(靳继康, 夏慧敏, 张凤莲, 汪义丰, 有机化学, **2020**, *40*, 2185.)
- [10] Hari, D. P.; Schroll, P.; König, B. *J. Am. Chem. Soc.* **2012**, *134*, 2958.
- [11] (a) Huang, L.; Zhao, J. *RSC Adv.* **2013**, *3*, 23377.
(b) Marzo, L.; Ghosh, I.; Esteban, F.; Konig, B. *ACS Catal.* **2016**, *6*, 6780.
(c) Feng, Y.; Bu, X.; Huang, B.; Rong, C.; Dai, J.; Xu, J.; Xu, H. *Tetrahedron Lett.* **2017**, *58*, 1939.
(d) Kojima, M.; Oisaki, K.; Kanai, M. *Chem. Commun.* **2015**, *51*, 9718.
(e) Poelma, S. O.; Burnett, G. L.; Discekici, E. H.; Mattson, K. M.; Treat, N. J.; Luo, Y.; Hudson, Z. M.; Shankel, S. L.; Clark, P. G.; Kramer, J. W.; Hawker, C. J.; Read de Alaniz, J. *J. Org. Chem.* **2016**, *81*, 7155.
(f) Cui, L.; Matusaki, Y.; Tada, N.; Miura, T.; Uno, B.; Itoh, A. *Adv. Synth. Catal.* **2013**, *355*, 2203.
- [12] Vitale, A.; Bongiovanni, R.; Ameduri, B. *Chem. Rev.* **2015**, *115*, 8835.
- [13] (a) Feng, Y.; Zhang, Z.; Fu, Q.; Yao, Q.; Huang, H.; Shen, J.; Cui, X. *Chin. Chem. Lett.* **2020**, *31*, 58.
(b) Feng, Y.; Liu, Y.; Fu, Q.; Zou, Z.; Shen, J.; Cui, X. *Chin. Chem. Lett.* **2020**, *31*, 733.
(c) Feng, Y.; Wu, Z.; Chen, T.; Fu, Q.; You, Q.; Shen, J.; Cui, X. *Chin. Chem. Lett.* **2020**, *31*, 3263.
(d) Li, T.; Yang, Z.; Song, Z.; Chauvin, R.; Cui, X. *Org. Lett.* **2020**, *22*, 4078.
(e) Liu, Y.; Yang, Z.; Chauvin, R.; Fu, W.; Yao, Z.; Wang, L.; Cui, X. *Org. Lett.* **2020**, *22*, 5140.
(f) Yao, Z.; Lin, X.; Chauvin, R.; Wang, L.; Gras, E.; Cui, X. *Chin. Chem. Lett.* **2020**, *31*, 3250.
(g) Song, Z.; Yang, Z.; Wang, P.; Shi, Z.; Li, T.; Cui, X. *Org. Lett.* **2020**, *22*, 6272.
(h) Wu, Y.; Pi, C.; Cui, X.; Wu, Y. *Org. Lett.* **2020**, *22*, 361.
(i) Ren, J.; Yan, X.; Cui, X.; Pi, C.; Wu, Y.; Cui, X. *Green Chem.* **2020**, *22*, 265.
(j) He, Y.; Pi, C.; Wu, Y.; Cui, X. *Chin. Chem. Lett.* **2020**, *31*, 396.
(k) Huang, Y.; Pi, C.; Cui, X.; Wu, Y. *Adv. Synth. Catal.* **2020**, *362*, 1385.
(l) Wan, T.; Pi, C.; Wu, Y.; Cui, X. *Org. Lett.* **2020**, *22*, 6484.
(m) Yu, H.; Pi, C.; Wang, Y.; Cui, X.; Wu, Y. *Chin. J. Org. Chem.* **2018**, *38*, 124 (in Chinese).
- (余海洋, 皮超, 王勇, 崔秀灵, 吴养洁, 有机化学, **2018**, *38*, 124.)
(n) Shi, Z.-J.; Wang, L.-H.; Cui, X. *Chin. J. Org. Chem.* **2019**, *39*, 1596 (in Chinese).
(施兆江, 王连会, 崔秀灵, 有机化学, **2019**, *39*, 1596.)
- [14] (a) Li, Z.; Li, D.; Wang, G. *J. Org. Chem.* **2013**, *78*, 10414.
(b) Li, H.; Li, P.; Wang, L. *Chem. Commun.* **2013**, *49*, 9170.
(c) Li, H.; Li, P.; Wang, L. *Chem.-Eur. J.* **2013**, *19*, 14432.
(d) Wang, G.; Shang, R.; Cheng, W.; Fu, Y. *Org. Lett.* **2015**, *17*, 4830.
(e) Ji, W.; Tan, H.; Wang, M.; Li, P.; Wang, L. *Chem. Commun.* **2016**, *52*, 1462.
(f) Li, C.; Xu, G.; Xu, P. *Org. Lett.* **2017**, *19*, 512.
(g) Xu, S.; Chen, J.; Liu, D.; Bao, Y.; Liang, Y.; Xu, P. *Org. Chem. Front.* **2017**, *4*, 1331.
(h) Liu, Y.; Wang, Q.; Zhou, C.; Xiong, B.; Zhang, P.; Kang, S.; Yang, C.; Tang, K. *Tetrahedron Lett.* **2018**, *59*, 2038.
- [15] (a) Sharma, U. K.; Gemoets, H. P. L.; Schroder, F.; Noel, T.; Eycken, E. V. *ACS Catal.* **2017**, *7*, 3818.
(b) Zhang, L.; Zhang, G.; Li, Y.; Wang, S.; Lei, A. *Chem. Commun.* **2018**, *54*, 5744.
(c) Chatgililoglu, C.; Crich, D.; Komatsu, M.; Ryu, I. *Chem. Rev.* **1999**, *99*, 1991.
(d) Yu, X.; Chen, J.; Chen, H.; Xiao, W.; Chen, J. *Org. Lett.* **2020**, *22*, 2333.
(e) Zhao, Q.; Zhou, X.; Xu, S.; Wu, Y.; Xiao, W.; Chen, J. *Org. Lett.* **2020**, *22*, 2470.
(f) Liang, D.; Tan, L.; Xiao, W.; Chen, J. *Chem. Commun.* **2020**, *56*, 3777.
(g) Leifert, D.; Daniliuc, C. G.; Studer, A. *Org. Lett.* **2013**, *15*, 6286.
(h) Wertz, S.; Leifert, D.; Studer, A. *Org. Lett.* **2013**, *15*, 928.
- [16] (a) Yi, R.; He, W. *Chin. J. Org. Chem.* **2021**, *41*, 1267 (in Chinese).
(易荣楠, 何卫民, 有机化学, **2021**, *41*, 1267.)
(b) Liu, R.; Yang, M.; Qiu, G.; Zhang, L.; Wang, Y.; Luo, J. *Chin. J. Org. Chem.* **2020**, *40*, 2071 (in Chinese).
(刘仁志, 杨民, 邱观音生, 张莲鹏, 王玉超, 罗劲, 有机化学, **2020**, *40*, 2071.)
(c) Peng, S.; Lin, Y.; He, W. *Chin. J. Org. Chem.* **2020**, *40*, 541 (in Chinese).
(彭莎, 林英武, 何卫民, 有机化学, **2020**, *40*, 541.)
(d) Yuan, X.; Yang, G.; Yu, B. *Chin. J. Org. Chem.* **2020**, *40*, 3620 (in Chinese).
(袁晓亚, 杨国平, 於兵, 有机化学, **2020**, *40*, 3620.)
(e) Wu, Y.; Chen, J.; Ning, J.; Jiang, X.; Deng, J.; Deng, Y.; Xu, R.; He, W. *Green Chem.* **2021**, *23*, 3950.
(f) Jiang, J.; Xiao, F.; He, W.; Wang, L. *Chin. Chem. Lett.* **2021**, *32*, 1637.
(g) He, W.; Gao, L.; Chen, X.; Wu, Z.; Huang, Y.; Cao, Z.; Xu, X.; He, W. *Chin. Chem. Lett.* **2020**, *31*, 1895.
- [17] Tsukamoto, H.; Kondo, Y. *Org. Lett.* **2007**, *21*, 4227.
- [18] Feng, Y.; Zhang, H.; Yu, Y.; Yang, L.; Cui, X. *Eur. J. Org. Chem.* **2019**, *16*, 2740.
- [19] Zhang, J.; Yang, F.; Wu, Y. *Appl. Organomet. Chem.* **2011**, *25*, 675.
- [20] Jafarpour, F.; Azizzade, M.; Golpazir-Sorkheh, Y.; Navid, H.; Rajai-Daryasarei, S. *J. Org. Chem.* **2020**, *85*, 8287.
- [21] Curtin, D. Y.; Johnson, H. W., Jr.; Steiner, E. G. *J. Am. Chem. Soc.* **1955**, *77*, 4566.

(Cheng, F.)