1,4-二氮杂二环[2.2.2]辛烷促进三酮单水合物和活泼炔通过连续碳-碳/碳-氧键断裂的加成反应

张金凤 叶如意 李子阳 孟令国*

(淮北师范大学化学与材料科学学院 安徽淮北 235000)

摘要 在无金属和温和反应条件下, 发展了新颖的三酮单水合物与活泼炔通过连续碳-碳/碳-氧键断裂的加成反应, 且反应可高效地在 1 min 内完成. 多样的三酮单水合物均可在 1,4-二氮杂二环[2.2.2]辛烷(DABCO)的促进下, 通过这种转变与活泼炔反应, 生成多种酯类化合物.

关键词 碳-碳键断裂; 碳-氧键断裂; 加成; 促进

1,4-Diazabicyclo[2.2.2]octane-Promoted Addition Reaction of Tricarbonyl Monohydrates with Active Alkynes via Consecutive C—C/C—O Cleavage

Zhang, Jinfeng Ye, Ruyi Li, Ziyang Meng, Lingguo*

(School of Chemistry and Materials Science, Huaibei Normal University, Huaibei, Anhui 235000)

Abstract A novel addition reaction of vicinal tricarbonyl monohydrates with active alkynes proceeded via rapid consecutive C—C/C—O cleavage processes under metal-free and mild reaction conditions, and the transformation was completed within only 1 min. Diverse tricarbonyl monohydrates could react with active alkynes to produce carboxylate esters via these transformations in the presence of 1,4-diazabicyclo[2.2.2]octane (DABCO).

Keywords C—C cleavage; C—O cleavage; addition; promoter

1 Introduction

Chemical bond cleavage has emerged as an important means of bond formation, is fundamental research areas in organic synthesis, and has proven to be an efficient method for using molecular rearrangement to construct many organic compounds.^[1] Among the various types of chemical bond cleavages, C—C bond cleavage has received more attention in recent years as a direct means of constructing novel molecular skeletons.^[2] As a result, efficient promotion of C—C bond cleavage has attracted considerable interest, and several methods for this transformation have been devised. The mainstream method employed to date to achieve C—C bond cleavage is the use of transitional metals as assistants,^[3] such as palladium,^[4] rhodium^[5] and copper^[6] *etc.* However, the difficulty of removing heavy metals and releasing environmental waste has brought problems to the pharmaceutical industry and has limited the

applications of heavy metals in organic synthesis. Recently, C—C bond cleavage was achieved in the absence of heavy metals by using an oxidative system^[7] or photoassistance conditions^[8]. Despite the significant progress that has been made in this field, finding a simple method to promote C—C bond cleavage remains a significant challenge.

Vicinal triketones and their monohydrates are raw materials that can be easily obtained and employed to achieve diverse synthetic transformations via the C—C bond cleavage process. For example, Schofield *et al.*^[9a] reported the C—C cleavage of vicinal tricarbonyl compounds in the presence of iron salts (Scheme 1a). A similar chemical bond cleavage process can also be realized in the presence of nickel salts,^[9b] phenyl magnesium bromide,^[10] selenium^[11] or aluminium chloride.^[12] Alternative methods, such as the use of I₂,^[13] peroxide^[14] and irradiation conditions^[15] are also powerful tactics for inducing the cleavage process. Note that all the above mentioned reported methods have

* Corresponding author. E-mail: milig@126.com

Received April 8, 2021; revised June 1, 2021; published online August 17, 2021.

Project supported by the Natural Science Foundation of Educational Committee of Anhui Province (Nos. KJ2020A0045, KJ2020B01).

安徽高校自然科学基金(Nos. KJ2020A0045, KJ2020B01)资助项目.

Scheme 1 Chemical bond cleavage reaction of tricarbonyl monohydrate

drawbacks, such as expensive reagents, heavy metal pollution, harsh conditions, or long reaction times. Therefore, a facile and more efficient method for the C—C bond cleavage of triketones or their derivatives is highly desirable.

Tertiary amine or phosphine catalyzed reactions based on active alkynes have become an active research area.^[16] However, few studies have been performed on chemical bond cleavage in this field.^[17] In recent years, we have developed efficient methods for the direct cleavage of C—N and N—N bonds by using tertiary amines as promoters.^[18] In continuation of our interest in applying this synthetic strategy to novel bond cleavage reactions, we further investigated the potential reaction between vicinal tricarbonyl monohydrates with alkynyl ketones in the presence of a tertiary amine. An addition reaction of vicinal tricarbonyl monohydrates with alkynyl ketones via consecutive C—C/C—O cleavage promoted by 1,4-diazabicyclo[2.2.2]octane (DABCO) was found. A series of carboxylate esters were obtained under the above mentioned mild conditions (Scheme 1b). The versatile reactivity of carboxylate esters has been utilized to synthesize many compounds.^[19] Herein, we report an alternative method for the promotion of C—C bond cleavage of vicinal tricarbonyl monohydrates in the presence of DABCO and alkynyl ketones.

2 Results and discussion

Our initial investigation focused on the optimization of the conditions for the reaction of 1,3-bis(4-chlorophenyl)-

2,2-dihydroxypropane-1,3-dione (**1a**, 0.20 mmol) with 1-phenylprop-2-yn-1-one (**2a**, 0.40 mmol) using 0.20 equiv. of DABCO in CH₂Cl₂ at room temperature under stirring for 30 min (Table 1). Surprisingly, a rearrangement reaction was found between **1a** and **2a** via consecutive C—C and C—O bond cleavage to afford an unexpected carboxylate ester product, (*E*)-3-oxo-3-phenylprop-1-en-1-yl 4-chlorobenzoate in 25% isolated yield (Table 1, Entry 1). Efforts were then made to improve the yield of this novel reaction under different conditions. The yield was improved by increasing the quantity of DABCO, where 0.50 equiv. of DABCO based on vicinal tricarbonyl monohydrate was found to be optimal. The yield of the product was decreased when 1.0 equiv. of DABCO was used (Table 1, Entries 2~4). Unsatisfactory results were obtained by changing the reaction temperature (Table 1, Entries 5 and 6). The results presented in Table 1 show that the choice of base for the reaction was clearly important. Among a wide range of candidates, DABCO was found to be a powerful base for promoting the reaction. Lower yields were observed upon replacing DABCO with Et₃N, pyridine or DBU. The desired product was not formed when other bases, such as DMAP, HMTA, PPh₃, K₂CO₃ and KOBu^t were added to the reaction. The effect of the solvent on the reaction was also examined, and 12% and 17% yields of the desired products were obtained using CHCl₃ and THF as solvents, respectively. When the solvent was switched to CCl₄, CH₃CN, CS₂, toluene or EtOAc, only a trace amount of product was observed by thin layer chromatography analysis. Using *N,N*-dimethylformamide (DMF) as a solvent afforded no product and was therefore not suitable for the reaction.

Table 1 Optimization of reaction conditions^a

Entry	Solvent	Base	Yield ^b
1	CH ₂ Cl ₂	DABCO	25 ^c
2	CH ₂ Cl ₂	DABCO	36 ^d
3	CH ₂ Cl ₂	DABCO	45
4	CH ₂ Cl ₂	DABCO	18 ^e
5	CH ₂ Cl ₂	DABCO	<10 ^f
6	CH ₂ Cl ₂	DABCO	Complex ^g
7	CH ₂ Cl ₂	Et ₃ N	13
8	CH ₂ Cl ₂	Pyridine	Trace
9	CH ₂ Cl ₂	DBU	Trace
10	CH ₂ Cl ₂	DMAP	ND ^h
11	CH ₂ Cl ₂	HMTA	NR ⁱ
12	CH ₂ Cl ₂	PPh ₃	ND ^h
13	CH ₂ Cl ₂	K ₂ CO ₃	NR ⁱ
14	CH ₂ Cl ₂	KOBu ^t	NR ⁱ
15	CHCl ₃	DABCO	17
16	THF	DABCO	12
17	CCl ₄	DABCO	Trace
18	CH ₃ CN	DABCO	Trace
19	CS ₂	DABCO	Trace
20	Toluene	DABCO	Trace
21	EtOAc	DABCO	Trace
22	DMF	DABCO	ND ^h

^a **1a** (0.20 mmol), **2a** (0.40 mmol), base (0.10 mmol), solvent (1.5 mL), r.t., in air, 30 min; ^b Isolated yield; ^c DABCO (0.20 equiv.); ^d DABCO (0.30 equiv.); ^e DABCO (1.0 equiv.); ^f At 0 °C; ^g At reflux; ^h ND=No desired product detected; ⁱ NR=No reaction observed. DABCO=1,4-diazabicyclo[2.2.2]octane, DBU=1,8-diazabicyclo[5.4.0]undec-7-ene, DMAP=4-dimethylaminopyridine, HMTA=hexamethylene tetramine.

Having identified the optimal reaction conditions, the proposed approach was applied to the consecutive cleavage of chemical bonds in different symmetrical vicinal tricarbonyl monohydrates (**1**) by reaction with 1-phenylprop-2-yn-1-one (**2a**), and the results are shown in Table 2. A variety of aromatic tricarbonyl monohydrates were found to react smoothly with **2a** to produce the desired carboxylate esters in moderate to good yields. Aromatic tricarbonyl monohydrates with an electron-donating group on the aromatic ring afforded clearly higher yields than those with an electron-withdrawing group on the aromatic ring. For example, substrate **1** with an electron-donating group, such as CH₃ or CH₃O at the *para*-position of the benzene ring, reacted with **2a** to provide the desired products **3ba** and **3ca** in 85% and 65% yields, respectively. By comparison, **1** with an electron-withdrawing group, such as F, Cl or Br at the *para*-position of the benzene ring, afforded the corresponding products in 41%~46% yields. The *ortho*-substituted

tricarbonyl monohydrate was tolerated, and the corresponding product **3ga** was obtained in 70% yield. Treatment of 1,3-bis(3-bromophenyl)-2,2-dihydroxypropane-1,3-dione (**1h**) with **2a** afforded the desired product **3ha** in 72% yield. Notably, a heteroaromatic symmetrical vicinal tricarbonyl monohydrate could also be converted to desired product (**3ia**) in moderate yield. Note that all the reactions presented in Table 2 were efficiently completed within 1 to 30 min.

Table 2 DABCO-promoted addition reaction of symmetrical vicinal tricarbonyl monohydrates **1** with **2a**^a

^a Reaction conditions: **1** (0.20 mmol), **2a** (0.40 mmol), DABCO (0.10 mmol), CH₂Cl₂ (1.5 mL), r. t., in air, 1~30 min. ^b Isolated yield. ^c The reaction was stirred for 30 min. ^d The reaction was stirred for 1 min.

Subsequently, other alkynyl ketones were employed in the reaction and the results are shown in Table 3. It was gratifying to find that a variety of substituted alkynyl ketones rapidly reacted with diverse tricarbonyl monohydrates in satisfactory yields regardless of the different substitutions on the aromatic rings of **2**. Substrates with CH₃O group on the benzene ring produce higher yields than those with F or Cl group on the benzene ring (**3bb** vs **3bc**, **3bd**). There was no discernible product formation using 1-(4-nitrophenyl)-prop-2-yn-1-one (**2e**) as the substrate. 1-(Naphthalen-2-yl)-prop-2-yn-1-one was also converted to the desired product

(**3bf**) in excellent yield. Furthermore, substrates **1g** and **1d**, also reacted with diverse alkynyl ketones to generate the corresponding products (**3gg** ~ **gi**, **3db**) in 45% ~ 71% yields. However, other electron-deficient olefins or alkynes, such as ethyl propiolate, propiolamide and chalcone, failed to react with the tricarbonyl monohydrates to afford the desired addition product.

Table 2 Scope of alkynyl ketones^a

^a Reaction conditions: **1** (0.20 mmol), **2** (0.40 mmol), DABCO (0.10 mmol), CH₂Cl₂ (1.5 mL), r.t., in air, 10 min. ^b Isolated yield. ^c The reaction was stirred for 30 min. ^d The reaction was stirred for 1 min.

Last, to address the regioselectivity of the reaction, a series of intramolecular competition reactions were explored (Scheme 2). Among the unsymmetrical vicinal tricarbonyl monohydrates, 1-(4-chlorophenyl)-2,2-dihydroxy-3-(4-methoxyphenyl)propane-1,3-dione reacted with **2a**, affording

a mixture of **3aa** (26% yield) and **3ca** (20% yield). Note that the reactions of 1,3-aromatic/aliphatic and 1,3-aromatic/ester tricarbonyl monohydrates, such as 2,2-dihydroxy-1-phenylbutane-1,3-dione and ethyl 2,2-dihydroxy-3-oxo-3-phenylpropanoate, exhibited excellent selectivity, although the corresponding product yields were low. Disappointingly, the addition of 2,2-dihydroxy-1*H*-indene-1,3(2*H*)dione (**1m**) made the reaction complex, with a poor result. Notably, impure raw material **1d**, which contained 6/7 tricarbonyl form **1d'**, could also be converted to expected product **3da** in 46% yield.

Table 3 DABCO-promoted the reactions of other vicinal tricarbonyl monohydrates **1** with **2a**^a

^a Reaction conditions: **1** (0.20 mmol), **2a** (0.40 mmol), DABCO (0.10 mmol), CH₂Cl₂ (1.5 mL), r.t., in air, 30 min. ^b Isolated yields. ^c Containing 1/3 tricarbonyl form. ^d Containing 6/7 tricarbonyl form. ^e ND=No desired product was detected.

Based on the above mentioned observations, the mechanism depicted in Scheme 3 was proposed. The reaction was triggered by the reaction of DABCO with **2a** to produce zwitterion **A**, which deprotonated 2,2-dihydroxy-1,3-diphenylpropane-1,3-dione (**1d**) to generate **B** and **C** (Scheme 3, path 1). The C—C bond cleavage reaction also proceeded using an equilibrium mixture of **1d** and **1d'** as substrates, and intermediate **C** could also be obtained by another process (Scheme 3, path 2), whereby **1d** was easily converted into intermediate triketone **1d'** by losing H₂O, which was

Scheme 3 Plausible mechanism

deprotonated by **A** to afford a hydroxide anion. The hydroxide anion reacted with **1d'** through a nucleophilic addition to produce the same intermediate **C** as path 1. It should be noted that we do not know which path (path 1 or path 2) was taken to produce **C** due to the fast intermolecular transformation between **1d** and **1d'**. Subsequently, intermediate **C** underwent an intramolecular nucleophilic attack of the oxygen anion to the vicinal carbonyl to generate **D**, followed by a C—C bond cleavage process to give **E**. Finally, intermediate **E** underwent proton transfer and C—O cleavage to form a benzoate anion, which reacted with **B** to afford the desired product **3da** by Michael addition and elimination of DABCO. Note, however, that the concomitant generative product 2-oxo-2-phenylacetaldehyde **G** or its monohydrate **G'** could not be obtained from the reaction system, which can be explained using the results of a control experiment shown in Scheme 4.

Scheme 4 Control experiments

3 Conclusions

In summary, a novel and interesting addition reaction was reported via rapid consecutive C—C and C—O bonds cleavage processes promoted by DABCO under metal-free conditions. Note that this direct C—C bond cleavage constitutes a novel protocol for the construction of molecular skeletons, which could be very valuable and efficient in organic chemistry.

4 Experimental section

4.1 Instruments and reagents

Chemicals and analytical grade solvents were purchased from commercial suppliers and used without further purification unless otherwise stated. Chromatographic purifica-

tion was performed on silica gel (100~200 mesh) and analytical thin layer chromatography (TLC) on silica gel 60-F₂₅₄ (Qindao), which was detected by fluorescence. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra were measured on a Bruker AC 400 spectrometer with CDCl₃ as solvent and recorded relative to internal tetramethylsilane standard. High resolution mass spectra (HRMS) were obtained with a Micromass GCT-TOF mass spectrometer. Melting points were determined on a digital melting point apparatus and temperatures were uncorrected.

4.2 General procedure for the synthesis of 3

To a solution of tricarbonyl monohydrates (0.2 mmol) with active alkynes (0.4 mmol) in CH₂Cl₂ (1.5 mL) was added DABCO (0.1 mmol). The mixture was stirred at room temperature for approximately 1 min in a reaction flask. The solvent was removed *in vacuo* and residue was purified by column chromatography on silica gel to afford the desired products.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl 4-chlorobenzoate (**3aa**): Pale yellow solid. m.p. 90~92 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.64 (d, *J*=12.4 Hz, 1H), 8.11 (d, *J*=7.6 Hz, 2H), 7.97 (d, *J*=7.6 Hz, 2H), 7.62 (t, *J*=7.6 Hz, 1H), 7.51~7.50 (m, 4H), 7.01 (d, *J*=12.4 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 190.0, 161.8, 150.2, 140.9, 137.7, 132.9, 131.5, 129.1, 128.5, 128.2, 126.1, 110.6. HRMS (EI) calcd for C₁₆H₁₁ClO₃ [M]⁺: 286.0397; found 286.0401.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl 4-methylbenzoate (**3ba**): Pale yellow solid. m.p. 85~87 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.67 (d, *J*=12.4 Hz, 1H), 8.06 (d, *J*=7.6 Hz, 2H), 7.97 (d, *J*=7.6 Hz, 2H), 7.61 (t, *J*=7.2 Hz, 1H), 7.52 (t, *J*=7.2 Hz, 2H), 7.33 (d, *J*=7.6 Hz, 2H), 6.98 (d, *J*=12.4 Hz, 1H), 2.45 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 190.3, 162.6, 150.7, 145.3, 137.8, 132.8, 130.3, 129.4, 128.5, 128.2, 124.9, 110.1, 21.6. HRMS (EI) calcd for C₁₇H₁₄O₃ [M]⁺: 266.0943; found 266.0947.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl 4-methoxybenzoate (**3ca**): Pale yellow solid. m.p. 126~128 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.67 (d, *J*=12.4 Hz, 1H), 8.13 (d, *J*=8.8 Hz, 2H), 7.97 (d, *J*=7.2 Hz, 2H), 7.61 (t, *J*=7.2 Hz, 1H), 7.52 (t, *J*=7.6 Hz, 2H), 7.01 (d, *J*=8.8 Hz, 2H), 6.97 (d, *J*=12.4 Hz, 1H), 3.91 (s, 3H); ¹³C NMR (100 MHz, CDCl₃)

δ : 190.4, 164.4, 162.2, 150.9, 137.9, 132.7, 132.5, 128.5, 128.2, 119.8, 114.0, 109.9, 55.4. HRMS (EI) calcd for $C_{17}H_{14}O_4$ $[M]^+$: 282.0892; found 282.0890.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl benzoate (**3da**): White solid. m.p. 73~75 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.68 (d, $J=12.0$ Hz, 1H), 8.18 (d, $J=8.0$ Hz, 2H), 7.98 (d, $J=7.6$ Hz, 2H), 7.69 (t, $J=7.2$ Hz, 1H), 7.62 (t, $J=7.2$ Hz, 1H), 7.55~7.49 (m, 4H), 7.01 (d, $J=12.0$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 190.2, 162.6, 150.5, 137.7, 134.3, 132.9, 130.2, 128.6, 128.5, 128.2, 127.6, 110.3. HRMS (EI) calcd for $C_{16}H_{12}O_3$ $[M]^+$: 252.0786; found 252.0790.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl 4-fluorobenzoate (**3ea**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.65 (d, $J=12.4$ Hz, 1H), 8.22 (dd, $J=5.6, 8.0$ Hz, 2H), 7.97 (d, $J=7.2$ Hz, 2H), 7.62 (t, $J=7.2$ Hz, 1H), 7.53 (t, $J=7.6$ Hz, 2H), 7.23 (t, $J=8.4$ Hz, 2H), 7.01 (d, $J=12.4$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 190.1, 167.7 (d, $J=255.3$ Hz), 161.6, 150.3, 137.7, 133.0, 132.9, 128.5, 128.2 (d, $J=4.1$ Hz), 124.0 (d, $J=3.0$ Hz), 116.1 (d, $J=22.1$ Hz), 110.5. HRMS (EI) calcd for $C_{16}H_{11}FO_3$ $[M]^+$: 270.0692; found 270.0691.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl 4-bromobenzoate (**3fa**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.64 (d, $J=12.4$ Hz, 1H), 8.03 (d, $J=8.4$ Hz, 2H), 7.97 (d, $J=7.2$ Hz, 2H), 7.69 (d, $J=8.4$ Hz, 2H), 7.62 (t, $J=7.6$ Hz, 1H), 7.53 (t, $J=7.6$ Hz, 2H), 7.01 (d, $J=12.4$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 190.0, 162.0, 150.2, 137.6, 132.9, 132.1, 131.6, 129.7, 128.5, 128.2, 126.6, 110.6. HRMS (EI) calcd for $C_{16}H_{11}BrO_3$ $[M]^+$: 329.9892; found 329.9897.

(*E*)-3-oxo-3-Phenylprop-1-en-1-yl 2-methylbenzoate (**3ga**): Yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.68 (d, $J=12.0$ Hz, 1H), 8.11 (d, $J=7.6$ Hz, 1H), 7.97 (d, $J=7.2$ Hz, 2H), 7.61 (t, $J=7.2$ Hz, 1H), 7.52~7.49 (m, 3H), 7.35~7.32 (m, 2H), 6.97 (d, $J=12.0$ Hz, 1H), 2.68 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 190.3, 162.7, 150.6, 142.0, 137.8, 133.4, 132.8, 132.0, 131.1, 128.5, 128.2, 126.6, 125.9, 110.1, 21.8. HRMS (EI) calcd for $C_{17}H_{14}O_3$ $[M]^+$: 266.0943; found 266.0946.

(*E*)-3-oxo-3-phenylprop-1-en-1-yl 3-bromobenzoate (**3ha**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.64 (d, $J=12.0$ Hz, 1H), 8.30 (s, 1H), 8.11 (d, $J=7.6$ Hz, 1H), 7.98 (d, $J=7.2$ Hz, 2H), 7.81 (d, $J=8.0$ Hz, 1H), 7.62 (t, $J=7.2$ Hz, 1H), 7.53 (t, $J=7.6$ Hz, 2H), 7.43 (t, $J=7.6$ Hz, 1H), 7.04 (d, $J=12.0$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 189.9, 161.4, 150.1, 137.6, 137.2, 133.1, 133.0, 130.2, 129.6, 128.8, 128.5, 128.2, 122.7, 110.7. HRMS (EI) calcd for $C_{16}H_{11}BrO_3$ $[M]^+$: 329.9892; found 329.9891.

(*E*)-3-oxo-3-phenylprop-1-en-1-yl thiophene-2-carboxylate (**3ia**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.61 (d, $J=12.0$ Hz, 1H), 8.01 (d, $J=3.2$ Hz, 1H), 7.97 (d, $J=7.2$ Hz, 2H), 7.74 (d, $J=4.8$ Hz, 1H), 7.62 (t, $J=7.2$ Hz, 1H), 7.53 (t, $J=7.2$ Hz, 2H), 7.22~7.20 (m, 1H), 6.99 (d, $J=12.0$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 190.0, 158.0, 150.2, 137.7, 135.6, 134.7, 132.9, 130.8, 128.5, 128.2, 110.2. HRMS (EI) calcd for $C_{14}H_{10}O_3S$ $[M]^+$:

258.0351; found 258.0354.

(*E*)-3-oxo-3-phenylprop-1-en-1-yl 2-methylbenzoate (**3bb**): Pale yellow solid. m.p. 90~91 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.65 (d, $J=12.0$ Hz, 1H), 8.05 (d, $J=8.0$ Hz, 2H), 7.98 (d, $J=8.4$ Hz, 2H), 7.32 (d, $J=8.4$ Hz, 2H), 6.99~6.96 (m, 3H), 3.89 (s, 3H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 188.5, 163.4, 162.7, 150.0, 145.2, 130.7, 130.5, 130.2, 129.3, 125.0, 113.7, 109.8, 55.3, 21.7. HRMS (EI) calcd for $C_{18}H_{16}O_4$ $[M]^+$: 296.1049; found 296.1051.

(*E*)-3-(4-Fluorophenyl)-3-oxoprop-1-en-1-yl 4-methylbenzoate (**3bc**): White solid. m.p. 92~94 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.67 (d, $J=12.4$ Hz, 1H), 8.05~7.98 (m, 4H), 7.33 (d, $J=8.0$ Hz, 2H), 7.19 (t, $J=8.4$ Hz, 2H), 6.96 (d, $J=12.4$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 188.5, 166.8 (d, $J=254.0$ Hz), 162.5, 150.8, 145.5, 134.1 (d, $J=3.0$ Hz), 130.8 (d, $J=9.3$ Hz), 130.3, 129.4, 124.8, 115.7 (d, $J=21.7$ Hz), 109.6, 21.7. HRMS (EI) calcd for $C_{17}H_{13}FO_3$ $[M]^+$: 284.0849; found 284.0850.

(*E*)-3-(4-Chlorophenyl)-3-oxoprop-1-en-1-yl 4-methylbenzoate (**3bd**): Pale yellow solid. m.p. 115~117 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.68 (d, $J=12.4$ Hz, 1H), 8.06 (d, $J=8.4$ Hz, 2H), 7.92 (d, $J=8.4$ Hz, 2H), 7.49 (d, $J=8.4$ Hz, 2H), 7.33 (d, $J=8.4$ Hz, 2H), 6.95 (d, $J=12.4$ Hz, 1H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 188.9, 162.5, 151.0, 145.5, 139.3, 136.1, 130.3, 129.6, 129.4, 128.8, 124.7, 109.6, 21.7. HRMS (EI) calcd for $C_{17}H_{13}ClO_3$ $[M]^+$: 300.0553; found 300.0554.

(*E*)-3-(Naphthalen-2-yl)-3-oxoprop-1-en-1-yl 4-methylbenzoate (**3bf**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.49 (d, $J=12.4$ Hz, 1H), 8.36 (d, $J=8.0$ Hz, 1H), 8.04~7.99 (m, 3H), 7.92 (d, $J=7.6$ Hz, 1H), 7.76 (d, $J=7.2$ Hz, 1H), 7.59~7.51 (m, 3H), 7.32 (d, $J=8.0$ Hz, 2H), 6.78 (d, $J=12.4$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 195.0, 162.5, 151.9, 145.5, 136.4, 133.7, 131.7, 130.3, 130.2, 129.4, 128.3, 127.4, 127.1, 126.4, 125.4, 124.7, 124.3, 115.3, 21.7. HRMS (EI) calcd for $C_{21}H_{16}O_3$ $[M]^+$: 316.1099; found 316.1101.

(*E*)-3-(2-Chlorophenyl)-3-oxoprop-1-en-1-yl 2-methylbenzoate (**3gg**): Yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.34 (d, $J=12.4$ Hz, 1H), 8.09 (d, $J=8.0$ Hz, 1H), 7.51~7.31 (m, 7H), 6.58 (d, $J=12.4$ Hz, 1H), 2.64 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 193.2, 162.4, 152.4, 142.2, 138.5, 133.6, 132.1, 131.4, 131.2, 131.0, 130.2, 128.9, 126.7, 126.1, 126.0, 114.7, 21.8. HRMS (EI) calcd for $C_{17}H_{13}ClO_3$ $[M]^+$: 300.0553; found 300.0556.

(*E*)-3-(4-Bromophenyl)-3-oxoprop-1-en-1-yl 2-methylbenzoate (**3gh**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.68 (d, $J=12.4$ Hz, 1H), 8.10 (d, $J=8.0$ Hz, 1H), 7.83 (d, $J=8.4$ Hz, 2H), 7.65 (d, $J=8.4$ Hz, 2H), 7.53 (t, $J=7.6$ Hz, 1H), 7.35~7.32 (m, 2H), 6.92 (d, $J=12.4$ Hz, 1H), 2.67 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 189.1, 162.5, 151.0, 142.1, 136.5, 133.5, 132.1, 131.8, 131.1, 129.7, 127.9, 126.4, 125.9, 109.5, 21.8. HRMS (EI) calcd for $C_{17}H_{13}BrO_3$ $[M]^+$: 344.0048; found 344.0051.

(*E*)-3-(3-Bromophenyl)-3-oxoprop-1-en-1-yl 2-methylbenzoate (**3gi**): Pale yellow oil. 1H NMR (400 MHz, $CDCl_3$)

δ : 8.69 (d, $J=12.4$ Hz, 1H), 8.11 (d, $J=8.0$ Hz, 1H), 8.09 (s, 1H), 7.88 (d, $J=7.6$ Hz, 1H), 7.72 (d, $J=7.6$ Hz, 1H), 7.54 (t, $J=7.6$ Hz, 1H), 7.40~7.32 (m, 3H), 6.92 (d, $J=12.4$ Hz, 1H), 2.68 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 188.8, 162.5, 151.2, 142.1, 139.5, 135.6, 133.5, 132.1, 131.2, 131.1, 130.1, 126.6, 126.3, 125.9, 122.9, 109.4, 21.8. HRMS (EI) calcd for $\text{C}_{17}\text{H}_{13}\text{BrO}_3$ $[\text{M}]^+$: 344.0048; found 344.0044.

(*E*)-3-(4-Methoxyphenyl)-3-oxoprop-1-en-1-yl benzoate (**3db**): Pale yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.66 (d, $J=12.4$ Hz, 1H), 8.18 (d, $J=7.6$ Hz, 2H), 7.99 (d, $J=8.0$ Hz, 2H), 7.69 (t, $J=7.6$ Hz, 1H), 7.54 (t, $J=7.6$ Hz, 2H), 7.02~6.97 (m, 3H), 3.89 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 188.4, 163.5, 162.7, 149.8, 134.2, 130.7, 130.5, 130.2, 130.0, 128.6, 113.7, 110.1, 55.3. HRMS (EI) calcd for $\text{C}_{17}\text{H}_{14}\text{O}_4$ $[\text{M}]^+$: 282.0892; found 282.0889.

Supporting Information ^1H NMR and ^{13}C NMR spectra for compound **3**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Fuji, K. Bond-Cleavage Reactions with Hard Acid and Soft Nucleophile Systems, *Advances in Chemistry, ACS Symposium Series*, American Chemical Society, Washington, DC, **1987**, Vol. 215, p. 219.
- [2] For selected reviews, see:
 - (a) van der Boom, M. E.; Milstein, D. *Chem. Rev.* **2003**, *103*, 1759.
 - (b) Tobisu, M.; Chatani, N. *Chem. Soc. Rev.* **2008**, *37*, 300.
- [3] For selected reviews, see:
 - (a) Lu, H.; Yu, T.-Y.; Xu, P.-F.; Wei, H. *Chem. Rev.* **2021**, *121*, 365.
 - (b) Lutz, M. D. R.; Morandi, B. *Chem. Rev.* **2021**, *121*, 300.
 - (c) McDonald, T. R.; Mills, R. M.; West, M. S.; Rousseaux, S. A. L. *Chem. Rev.* **2021**, *121*, 3.
 - (d) Jun, C.-H. *Chem. Soc. Rev.* **2004**, *33*, 610.
 - (e) Wu, J.; Zhu, J.; Li, H.; Wu, C.; Shen, R.; Yu, L. *Chin. J. Org. Chem.* **2019**, *39*, 573 (in Chinese).
 (吴锦雯, 朱佳雯, 李慧, 吴春雷, 沈润溥, 余乐茂, 有机化学, **2019**, *39*, 573.)
- [4] For selected examples, see:
 - (a) Suginome, M.; Matsuda, T.; Ito, Y. *J. Am. Chem. Soc.* **2000**, *122*, 11015.
 - (b) Youn, S. W.; Kim, B. S.; Jagdale, A. R. *J. Am. Chem. Soc.* **2012**, *134*, 11308.
 - (c) Zhou, M.-B.; Luo, M.-J.; Hu, M.; Li, J.-H. *Chin. J. Chem.* **2020**, *38*, 553.
- [5] For selected examples, see:
 - (a) Onodera, S.; Ishikawa, S.; Kochi, T.; Fakiuchi, F. *J. Am. Chem. Soc.* **2018**, *140*, 9788.
 - (b) Chatani, N.; Le, Y.; Kakiuchi, F.; Murai, S. *J. Am. Chem. Soc.* **1999**, *121*, 8645.
 - (c) Cheng, C.-W.; Kuo, Y.-C.; Chang, S.-H.; Lin, Y.-C.; Liu, Y.-H.; Wang, Y. *J. Am. Chem. Soc.* **2007**, *129*, 14974.
- [6] For selected examples, see:
 - (a) Brendel, M.; Sakhare, P. R.; Dahiya, G.; Subramanian, P.; Kaliappan, K. P. *J. Org. Chem.* **2020**, *85*, 8102.
 - (b) Fan, W.; Yang, Y.; Lei, J.; Jiang, Q.; Zhou, W. *J. Org. Chem.* **2015**, *80*, 8782.
 - (c) Zhu, Y.; Yan, H.; Lu, L.; Liu, D.; Rong, G.; Mao, J. *J. Org. Chem.* **2013**, *78*, 9898.
 - (d) Zhang, L.; Bi, X.; Guan, X.; Li, X.; Liu, Q. *Angew. Chem., Int. Ed.* **2013**, *52*, 11303.
- [7] For selected examples, see:
 - (a) Wang, T.; Jiao, N. *J. Am. Chem. Soc.* **2013**, *135*, 11692.
 - (b) Liang, D.; He, Y.; Liu, L.; Zhu, Q. *Org. Lett.* **2013**, *15*, 3476.
- [8] For selected examples, see:
 - (a) Ji, T.; Chen, X.-Y.; Huang, L.; Rueping, M. *Org. Lett.* **2020**, *22*, 2579.
 - (b) Zhu, S.; Das, A.; Bui, L.; Zhou, H.; Curran, D. P.; Rueping, M. *J. Am. Chem. Soc.* **2013**, *135*, 1823.
 - (c) Zou, S.; Li, R.; Kobayashi, H.; Liu, J.; Fan, J. *Chem. Commun.* **2013**, *49*, 1906.
 - (d) Song, C.; Shen, X.; Yu, F.; He, Y.; Yu, S. *Chin. J. Org. Chem.* **2020**, *40*, 3748 (in Chinese).
 (宋常华, 沈许, 于芳, 何宇鹏, 俞寿云, 有机化学, **2020**, *40*, 3748.)
- (e) Zhao, B.; Tan, H.; Chen, C.; Jiao, N.; Shi, Z. *Chin. J. Chem.* **2018**, *36*, 995.
- [9] (a) Mecinovic, J.; Hamed, R. B.; Schofield, C. J. *Angew. Chem., Int. Ed.* **2009**, *48*, 2796.
- (b) Allpress, C. J.; Grubel, K.; Szajna-Fuller, E.; Arif, A. M.; Berreau, L. M. *J. Am. Chem. Soc.* **2013**, *135*, 659.
- [10] Kohler, E.; Erickson, J. L. E. *J. Am. Chem. Soc.* **1931**, *53*, 2301.
- [11] Barany, H. C.; Pianka, M.; Waters, W. A.; Schöenberg, A.; Moubasher, R.; Mostafa, A. *J. Chem. Soc.* **1946**, 965.
- [12] Schöenberg, A.; Azzam, R. C. *J. Chem. Soc.* **1939**, 1428.
- [13] Tada, N.; Shomura, M.; Cui, L.; Nobuta, T.; Miura, T.; Itoh, A. *Synlett* **2011**, 2896.
- [14] Berreau, L. M.; Grubel, K.; Allpress, C. J.; Borowski, T.; Wikstrom, J. P.; Germain, M. E.; Rybak-Akimova, E. V.; Tierney, D. L. *Inorg. Chem.* **2011**, *50*, 1047.
- [15] Schöenberg, A.; Mustafa, A. *J. Chem. Soc.* **1947**, 997.
- [16] For selected recent reviews see:
 - (a) Guo, H.; Fan, Y.; Sun, Z.; Wu, Y.; Kwon, O. *Chem. Rev.* **2018**, *118*, 10049.
 - (b) Lu, X.; Zhang, C.; Xu, Z. *Acc. Chem. Res.* **2001**, *34*, 535.
 - (c) Ye, L.-W.; Zhou, J.; Tang, Y. *Chem. Soc. Rev.* **2008**, *37*, 1140.
- [17] (a) Nair, V.; Pillai, A. N.; Menon, R. S.; Suresh, E. *Org. Lett.* **2005**, *7*, 1189.
- (b) Guo, H.; Xu, Q.; Kwon, O. *J. Am. Chem. Soc.* **2009**, *131*, 6318.
- (c) Wu, J.-Y.; Luo, Z.-B.; Dai, L.-X.; Hou, X.-L. *J. Org. Chem.* **2008**, *73*, 9137.
- (d) Garima; Srivastava, V. P.; Yadav, L. D. S. *Green Chem.* **2010**, *12*, 1460.
- (e) Meng, L.-G.; Hu, B.; Wu, Q.-P.; Liang, M.; Xue, S. *Chem. Commun.* **2009**, 6089.
- [18] (a) Meng, L.-G.; Wang, L. *Chem. Commun.* **2012**, *48*, 3242.
- (b) Meng, L.-G.; Wang, L. *Adv. Synth. Catal.* **2013**, *355*, 2967.
- (c) Meng, L.-G.; Ge, N.-L.; Yang, M.-M.; Wang, L. *Eur. J. Org. Chem.* **2011**, 3403.
- [19] (a) Stang, P. J. *Acc. Chem. Res.* **1991**, *24*, 304.
- (b) Ackermann, L. *Chem. Rev.* **2011**, *111*, 1315.

(Fan, Y.)