铁(0)促进的醛与溴代硝基甲烷的 Henry 型反应高效合成 2-硝基-1-醇

张斯旋^{†,a} 李祥瑞^{†,a} 李文欣^a 饶卫东^b 葛丹华^{*,a}
沈志良^a 褚雪强^{*,a}

(^a 南京工业大学化学与分子工程学院 先进化学制造研究院 化学实验教学中心 南京 211816)

(^b 南京林业大学化工学院 江苏省生物质绿色燃料与化学品重点实验室 南京 210037)

摘要 以廉价易得、商品化的铁粉为反应促进剂,研究了溴代硝基甲烷与各种醛的 Henry 型反应,高效地合成了 2-硝基-1-醇。该反应在 PbCl₂ 和四丁基硫酸氢铵(TBAHS)存在下有效地进行,以中等至良好的产率得到目标产物,并且该反应应具有广泛的官能团耐受性。

关键词 铁; 二氯化铅; Henry 型反应; 溴代硝基甲烷; 2-硝基-1-醇

Iron(0)-Mediated Henry-Type Reaction of Bromonitromethane with Aldehydes for the Efficient Synthesis of 2-Nitro-alkan-1-ols

Zhang, Sixuan^{†,a} Li, Xiangrui^{†,a} Li, Wenxin^a Rao, Weidong^b Ge, Danhua^{*,a}
Shen, Zhiliang^a Chu, Xueqiang^{*,a}

(^a Chemical Experiment Teaching Center, Institute of Advanced Synthesis, School of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816)

(^b Jiangsu Key Laboratory of Biomass-based Green Fuels and Chemicals, College of Chemical Engineering, Nanjing Forestry University, Nanjing 210037)

Abstract The Henry-type reaction of bromonitromethane with various aldehydes for the efficient synthesis of 2-nitro-alkan-1-ols by using inexpensive and commercial iron powder as reaction mediator was developed. The reaction proceeded efficiently in the presence of PbCl₂ and tetrabutylammonium hydrogen sulfate (TBAHS) to produce the desired products in moderate to good yield with wide functional group tolerance.

Keywords iron, lead dichloride; Henry-type reaction; bromonitromethane; 2-nitro-alkan-1-ol

1 Introduction

2-Nitro-alkan-1-ols serve as versatile synthetic intermediates in organic synthesis and could be readily transformed into amino alcohols, amino sugars, nitroketones, nitroalkenes, and other important molecules.^[1] Conventionally, 2-nitro-alkan-1-ol could be prepared by the base-mediated Henry reaction (also known as nitro-aldol reaction) of carbonyl compounds with nitroalkanes.^[2-4] In recent decades, metal-mediated Henry-type reaction of carbonyl compounds with bromonitromethanes also serves as an attractive alternative to the conventional Henry reaction where normally bases are required. For instance, several metal

mediators, including SmI₂,^[5] In,^[6] Zn,^[7] and SnCl₂^[8] were reported to effectively promote the Henry-type reaction of aldehydes with bromonitromethanes. However, the metal mediators used for the aforementioned Henry-type reaction are more or less air-sensitive, expensive, and toxic. Thus, the development of a Henry-type reaction by using an inexpensive and environmentally benign metal mediator is still highly desirable.

Among the various metals in Periodic Table, iron(0) is definitely one of the most abundant and inexpensive metals. In addition, iron(0) and most of its salts are less toxic and thus are environmentally benign. Nonetheless, the utility of

* Corresponding authors. E-mail: xueqiangchu@njtech.edu.cn; gedanhua@njtech.edu.cn.

Received July 23, 2021; revised August 27, 2021; published online August 28, 2021.

Project supported by the Nanjing Tech University (Start-up Grant Nos. 39837118, 39837146), the National Natural Science Foundation of China (No. 22001121), and the Natural Science Foundation of Jiangsu Province (No. BK20180690).

南京工业大学科研启动基金(Nos. 39837118, 39837146)、国家自然科学基金(No. 22001121)、江苏省自然科学基金(No. BK20180690)资助项目。

[†] 共同第一作者(These authors contributed equally to this work).

iron(0) powder in synthetic organic chemistry has not been widely exploited,^[9] presumably owing to the lack of effective methods for the activation of commercial iron powder since it contains an oxide layer on its surface. In continuation of our task in developing iron(0)-mediated organic transformations^[9d,9e] herein we report an efficient Henry-type reaction of bromonitromethane with various aldehydes by using inexpensive, non-toxic, and commercial iron(0) powder as reaction mediator. The reactions proceeded efficiently in the presence of PbCl₂ and tetrabutylammonium hydrogen sulfate (TBAHS) to produce the desired 2-nitro-alkan-1-ols in moderate to good yield with wide functional group tolerance, serving as an attractive alternative to existing methods.

2 Results and discussion

Initially, the Henry-type reaction of 4-chlorobenzaldehyde (**1a**) with bromonitromethane (**2a**) was performed in acetonitrile at 60 °C for 24 h in the presence of commercial iron powder (3 equiv.) and a series of catalysts (0.2 equiv.). As shown in Table 1, the Henry-type reaction could not occur in the absence of any catalyst (Entry 1). Interestingly, varying reaction performance was observed when the reaction was performed by employing various metallic salts as reaction catalyst. Among the different catalysts surveyed, such as InCl₃,^[10-11] BiCl₃,^[10,12] CrCl₃,^[13] FeCl₃,^[14] PbCl₂,^[10,15] CuX,^[16] NH₄I,^[17] and LiI,^[18] PbCl₂ emerged to be the most efficient catalyst for the reaction, leading to the corresponding product **3a** in an acceptable yield of 58% (53% isolated yield; Entry 11).

Table 1 Optimization of reaction conditions by using various catalysts^a

Entry	Catalyst	Yield ^b /%	Entry	Catalyst	Yield ^b /%
1	—	0	10	FeBr ₂	26
2	InCl ₃	25	11	PbCl ₂	58 (53) ^c
3	BiCl ₃	28	12	PbBr ₂	32
4	MnCl ₂	48	13	PbI ₂	28
5	CrCl ₃	24	14	CuBr	38
6	ZnCl ₂	46	15	CuI	26
7	FeCl ₃	34	16	NaI	30
8	FeCl ₂	22	17	NH ₄ I	10
9	FeBr ₃	26	18	LiI	30

^a Unless otherwise noted, the reactions were performed at 60 °C for 24 h by using aldehyde **1a** (0.3 mmol), bromonitromethane (**2a**, 0.9 mmol), Fe (0.9 mmol), and catalyst (0.2 equiv.) in CH₃CN (1 mL). ^b Yields were determined by NMR analysis of crude reaction mixture after work-up by using 1,4-dimethoxybenzene as an internal standard. ^c Isolated yield.

Next, we continued to optimize the reaction conditions by employing various additives. As shown in Table 2, of the several additives screened (Entries 1~4), tetrabutylammonium hydrogen sulfate (TBAHS)^[19] was found to be the additive of choice, affording the desired 2-nitro-alkan-1-ol **3a** in an improved NMR yield of 78% (74% isolated yield; Entry 3). Interestingly, the use of other acidic additives, such as KHSO₄ and TsOH, also led to the product **3a** in acceptable yield (67% NMR yield, Entries 5~6). However, when the reaction was performed by using either 2 equiv. of iron as mediator or 2 equiv. of bromonitromethane as substrate, the corresponding product **3a** was only obtained in 22% NMR yield and 18% NMR yield, respectively (Entries 7~8). In addition, the use of other metal to replace iron as mediator under the optimized reaction conditions (shown in Entry 3) resulted in sharply decreased product yield (Mn: 9%, Al: <5%, Pb: <5%, Ga: <5%, Bi: 14%). Moreover, survey of a series of reaction solvents showed that CH₃CN was still the solvent of choice (THF: 22%, DMF: 56%, DMSO: 37%, MeOH: 13%, DCE: 13%, DME: 38%). With the encouraging result in hand (see Entry 3), we proceeded to study the substrate scope of the reaction by using various aldehydes as starting materials.

Table 2 Optimization of reaction conditions by using various additives^a

Entry	Additive	Yield ^b /%
1	TBAB	32
2	TBAI	<5
3	TBAHS	78 (74) ^c
4	CTMAB	20
5	KHSO ₄	67
6	TsOH	67
7	TBAHS	22 ^d
8	TBAHS	18 ^e

^a The reactions were performed at 60 °C for 24 h by using aldehyde **1a** (0.3 mmol), bromonitromethane (**2a**, 0.9 mmol), Fe (0.9 mmol), PbCl₂ (0.2 equiv.), and additive (1 equiv.) in MeCN (1 mL). ^b Yields were determined by NMR analysis of crude reaction mixture after work-up by using 1,4-dimethoxybenzene as an internal standard. ^c Isolated yield. ^d Using 2 equiv. of iron powder. ^e Using 2 equiv. of **2a**. TBAHS=tetrabutylammonium hydrogen sulfate, CTMAB=hexadecyl trimethyl ammonium bromide.

As outlined in Table 3, various aldehydes **1b**~**1r** were capable of effectively undergoing iron-mediated Henry-type reaction with bromonitromethane (**2a**) in the presence of lead dichloride and TBAHS under the well-established

Table 3 Substrates scope study by employing different aldehydes^a

$ \begin{array}{c} \text{R}-\text{CHO} \\ \text{1b} \sim \text{1r} \end{array} + \text{O}_2\text{N}-\text{CH}_2-\text{Br} \xrightarrow[\text{CH}_3\text{CN}, 60^\circ\text{C}, 24\text{ h}]{\text{Fe (3 equiv.)}, \text{PbCl}_2 (0.2 \text{ equiv.}), \text{TBAHS (1 equiv.)}} \begin{array}{c} \text{R}-\text{CH}(\text{OH})-\text{CH}_2-\text{NO}_2 \\ \text{3b} \sim \text{3r} \end{array} $							
Entry	Aldehyde	Product	Yield ^b /%	Entry	Aldehyde	Product	Yield ^b /%
1		3b	68	10		3k	64
2		3c	63	11		3l	41
3		3d	75	12		3m	36
4		3e	69	13		3n	49
5		3f	65	14		3o	55
6		3g	60	15		3p	45
7		3h	66	16		3q	43
8		3i	85	17		3r	47 ^{c,d}
9		3j	67				

^a Unless otherwise noted, the reactions were performed at 60 °C for 24 h by using aldehyde **1b**~**1r** (0.3 mmol), bromonitromethane (**2a**, 0.9 mmol), Fe (0.9 mmol), PbCl₂ (0.05 mmol), and TBAHS (0.3 mmol) in CH₃CN (1 mL). ^b Isolated yield. ^c 50:50 *dr*. ^d With around 10% impurity which could not be purified by silica gel column chromatography.

reaction conditions, leading to the expected products **3b**~**3r** in moderate to good yields. In most cases, aryl aldehydes containing electron-withdrawing group (Entries 1~10) showed improved reactivities than aryl aldehydes possessing electron-donating group (Entries 11~12). It should be mentioned that the nitro group embedded in substrate **1b**, which has the tendency to be reduced by iron metal^[9a-9c], remained intact in the present protocol. Analogously, heterocyclic aldehyde **1o** bearing thienyl moiety, which might be sensitive to acidic conditions, worked equally well under the optimal conditions to furnish the desired product **3o** in 55% yield (Entry 14). In a same manner, aliphatic aldehydes **1q**~**1r** were proven to be appropriate candidates, although slightly reduced product yields were obtained (Entries 16~17). Remarkably, the mild reaction conditions allowed the presence of many important functional groups in the aldehydes, including nitro, methoxycarbonyl, trifluoromethyl, trifluoromethoxy, cyano, fluoro, chloro, bromo, iodo, and alkynyl, which could potentially be kept for late-stage derivatization. However, when 2-bromo-2-nitropropane was used as substrate, the reaction could not proceed under the established reaction conditions, presumably because of the low reactivity or steric hindrance of the substrate. Additionally, when less reactive ketone (e.g., acetophenone) was employed as starting material, the reaction could not occur either.

The mildness of the reaction conditions also enabled the reaction of substrate **2a** with α,β -unsaturated aldehyde **1s** to regioselectively take place at the aldehyde rather than the alkene, giving rise to the corresponding product **3s** in 56% yield (Scheme 1a). When substrate **1t** containing both formyl and acetyl groups was subjected to the optimized reaction conditions, it was found that the reaction occurred regioselectively at the more reactive formyl group, leaving the relatively less reactive acetyl functionality untouched (Scheme 1b).

Scheme 1 Regioselective reaction with substrates **1s** and **1t**

Mechanistically, the reaction might occur via a single electron transfer (SET) from iron metal to bromonitromethane to form a nitromethyl radical, which then attacks the aldehyde to give the desired product of 2-nitro-alkan-1-ol (after another SET from iron to the *in situ* formed alkoxy radical to convert it to alkoxy anion followed by protonation). Alternatively, the reaction might proceed via the formation of a nitromethyliron compound (from bromonitromethane **2a**) which behaves as a nucleophile to react with aldehyde; subsequent protonation of the formed alkoxy anion would also give the final product. But other possibilities could also not be ruled out at the current stage. As for the role of PbCl₂ and TBAHS, currently it is unknown. It should be mentioned that the accelerating effect of lead(II) salt in promoting either metal insertion reaction or metal-mediated organic transformation has been reported in several cases^[10,15]. In the present reaction, PbCl₂ might also act as a catalyst to activate the iron metal. Analogously, the slightly acidic TBAHS could also help to activate the iron metal by removing the oxide layer from its surface. Besides, we conjecture that PbCl₂ and TBAHS might also serve as Lewis acid and protic acid to coordinate with the oxygen atom of the carbonyl substrate to increase the electrophilicity/reactivity of the aldehyde substrate.

3 Conclusions

In a nutshell, a facile method for the synthesis of 2-nitro-alkan-1-ols via the Henry-type reaction of bromonitromethane with various aldehydes by using inexpensive and commercial iron powder as reaction mediator was developed. The reactions proceeded effectively in the presence of PbCl₂ and TBAHS to afford the corresponding products in moderate to good yield with broad functional group compatibility, serving as an attractive alternative to existing methods.

4 Experimental section

4.1 General

Unless otherwise stated, all reagents were purchased from commercial suppliers and used without further purification. All reactions were carried out under air using undistilled solvent, without the need of precautions to exclude air and moisture. IR spectra were recorded on a FT-IR spectrophotometer using KBr optics. ¹H NMR and ¹³C NMR spectra were recorded in CDCl₃ on a Bruker Avance or a Jeol 400 MHz spectrometers. Tetramethylsilane (TMS) served as an internal standard for ¹H NMR and ¹³C NMR. High resolution mass spectra (HRMS) were obtained using a commercial apparatus (ESI or EI Source).

4.2 Typical procedures for the Henry-type reaction for the synthesis of 2-nitroalkan-1-ol

To a 10 mL Schlenk flask was added iron powder (50 mg, 0.9 mmol) and CH₃CN (1 mL), and the iron powder was sequentially activated by using 1,2-dibromoethane (0.10 mmol) and TMSCl (0.10 mmol). After cooling down, al-

dehyde (0.3 mmol), bromo(nitro)methane (126 mg, 0.9 mmol), PbCl_2 (17 mg, 0.06 mmol), and TBAHS (101 mg, 0.3 mmol) were added. The reaction mixture was vigorously stirred at 60 °C for 24 h under nitrogen atmosphere, followed by quenching with sat. NH_4Cl solution (20 mL) and extraction with ethyl acetate (20 mL $\times$ 3). The combined extracts were washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated *in vacuo*. Purification by silica gel column chromatography using ethyl acetate/petroleum ether as eluent gave the analytically pure product.

1-(4-Chlorophenyl)-2-nitroethan-1-ol (**3a**): 44.6 mg, 74% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.41~7.31 (m, 4H), 5.44 (dt, $J=9.5, 3.1$ Hz, 1H), 4.57 (dd, $J=13.5, 9.5$ Hz, 1H), 4.48 (dd, $J=13.5, 3.2$ Hz, 1H), 3.07 (d, $J=3.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 136.5, 134.8, 129.2, 127.3, 80.9, 70.2; IR (KBr) ν : 3447, 1623, 1595, 1553, 1380, 1332, 910 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{ClNO}_3$ $[\text{M}+\text{H}]^+$ 202.0265, found 202.0271.

2-Nitro-1-(4-nitrophenyl)ethan-1-ol (**3b**): 43.3 mg, 68% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 8.26 (d, $J=8.8$ Hz, 2H), 7.62 (d, $J=8.8$ Hz, 2H), 5.60 (dd, $J=8.4, 3.9$ Hz, 1H), 4.62~4.55 (m, 2H), 2.59 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 148.0, 145.0, 126.9, 124.1, 80.6, 69.9; IR (KBr) ν : 2920, 1557, 1520, 1350, 1085, 856, 755 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{N}_2\text{O}_5$ $[\text{M}+\text{H}]^+$ 213.0506, found 213.0511.

Methyl 4-(1-hydroxy-2-nitroethyl)benzoate (**3c**): 42.6 mg, 63% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 8.03 (d, $J=8.4$ Hz, 2H), 7.48 (d, $J=8.4$ Hz, 2H), 5.53 (dd, $J=9.3, 3.2$ Hz, 1H), 4.59 (dd, $J=13.4, 9.3$ Hz, 1H), 4.52 (dd, $J=13.4, 3.3$ Hz, 1H), 3.91 (s, 3H), 3.34 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.6, 143.0, 130.5, 130.2, 125.9, 80.9, 70.5, 52.3; IR (KBr) ν : 3326, 1719, 1558, 1431, 1282, 1110, 767 cm^{-1} . HRMS calcd for $\text{C}_{10}\text{H}_{12}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 226.0710, found 226.0715.

2-Nitro-1-(4-(trifluoromethyl)phenyl)ethan-1-ol (**3d**): 52.9 mg, 75% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.67 (d, $J=8.0$ Hz, 2H), 7.55 (d, $J=8.1$ Hz, 2H), 5.55 (dt, $J=9.0, 3.5$ Hz, 1H), 4.60 (dd, $J=13.6, 9.1$ Hz, 1H), 4.53 (dd, $J=13.6, 3.4$ Hz, 1H), 3.14 (d, $J=3.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 141.9 (q, $J=1.3$ Hz), 131.1 (q, $J=32.4$ Hz), 126.3, 126.0 (q, $J=3.8$ Hz), 123.8 (q, $J=270.2$ Hz), 80.9, 70.3; ^{19}F NMR (376 MHz, CDCl_3) δ : -62.62 (s, 3F); IR (KBr) ν : 1551, 1380, 1328, 1169, 1126, 1069, 844 cm^{-1} . HRMS calcd for $\text{C}_9\text{H}_9\text{F}_3\text{NO}_3$ $[\text{M}+\text{H}]^+$ 236.0529, found 236.0535.

2-Nitro-1-(4-(trifluoromethoxy)phenyl)ethan-1-ol (**3e**): 52.0 mg, 69% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.47~7.43 (m, 2H), 7.28~7.24 (m, 2H), 5.49 (dt, $J=9.5, 3.4$ Hz, 1H), 4.59 (dd, $J=13.5, 9.4$ Hz, 1H), 4.51 (dd, $J=13.5, 3.1$ Hz, 1H), 3.05 (d, $J=3.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 149.6 (q, $J=2.0$ Hz), 136.8, 127.6, 121.7~121.6 (m), 120.5 (q, $J=257.6$ Hz), 81.1, 70.3; ^{19}F NMR (376 MHz, CDCl_3) δ : -57.78 (s, 3F); IR (KBr) ν : 3585, 1557, 1510, 1271, 1163, 1075, 856 cm^{-1} . HRMS calcd for $\text{C}_9\text{H}_9\text{F}_3\text{NO}_4$ $[\text{M}+\text{H}]^+$ 252.0478, found

252.0484.

4-(1-Hydroxy-2-nitroethyl)benzonitrile (**3f**): 37.5 mg, 65% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.71 (d, $J=8.6$ Hz, 2H), 7.56 (d, $J=8.0$ Hz, 2H), 5.57~5.53 (m, 1H), 4.64~4.50 (m, 2H), 3.21 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 143.1, 132.8, 126.7, 118.2, 112.7, 80.6, 70.1; IR (KBr) ν : 3435, 2239, 1552, 1410, 1376, 1085, 840 cm^{-1} . HRMS calcd for $\text{C}_9\text{H}_9\text{N}_2\text{O}_3$ $[\text{M}+\text{H}]^+$ 193.0608, found 193.0613.

1-(4-Fluorophenyl)-2-nitroethan-1-ol (**3g**): 33.3 mg, 60% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.42~7.38 (m, 2H), 7.13~7.08 (m, 2H), 5.46 (dd, $J=9.5, 3.2$ Hz, 1H), 4.59 (dd, $J=13.4, 9.5$ Hz, 1H), 4.49 (dd, $J=13.4, 3.1$ Hz, 1H), 2.94 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.8 (d, $J=246.1$ Hz), 133.9 (d, $J=3.3$ Hz), 127.7 (d, $J=8.3$ Hz), 116.0 (d, $J=21.6$ Hz), 81.1, 70.3; ^{19}F NMR (376 MHz, CDCl_3) δ : -112.35 (s, 1F); IR (KBr) ν : 3566, 1606, 1557, 1380, 1226, 1079, 839 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{FNO}_3$ $[\text{M}+\text{H}]^+$ 186.0561, found 186.0566.

1-(4-Bromophenyl)-2-nitroethan-1-ol (**3h**): 48.7 mg, 66% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.53 (d, $J=8.5$ Hz, 2H), 7.28 (d, $J=8.5$ Hz, 2H), 5.43 (dd, $J=9.4, 3.2$ Hz, 1H), 4.57 (dd, $J=13.5, 9.4$ Hz, 1H), 4.48 (dd, $J=13.5, 3.2$ Hz, 1H), 2.99 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 137.0, 132.1, 127.6, 122.9, 80.9, 70.3; IR (KBr) ν : 3373, 1553, 1489, 1385, 1074, 1010, 823 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{BrNO}_3$ $[\text{M}+\text{H}]^+$ 245.9760, found 245.9766.

1-(2-Bromophenyl)-2-nitroethan-1-ol (**3i**): 62.7 mg, 85% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.66 (dd, $J=7.8, 1.6$ Hz, 1H), 7.57 (dd, $J=8.0, 1.3$ Hz, 1H), 7.41 (td, $J=7.6, 1.2$ Hz, 1H), 7.24 (td, $J=7.7, 1.8$ Hz, 1H), 5.81 (d, $J=9.3$ Hz, 1H), 4.69 (dd, $J=13.6, 2.3$ Hz, 1H), 4.43 (dd, $J=13.6, 9.7$ Hz, 1H), 3.02 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 137.0, 132.9, 130.2, 128.2, 127.7, 121.4, 79.3, 69.9; IR (KBr) ν : 2915, 1556, 1468, 1377, 1340, 1024, 759 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{BrNO}_3$ $[\text{M}+\text{H}]^+$ 245.9760, found 245.9766.

1-(3-Bromophenyl)-2-nitroethan-1-ol (**3j**): 49.5 mg, 67% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.58 (t, $J=1.9$ Hz, 1H), 7.49 (dt, $J=7.7, 1.6$ Hz, 1H), 7.34~7.25 (m, 2H), 5.44 (dd, $J=9.3, 3.2$ Hz, 1H), 4.57 (dd, $J=13.5, 9.4$ Hz, 1H), 4.50 (dd, $J=13.5, 3.2$ Hz, 1H), 3.02 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 140.2, 132.0, 130.6, 129.1, 124.5, 123.1, 80.9, 70.2; IR (KBr) ν : 1555, 1378, 1341, 1194, 1073, 899, 789 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{BrNO}_3$ $[\text{M}+\text{H}]^+$ 245.9760, found 245.9766.

1-(4-Iodophenyl)-2-nitroethan-1-ol (**3k**): 56.3 mg, 64% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.74 (d, $J=8.0$ Hz, 2H), 7.16 (d, $J=8.1$ Hz, 2H), 5.42 (dd, $J=9.4, 3.0$ Hz, 1H), 4.56 (dd, $J=13.5, 9.4$ Hz, 1H), 4.48 (dd, $J=13.5, 3.1$ Hz, 1H), 2.97 (brs, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 138.1, 137.7, 127.8, 94.6, 80.9, 70.4; IR (KBr) ν : 3373, 1550, 1485, 1383, 1069, 1005, 822 cm^{-1} . HRMS calcd for $\text{C}_8\text{H}_9\text{INO}_3$ $[\text{M}+\text{H}]^+$ 293.9622, found 293.9627.

2-Nitro-1-(*p*-tolyl)ethan-1-ol (**3l**): 22.3 mg, 41% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ : 7.26 (d, $J=8.3$

H₂, 2H), 7.19 (d, $J=7.9$ Hz, 2H), 5.38 (dt, $J=9.6, 3.3$ Hz, 1H), 4.57 (dd, $J=13.2, 9.6$ Hz, 1H), 4.45 (dd, $J=13.2, 3.1$ Hz, 1H), 2.95 (d, $J=3.8$ Hz, 1H), 2.35 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 138.9, 135.1, 129.7, 125.8, 81.2, 70.9, 21.1; IR (KBr) ν : 3500, 1557, 1420, 1374, 1283, 1081, 723 cm⁻¹. HRMS calcd for C₉H₁₂NO₃ [M+H]⁺ 182.0812, found 182.0817.

1-(4-Methoxyphenyl)-2-nitroethan-1-ol (**3m**): 21.3 mg, 36% yield, colorless oil; ¹H NMR (400 MHz, CDCl₃) δ : 7.31 (d, $J=8.7$ Hz, 2H), 6.91 (d, $J=8.7$ Hz, 2H), 5.40 (dd, $J=9.6, 3.0$ Hz, 1H), 4.60 (dd, $J=13.1, 9.7$ Hz, 1H), 4.47 (dd, $J=13.2, 3.1$ Hz, 1H), 3.81 (s, 3H), 2.94 (brs, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 159.9, 130.2, 127.2, 114.3, 81.2, 70.6, 55.3; IR (KBr) ν : 3463, 2840, 1612, 1555, 1513, 1252, 833 cm⁻¹. HRMS calcd for C₉H₁₂NO₄ [M+H]⁺ 198.0761, found 198.0766.

2-Nitro-1-(4-((trimethylsilyl)ethynyl)phenyl)ethan-1-ol (**3n**): 38.7 mg, 49% yield, colorless oil; ¹H NMR (400 MHz, CDCl₃) δ : 7.49 (d, $J=8.5$ Hz, 2H), 7.34 (d, $J=8.1$ Hz, 2H), 5.46 (dd, $J=9.4, 3.1$ Hz, 1H), 4.57 (dd, $J=13.4, 9.5$ Hz, 1H), 4.49 (dd, $J=13.5, 3.1$ Hz, 1H), 2.91 (brs, 1H), 0.25 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ : 138.1, 132.5, 125.8, 123.8, 104.1, 95.3, 80.9, 70.6, -0.1; IR (KBr) ν : 3529, 2958, 2156, 1553, 1384, 1076, 867 cm⁻¹. HRMS calcd for C₁₃H₁₈NO₃Si [M+H]⁺ 264.1050, found 264.1056.

2-Nitro-1-(thiophen-2-yl)ethan-1-ol (**3o**): 28.6 mg, 55% yield, colorless oil; ¹H NMR (400 MHz, CDCl₃) δ : 7.34 (d, $J=5.0$ Hz, 1H), 7.07 (d, $J=3.5$ Hz, 1H), 7.02 (dd, $J=5.0, 3.6$ Hz, 1H), 5.73 (d, $J=9.2$ Hz, 1H), 4.72 (dd, $J=13.4, 9.4$ Hz, 1H), 4.61 (dd, $J=13.4, 3.2$ Hz, 1H), 3.12 (d, $J=4.1$ Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 141.2, 127.2, 126.8, 125.1, 80.8, 67.1; IR (KBr) ν : 3545, 1556, 1380, 1329, 1200, 1081, 711 cm⁻¹. HRMS calcd for C₆H₈NO₃S [M+H]⁺ 174.0219, found 174.0225.

1-(Naphthalen-1-yl)-2-nitroethan-1-ol (**3p**): 29.3 mg, 45% yield, colorless oil; ¹H NMR (400 MHz, CDCl₃) δ : 8.04 (d, $J=8.6$ Hz, 1H), 7.94~7.90 (m, 1H), 7.87 (d, $J=8.2$ Hz, 1H), 7.77 (d, $J=7.2$ Hz, 1H), 7.63~7.50 (m, 3H), 6.27 (dd, $J=8.6, 3.6$ Hz, 1H), 4.73~4.62 (m, 2H), 2.92 (brs, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 133.7, 133.5, 129.5, 129.4, 129.3, 127.1, 126.1, 125.5, 123.8, 121.8, 80.8, 68.3; IR (KBr) ν : 3480, 1572, 1550, 1381, 1225, 1100, 773 cm⁻¹. HRMS calcd for C₁₂H₁₂NO₃ [M+H]⁺ 218.0812, found 218.0817.

1-Nitro-4-phenylbutan-2-ol (**3q**): 25.2 mg, 43% yield, colorless oil; ¹H NMR (400 MHz, CDCl₃) δ : 7.34~7.30 (m, 2H), 7.25~7.20 (m, 3H), 4.43~4.39 (m, 2H), 4.36~4.27 (m, 1H), 2.90~2.83 (m, 1H), 2.80~2.71 (m, 2H), 1.94~1.74 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ : 140.6, 128.6, 128.4, 126.3, 80.5, 67.7, 35.1, 31.3; IR (KBr) ν : 3391, 1558, 1454, 1203, 1102, 880, 704 cm⁻¹. HRMS calcd for C₁₀H₁₄NO₃ [M+H]⁺ 196.0968, found 196.0974.

3,7-Dimethyl-1-nitrooct-6-en-2-ol (**3r**): 28.4 mg, 47% yield, 50 : 50 *dr*, white solid; ¹H NMR (400 MHz, CDCl₃) δ : 5.07 (t, $J=6.8$ Hz, 1H), 4.45~4.31 (m, 2H), 2.58 (brs, 1H), 1.99~1.95 (m, 2H), 1.68 (s, 3H), 1.60 (s, 3H), 1.46~1.30 (m, 2H), 1.28~1.12 (m, 2H), 0.95 (t, $J=7.1$ Hz,

3H); ¹³C NMR (100 MHz, CDCl₃) δ : 131.7, 131.6, 124.2, 124.2, 81.1, 80.7, 66.5, 60.9, 40.8, 40.5, 37.5, 36.2, 28.8, 28.4, 25.3, 25.2, 20.0, 18.0, 17.7, 17.7; IR (KBr) ν : 2964, 2920, 1556, 1455, 1383, 1095, 1052 cm⁻¹. HRMS calcd for C₁₁H₂₂NO₃ [M+H]⁺ 202.1438, found 202.1444.

(*E*)-1-Nitro-4-phenylbut-3-en-2-ol (**3s**): 32.5 mg, 56% yield, colorless oil; ¹H NMR (400 MHz, CDCl₃) δ : 7.40~7.29 (m, 5H), 6.83~6.75 (m, 1H), 6.15 (dd, $J=15.9, 6.3$ Hz, 1H), 5.06 (q, $J=5.7$ Hz, 1H), 4.55~4.48 (m, 2H), 2.75 (brs, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 135.4, 133.6, 128.7, 128.5, 126.7, 124.8, 79.8, 69.6; IR (KBr) ν : 3447, 1623, 1554, 1495, 1380, 1332, 970 cm⁻¹. HRMS calcd for C₁₀H₁₂NO₃ [M+H]⁺ 194.0812, found 194.0817.

1-(4'-(1-Hydroxy-2-nitroethyl)-[1,1'-biphenyl]-4-yl)ethan-1-one (**3t**): 52.2 mg, 61% yield, white solid, m.p. 129.8~131.0 °C; ¹H NMR (400 MHz, CDCl₃) δ : 8.03 (d, $J=8.3$ Hz, 2H), 7.67 (dd, $J=8.4, 2.4$ Hz, 4H), 7.52 (d, $J=8.2$ Hz, 2H), 5.55 (dd, $J=9.4, 3.0$ Hz, 1H), 4.66 (dd, $J=13.4, 9.5$ Hz, 1H), 4.57 (dd, $J=13.4, 3.0$ Hz, 1H), 2.64 (s, 3H), 1.68 (brs, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 197.8, 144.8, 140.5, 138.1, 136.1, 129.0, 127.9, 127.2, 126.6, 81.1, 70.6, 26.7; IR (KBr) ν : 3347, 2918, 1668, 1549, 1381, 1274, 819 cm⁻¹. HRMS calcd for C₁₆H₁₆NO₄ [M+H]⁺ 286.1074, found 286.1079.

Supporting Information ¹H NMR, ¹⁹F NMR, and ¹³C NMR spectra of products **3a**~**3t**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Ono, N. In *The Nitro Group in Organic Synthesis*, Wiley-VCH, New York, **2001**, p. 30.
(b) Norman, B. H.; Morris, M. L. *Tetrahedron Lett.* **1992**, 33, 6803.
(c) Kawabata, T.; Kiryu, Y.; Sugiyure, Y.; Fujii, K. *Tetrahedron Lett.* **1993**, 34, 5127.
(d) Sasai, H.; Kim, W.-S.; Suzuki, T.; Shibasaki, M. *Tetrahedron Lett.* **1994**, 35, 6123.
(e) Corey, E. J.; Zhang, F.-Y. *Angew. Chem. Int. Ed.* **1999**, 38, 1931.
(f) Grembecka, J.; Kafarski, P. *Mini Rev. Med. Chem.* **2001**, 1, 133.
- [2] Henry, L. *Bull. Soc. Chim. Fr.* **1895**, 13, 999.
- [3] (a) Rosini, G.; Ballini, R. *Synthesis* **1988**, 833.
(b) Shvekhgeimer, M. A. *Russ. Chem. Rev.* **1998**, 67, 35.
(c) Luzzio, F. A. *Tetrahedron* **2001**, 57, 915.
- [4] (a) Concellon, J. M.; Rodriguez-Solla, H.; Concellon, C.; Garcia-Granda, S.; Diaz, M. R. *Org. Lett.* **2006**, 8, 5979.
(b) Alcaide, B.; Almendros, P.; Luna, A.; Torres, M. R. *Org. Biomol. Chem.* **2008**, 6, 1635.
(c) Blay, G.; Hernandez-Olmos, V.; Pedro, J. R. *Chem. Commun.* **2008**, 4840.
(d) Leighty, M. W.; Shen, B.; Johnston, J. N. *J. Am. Chem. Soc.* **2012**, 134, 15233.
- [5] (a) Mao, P.; Yang, L.; Xiao, Y.; Yuan, J.; Mai, W.; Gao, J.; Zhang, X. *Chin. J. Org. Chem.* **2019**, 39, 443.
(f) Hu, Z.; Jiang, G.; Zhu, Z. Gong, B.; Xie, Z.; Le, Z. *Chin. J. Org. Chem.* **2021**, 41, 325.
- [6] (a) Concellon, J. M.; Rodriguez-Solla, H.; Concellon, C. *J. Org. Chem.* **2006**, 71, 7919.
(b) Soengas, R. G.; Estévez, A. M. *Eur. J. Org. Chem.* **2010**, 5190.
(c) Soengas, R. G.; Estévez, A. M. *Synlett* **2010**, 2625.
(d) Soengas, R. G.; Estévez, A. M. *Tetrahedron Lett.* **2012**, 53, 570.

- [7] Soengas, R. G.; Silva, A. M. S. *Synlett* **2012**, 873.
- [8] Mahasneh, A. S. Z. *Naturforsch.* **2005**, 60b, 416.
- [9] (a) Liu, Y.; Lu, Y.; Prashad, M.; Repic, O.; Blacklock, T. J. *Adv. Synth. Catal.* **2005**, 347, 217.
(b) Gao, G.; Tao, Y.; Jiang, J. *Green Chem.* **2008**, 10, 439.
(c) Dey, R.; Mukherjee, N.; Ahammed, S.; Ranu, B. C. *Chem. Commun.* **2012**, 48, 7982.
(d) Liu, X.-Y.; Cheng, B.-Q.; Guo, Y.-C.; Chu, X.-Q.; Rao, W.; Loh, T.-P.; Shen, Z.-L. *Org. Chem. Front.* **2019**, 6, 1581.
(e) Liu, X.-Y.; Li, X.-R.; Zhang, C.; Chu, X.-Q.; Rao, W.; Loh, T.-P.; Shen, Z.-L. *Org. Lett.* **2019**, 21, 5873.
(f) Chan, T. C.; Lau, C. P.; Chan, T. H. *Tetrahedron Lett.* **2004**, 45, 4189.
- [10] (a) Blümke, T. D.; Chen, Y.-H.; Peng, Z.; Knochel, P. *Nat. Chem.* **2010**, 2, 313.
(b) Chen, B.-Z.; Wang, C.-X.; Jing, Z.-H.; Chu, X.-Q.; Loh, T.-P.; Shen, Z.-L. *Org. Chem. Front.* **2019**, 6, 313.
- [11] (a) Blümke, T. D.; Klatt, T.; Koszinowski, K.; Knochel, P. *Angew. Chem. Int. Ed.* **2012**, 51, 9926.
(b) Takai, K.; Ikawa, Y. *Org. Lett.* **2002**, 4, 1727.
(c) Yun, J.-J.; Zhi, M.-L.; Shi, W.-X.; Chu, X.-Q.; Shen, Z.-L.; Loh, T.-P.; *Adv. Synth. Catal.* **2018**, 360, 2632.
(d) Shen, L.; Zhao, K.; Doitomi, K.; Ganguly, R.; Li, Y.-X.; Shen, Z.-L.; Hirao, H.; Loh, T.-P. *J. Am. Chem. Soc.* **2017**, 139, 13570.
- [12] (a) Ollevier, T. *Org. Biomol. Chem.* **2013**, 11, 2740.
(b) Wu, Z.; Feng, X.-X.; Wang, Q.-D.; Yun, J.-J.; Rao, W.; Yang, J.-M.; Shen, Z.-L. *Chin. Chem. Lett.* **2020**, 31, 1297.
- [13] (a) Chen, C.; Liu, P.; Luo, M.; Zeng, X. *ACS Catal.* **2018**, 8, 5864.
(b) Li, Y.; Deng, G.; Zeng, X. *Organometallics* **2016**, 35, 747.
(c) Yun, J.-J.; Liu, X.-Y.; Deng, W.; Chu, X.-Q.; Shen, Z.-L.; Loh, T.-P. *J. Org. Chem.* **2018**, 83, 10898.
- [14] (a) Enthaler, S.; Junge, K.; Beller, M. *Angew. Chem. Int. Ed.* **2008**, 47, 3317.
(b) Bauer, I.; Knölker, H.-J. *Chem. Rev.* **2015**, 115, 3170.
(c) Xiong, H.; Ramkumar, N.; Chiou, M.-F.; Jian, W.; Li, Y.; Su, J.-H.; Zhang, X.; Bao, H. *Nat. Commun.* **2019**, 10, 122.
(d) Wei, R.; Xiong, H.; Ye, C.; Li, Y.; Bao, H. *Org. Lett.* **2020**, 22, 3195.
(e) Deng, W.; Ye, C.; Li, Y.; Li, D.; Bao, H. *Org. Lett.* **2019**, 21, 261.
(f) He, Z.; Fan, M.; Xu, J.; Hu, Y.; Wang, L.; Wu, X.; Xia, C.; Liu, C. *Chin. J. Org. Chem.* **2019**, 39, 3438.
- [15] (a) Takai, K.; Kakiuchi, T.; Utimoto, K. *J. Org. Chem.* **1994**, 59, 2671.
(b) Cheng, B.-Q.; Zhao, S.-W.; Song, X.-D.; Chu, X.-Q.; Rao, W.; Loh, T.-P.; Shen, Z.-L. *J. Org. Chem.* **2019**, 84, 5348.
- [16] (a) Cheng, B.-Q.; Zhang, S.-X.; Cui, Y.-Y.; Chu, X.-Q.; Rao, W.; Xu, H.; Han, G.-Z.; Shen, Z.-L. *Org. Lett.* **2020**, 22, 5456.
(b) Chu, X.-Q.; Cheng, B.-Q.; Zhang, Y.-W.; Ge, D.; Shen, Z.-L.; Loh, T.-P. *Chem. Commun.* **2018**, 54, 2615.
(c) Wang, P.; Chen, B.-Z.; Guo, Y.-C.; Rao, W.; Shen, Z.-L. *Tetrahedron Lett.* **2019**, 60, 151288.
(d) Wu, L.-H.; Zhao, K.; Shen, Z.-L.; Loh, T.-P. *Org. Chem. Front.* **2017**, 4, 1872.
(e) Liu, B.; Xu, X.; Huang, L.; Feng, H. *Chin. J. Org. Chem.* **2020**, 40, 1290.
(f) Huang, S.; Nie, Y.; Yang, J.; Zheng, Z.; Cao, J.; Xu, Z.; Xu, L. *Chin. J. Org. Chem.* **2020**, 40, 2018.
- [17] Wu, Z.; Feng, X.-X.; Wang, Q.-D.; Liu, X.-Y.; Rao, W.; Yang, J.-M.; Shen, Z.-L. *Chin. Chem. Lett.* **2020**, 31, 391.
- [18] (a) Chen, B.-Z.; Zhi, M.-L.; Wang, C.-X.; Chu, X.-Q.; Shen, Z.-L.; Loh, T.-P. *Org. Lett.* **2018**, 20, 1902.
(b) Liu, X.-Y.; Cheng, B.-Q.; Guo, Y.-C.; Chu, X.-Q.; Li, Y.-X.; Loh, T.-P.; Shen, Z.-L. *Adv. Synth. Catal.* **2019**, 361, 542.
- [19] (a) Tay, N. E. S.; Chen, W.; Levens, A.; Pistritto, V. A.; Huang, Z.; Wu, Z.; Li, Z.; Nicewicz, D. A. *Nat. Catal.* **2020**, 3, 734.
(b) Bhattacharyya, A.; Kavitha, C. V.; Ghorai, M. K.; *J. Org. Chem.* **2016**, 81, 643.
(c) Kirihaara, M.; Osugi, R.; Saito, K.; Adachi, K.; Yamazaki, K.; Matsushima, R.; Kimura, Y. *J. Org. Chem.* **2019**, 84, 8330.

(Li, L.; Fan, Y.)