可见光促进钯催化 C—H 键胺化反应合成咔唑醌衍生物的研究

韩 阳[†] 姜为超[†] 张 靖 彭进松* 陈春霞*

(东北林业大学化学化工与资源利用学院 哈尔滨 150040)

摘要 以 2-氨基-3-芳基萘-1,4-二酮为底物, 发展了一类可见光/钯催化的分子内 C—H 键氧化胺化反应. 以氧气作为氧化剂及醋酸钯作为催化剂, 在蓝色 LED 光照射下, 2-氨基-3-芳基萘-1,4-二酮底物在室温下被转化为咔唑醌衍生物, 反应应具有良好的产率、官能团耐受性和优异的区域选择性.

关键词 光促进; 钯催化; 碳-氢键胺化; 咔唑醌

Visible-Light-Promoted Palladium-Catalyzed C—H Amination for the Synthesis of Carbazolequinones

Han, Yang[†] Jiang, Weichao[†] Zhang, Jing Peng, Jinsong* Chen, Chunxia*

(College of Chemistry, Chemical Engineering and Resource Utilization, Northeast Forestry University, Harbin 150040)

Abstract An intramolecular palladium-catalyzed cross-dehydrogenative C—H amination under blue LED irradiation at room temperature was developed. With oxygen as the external oxidant, a broad range of functional groups was compatible with the aromatic C—H amination of 2-amino-3-arylnaphthalene-1,4-diones, and diverse carbazolequinones can be obtained with good yields and excellent regioselectivity.

Keywords light-promoted; palladium-catalyzed; C—H amination; carbazolequinones

1 Introduction

The carbazolequinone motif is widely found in a diverse array of bioactive natural products (Figure 1), many carbazolequinone alkaloids (such as ellipticine quinone,^[1] murrayaquinones,^[2] calothrixins,^[3] and koeniginequinones^[4]) have exhibited various biological activities to act as anticancer, anti-inflammatory, antibacterial, and anti-infective properties agents. In addition, they are also valuable intermediates for the synthesis of carbazoles, carbazolequinols, and other alkaloids.^[5] Carbazolequinone alkaloids are the focus of intensive research due to their potential in the field of medicinal chemistry, therefore, much effort has been focused on the synthetic methods of carbazolequinone derivatives.^[6]

Using functionalized indoles or arylamines as precursors, various annulation strategies have been developed for the construction of carbazolequinone skeleton through cross-coupling reaction,^[6c-6h,7] radical chemistry,^[8] and metal-catalyzed^[9] or metal-free^[10] cascade reaction for the

formation of C—C or C—N bond. Over the last decades, transition metal-catalyzed C—H bond functionalization has emerged as popular and powerful tools for the assembly of complex polycyclic structures. Following this tendency, several attractive palladium-catalyzed C—H functionalization protocols have been discovered for the synthesis of carbazolequinones from the corresponding anilinoquinones (Scheme 1, paths a~c).^[6c-6h,7] The above methods have a number of drawbacks, such as the use of a stoichiometric amount of palladium or highly toxic and dangerous reagents, harsh reaction conditions, and the use of a large amount of copper salts as the terminal oxidant. Thus, novel synthetic approaches with enhanced reaction efficiency, milder and greener reaction conditions are still desirable.

Recently, photochemical C—H amination has emerged as an efficient and selective alternative for the construction of C—N bond by photoredox catalysis.^[11] This chemistry can be accomplished with the help of air or oxygen as the

* Corresponding authors. E-mail: ccx0109@nefu.edu.cn; jspeng1998@nefu.edu.cn

Received April 17, 2021; revised July 14, 2021; published online September 2, 2021.

Project supported by the Natural Science Foundation of Heilongjiang Province (No. ZD2021C001), and the Fundamental Research Funds for the Central Universities (No. 2572019CG06).

黑龙江省自然科学基金(No. ZD2021C001)和中央高校基本科研基金(No. 2572019CG06)资助项目.

[†] 共同第一作者(These authors contributed equally to this work).

Figure 1 Selected examples of biologically active carbazolequinones

Scheme 1 Palladium-catalyzed synthesis of carbazolequinones

green oxidant.^[12] Various photocatalytic C—N forming reactions have been applied in the construction of polycyclic structures.^[13] For instance, Cho and You *et al.*^[13b] reported a novel intramolecular C—N coupling reaction by merging Pd and photoredox catalysis to construct the *N*-substituted carbazole framework in 2015. Unfortunately,

2-aminobiaryl substrates did not undergo the C—H amination. This lack of reactivity was due to reductive quenching of the photoexcited catalysts by “free” aminobiaryls (Scheme 1d).

Although visible light-mediated metal-catalyzed C—H amination processes using dual metal/photoredox catalysts

have recently been reported,^[11] there are no examples of visible light-mediated, palladium-catalyzed C—H amination without photosensitizers. Inspired by Cho and You's study^[13b] and experience of photoredox catalysis for direct amination of aromatic C—H bonds with free amines,^[11,14] we designed a photosensitizer-free palladium-catalyzed protocol for the synthesis of carbazolequinones where an intramolecular C—H bond amination would take place by visible light irradiation of a solution of a “free” 2-amino-3-phenylnaphthalene-1,4-dione under an oxygen atmosphere (Scheme 1e).

2 Results and discussion

2-Amino-3-phenylnaphthalene-1,4-dione (**1a**), readily derived from the reaction of 2-aminonaphthoquinone and phenyl boronic acid in the presence of $\text{Mn}(\text{OAc})_3 \cdot 2\text{H}_2\text{O}$,^[15] was selected as a model substrate for visible light-promoted Pd-catalyzed C—H amination to construct the carbazolequinone (**2a**) (Table 1).

Table 1 Visible light-mediated Pd-catalyzed C—H amination condition optimization^a

Entry	Pd	LEDs	Solvent	Yield ^b /%
1	$\text{Pd}(\text{OAc})_2$	—	DMSO	0
2	—	Blue (4 W)	DMSO	0
3 ^c	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	DMSO	74
4	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	NMP	11
5	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	Methanol	34
6	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	DCM	0
7	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	DCE	0
8	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	THF	0
9	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	1,4-Dioxane	0
10	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	DMF	0
11	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	Acetone	0
12	$\text{Pd}(\text{OAc})_2$	Blue (4 W)	Toluene	0
13	PdCl_2	Blue (4 W)	DMSO	Trace
14	$\text{Pd}(\text{CF}_3\text{COO})_2$	Blue (4 W)	DMSO	10
15	$\text{Pd}_2(\text{dba})_3$	Blue (4 W)	DMSO	Trace
16	$\text{Pd}(\text{dppf})\text{Cl}_2$	Blue (4 W)	DMSO	Trace
17	$\text{Pd}(\text{PPh}_3)_4$	Blue (4 W)	DMSO	Trace
18	$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$	Blue (4 W)	DMSO	Trace
19 ^c	$\text{Pd}(\text{OAc})_2$	Blue (2 W)	DMSO	22
20 ^c	$\text{Pd}(\text{OAc})_2$	Blue (6 W)	DMSO	21
21 ^c	$\text{Pd}(\text{OAc})_2$	Blue (8 W)	DMSO	16
22 ^c	$\text{Pd}(\text{OAc})_2$	Blue (10 W)	DMSO	36
23	$\text{Pd}(\text{OAc})_2$	Purple (4 W)	DMSO	11
24	$\text{Pd}(\text{OAc})_2$	Green (4 W)	DMSO	18

^a Reaction conditions: **1a** (0.2 mmol), palladium catalyst (0.02 mmol, 10 mol%), solvent (1.0 mL), O_2 (101 kPa), 25–30 °C, 24 h. ^b Isolated yields after chromatography. ^c Under the same wavelength.

At first, the blank experiment (without the visible light irradiation or palladium catalyst) of **1a** was examined in dimethyl sulfoxide (DMSO) in the presence of oxygen at 25–30 °C for 24 h, and no desired product **2a** was obtained (Entries 1 and 2, Table 1). Then the model reaction was examined in the presence of $\text{Pd}(\text{OAc})_2$ (10 mol%) under irradiation with blue LEDs (4 W) in DMSO, and fortunately, the desired product **2a** was indeed formed in 74% yield (Entry 3, Table 1). Next, different solvents were evaluated using $\text{Pd}(\text{OAc})_2$ as the catalyst and 4 W blue LEDs light source (Entries 4–12, Table 1). *N*-Methylpyrrolidone (NMP) and methanol were inferior to DMSO, affording the product **2a** in low yields (11% and 34% yields, respectively, Entries 4 and 5, Table 1). Other solvents, such as dichloromethane (DCM), 1,2-dichloroethane (DCE), tetrahydrofuran (THF), 1,4-dioxane, *N,N*-dimethyl formamide (DMF), acetone and toluene, gave no desired product (Entries 6–12, Table 1).

The effect of palladium catalysts bearing various ligands, such as PdCl_2 , $\text{Pd}(\text{CF}_3\text{COO})_2$, $\text{Pd}_2(\text{dba})_3$, $\text{Pd}(\text{dppf})\text{Cl}_2$, $\text{Pd}(\text{PPh}_3)_4$, and $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$, was then investigated on the reaction (Entries 13–18, Table 1). Compared to $\text{Pd}(\text{OAc})_2$, the use of $\text{Pd}(\text{CF}_3\text{COO})_2$ gave a lower yield (Entry 14, Table 1). With a palladium(0) or palladium(II) complex possessing a phosphine ligand, the product **2a** was formed only in a trace quantity (Entries 15–18, Table 1). Finally, a set of experiments were carried out to reveal the crucial role of the LEDs light source (power and wave length). The results showed that 4 W blue LEDs provided better yield than blue LEDs (2, 6 and 10 W), green LEDs (4 W), and purple LEDs (4 W) (Entries 19–24, Table 1). The yield of **2a** was depended on both decomposition and conversion rate of **1a**. Under the same wavelength, the rate of substrate decomposition increased along with the increase of light power, leading to a decrease in the yield of **2a**. However, 10 W blue LEDs provided better yield than blue LEDs (2, 6, 8 W), which was related to higher conversion of **1a**.

Having established the feasibility of this Pd-catalyzed intramolecular C—H amination under visible light irradiation (Entry 3, Table 1), we next focused on the expansion of the substrate scope by investigating a range of 2-amino-3-arylnaphthalene-1,4-diones (Table 2). For 3-aryl moiety of substrates **1**, a variety of substituents (F, Cl, Br, I, CF_3 , alkyl, alkenyl, aryl, alkoxy, phenoxy, and methylthio) were applicable, and both electron-withdrawing (**1b**–**1f**, **1r**–**1t**, and **1v**) and electron-donating (**1g**–**1q**, **1u**, **1w**–**1y**, **1bb**, and **1bc**) substituents can be incorporated at the *para*-, *meta*-, or *ortho*-position, affording carbazolequinone products **2b**–**2z**, and **2ba**–**2bc** in 15%–93% yields. The most of halo-substituted 2-aminonaphthoquinone substrates can smoothly be converted to the desired products in good yields (46%–77%), however, the use of aryl iodide **1e** to effect this transformation afforded an inferior result than fluoro, chloro, or bromo analogues, only 23% isolated yield was obtained.

Table 2 Visible-light-induced Pd-catalyzed synthesis of carbazolequinones^{a,b}

^a Reaction conditions: **1** (0.2 mmol), $\text{Pd}(\text{OAc})_2$ (0.02 mmol, 10 mol%), 4 W blue LEDs, DMSO (1.0 mL), O_2 (101 kPa), 25 ~ 30 °C, 24 h. ^b Isolated yields after chromatography. ^c One millimole scale.

For hydrocarbon substitutions, aliphatic primary, secondary, and tertiary alkyl-substituted substrates **1g**~**1i** were generally more reactive than those bearing an alkenyl (**1j**) or aromatic substituent (**1k**). For various alkoxy substitutions, methoxy substrate **1l** afforded a corresponding product **2l** in 93% yield, however, other analogues **1m**~**1p** generated the desired products **2m**~**2p** in only moderate yields (35%~38%). Regioselectivity issues may po-

tentially exist in the C—H amination process when functional groups are selectively introduced at the *meta* positions of the aryl moiety. The influence of sterics and electronics of the substituent (F, Cl, Br, CF_3 , Me, OMe, and OCF_3) on the regioselectivity was then examined. The incorporation of the above groups did not hamper the reaction, and the corresponding products **2r**~**2x** can be obtained in 31%~89% yields. Interestingly, although there

are two possible C—H activation sites, the amination reaction in all the examined cases occurred only at the most sterically accessible site to give single regioisomers. Similarly, β -naphthyl substrate **1z** also provided a single product **2z** through β -site specific C—H amination. Disubstituted substrates (dimethyl and dimethoxy, **1bb** and **1bc**) were finally investigated, the corresponding products (**2bb** and **2bc**) were obtained in 80% and 59% yield respectively.

To gain insight into the mechanism of the reaction, some designed control experiments were conducted under standard reaction conditions (Scheme 2). The light on/off experiment was firstly carried out (Scheme 2a), and the result showed that the conversion of **1a** ceased upon removal from irradiation and continued upon additional irradiation, and the continuous light irradiation was essential for the completion of the reaction. The above result indicated that the reaction may be amenable to photo-induced palladium catalysis rather than just photo-initiated chain process. UV/Vis absorption spectra studies were then conducted (Scheme 2b), the absorbance of **1a** gradually increased at the wavelength of 465 nm with the introduction of palladium acetate and the radiation of the blue LEDs, proving that **1a** can be gradually excited. At a wavelength of 378 nm, the absorbance of the mixture was similar to that of the product **2a**, indicating a new substance had been formed. Finally, when two equivalents of 2,2,6,6-tetram-

ethylpiperidinoxy (TEMPO) were added into the reaction mixture under standard reaction conditions, the reaction was shut down and the substrate **1a** was completely recovered from the reaction mixture (Scheme 2c), indicating that radical species was abolished by radical scavenger TEMPO.

On the basis of the above results, a possible catalytic cycle involved in the intramolecular C—H amination process is outlined in Scheme 3. The initial step involved the coordination of 2-amino-3-arylnaphthalene-1,4-dione **1** to the Pd(II) center to form the complex **3**, which was then excited into Pd^{II}* species **3*** under blue LED irradiation. Next, the superoxide anion radical was generated from O₂ by oxidative quenching of the excited state of Pd^{II}* **3*** to Pd^{III} species **4** (Scheme 3, cycle A).^[16] In the catalytic cycle B, an intermediate complex **5** was generated from the deprotonation of N—H and the dissociation of acetate anion of Pd(II) complex **3**.^[13b] Subsequently, the aromatic ring reacted with the Pd(II) center to afford a six-membered palladacycle **6** via a concerted metalation-deprotonation (CMD) transition state.^[17] Palladacycle intermediate **6** underwent C—N bond-forming reductive elimination to afford the desired product **2** and a Pd⁰ species **7**, and the latter was finally oxidized to generate a Pd^{II} species by Pd^{III} species **4** and the superoxide anion radical in the photoredox process.

Scheme 2 Investigation of the reaction mechanism

Scheme 3 Proposed mechanism for visible-light-promoted Pd-catalyzed C—H amination

3 Conclusions

In conclusion, we have developed a mild and efficient method for the synthesis of diverse carbazolequinones, which involves an intramolecular aromatic C—H amination of 2-amino-3-arylnaphthalene-1,4-dione derivatives. The desired catalytic process is modulated by blue LED light, and does not require additional photosensitizers, and two distinct catalytic principles can operate in the presence of a single palladium catalyst. Furthermore, the above described synthetic strategy combines photocatalysis and transition-metal catalysis, avoiding the use of potentially toxic chemical additives and strong oxidants. Further investigations of other visible-light-induced metal-catalyzed reactions without photosensitizers are ongoing in our group.

4 Experimental section

4.1 General information

Chemicals were all purchased from commercial supplies and used without further purification unless otherwise stated. Solvents were dried and purified according to the standard procedures before use. Reactions were monitored by analytical thin-layer chromatography (TLC). All reactions were conducted in dried glassware. Purification of reaction products was done by flash chromatography with 230~400 mesh silica gel. All the 2-amino-3-arylnaphthalene-1,4-dione substrates were prepared according to literature methods.^[15] **1h**, **1n**, and **1x** are new compounds, other substrates are known compounds, and the spectroscopic and physical data are matched with those from the literature. “RLH-18 8-position Photo Reaction System” was

used, which manufactured by Beijing Rogertech Co. Ltd. The wavelength of blue light is 453.6 nm, peak width at half-height is 20.4 nm. Irradiation vessel is borosilicate glass test tube, LED irradiate through a high-reflection channel to the test tube, and path length is 2 cm. Melting points were determined on a melting point apparatus in open capillaries and were uncorrected. ¹H NMR spectra were recorded on a 500 MHz spectrometer, and ¹³C NMR spectra were recorded at 126 MHz. Unless otherwise stated, deuterated dimethyl sulfoxide (DMSO-*d*₆) was used as a solvent. Chemical shifts (δ) were referenced to tetramethylsilane (TMS). Chemical shifts for carbon resonances were referenced to the carbon resonance of the solvent DMSO-*d*₆ (δ 39.52). High-resolution mass spectra were recorded on a BIO TOF Q mass spectrometer equipped with an electrospray ion source (ESI), operated in the positive (**1h**, **1x**, **2h**, **2i**, **2k**, **2m**, **2o**~**2q**, **2y**, **2ba**~**2bc**) or negative mode (**1n**, **2d**, **2e**, **2j**, **2n**, **2t**, **2x**, **2z**).

4.2 Preparation of 2-amino-3-arylnaphthalene-1,4-diones **1**

The reaction was carried out using 2-aminonaphthoquinone (80 mg, 0.462 mmol), arylboronic acid (0.693 mmol, 1.5 equiv.), Mn(OAc)₃·2H₂O (248 mg, 0.925 mmol, 2.0 equiv.) in DMSO (4 mL) at 100 °C for 24 h. The substrates **1h**, **1n**, and **1x** were obtained as orange red solid after passing through a short silica gel column, eluting with 15% ethyl acetate/petroleum ether.^[15]

2-([1,1'-Biphenyl]-4-yl)-3-aminonaphthalene-1,4-dione (**1h**): Yield 62% (93.1 mg, Eluent: EtOAc/petroleum ether, *V*: *V*=1:7); red solid, m.p. 237~240 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 8.06~7.99 (m, 2H), 7.85 (td, *J*=7.6, 1.3 Hz, 1H), 7.79~7.72 (m, 5H), 7.51 (t, *J*=7.7 Hz, 2H),

7.41~7.38 (m, 3H), 6.67 (br, 2H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 182.2, 180.7, 146.8, 140.5, 139.4, 135.3, 133.4, 133.0, 132.8, 131.6, 130.7, 129.5, 127.9, 127.07, 127.06, 126.2, 126.0, 114.6; IR (KBr) ν : 3481, 3352, 1676, 1574, 1434, 1385, 1348, 1022, 726, 699 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{15}\text{NNaO}_2$ $[\text{M} + \text{Na}]^+$ 348.0995, found 348.0995.

2-Amino-3-(4-ethoxyphenyl)naphthalene-1,4-dione (**1n**): Yield 42% (56.8 mg, Eluent: EtOAc/petroleum ether, $V: V=1:7$); dark red solid, m.p. 212~225 $^\circ\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 8.00~7.96 (m, 2H), 7.84~7.81 (m, 1H), 7.76~7.72 (m, 1H), 7.24~7.15 (m, 2H), 7.03~6.96 (m, 2H), 6.49 (br, 2H), 4.07 (q, $J=7.0$ Hz, 2H), 1.36 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 182.2, 180.8, 158.1, 146.7, 135.1, 133.3, 132.7, 132.1, 130.7, 126.2, 125.9, 125.5, 115.0, 114.7, 63.4, 15.2; IR (KBr) ν : 3460, 3350, 1671, 1557, 1346, 1249, 1046, 1001, 723 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{NO}_3$ $[\text{M}-\text{H}]^-$ 292.0979, found 292.0980.

2-Amino-3-[3-(trifluoromethoxy)phenyl]naphthalene-1,4-dione (**1x**): Yield 64% (98.5 mg, Eluent: EtOAc/petroleum ether, $V: V=1:7$); red solid, m.p. 156~160 $^\circ\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 8.02 (d, $J=7.5$ Hz, 1H), 7.99 (d, $J=7.5$ Hz, 1H), 7.85 (t, $J=7.4$ Hz, 1H), 7.76 (t, $J=7.4$ Hz, 1H), 7.58 (t, $J=7.9$ Hz, 1H), 7.33 (t, $J=9.0$ Hz, 2H), 7.24 (s, 1H), 6.87 (br, 2H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 182.1, 180.3, 148.9, 147.1, 136.2, 135.3, 133.3, 132.8, 130.7, 130.2, 126.2, 126.1, 126.0, 123.7, 121.7, 120.1, 113.2; IR (KBr) ν : 3492, 3355, 1678, 1601, 1567, 1245, 1006, 726, 698 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{10}\text{F}_3\text{NO}_3\text{Na}$ $[\text{M}-\text{H}]^+$ 356.0505, found 356.0505.

4.3 General procedure for the synthesis of carbazolequinones **2**

A 10 mL Schlenk tube equipped with a magnetic stirring bar was charged with 2-amino-3-arylnaphthalene-1,4-dione (**1**, 0.2 mmol, 1.0 equiv.), and then $\text{Pd}(\text{OAc})_2$ (0.02 mmol, 4.5 mg) was added. Finally, DMSO (1.0 mL) was added to the mixture via syringe at room temperature under 101 kPa of O_2 . The tube was sealed and irradiated with 4 W blue LED for 24 h at room temperature. The mixture was cooled to room temperature, quenched with water (5 mL), and diluted with dichloromethane (5 mL). The layers were separated, and the aqueous layer was extracted with 5 mL $\times$ 2 of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. The residue was purified by flash chromatography to afford the product carbazolequinone **2**, eluting with 15%~30% ethyl acetate/petroleum ethers.

5*H*-Benzo[*b*]carbazole-6,11-dione (**2a**): The reaction was carried out on a 1.0 mmol scale. Yield 52% (120.8 mg, Eluent: EtOAc/petroleum ether, $V: V=1:3$); red solid, m.p. 301~303 $^\circ\text{C}$ (lit.^[7f] 301~303 $^\circ\text{C}$); ^1H NMR (500 MHz, DMSO- d_6) δ : 13.07 (s, 1H), 8.21 (d, $J=8.0$ Hz, 1H), 8.11 (t, $J=8.4$ Hz, 2H), 7.83~7.79 (m, 2H), 7.59 (d, $J=8.3$ Hz, 1H), 7.44 (t, $J=7.6$ Hz, 1H), 7.86 (t, $J=7.5$ Hz,

1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.8, 178.0, 138.7, 137.6, 134.6, 134.5, 133.6, 133.1, 127.4, 126.5, 126.4, 124.4, 124.3, 122.8, 117.8, 114.3.

3-Fluoro-5*H*-benzo[*b*]carbazole-6,11-dione (**2b**): Yield 54% (28.6 mg, Eluent: EtOAc/petroleum ether, $V: V=1:6$); orange solid, m.p. 260~262 $^\circ\text{C}$ (lit.^[7f] 262~264 $^\circ\text{C}$); ^1H NMR (500 MHz, DMSO- d_6) δ : 13.07 (s, 1H), 8.14~8.11 (m, 1H), 8.05~8.02 (m, 2H), 7.83~7.76 (m, 2H), 7.25~7.17 (m, 2H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.7, 177.6, 161.9 (d, $J_{\text{C-F}}=242.5$ Hz), 139.0 (d, $J_{\text{C-F}}=13.0$ Hz), 138.4 (d, $J_{\text{C-F}}=2.7$ Hz), 134.6, 134.2, 133.7, 132.9, 126.5, 126.4, 124.5 (d, $J_{\text{C-F}}=10.4$ Hz), 121.1, 117.9, 113.5 (d, $J_{\text{C-F}}=25.0$ Hz), 100.0 (d, $J_{\text{C-F}}=25.9$ Hz); ^{19}F NMR (376 MHz, DMSO) δ : -118.07.

3-Chloro-5*H*-benzo[*b*]carbazole-6,11-dione (**2c**):^[18] Yield 77% (40.7 mg, Eluent: EtOAc/petroleum ether, $V: V=1:6$); orange-red solid, m.p. 370~372 $^\circ\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.17 (s, 1H), 8.16 (d, $J=8.6$ Hz, 1H), 8.09 (t, $J=6.6$ Hz, 2H), 7.88~7.80 (m, 2H), 7.56 (s, 1H), 7.38 (d, $J=8.5$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.7, 177.9, 139.0, 138.4, 134.8, 134.3, 133.8, 133.0, 131.8, 126.6, 126.5, 124.9, 124.3, 123.0, 117.7, 113.7.

3-Bromo-5*H*-benzo[*b*]carbazole-6,11-dione (**2d**): Yield 46% (30.1 mg, Eluent: EtOAc/petroleum ether, $V: V=1:6$); orange-red solid, m.p. 234~236 $^\circ\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.18 (s, 1H), 8.13~8.09 (m, 3H), 7.89~7.82 (m, 2H), 7.73 (s, 1H), 7.50 (d, $J=8.6$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.7, 177.9, 139.4, 138.3, 134.8, 134.4, 133.8, 133.1, 127.5, 126.62, 126.57, 124.6, 123.2, 120.1, 117.8, 116.8; IR (KBr) ν : 3417, 1594, 1563, 1444, 1424, 1351, 1131, 726 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{16}\text{H}_7\text{BrNO}_2$ $[\text{M}-\text{H}]^-$ 323.9666, found 323.9665.

3-Iodo-5*H*-benzo[*b*]carbazole-6,11-dione (**2e**): Yield 23% (17.1 mg, Eluent: EtOAc/petroleum ether, $V: V=1:5$); orange solid, m.p. 180~182 $^\circ\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.15 (s, 1H), 8.12 (t, $J=6.6$ Hz, 2H), 8.01 (d, $J=8.4$ Hz, 1H), 7.95 (s, 1H), 7.89~7.82 (m, 2H), 7.67 (d, $J=8.4$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.8, 178.0, 139.8, 137.9, 134.9, 134.5, 133.9, 133.1, 132.9, 126.7, 126.6, 124.6, 123.5, 122.8, 117.8, 92.7; IR (KBr) ν : 3237, 1724, 1643, 1288, 1146, 1073, 753, 710 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{16}\text{H}_7\text{INO}_2$ $[\text{M}-\text{H}]^-$ 371.9527, found 371.9527.

3-(Trifluoromethyl)-5*H*-benzo[*b*]carbazole-6,11-dione (**2f**): Yield 81% (51.5 mg, Eluent: EtOAc/petroleum ether, $V: V=1:5$); yellow solid, m.p. 268~270 $^\circ\text{C}$ (lit.^[7f] 268~270 $^\circ\text{C}$); ^1H NMR (500 MHz, DMSO- d_6) δ : 13.44 (s, 1H), 8.45 (s, 1H), 8.10 (t, $J=6.3$ Hz, 2H), 7.90~7.82 (m, 2H), 7.75 (q, $J=10.0$ Hz, 2H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.7, 178.1, 139.9, 137.4, 137.1, 135.0, 134.4, 133.9, 133.0, 127.1, 126.72, 126.71, 124.9 (q, $J_{\text{C-F}}=272.2$ Hz), 124.0, 120.3 (q, $J_{\text{C-F}}=2.5$ Hz), 117.4, 111.6 (q, $J_{\text{C-F}}=3.8$ Hz); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -59.72.

3-Methyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2g**): Yield

91% (47.7 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); orange solid, m.p. 271~273 °C (lit.^[7] 271~274 °C); ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.92 (s, 1H), 8.09~8.05 (m, 3H), 7.85~7.78 (m, 2H), 7.35 (s, 1H), 7.18 (d, $J=8.1$ Hz, 1H), 2.44 (s, 3H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.9, 177.9, 139.2, 137.4, 137.3, 134.6, 134.5, 133.6, 133.2, 126.5, 126.4, 122.5, 122.3, 118.0, 113.7, 22.1.

3-Ethyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2h**): Yield 39% (21.3 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); red solid, m.p. 243~245 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.96 (s, 1H), 8.11~8.09 (m, 3H), 7.88~7.79 (m, 2H), 7.38 (s, 1H), 7.25 (d, $J=8.2$ Hz, 1H), 2.76 (q, $J=7.5$ Hz, 2H), 1.25 (t, $J=7.6$ Hz, 3H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.9, 177.9, 143.9, 139.3, 137.4, 134.6, 134.5, 133.7, 133.2, 126.5, 126.4, 125.5, 122.7, 122.5, 118.0, 112.5, 29.1, 16.2; IR (KBr) ν : 3425, 3298, 1663, 1635, 1590, 1531, 1466, 1400, 1228, 715 cm⁻¹. HRMS (ESI) calcd for C₁₈H₁₄NO₂ [M+H]⁺ 276.1019, found 276.1023.

3-(*tert*-Butyl)-5*H*-benzo[*b*]carbazole-6,11-dione (**2i**): Yield 48% (27.9 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); orange solid, m.p. 292~294 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.93 (s, 1H), 8.10 (dd, $J=13.9$, 7.6 Hz, 3H), 7.83 (dd, $J=14.0$, 7.4 Hz, 2H), 7.58~7.22 (m, 2H), 1.36 (s, 9H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.8, 177.8, 150.7, 139.1, 137.5, 134.6, 134.5, 133.7, 133.2, 126.5, 126.4, 123.2, 122.4, 122.2, 117.9, 109.7, 35.4, 31.6; IR (KBr) ν : 3241, 1650, 1590, 1525, 1469, 1398, 1333, 1228, 762 cm⁻¹. HRMS (ESI) calcd for C₂₀H₁₈NO₂ [M+H]⁺ 304.1332, found 304.1333.

3-Vinyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2j**): Yield 31% (16.4 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); orange solid, m.p. 273~277 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 13.12 (s, 1H), 8.17 (d, $J=8.4$ Hz, 1H), 8.12 (t, $J=7.1$ Hz, 2H), 7.89~7.82 (m, 2H), 7.66~7.53 (m, 2H), 6.91 (dd, $J=17.6$, 10.9 Hz, 1H), 5.93 (d, $J=17.6$ Hz, 1H), 5.36 (d, $J=11.0$ Hz, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.9, 177.8, 139.1, 138.1, 137.3, 136.7, 134.7, 134.5, 133.8, 133.2, 126.6, 126.5, 124.2, 122.9, 122.5, 115.4, 115.3, 112.2; IR (KBr) ν : 2975, 2723, 1617, 1459, 1376, 1161, 1042, 950, 722 cm⁻¹. HRMS (ESI) calcd for C₁₈H₁₀NO₂ [M-H]⁻ 272.0717, found 272.0717.

3-Phenyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2k**): Yield 28% (18.1 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); red solid, m.p. 227~230 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 13.16 (s, 1H), 8.27 (d, $J=8.4$ Hz, 1H), 8.12 (t, $J=7.7$ Hz, 2H), 7.88~7.81 (m, 2H), 7.76 (s, 1H), 7.73 (d, $J=7.5$ Hz, 2H), 7.69 (d, $J=8.4$ Hz, 1H), 7.52 (t, $J=7.6$ Hz, 2H), 7.43 (d, $J=7.3$ Hz, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.9, 177.9, 140.7, 139.7, 139.4, 138.2, 134.7, 134.5, 133.7, 133.2, 129.6, 128.2, 127.5, 126.6, 126.5, 124.1, 123.7, 123.3, 117.9, 111.8; IR (KBr) ν : 3431, 3258, 1651, 1563, 1385, 1152, 1004, 762 cm⁻¹. HRMS (ESI) calcd for C₂₂H₁₄NO₂ [M+H]⁺ 324.1019, found 324.1018.

3-Methoxy-5*H*-benzo[*b*]carbazole-6,11-dione (**2l**):^[18] Yield 93% (51.5 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); red solid, m.p. 288~290 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.92 (s, 1H), 8.10~8.06 (m, 3H), 7.88~7.77 (m, 2H), 7.02 (d, $J=8.9$ Hz, 1H), 6.98 (s, 1H), 3.85 (s, 3H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 181.0, 177.3, 159.9, 140.3, 136.8, 134.5, 134.4, 133.7, 133.2, 126.5, 126.4, 123.7, 118.5, 118.4, 116.0, 95.5, 55.9.

3-(Trifluoromethoxy)-5*H*-benzo[*b*]carbazole-6,11-dione (**2m**): Yield 36% (23.8 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); orange solid, m.p. 281~283 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 13.28 (s, 1H), 8.27 (d, $J=8.7$ Hz, 1H), 8.10 (t, $J=6.8$ Hz, 2H), 7.88~7.81 (m, 2H), 7.48 (s, 1H), 7.35 (d, $J=8.7$ Hz, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.7, 177.8, 147.5, 139.1, 138.5, 134.9, 134.3, 133.9, 133.0, 126.62, 126.6, 124.5, 123.1, 120.6 (q, $J_{C-F}=257$ Hz), 118.2, 117.6, 106.5; IR (KBr) ν : 3280, 1656, 1581, 1532, 1469, 1280, 1168, 1137, 715 cm⁻¹. HRMS (ESI) calcd for C₁₇H₉F₃NO₃ [M+H]⁺ 332.0529, found 332.0529; ¹⁹F NMR (376 MHz, DMSO) δ : -56.98.

3-Ethoxy-5*H*-benzo[*b*]carbazole-6,11-dione (**2n**): Yield 38% (23.7 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); red solid, m.p. 215~217 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.84 (s, 1H), 8.08~8.02 (m, 3H), 7.87~7.76 (m, 2H), 7.01~6.90 (m, 2H), 4.08 (q, $J=6.8$ Hz, 2H), 1.38 (t, $J=6.8$ Hz, 3H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 181.0, 177.2, 159.1, 140.3, 136.8, 134.4, 133.7, 133.2, 126.4, 126.3, 123.7, 118.5, 118.4, 116.2, 96.1, 79.6, 63.9, 15.1; IR (KBr) ν : 3229, 1644, 1633, 1434, 1261, 1170, 1045, 1012, 708 cm⁻¹. HRMS (ESI) calcd for C₁₈H₁₂NO₃ [M-H]⁻ 290.0823, found 290.0824.

3-Isopropoxy-5*H*-benzo[*b*]carbazole-6,11-dione (**2o**): Yield 36% (22.2 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); red solid, m.p. 277~279 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.81 (s, 1H), 8.10~8.02 (m, 3H), 7.85~7.78 (m, 2H), 6.99~6.97 (m, 2H), 4.71~4.65 (m, 1H), 1.33 (d, $J=6.0$ Hz, 6H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 181.0, 177.3, 157.9, 140.3, 136.9, 134.4, 134.3, 133.7, 133.2, 126.4, 126.3, 123.8, 118.5, 118.4, 116.9, 97.7, 70.3, 22.2; IR (KBr) ν : 3235, 1644, 1619, 1526, 1436, 1260, 1010, 824, 733 cm⁻¹. HRMS (ESI) calcd for C₁₉H₁₆NO₃ [M+H]⁺ 306.1125, found 306.1128.

3-Phenyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2p**): Yield 35% (23.7 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); red solid, m.p. 258~260 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 12.91 (s, 1H), 8.19 (d, $J=8.7$ Hz, 1H), 8.12~8.06 (m, 2H), 7.88~7.80 (m, 2H), 7.45 (t, $J=7.9$ Hz, 2H), 7.22 (t, $J=7.4$ Hz, 1H), 7.15~7.11 (m, 3H), 7.02 (d, $J=1.9$ Hz, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ : 180.9, 177.5, 157.1, 156.8, 139.7, 137.8, 134.7, 134.4, 133.8, 133.1, 130.7, 126.54, 126.47, 124.5, 124.4, 120.4, 119.7, 118.1, 117.5, 102.1; IR (KBr) ν : 3415, 3241, 1643, 1567, 1528, 1433, 1261, 1139, 1011, 727 cm⁻¹. HRMS (ESI) calcd for C₂₂H₁₄NO₃ [M+H]⁺ 340.0968, found 340.0964.

3-(Methylthio)-5*H*-benzo[*b*]carbazole-6,11-dione (**2q**):

Yield 54% (31.4 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); red solid, m.p. 263~265 °C; ^1H NMR (500 MHz, DMSO- d_6) δ : 12.92 (s, 1H), 8.07~8.04 (m, 3H), 7.84~7.78 (m, 2H), 7.35 (s, 1H), 7.24 (d, $J=8.5$ Hz, 1H), 2.54 (s, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.8, 177.6, 139.4, 138.2, 137.2, 134.6, 134.4, 133.7, 133.2, 126.5, 126.4, 123.6, 123.0, 121.8, 118.1, 109.8, 15.4; IR (KBr) ν : 3216, 1648, 1562, 1528, 1427, 1332, 1232, 1006, 711 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{12}\text{NO}_2\text{S} [\text{M}+\text{H}]^+$ 294.0583, found 294.0582.

2-Fluoro-5*H*-benzo[*b*]carbazole-6,11-dione (**2r**): Yield 31% (16.4 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); yellow solid, m.p. 260~262 °C (lit.^[7] 260~262 °C); ^1H NMR (500 MHz, DMSO- d_6) δ : 13.22 (s, 1H), 8.11 (t, $J=6.7$ Hz, 2H), 7.89~7.81 (m, 3H), 7.61 (dd, $J=9.0, 4.4$ Hz, 1H), 7.35~7.32 (m, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.6, 177.9, 159.9 (d, $J_{\text{C-F}}=239$ Hz), 138.9, 135.4, 134.8, 134.5, 133.8, 133.1, 126.6, 124.7 (d, $J_{\text{C-F}}=11$ Hz), 117.8 (d, $J_{\text{C-F}}=5.4$ Hz), 116.4 (d, $J_{\text{C-F}}=27$ Hz), 116.1 (d, $J_{\text{C-F}}=10$ Hz), 107.2, 107.0; ^{19}F NMR (376 MHz, DMSO) δ : -115.13.

2-Chloro-5*H*-benzo[*b*]carbazole-6,11-dione (**2s**): Yield 62% (34.8 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); yellow solid, m.p. 318~320 °C (lit.^[7] 318~320 °C); ^1H NMR (500 MHz, DMSO- d_6) δ : 13.27 (s, 1H), 8.14 (d, $J=1.8$ Hz, 1H), 8.13~8.08 (m, 2H), 7.89~7.81 (m, 2H), 7.61 (d, $J=8.8$ Hz, 1H), 7.47 (dd, $J=8.8, 2.1$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.6, 178.0, 138.7, 137.1, 134.9, 134.4, 133.9, 133.1, 129.0, 127.6, 126.6, 125.2, 121.6, 117.2, 116.2.

2-Bromo-5*H*-benzo[*b*]carbazole-6,11-dione (**2t**): Yield 40% (25.8 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); yellow solid, m.p. 251~255 °C; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.26 (s, 1H), 8.29 (s, 1H), 8.10 (t, $J=6.7$ Hz, 2H), 7.89~7.81 (m, 2H), 7.56 (q, $J=8.8$ Hz, 2H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.7, 177.9, 137.3, 134.9, 134.4, 133.9, 133.1, 130.1, 130.0, 126.6, 125.9, 124.7, 117.0, 116.5; IR (KBr) ν : 3427, 1635, 1568, 1385, 1012, 877, 710 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{16}\text{H}_7\text{BrNO}_2 [\text{M}-\text{H}]^-$ 323.9666, found 323.9667.

2-Methyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2u**): Yield 55% (28.7 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); yellow solid, m.p. 258~260 °C (lit.^[6,7] 258~260 °C); ^1H NMR (500 MHz, DMSO- d_6) δ : 12.95 (s, 1H), 8.07 (t, $J=8.2$ Hz, 2H), 7.96 (s, 1H), 7.83 (t, $J=7.3$ Hz, 1H), 7.79 (t, $J=7.3$ Hz, 1H), 7.45 (d, $J=8.4$ Hz, 1H), 7.24 (d, $J=8.4$ Hz, 1H), 2.44 (s, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.7, 177.9, 137.5, 137.1, 134.6, 134.58, 133.8, 133.6, 133.1, 129.3, 126.5, 126.4, 124.7, 122.1, 117.4, 114.0, 21.8.

3-(Trifluoromethyl)-5*H*-benzo[*b*]carbazole-6,11-dione (**2v**): Yield 40% (25.2 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); orange solid, m.p. 268~270 °C (lit.^[7,18] 268~270 °C); ^1H NMR (500 MHz, DMSO- d_6) δ : 13.40 (s, 1H), 8.31 (d, $J=8.4$ Hz, 1H), 8.08~8.05 (m, 2H), 7.86~7.70 (m, 3H), 7.60 (d, $J=8.4$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.6, 178.1, 139.8, 137.4, 135.0,

134.4, 133.9, 133.0, 127.3, 126.7, 126.6, 126.5 (128.1, 127.0, 126.0, q, $J_{\text{C-F}}=131$ Hz), 124.0, 123.8, 120.3 (d, $J_{\text{C-F}}=3.8$ Hz), 117.4, 111.6 (d, $J_{\text{C-F}}=5.1$ Hz); ^{19}F NMR (376 MHz, DMSO) δ : -60.13; IR (KBr) ν : 3240, 1656, 1594, 1403, 1341, 1230, 1112, 709 cm^{-1} .

2-Methoxy-5*H*-benzo[*b*]carbazole-6,11-dione (**2w**): Yield 89% (48.9 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); orange solid, m.p. 296~298 °C (lit.^[6] 297~298 °C); ^1H NMR (500 MHz, DMSO- d_6) δ : 12.97 (s, 1H), 8.06 (t, $J=7.1$ Hz, 2H), 7.82 (t, $J=7.3$ Hz, 1H), 7.78 (t, $J=7.3$ Hz, 1H), 7.56 (s, 1H), 7.45 (d, $J=9.0$ Hz, 1H), 7.07~7.02 (m, 1H), 3.84 (s, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.6, 177.6, 157.4, 137.4, 134.6, 134.5, 133.9, 133.5, 133.2, 126.5, 126.4, 125.3, 118.8, 117.5, 115.4, 102.5, 55.8.

3-(Trifluoromethoxy)-5*H*-benzo[*b*]carbazole-6,11-dione (**2x**): Yield 62% (41.1 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); orange solid, m.p. 323~326 °C; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.31 (s, 1H), 8.10 (d, $J=2.8$ Hz, 2H), 8.04 (s, 1H), 7.88~7.81 (m, 2H), 7.69 (d, $J=8.9$ Hz, 1H), 7.43 (d, $J=8.6$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.6, 178.0, 145.4, 139.4, 137.1, 134.9, 134.4, 133.9, 133.0, 126.7, 126.6, 124.4, 121.2, 120.8 (123.8, 121.8, 119.7, 117.7, q, $J_{\text{C-F}}=257$ Hz), 117.9, 116.2, 114.3; ^{19}F NMR (376 MHz, DMSO) δ : -56.78; IR (KBr) ν : 3296, 1657, 1587, 1531, 1482, 1390, 1255, 1209, 1015, 715 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{17}\text{H}_7\text{F}_3\text{NO}_2 [\text{M}-\text{H}]^-$ 314.0434, found 314.0433.

1-Methyl-5*H*-benzo[*b*]carbazole-6,11-dione (**2y**): Yield 34% (17.8 mg, Eluent: EtOAc/petroleum ether, $V:V=1:3$); orange solid, m.p. 321~323 °C; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.07 (s, 1H), 8.15 (t, $J=8.1$ Hz, 1H), 8.08 (t, $J=8.2$ Hz, 1H), 7.89~7.78 (m, 2H), 7.41 (t, $J=8.5$ Hz, 1H), 7.34~7.29 (m, 1H), 7.10 (t, $J=8.0$ Hz, 1H), 2.93 (d, $J=9.1$ Hz, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 179.8, 178.2, 139.4, 138.3, 135.2, 134.9, 134.2, 133.5, 132.4, 127.6, 127.3, 126.0, 125.6, 124.4, 119.0, 111.7, 23.4; IR (KBr) ν : 3427, 2955, 1565, 1441, 1426, 1385, 745, 712 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{12}\text{NO}_2 [\text{M}+\text{H}]^+$ 262.0863, found 262.0864.

7*H*-Dibenzo[*a,h*]carbazole-7,12(13*H*)-dione (**2z**): Yield 22% (17.8 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); red solid, m.p. 291~293 °C; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.60 (s, 1H), 9.95 (d, $J=8.4$ Hz, 1H), 8.28 (d, $J=7.6$ Hz, 1H), 8.14 (d, $J=7.4$ Hz, 1H), 8.04 (d, $J=7.9$ Hz, 1H), 7.96 (d, $J=8.9$ Hz, 1H), 7.91 (t, $J=7.5$ Hz, 1H), 7.86 (t, $J=7.3$ Hz, 1H), 7.75~7.71 (m, 2H), 7.61 (t, $J=7.4$ Hz, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.5, 177.3, 137.2, 136.7, 135.1, 134.7, 133.7, 132.7, 131.3, 129.8, 129.4, 128.9, 127.6, 127.5, 127.3, 126.1, 126.0, 120.8, 120.5, 114.4; IR (KBr) ν : 3275, 1646, 1593, 1522, 717, 678 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{12}\text{NO}_2 [\text{M}+\text{H}]^+$ 298.0863, found 298.0860.

7*H*-Dibenzo[*b,g*]carbazole-8,13-dione (**2ba**): Yield 15% (8.91 mg, Eluent: EtOAc/petroleum ether, $V:V=1:5$); red solid, m.p. 336~338 °C; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.88 (s, 1H), 8.80 (d, $J=8.1$ Hz, 1H), 8.23

(d, $J=8.7$ Hz, 1H), 8.18~8.11 (m, 2H), 8.04 (d, $J=7.9$ Hz, 1H), 7.88~7.79 (m, 3H), 7.70~7.67 (m, 1H), 7.64~7.60 (m, 1H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 181.5, 177.1, 136.0, 135.2, 134.5, 134.4, 133.9, 133.5, 132.6, 129.3, 127.4, 127.2, 126.6, 125.9, 122.9, 122.6, 121.4, 120.7, 119.3; IR (KBr) ν : 3235, 1638, 1585, 1510, 1432, 1263, 1186, 808, 713 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{12}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 298.0863, found 298.0862.

2,4-Dimethyl-5H-benzo[*b*]carbazole-6,11-dione (2bb): Yield 80% (36.2 mg, Eluent: EtOAc/petroleum ether, $V: V=1:5$); red solid, m.p. 343~345 $^{\circ}\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 12.89 (s, 1H), 8.10 (d, $J=8.3$ Hz, 2H), 7.86~7.78 (m, 3H), 7.04 (s, 1H), 2.53 (s, 3H), 2.40 (s, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.8, 177.8, 137.5, 136.9, 134.5, 133.9, 133.6, 133.2, 129.8, 126.5, 126.4, 124.7, 123.9, 119.5, 117.8, 21.7, 17.4; IR (KBr) ν : 3346, 2923, 1719, 1433, 1381, 1251, 1007, 716, 529 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 276.1019, found 276.1022.

2,4-Dimethoxy-5H-benzo[*b*]carbazole-6,11-dione (2bc): Yield 59% (36.2 mg, Eluent: EtOAc/petroleum ether, $V: V=1:5$); dark red solid, m.p. 311~313 $^{\circ}\text{C}$; ^1H NMR (500 MHz, DMSO- d_6) δ : 13.21 (s, 1H), 8.09 (t, $J=7.5$ Hz, 2H), 7.86~7.79 (m, 2H), 7.21 (s, 1H), 6.61 (s, 1H), 3.94 (s, 3H), 3.86 (s, 3H); ^{13}C NMR (126 MHz, DMSO- d_6) δ : 180.9, 177.3, 158.7, 148.6, 137.0, 134.6, 134.5, 133.6, 133.4, 126.5, 126.4, 126.1, 125.3, 118.0, 99.7, 94.2, 56.3, 56.0; IR (KBr) ν : 3248, 1644, 1595, 1436, 1290, 1270, 1145, 964, 712 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 308.0917, found 308.0916.

Supporting Information ^1H NMR and ^{13}C NMR spectra of **1h**, **1n**, **1x**, **2a**~**2bc** and HRMS spectra of **1h**, **1n**, **1x**, **2d**, **2e**, **2h**, **2j**, **2k**, **2m**~**2q**, **2t**, **2x**~**2bc** are available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- (a) Bernardo, P. H.; Chai, C. L. L.; Heath, G. A.; Mahon, P. J.; Smith, G. D.; Waring, P.; Wilkes, B. A. *J. Med. Chem.* **2004**, *47*, 4958.
(b) Moon, Y.; Jeong, Y.; Kook, D.; Hong, S. *Org. Biomol. Chem.* **2015**, *13*, 3918.
- Furukawa, H.; Wu, T. S.; Ohta, T.; Kuoh, C. S. *Chem. Pharm. Bull.* **1985**, *33*, 4132.
- Rickards, R. W.; Rothschild, J. M.; Willis, A. C.; de Chazal, N. M.; Kirk, J.; Kirk, K.; Saliba, K. J.; Smith, G. D. *Tetrahedron* **1999**, *55*, 13513.
- Saha, C. K.; Chowdhury, B. *Phytochemistry* **1998**, *48*, 363.
- (a) Watanabe, M.; Snieckus, V. *J. Am. Chem. Soc.* **1980**, *102*, 1457.
(b) Sissouma, D.; Collet, S. C.; Guingant, A. Y. *Synlett* **2004**, 2612.
- (a) Schmidt, A. W.; Reddy, K. R.; Knolker, H. J. *Chem. Rev.* **2012**, *112*, 3193.
(b) Norman, A. R.; Norcott, P.; McErlean, C. S. P. *Tetrahedron Lett.* **2016**, *57*, 4001.
(c) Sridharan, V.; Martín, M. A.; Menéndez, J. C. *Eur. J. Org. Chem.* **2009**, 4614.
(d) Bolibrukh, K.; Khoumeri, O.; Polovkovych, S.; Novikov, V.; Terme, T.; Vanelle, P. *Synlett* **2014**, 2765.
- (e) Kaliyaperumal, S. A.; Banerjee, S.; Kumar, U. K. S. *Org. Biomol. Chem.* **2014**, *12*, 6105.
- Ramkumar, N.; Nagarajan, R. *Org. Biomol. Chem.* **2015**, *13*, 11046.
- Ramkumar, N.; Nagarajan, R. *RSC Adv.* **2015**, *5*, 87838.
- Do, H. H.; Ejaz, S. A.; Molenda, R.; Ohlendorf, L.; Villinger, A.; Khan, S. U.; Lecka, J.; Sevigny, J.; Iqbal, J.; Ehlers, P.; Langer, P. *ChemistrySelect* **2019**, *4*, 2545.
- Abe, T.; Ikeda, T.; Yanada, R.; Ishikura, M. *Org. Lett.* **2011**, *13*, 3356.
- Indumathi, T.; Fronczek, F. R.; Prasad, K. J. R. *Tetrahedron Lett.* **2014**, *55*, 5361.
- Dethe, D. H.; Murhade, G. M. *Eur. J. Org. Chem.* **2014**, 6953.
- Zhang, K.; Xu, H.; Liu, Z.; Song, C. *Chin. J. Org. Chem.* **2019**, *39*, 1142 (in Chinese).
(张坤, 徐红进, 刘咨博, 宋传君, 有机化学, **2019**, *39*, 1142.)
- (a) Moon, Y.; Jeong, Y.; Kook, D.; Hong, S. *Org. Biomol. Chem.* **2015**, *13*, 3918.
(b) Sieveking, I.; Thomas, P.; Estévez, J. C.; Quiñones, N.; Cuéllar, M. A.; Villena, J.; Espinosa-Bustos, C.; Fierro, A.; Tapia, R. A.; Maya, J. D.; López-Muñoz, R.; Cassels, B. K.; Estévez, R. J.; Salas, C. O. *Med. Chem.* **2014**, *22*, 4609.
(c) Mandal, A.; Mondal, S. K.; Jana, A.; Manna, S. K.; Ali, S. A.; Samanta, S. *J. Heterocycl. Chem.* **2017**, *54*, 2529.
(d) Suematsu, N.; Ninomiya, M.; Sugiyama, H.; Udagawa, T.; Tanaka, K.; Koketsu, M. *Bioorg. Med. Chem. Lett.* **2019**, *29*, 2243.
(e) Chen, X.-L.; Dong, Y.; He, S.; Zhang, R.; Zhang, H.; Tang, L.; Zhang, X.-M.; Wang, J.-Y. *Synlett* **2019**, *30*, 615.
(f) Li, X.; Song, Z.; Chen, X.; Cai, Y.; Liu, Y.; Chen, C.; Peng, J. *Chin. J. Org. Chem.* **2020**, *40*, 950 (in Chinese).
(李雪, 宋子锐, 陈鑫, 蔡铁超, 刘雅婕, 陈春霞, 彭进松, 有机化学, **2020**, *40*, 950.)
- Xu, S.; Nguyen, T.; Pomilio, I.; Vitale, M. C.; Velu, S. E. *Tetrahedron* **2014**, *70*, 5928.
- Nishiyama, T.; Choshi, T.; Kitano, K.; Hibino, S. *Tetrahedron Lett.* **2011**, *52*, 3876.
- (a) Guo, J.; Kiran, I. N. C.; Reddy, R. S.; Gao, J.; Tang, M.; Liu, Y.; He, Y. *Org. Lett.* **2016**, *18*, 2499.
(b) Guo, J.; Kiran, I. N. C.; Gao, J.; Reddy, R. S.; He, Y. *Tetrahedron Lett.* **2016**, *57*, 3481.
(c) Banne, S.; Reddy, D. P.; Li, W.; Wang, C.; Guo, J.; He, Y. *Org. Lett.* **2017**, *19*, 4996.
- (a) Chan, C. M.; Chow, Y. C.; Yu, W. Y. *Synthesis* **2020**, *52*, 2899.
(b) Yang, Y. F.; Zhang, D. Q.; Vessally, E. *Top. Curr. Chem.* **2020**, *378*, 37.
- Selected reviews:
(a) Cheng, X.; Hu, X.; Lu, Z. *Chin. J. Org. Chem.* **2017**, *37*, 251 (in Chinese).
(程晓恺, 胡新根, 陆展, 有机化学, **2017**, *37*, 251.)
(b) Zhang, Y.; Schilling, W.; Das, S. *ChemSusChem* **2019**, *12*, 2898.
(c) Zhang, X.; Rakesh, K. P.; Ravindar, L.; Qin, H. L. *Green Chem.* **2018**, *20*, 4790.
- Selected examples, see:
(a) Maity, S.; Zheng, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 9562.
(b) Choi, S.; Chatterjee, T.; Choi, W. J.; You, Y.; Cho, E. J. *S ACS Catal.* **2015**, *5*, 4796.
(c) Martínez, C.; Bosnidou, A. E.; Allmendinger, S.; Muñiz, K. *Chem. Eur. J.* **2016**, *22*, 9929.
(d) Moon, Y.; Jang, E.; Choi, S.; Hong, S. *Org. Lett.* **2018**, *20*, 240.
(e) Wang, N.; Gu, Q.-S.; Li, Z.-L.; Li, Z.; Guo, Y.-L.; Guo, Z.; Liu, X.-Y. *Angew. Chem., Int. Ed.* **2018**, *57*, 14225.
- Selected examples, see:

- (a) Margrey, K. A.; McManus, J. B.; Bonazzi, S.; Zecri, F.; Nicewicz, D. A. *J. Am. Chem. Soc.* **2017**, *139*, 11288.
- (b) Ruffoni, A.; Juliá, F.; Svejstrup, T. D.; McMillan, A. J.; Douglas, J. J.; Leonori, D. *Nat. Chem.* **2019**, *11*, 426.
- (c) Kaur, S.; Kumar, M.; Bhalla, V. *Green Chem.* **2016**, *18*, 5870.
- (d) Zheng, Y.-W.; Chen, B.; Ye, P.; Feng, K.; Wang, W.; Meng, Q.-Y.; Wu, L.-Z.; Tung, C.-H. *J. Am. Chem. Soc.* **2016**, *138*, 10080.
- (e) Zhao, F.; Yang, Q.; Zhang, J.; Shi, W.; Hu, H.; Liang, F.; Wei, W.; Zhou, S. *Org. Lett.* **2018**, *20*, 7753.
- (f) Engl, P. S.; Häring, A. P.; Berger, F.; Berger, G.; Pérez-Bitrián, A.; Ritter, T. *J. Am. Chem. Soc.* **2019**, *141*, 13346.
- [15] Ashok, P.; Ilangoan, A. *Tetrahedron Lett.* **2018**, *59*, 438.
- [16] Hsu, Y. C.; Wang, V. C. C.; Au-Yeung, K. C.; Tsai, C. Y.; Chang, C. C.; Lin, B. C.; Chan, Y. T.; Hsu, C. P.; Yap, G. P. A.; Jurca, T.; Ong, T. G. *Angew. Chem., Int. Ed.* **2018**, *57*, 4622.
- [17] (a) Campeau, L. C.; Parisien, M.; Jean, A.; Fagnou, K. *J. Am. Chem. Soc.* **2006**, *128*, 581.
- (b) Mota, A. J.; Dedieu, A.; Bour, C.; Suffert, J. *J. Am. Chem. Soc.* **2005**, *127*, 7171.
- (c) Garcia-Cuadrado, D.; Braga, A. A. C.; Maseras, F.; Echavarren, A. M. *J. Am. Chem. Soc.* **2006**, *128*, 1066.
- [18] Lisboa, C. D.; de Lucas, N. C.; Garden, S. J. *Dyes Pigm.* **2016**, *134*, 618.

(Lu, Y.)