铈催化定位基促进芳环 C—H 键和碳酸亚乙烯酯的缩合得到异香豆素

王璐^a 邵莺^a 陈帆^b 钱鹏程^{*,b} 成江^{*,a,b}^(a) 常州大学石油化工学院 江苏常州 213164^(b) 温州大学化学与材料工程学院 浙江温州 325035

摘要 异香豆素是一种重要的药物分子, 目前已经开发了许多合成途径, 主要集中在苯甲酸或其他物质与炔烃的环化. 亚磺酰基苯甲酰胺在螯合辅助的芳烃邻位 C—H 官能化中可以轻松地制备, 碳酸亚乙烯酯可以建立无氧化剂的催化体系, 使碳酸作为唯一的副产物. 在铈催化剂的存在下, 亚磺酰基苯甲酰胺和碳酸亚乙烯酯之间形成了环化, 通过导向基团辅助的邻位 C—H 偶联和分子内醇解, 以中等至良好的产率获得未取代的异香豆素.

关键词 铈; 亚磺酰基苯甲酰胺; 碳酸亚乙烯酯

Rhodium-Catalyzed Directing Group-Assisted Annulation of Arene C—H Bond with Vinylene Carbonate toward Isocoumarins

Wang, Lu^a Shao, Ying^a Chen, Fan^b Qian, Pengcheng^{*,b} Cheng, Jiang^{*,a,b}^(a) School of Petrochemical Engineering, Changzhou University, Changzhou, Jiangsu 213164^(b) College of Chemistry & Materials Engineering, Wenzhou University, Wenzhou, Zhejiang 325035

Abstract Isocoumarins are an important pharmaceutical compounds. Many synthetic pathways have been developed, mainly focusing on the cyclization of benzoic acid or other with alkynes. Sulfinylidene benzamides were facilely prepared and developed in the chelation-assisted arene *ortho*-C—H functionalization, and vinylene carbonate can establish an oxidant-free catalytic system, giving carbonic acid as a sole side product. In the presence of rhodium catalyst, an annulation between sulfinylidene benzamides and vinylene carbonate was developed, proceeding with directing group-assisted *ortho*-C—H coupling and intramolecular alcoholysis toward nonsubstituted isocoumarins in moderate to good yields.

Keywords rhodium; sulfinylidene benzamides; vinylene carbonate

Isocoumarins are ubiquitous in pharmaceutical chemistry,^[1] serving as antimicrobial,^[2] antifungal,^[3] anticancer,^[4] anti-HIV,^[5] anti-inflammatory^[6] and anti-allergy^[6] reagents (Scheme 1). As a result, many synthetic pathways have been developed,^[7] which mainly focused on the annulation of benzoic acid^[8] or others^[9] with alkynes (Scheme 2a). Intriguingly, the access of nonsubstituted isocoumarins was rarely reported, which was at least partially due to the lack of catalytic system using ethyne equivalent. Recently, Miura *et al.*^[10–11] reported a concise synthesis of isocoumarins whereby Rh-catalyzed direct vinylene annulation (Scheme 2b).

Sulfinylidene benzamides were facilely prepared and developed in the chelation-assisted arene *ortho*-C—H fun-

ctionalization.^[12] Generally, further step was required for the transformation of sulfinylidene benzamides into useful functional groups. Vinylene carbonate can establish an oxidant-free catalytic system, giving carbonic acid as a sole side product. We envisaged the cyclization would take place between sulfinylidene benzamides and the new introduced functional groups. Herein, we wish to report the rhodium-catalyzed annulation of sulfinylidene benzamides and vinylene carbonate toward isocoumarins (Scheme 2c). In this reaction, with the aid of a transformable sulfoximine directing group, the construction of C—O and C—C bonds occurs in one step in the presence of rhodium catalyst. Compared to Miura's work,^[10] vinylene carbonate is not served as ethyne surrogate in this procedure, which extends

* Corresponding authors. E-mail: jiangcheng@cczu.edu.cn, qpc@wzu.edu.cn

Received June 10, 2021; revised July 14, 2021; published online September 2, 2021.

Project supported by the National Natural Science Foundation of China (No. 21971025), the "Innovation & Entrepreneurship Talents" Introduction Plan of Jiangsu Province, the Jiangsu Key Laboratory of Advanced Catalytic Materials & Technology (No. BM2012110), the Foundation of Wenzhou Science & Technology Bureau (No. W20170003), and the Advanced Catalysis and Green Manufacturing Collaborative Innovation Center.

国家自然科学基金(No. 21971025)、江苏省“创新创业人才”引进计划、江苏省先进催化材料与技术重点实验室(No. BM2012110)、温州市科技局基金(No. W20170003)、先进催化与绿色制造协同创新中心资助项目.

the application for vinylene carbonate.

Scheme 1 Representative isocoumarins compounds.

Scheme 2 Annulation of benzoic acid and its derivatives toward isocoumarins.

2 Results and discussion

We began our investigation with the reaction of *N*-(4-methylbenzoyl)methylphenyl sulfoximine (**1a**) (0.10 mmol) and vinylene carbonate **2a** (0.40 mmol) in the presence of $\text{Cp}^*\text{Rh}(\text{OAc})_2 \cdot \text{H}_2\text{O}$, AgSbF_6 and $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ in 1,4-dioxane (1.5 mL) at 120 °C under air. Disappointingly, serving $\text{Cp}^*\text{Rh}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ or $[\text{Cp}^*\text{IrCl}_2]_2$ as catalyst resulted in poor yields (<5%, Table 1, Entries 1 and 2). To our delight, the product **3a** was isolated in 15% yield in the presence of $[\text{Cp}^*\text{RhCl}_2]_2$ (Entry 3). However, no reaction took place in the absence of $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (Entry 4). In this reaction, $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ served as an important additive, which may enhance the reaction efficiency by promoting the oxidation of silver salt. Among the Ag(I) tested, AgNTf_2 (56%, Entry 6) was superior to AgSbF_6 (15%, Entry 3) and AgBF_4 (22%, Entry 5). Notably, it resulted in lower efficiency under either nitrogen (49%) or oxygen (51%) atmosphere (Entry 6). Additive 1, such as $\text{Cu}(\text{OTf})_2$ (39%, Entry 7), CuI (45%, Entry 8), $\text{Zn}(\text{OAc})_2 \cdot$

$2\text{H}_2\text{O}$ (<5%, Entry 9) and $\text{Zn}(\text{OTf})_2$ (58%, Entry 10) did not improve the reaction efficiency significantly. Next, the reaction efficiency was further tested by introducing additive 2 (HOAc , $\text{Zn}(\text{OTf})_2$, ZnCl_2 , ZnBr_2 or MgCl_2) respectively in the presence of $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$, and additive 2 $\text{Zn}(\text{OTf})_2$ was the best (78%, Entry 12). The solvent also had a profound effect in the reaction efficiency. 1,4-Dioxane (78%, Entry 12) was superior to PhCl (66%, Entry 14), 1,2-dichloroethane (DCE, 59%, Entry 15) and dichloromethane (DCM, 62%, Entry 16).

With the optimized conditions in hand, the substrate scope of the sulfinylidene benzamides was examined. As shown in Table 2, A series of functionalities, such as halogens (**3b**, 78%; **3c**, 86%; **3d**, 88%; **3n**, 47%; **3t**, 51%; **3s**, 40%), formyl (**3h**, 80%), trifluoromethyl (**3i**, 77%), alkyl (**3e**, 74%; **3g**, 66%; **3j**, 56%; **3l**, 52%; **3o**, 46%; **3p**, 57%), methoxy (**3f**, 82%; **3q**, 46%) and ester (**3k**, 68%) are compatible with the optimized reaction conditions. In addition, the electronic properties of functional groups have no obvious effect on reaction efficiency. The sulfinylidene benzamides with electron-withdrawing groups (**3b** ~ **3d**, **3h**, **3i**, **3k**) and the electron-donating groups (**3e**, **3f**, **3g**, **3j**) at the *para*-position provide isocoumarin derivatives in good yields. Importantly, various substrates with substituents at the *meta*- and *ortho*-positions worked well. Notably, the naphthalene substituted substrates ran smoothly during this reaction (**3u**, 56%; **3v**, 61%). The structure of **3v** was further confirmed by X-ray crystal analysis (CCDC 2051238). Disappointingly, heterocyclic substrates and substituted vinylene carbonate did not work under this procedure.

To improve the practicability of the procedure, the reaction was conducted in 1 mmol scale and **3a** was isolated in an acceptable 68% yield. To gain mechanistic insight into this reaction, the H/D exchange experiment was conducted. The result shows that 20% of *ortho*-H is replaced with D, leading to **1a-d2** (Scheme 3a). However, no D was incorporated in **1a** in the absence of $[\text{Cp}^*\text{RhCl}_2]_2$ (Scheme 3b). Hence, the cleavage of the *ortho*-C—H bond is probably reversible.

A proposed reaction mechanism for this transformation is illustrated in Scheme 4.^[10] Firstly, **1a** coordinates to the $[\text{Cp}^*\text{Rh}]^{2+}$ species, which facilitates the cleavage of the *ortho*-C—H bond of **1a** to form a five-membered rhodacycle **A**. Secondly, the insertion of vinylene carbonate in the Rh—C bond results in a seven-membered intermediate **B**. Then, complex **C** is generated via β -oxygen elimination and decarboxylation. Finally, intramolecular alcoholysis of complex **C** affords the product **3a**.^[11c]

3 Conclusion

In conclusion, we have developed a rhodium-catalyzed annulation of sulfinylidene benzamides with vinylene carbonate, whereby directing group-assisted *ortho*-C—H coupling and intramolecular alcoholysis leading to nonsubstituted isocoumarins in moderate to good yields.

Table 1 Selected results for screening the optimized reaction conditions^a

Entry	Catalyst	Oxidant	Additive 1	Additive 2	Yield ^b
1	Cp*Rh(OAc) ₂ •H ₂ O	AgSbF ₆	Cu(OAc) ₂ •H ₂ O	—	<5%
2	[Cp*RhCl ₂] ₂	AgSbF ₆	Cu(OAc) ₂ •H ₂ O	—	<5%
3	[Cp*RhCl ₂] ₂	AgSbF ₆	Cu(OAc) ₂ •H ₂ O	—	15%
4	[Cp*RhCl ₂] ₂	AgSbF ₆	—	—	NR
5	[Cp*RhCl ₂] ₂	AgBF ₄	Cu(OAc) ₂ •H ₂ O	—	22%
6	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	—	56%, 49% ^c , 51% ^d
7	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OTf) ₂	—	39%
8	[Cp*RhCl ₂] ₂	AgNTf ₂	CuI	—	45%
9	[Cp*RhCl ₂] ₂	AgNTf ₂	Zn(OAc) ₂ •H ₂ O	—	<5%
10	[Cp*RhCl ₂] ₂	AgNTf ₂	Zn(OTf) ₂	—	58%
11	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	HOAc	49%
12	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	Zn(OTf) ₂	78%, 71% ^e , 64% ^f
13	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	ZnCl ₂	54%, 65% ^g , 20% ^h
14	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	Zn(OTf) ₂	66% ⁱ
15	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	Zn(OTf) ₂	59% ^d
16	[Cp*RhCl ₂] ₂	AgNTf ₂	Cu(OAc) ₂ •H ₂ O	Zn(OTf) ₂	62% ^k

^a Reaction conditions: **1a** (0.10 mmol), **2a** (0.40 mmol), catalyst (10 mol%), oxidant (60 mol%), additive 1 (1.5 equiv.), additive 2 (20 mol%), solvent: 1,4-dioxane (1.5 mL), 120 °C, under air for 20 h; ^b isolated yields; ^c under N₂ for 20 h; ^d under O₂ for 20 h; ^e additive 2: Zn(OTf)₂ (60 mol%); ^f additive 2: Zn(OTf)₂ (0.1 mmol); ^g additive 2: ZnBr₂ (20 mol%); ^h additive 2: MgCl₂ (20 mol %); ⁱ solvent: PhCl (1.5 mL); ^j solvent: DCE (1.5 mL); ^k solvent: DCM (1.5 mL).

Table 2 Substrates scope of sulfonylidene benzamides

3b , 78%	3c , 86%	3d , 88%	3e , 74%	3f , 82%	3g , 66%	3h , 80%
3i , 77%	3j , 56%	3k , 68%	3l , 52%	3m , trace	3n , 47%	3o , 46%
3p , 57%	3q , 46%	3r , trace	3s , 40%	3t , 51%	3u , 56%	3v , 61%

^a Reaction conditions: **1a** (0.10 mmol), **2a** (0.40 mmol), [Cp*RhCl₂]₂ (10 mol%), AgNTf₂ (60 mol%), Cu(OAc)₂•H₂O (1.5 equiv.), Zn(OTf)₂ (20 mol%), 1,4-dioxane (1.5 mL), 120 °C, under air for 20 h. ^b Isolated yield.

Scheme 3 H/D exchange experiments

Scheme 4 A proposed reaction mechanism

4 Experimental section

4.1 General experimental information

Unless otherwise noted, all the reactions were carried out under oxygen atmosphere using standard Schlenk technique, and all chemicals were purchased from commercial suppliers and used without further purification. The 1H NMR spectra were recorded on a 300 MHz or 400 MHz NMR spectrometer. The ^{13}C NMR spectra were recorded at 75 MHz or 100 MHz. NMR experiments are reported in δ units, and were referenced to $CDCl_3$ (δ 7.26 or 77.0) as the internal standard. The coupling constants J are given in Hz. High-resolution mass spectra (HRMS) were obtained using a Bruker micro-TOF II focus spectrometer (ESI). Column chromatography was performed using EM Silica gel 60 (300~400 mesh), and the eluent was a mixture of petroleum ether (PE) and ethyl acetate (EA). All melting points were tested after recrystallization with CH_2Cl_2 using WRS-1B digital melting point instrument.

4.2 Experimental methods

4.2.1 Synthesis of compounds 1a~1v

A solution of *N'*-(3-dimethylaminopropyl)-*N*-ethylcarbamide hydrochloride salt (EDC•HCl) (2.0 equiv., 16 mmol, 3.05 g), 4-*N,N*-dimethylaminopyridine (DMAP) (2.2 equiv., 17.6 mmol, 2.15 g) and benzoic acids (1.1 equiv., 8.8 mmol, 1.07 g) in CH_2Cl_2 (40.0 mL) was stirred under an argon atmosphere. (Methylsulfonylimidoyl)benzene (1.0 equiv., 8 mmol, 1.24 g) was introduced dropwise at 0 °C; The resulting reaction mixture was stirred for about 1 h at 0 °C, then reacted for 12 h at room temperature. Upon complete consumption of (methylsulfonylimidoyl)benzene, the reaction mixture was acidified with hydrochloric acid (HCl , 2 mol•L⁻¹). The organic layer was separated, and the aqueous layer was extracted with CH_2Cl_2 (3 times). The combined extracts were washed with 10% aqueous $NaHCO_3$ and brine. The organic layer was dried over Na_2SO_4 . Solvent was filtered and evaporated under the reduced pressure. The crude residue was purified using column chromatography on silica gel using hexane/ethyl acetate ($V/V=5/1$) to give compound 1a as white solid (1.7 g, 82% yield).^[12,13] 1b~1v were prepared as above.

4.2.2 Synthesis of compounds 3a~3v

A 20 mL of Schlenk tube equipped with a stir bar was charged with *N*-(4-methylbenzoyl)methylphenyl sulfoximine (1a) (0.1 mmol, 1.0 equiv., 25.9 mg), vinylene carbonate (0.4 mmol, 4.0 equiv., 34.4 mg), $[Cp^*RhCl]_2$ (10 mol%, 8 mg), $AgNTf_2$ (23.3 mg, 60 mol%), $Cu(OAc)_2 \cdot H_2O$ (29.9 mg, 1.5 equiv.), $Zn(OTf)_2$ (7.3 mg, 20 mol%) and 1,4-dioxane (1.5 mL). The tube was sealed with a PTFE cap. The reaction mixture was stirred at 120 °C for 20 h under air in an oil bath. After the completion of the reaction, the mixture was then allowed to cool to room temperature. The organic solution was concentrated, and purified through silica gel column chromatography using a mixture of ethyl acetate (EtOAc) and petroleum ether (PE) as eluent to afford the corresponding products 3a (11.3 mg, 78% yield). 3b~3v were prepared as above.

1*H*-Isochromen-1-one (3a):^[14] 11.3 mg, 78% yield. Yellow oil. 1H NMR ($CDCl_3$, 300 MHz) δ : 8.30~8.27 (m, 1H), 7.75~7.69 (m, 1H), 7.55~7.48 (m, 1H), 7.45~7.41 (m, 1H), 7.28 (d, $J=5.6$ Hz, 1H), 6.51~6.49 (m, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ : 162.26, 144.69, 136.47, 134.81, 129.67, 128.60, 125.55, 121.87, 107.03.

6-Fluoro-1*H*-isochromen-1-one (3b):^[15] 12.7 mg, 78% yield. White solid, m.p. 157~161 °C; 1H NMR ($CDCl_3$, 300 MHz) δ : 8.32~8.27 (m, 1H), 7.29 (d, $J=5.6$ Hz, 1H), 7.23~7.16 (m, 1H), 7.10~7.04 (m, 1H), 6.46 (d, $J=5.6$ Hz, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ : 168.28, 163.09 (d, $J_{C-F}=269.5$ Hz), 145.95, 139.15 (d, $J_{C-F}=10.8$ Hz), 133.11 (d, $J_{C-F}=10.2$ Hz), 118.32 (d, $J_{C-F}=2.4$ Hz), 116.90 (d, $J_{C-F}=22.9$ Hz), 111.36 (d, $J_{C-F}=22.5$ Hz), 106.54 (d, $J_{C-F}=2.8$ Hz); ^{19}F NMR (75 MHz, $CDCl_3$) δ : 101.59.

6-Bromo-1*H*-isochromen-1-one (3c): 19.3 mg, 86% yield. Yellow solid, sublimation at 201 °C (lit.^[10]

sublimation at 207 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 8.16~8.11 (m, 1H), 7.66~7.62 (m, 1H), 7.62~7.60 (m, 1H), 7.30 (d, *J*=5.6 Hz, 1H), 6.44~6.42 (m, 1H); ¹³C NMR (75 MHz, CDCl₃) δ: 161.55, 145.94, 137.88, 131.99, 131.37, 130.26, 128.26, 120.55, 106.00.

6-Chloro-1*H*-isochromen-1-one (**3d**): 15.8 mg, 88% yield. White solid, m.p. 190~193 °C (lit.^[16] 192~193 °C); ¹H NMR (*d*₆-DMSO, 300 MHz) δ: 8.11 (d, *J*=8.4 Hz, 1H), 7.73 (d, *J*=8.0 Hz, 1H), 7.62~7.61 (m, 1H), 7.59 (d, *J*=8.0 Hz, 1H), 6.76 (d, *J*=5.6 Hz, 1H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 161.29, 147.37, 140.62, 138.50, 131.51, 129.35, 125.86, 120.14, 106.37.

6-Methyl-1*H*-isochromen-1-one (**3e**): 11.8 mg, 74% yield. White solid, m.p. 191~194 °C (lit.^[16] 191~193 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 8.09 (d, *J*=8.0 Hz, 1H), 7.26~7.23 (m, 1H), 7.18~7.14 (m, 1H), 7.14~7.13 (m, 1H), 6.36~6.34 (m, 1H), 2.39 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 162.36, 145.94, 144.76, 136.56, 129.94, 129.64, 125.58, 119.43, 106.98, 21.97.

6-Methoxy-1*H*-isochromen-1-one (**3f**): 14.4 mg, 82% yield. Yellow solid, m.p. 92~94 °C (lit.^[10] 89~91 °C); ¹H NMR (CDCl₃, 400 MHz) δ: 8.18 (d, *J*=8.8 Hz, 1H), 7.23 (d, *J*=5.6 Hz, 1H), 7.04~7.01 (m, 1H), 6.79 (d, *J*=9.6 Hz, 1H), 6.41 (d, *J*=5.6 Hz, 1H), 3.90 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 164.66, 162.05, 145.31, 138.81, 131.89, 116.81, 114.92, 107.73, 107.04, 55.71.

6-Ethyl-1*H*-isochromen-1-one (**3g**): 11.4 mg, 66% yield. White solid, m.p. 61~64 °C; ¹H NMR (CDCl₃, 300 MHz) δ: 8.18 (d, *J*=8.0 Hz, 1H), 7.36~7.33 (m, 1H), 7.26~7.23 (m, 1H), 7.23~7.22 (m, 1H), 6.46~6.42 (m, 1H), 2.79~2.71 (m, 2H), 1.30~1.22 (m, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 162.36, 152.04, 144.70, 136.67, 129.73, 128.85, 124.39, 119.61, 107.11, 29.19, 15.07. HRMS (APCI) calcd for C₁₁H₁₁O₂ [M+H]⁺ 175.0754, found 175.0748.

1-Oxo-1*H*-isochromene-6-carbaldehyde (**3h**): 13.9 mg, 80% yield. White solid, m.p. 175~177 °C (lit.^[10] 178~179 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 10.16 (s, 1H), 8.46 (d, *J*=8.0 Hz, 1H), 8.02~7.98 (m, 1H), 7.94 (d, *J*=5.6 Hz, 1H), 7.36 (d, *J*=5.6 Hz, 1H), 6.63~6.61 (m, 1H); ¹³C NMR (75 MHz, CDCl₃) δ: 191.16, 161.24, 145.79, 140.32, 136.99, 130.80, 128.24, 127.22, 125.75, 106.69.

6-(Trifluoromethyl)-1*H*-isochromen-1-one (**3i**): 16.4 mg, 77% yield. White solid, m.p. 152~155 °C (lit.^[10] 149~151 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 8.44~8.40 (m, 1H), 7.76~7.73 (m, 1H), 7.72~7.71 (m, 1H), 7.36 (d, *J*=5.6 Hz, 1H), 6.59~6.56 (m, 1H); ¹³C NMR (75 MHz, CDCl₃) δ: 160.98, 146.05, 136.79, 136.28 (d, *J*_{C-F}=33.0 Hz), 130.74, 124.92 (d, *J*_{C-F}=10.7 Hz), 124.91 (d, *J*_{C-F}=3.4 Hz), 124.33 (d, *J*_{C-F}=2.9 Hz), 122.78 (d, *J*_{C-F}=3.9 Hz), 106.48; ¹⁹F NMR (75 MHz, CDCl₃) δ: 63.45.

6-(*tert*-Butyl)-1*H*-isochromen-1-one (**3j**): 11.3 mg, 56% yield. Yellow oil. ¹H NMR (CDCl₃, 400 MHz) δ: 8.19 (d, *J*=8.4 Hz, 1H), 7.58~7.55 (m, 1H), 7.38 (d, *J*=5.6 Hz, 1H), 7.26~7.23 (m, 1H), 6.48 (d, *J*=5.6 Hz, 1H), 1.36 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ: 162.31, 158.85, 144.63, 136.45, 129.49, 126.52, 121.92, 119.42, 107.44, 35.43, 31.01. HRMS (APCI) calcd for C₁₃H₁₅O₂ [M+H]⁺

203.1067, found 203.1063.

Methyl 1-oxo-1*H*-isochromene-6-carboxylate (**3k**): 13.9 mg, 68% yield. White solid, m.p. 165~166 °C (lit.^[10] 167~169 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 8.32~8.28 (m, 1H), 8.09~8.06 (m, 2H), 7.29 (d, *J*=5.6 Hz, 1H), 6.55~6.53 (m, 1H), 3.95 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 165.66, 161.37, 145.40, 136.40, 135.57, 129.94, 128.77, 127.07, 124.74, 106.80, 52.75.

8-Methyl-1*H*-isochromen-1-one (**3l**): 8.3 mg, 52% yield. m.p. 116~118 °C (lit.^[10] 113~114 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 7.57~7.52 (m, 1H), 7.31~7.28 (m, 1H), 7.26~7.22 (m, 1H), 7.22~7.21 (m, 1H), 6.42 (d, *J*=4.6 Hz, 1H), 2.82 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.64, 144.39, 143.66, 138.06, 134.00, 131.47, 123.74, 120.30, 107.47, 23.38.

8-Chloro-1*H*-isochromen-1-one (**3n**): 8.2 mg, 47% yield. Pale white solid, m.p. 140~143 °C; ¹H NMR (CDCl₃, 300 MHz) δ: 7.59~7.53 (m, 1H), 7.53~7.50 (m, 1H), 7.31~7.28 (m, 1H), 7.26~7.24 (m, 1H), 6.44 (d, *J*=5.6 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ: 158.73, 145.35, 139.44, 137.09, 134.59, 131.52, 124.59, 118.87, 106.89. HRMS (APCI) calcd for C₉H₆ClO₂ [M+H]⁺ 181.0051, found 181.0051.

5,8-Dimethyl-1*H*-isochromen-1-one (**3o**):^[17] 8.0 mg, 46% yield. White solid, m.p. 149~150 °C; ¹H NMR (CDCl₃, 300 MHz) δ: 7.18 (d, *J*=5.6 Hz, 1H), 7.10~7.09 (m, 1H), 7.01 (d, *J*=5.6 Hz, 1H), 6.34 (d, *J*=5.6 Hz, 1H), 2.76 (s, 3H), 2.39 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 161.72, 144.92, 144.44, 143.43, 138.14, 132.71, 123.87, 117.83, 107.40, 23.22, 21.64.

7-Methyl-1*H*-isochromen-1-one (**3p**): 9.1 mg, 57% yield. White solid, m.p. 99~101 °C (lit.^[18] 95 °C); ¹H NMR (CDCl₃, 300 MHz) δ: 8.02~8.01 (m, 1H), 8.49~8.45 (m, 1H), 7.27 (d, *J*=8.0 Hz, 1H), 7.17 (d, *J*=5.6 Hz, 1H), 6.43~6.40 (m, 1H), 2.40 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 162.47, 143.88, 138.88, 136.04, 133.95, 129.30, 125.49, 121.68, 106.92, 21.42.

7-Methoxy-1*H*-isochromen-1-one (**3q**):^[15] 8.1 mg, 46% yield. White solid, m.p. 132~135 °C; ¹H NMR (CDCl₃, 300 MHz) δ: 7.86~7.82 (m, 1H), 7.45~7.40 (m, 1H), 7.23 (d, *J*=5.7 Hz, 1H), 7.15~7.12 (m, 1H), 6.85~6.83 (m, 1H), 3.91 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ: 162.16, 153.99, 143.91, 128.90, 126.99, 122.67, 120.88, 114.51, 101.49, 55.93.

7-Fluoro-1*H*-isochromen-1-one (**3s**):^[15] 6.6 mg, 40% yield. White solid, m.p. 147~149 °C; ¹H NMR (CDCl₃, 300 MHz) δ: 7.94~7.90 (m, 1H), 7.45~7.42 (m, 2H), 7.25 (d, *J*=5.7 Hz, 1H), 6.50 (d, *J*=5.6 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ: 162.03 (d, *J*_{C-F}=248.6 Hz), 161.35 (d, *J*_{C-F}=3.6 Hz), 144.02 (d, *J*_{C-F}=2.8 Hz), 132.94 (d, *J*_{C-F}=2.7 Hz), 127.92 (d, *J*_{C-F}=7.7 Hz), 123.52 (d, *J*_{C-F}=8.1 Hz), 123.18 (d, *J*_{C-F}=22.9 Hz), 115.22 (d, *J*_{C-F}=23.2 Hz), 106.26 (d, *J*_{C-F}=1.3 Hz).

7-Chloro-1*H*-isochromen-1-one (**3t**): 9.1 mg, 51% yield. Pale white solid, m.p. 110~116 °C (lit.^[16] 114~115 °C); ¹H NMR (CDCl₃, 400 MHz) δ: 8.26 (d, *J*=8.0 Hz, 1H), 7.68~7.65 (m, 1H), 7.39 (d, *J*=8.4 Hz, 1H), 7.28 (d, *J*=

5.6 Hz, 1H), 6.50~6.48 (m, 1 H); ^{13}C NMR (75 MHz, CDCl_3) δ : 161.05, 144.96, 135.17, 134.81, 134.40, 129.20, 127.11, 123.09, 106.32.

1*H*-Benzo[g]isochromen-1-one (**3u**):^[19] 10.9 mg, 56% yield. White solid, m.p. 148~149 °C; ^1H NMR (CDCl_3 , 300 MHz) δ : 8.91 (s, 1H), 8.04~8.00 (m, 1H), 7.93~7.88 (m, 1H), 7.85 (s, 1H), 7.67~7.61 (m, 1H), 7.59~7.53 (m, 1H), 7.22 (d, $J=5.6$ Hz, 1H), 6.61~6.58 (m, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ : 162.52, 143.22, 136.37, 132.52, 132.06, 131.22, 129.73, 129.46, 127.75, 126.83, 123.97, 120.07, 107.14.

1*H*-Benzo[h]isochromen-1-one (**3v**):^[10] 11.9 mg, 61% yield. White solid, m.p. 148~150 °C (lit.^[10] 145~146 °C); ^1H NMR (d_6 -DMSO, 300 MHz) δ : 9.55~9.51 (m, 1H), 8.33~8.30 (m, 1H), 8.07~8.04 (m, 1H), 7.81~7.80 (m, 1H), 7.78~7.75 (m, 1H), 7.69~7.67 (m, 1H), 7.67~7.64 (m, 1H), 6.96 (d, $J=11.2$ Hz, 1H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ : 161.35, 147.91, 139.85, 136.99, 133.13, 131.12, 129.81, 129.53, 127.35, 126.00, 124.41, 114.54, 108.04.

3m, **3r** in trace yield.

4.2.3 Synthesis of compounds **1a** and **1a-d₂** (the H/D exchange experiment)

A 20 mL of Schlenk tube equipped with a stir bar was charged with *N*-(4-methylbenzoyl)methylphenyl sulfoximine (0.1 mmol, 1.0 equiv., 25.9 mg), $[\text{Cp}^*\text{RhCl}]_2$ (10 mol%, 8 mg), AgNTf_2 (23.3 mg, 60 mol%), $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (29.9 mg, 1.5 equiv., 0.15 mmol), $\text{Zn}(\text{OTf})_2$ (7.3 mg, 20 mol%) and 1,4-dioxane (1.5 mL). The tube was sealed with a PTFE cap. The reaction mixture was stirred at 120 °C for 2 h under air in an oil bath. After the completion of the reaction, the mixture was then allowed to cool to room temperature and quenched with 1.0 mL of D_2O . Ethyl acetate (2.0 mL) were added, the layers were separated and the aqueous layer was washed with ethyl acetate (30 mL $\times$ 2). The organic layers were combined, washed with brine (5 mL) and dried over Na_2SO_4 . The residue was purified by flash column chromatography on silica gel with petroleum ether-EtOAc ($V:V=8:1$) as the eluent to give the product **1a** and **1a-d₂** (see Supporting Information).

Supporting Information X-Ray structure of **3v**, and ^1H NMR and ^{13}C NMR spectra of compounds **3a**~**3v**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Saeed, A. *Eur. J. Med. Chem.* **2016**, *116*, 290.
(b) Brasholz, M.; Soergel, S.; Azap, C.; Reissig, H.-U. *Eur. J. Org. Chem.* **2007**, 3801.
(c) Pochet, L.; Frederick, R.; Masereel, B. *Curr. Pharm. Des.* **2004**, *10*, 3781.
(d) Wang X.; Wedge, D. E.; Cutler, S. *J. Nat. Prod. Commun.* **2016**, *11*, 1595.
- [2] (a) Harunari, E.; Imada, C.; Igarashi, Y. *J. Nat. Prod.* **2019**, *82*, 1609.
(b) Das, P.; Babbar, P.; Malhotra, N.; Sharma, M.; Jachak, G. R.; Gonnade, R. G.; Shanmugam, D.; Harlos, K.; Yogavel, M.; Sharma, A.; Reddy, D. S. *J. Med. Chem.* **2018**, *61*, 5664.
- [3] (a) Hu, Z.; Wu, Z.; Su, Q.; Li, M.; Wu, S.; Meng, R.; Ding, W.; Li, C. *Bioorg. Chem.* **2020**, *104*, 104300.
(b) Zhou, J.; Zheng, L.; Hei, Z.; Li, W.; Wang, J.; Yu, B.; Fang, P. *ACS Chem. Biol.* **2020**, *15*, 1016.
(c) Guo, D.-D.; Zhang, W.-X.; Wang, Y.-Q. *Chin. J. Org. Chem.* **2019**, *39*, 2650 (in Chinese).
(郭冬冬, 张武霞, 王永强, 有机化学, **2019**, *39*, 2650.)
- [4] (a) Kumar, J. S.; Thirupataiah, B.; Medishetti, R.; Ray, A.; Bele, S. D.; Hossain, K. A.; Reddy, G. S.; Edwin, R. K.; Joseph, A.; Kumar, N.; Shenoy, G. G.; Rao, C. M.; Pal, M. *Eur. J. Med. Chem.* **2020**, *201*, 112335.
(b) Li, X.; Zhao, C.; Jing, S.; Sun, J.; Li, X.; Man, S.; Wang, Y.; Gao, W. *Bioorg. Med. Chem. Lett.* **2017**, *27*, 3595.
- [5] Hu, Z.-X.; Xue, Y.-B.; Bi, X.-B.; Zhang, J.-W.; Luo, Z.-W.; Li, X.-N.; Yao, G.-M.; Wang, J.-P.; Zhang, Y.-H. *Mar. Drugs* **2014**, *12*, 5563.
- [6] (a) Thirupataiah, B.; Reddy, G. S.; Ghule, S. S.; Kumar, J. S.; Mounika, G.; Hossain, K. A.; Mudgal, J.; Mathew, J. E.; Shenoy, G. G.; Parsa, K. V. L.; Pal, M. *Bioorg. Chem.* **2020**, *97*, 103691.
(b) Yang, Z.-N.; Su, B.-J.; Wang, Y.-Q.; Liao, H.-B.; Chen, Z.-F.; Liang, D. *J. Nat. Prod.* **2020**, *83*, 985.
(c) Gu, L.; Liu, C.; Fang, X.; Weng, Z.; Li, Z.; Tan, Y.; Tang, K. *Chin. J. Synth. Chem.* **2020**, *28*, 841 (in Chinese).
(辜玲慧, 刘长英, 方新月, 翁正云, 李喆宇, 谭玉强, 唐克慧, 合成化学, **2020**, *28*, 841.)
- [7] For reviews, please see:
(a) Chen, G.; Yu, Y.; Huang, X. *Synlett* **2018**, *29*, 2087.
(b) Saikia, P.; Gogoi, S. *Adv. Synth. Catal.* **2018**, *360*, 2063.
(c) Saeed, A.; Haroon, M.; Muhammad, F.; Larik, F. A.; Hesham, E.-S.; Channar, P. A. *J. Organomet. Chem.* **2017**, *834*, 88.
(d) Saeed, A.; Larik, F. A. *Chem. Heterocycl. Compd.* **2016**, *52*, 450.
(e) Ashraf, Z. *Chem. Heterocycl. Compd.* **2016**, *52*, 149.
(f) Pal, S.; Chatare, V.; Pal, M. *Curr. Org. Chem.* **2011**, *15*, 782.
- [8] For selected recently examples, please see:
(a) Liu, G.; Kuang, G.; Zhang, X.; Lu, N.; Fu, Y.; Peng, Y.; Zhou, Y. *Org. Lett.* **2019**, *21*, 3043.
(b) Jiang, G.; Li, J.; Zhu, C.; Wu, W.; Jiang, H. *Org. Lett.* **2017**, *19*, 4440.
(c) Mandal, R.; Sundararaju, B. *Org. Lett.* **2017**, *19*, 2544.
(d) Kudo, E.; Shibata, Y.; Yamazaki, M.; Masutomi, K.; Miyauchi, Y.; Fukui, M.; Sugiyama, H.; Uekusa, H.; Satoh, T.; Miura, M.; Tanaka, K. *Chem.-Eur. J.* **2016**, *22*, 14190.
(e) Guo, X.-X. *J. Org. Chem.* **2013**, *78*, 166.
- [9] With alcohol: (a) Luo, M.-J.; Hu, M.; Song, R.-J.; He, D.-L.; Li, J.-H. *Chem. Commun.* **2019**, *55*, 1124.
With aldehyde: (b) Tao, L.-M.; Li, C.-H.; Chen, J.; Liu, H. *J. Org. Chem.* **2019**, *84*, 6807.
With anhydride: (c) Prakash, R.; Shekarrao, K.; Gogoi, S.; Boruah, R. C. *Chem. Commun.* **2015**, *51*, 9972.
- [10] Mihara, G.; Ghosh, K.; Nishii, Y.; Miura, M. *Org. Lett.* **2020**, *22*, 5706.
- [11] For examples on the annulation of vinylene carbonate:
(a) Li, X.; Huang, T.; Song, Y.; Qi, Y.; Li, L.; Li, Y.; Xiao, Q.; Zhang, Y. *Org. Lett.* **2020**, *22*, 5925.
(b) Ghosh, K.; Nishii, Y.; Miura, M. *Org. Lett.* **2020**, *22*, 3547.
(c) Ghosh, K.; Nishii, Y.; Miura, M. *ACS Catal.* **2019**, *9*, 11455.
- [12] (a) Raghuvanshi, K.; Zell, D.; Ackermann, L. *Org. Lett.* **2017**, *19*, 1278.
(b) Shankar, M.; Ghosh, K.; Mukherjee, K.; Rit, R. K.; Sahoo, A.

- K. *Org. Lett.* **2018**, *20*, 5144.
- (c) Rit, R. K.; Ramu Yadav, M.; Sahoo, A. K. *Org. Lett.* **2012**, *14*, 3724.
- (d) Ramu Yadav, M.; Rit, R. K.; Sahoo, A. K. *Org. Lett.* **2013**, *15*, 1638.
- (e) Ramu Yadav, M.; Rit, R. K.; Sahoo, A. K. *Chem.-Eur. J.* **2012**, *18*, 5541.
- [13] (a) Yadav, M. R.; Rit, R. K.; Shankar, M.; Sahoo, A. K. *J. Org. Chem.* **2014**, *79*, 6123.
- (b) Yadav, M. R.; Shankar, M.; Ramesh, E.; Ghosh, K.; Sahoo, A. K. *Org. Lett.* **2015**, *17*, 886.
- [14] Santhi, J.; Baire, B. *Chem. Asian J.* **2019**, *14*, 3161.
- [15] Toure, M.; Jaime-Figueroa, S.; Burslem, G. M.; Crews, C. M. *Eur. J. Org. Chem.* **2016**, 4171.
- [16] Izumi, T.; Nishimoto, Y.; Kohei, K.; Kasahara, A. *J. Heterocycl. Chem.* **1990**, *27*, 1419.
- [17] Colonge, J.; Boide, P. C. R. *Hebd. Seances Acad. Sci.* **1954**, 239, 1047.
- [18] Kinder, M. A.; Kopf, J.; Margaretha, P. *Tetrahedron* **2000**, *56*, 6763.
- [19] Mal, D.; Bandyopadhyay, M.; Ghorai, S. K.; Datta, K. *Tetrahedron Lett.* **2000**, *41*, 3677.

(Li, L.; Fan, Y.)