以环丙烷限制构象的新型双酰胺的设计、合成、抗癌活性及计算分析

陈睿嘉^a 周 聪^a 逢锡文^a 刘佳君^b
顾玉诚^c 刘建文^b 李 忠^{*,a}^(a) 华东理工大学药学院 上海市化学生物学(芳香杂环)重点实验室 上海 200237)^(b) 华东理工大学药学院 生物反应器工程国家重点实验室&上海市新药设计重点实验室 上海 200237)^(c) 先正达 Jealott's Hill 国际研发中心 英国 RG42 6EY)

摘要 为了探索具有新型骨架的抗癌剂,设计了一系列的以环丙烷限制构象的双酰胺类化合物.大多数含有顺式环丙烷中心的外消旋体对四种肿瘤细胞系(MCF-7、BGC-823、HepG2 和 NCI-H460)具有良好的抑制活性.其中,消旋体顺式-*N*¹-(2-甲基-4-(全氟丙-2-基)苯基)-*N*²-(4-苯氧基苄基)环丙烷-1,2-二羧酰胺(**1-19**)对 MCF-7 的 IC₅₀ 值为(8.38±0.67) mg·L⁻¹,显示出可进一步优化为抗癌先导物的潜力.此外,甾醇硫酸酯酶(steryl-sulfatase, STS)被 PharmMapper 预测为可能的靶标蛋白,并由分子对接实验进行了验证.

关键词 双酰胺; 环丙烷; 抗癌活性; 靶标预测; 分子对接

Design, Synthesis, Anti-cancer Activities and Computational Analysis of Novel Diamides Conformationally Restricted by Cyclopropane

Chen, Ruijia^a Zhou, Cong^a Pang, Xiwen^a Liu, Jiajun^b
Gu, Yucheng^c Liu, Jianwen^b Li, Zhong^{*,a}^(a) Shanghai Key Laboratory of Chemical Biology, School of Pharmacy, East China University of Science and Technology, Shanghai 200237)^(b) State Key Laboratory of Bioreactor Engineering & Shanghai Key Laboratory of Drug Design, School of Pharmacy, East China University of Science and Technology, Shanghai 200237)^(c) Syngenta Jealott's Hill International Research Centre, RG42 6EY, UK)

Abstract A series of diamides conformationally restricted by cyclopropane were designed and synthesized. Most of the racemates containing *cis*-cyclopropane possessed promising inhibitory activity against four cancer cell lines (MCF-7, BGC-823, HepG2 and NCI-H460). In particular, racemate *cis*-*N*¹-(2-methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(4-phenoxybenzyl)cyclopropane-1,2-dicarboxamide (**1-19**) showed the potential for further optimization as an anti-cancer lead with the IC₅₀ value of (8.38±0.67) mg·L⁻¹ on MCF-7. Moreover, steryl-sulfatase (STS) was predicted as the potential target protein and molecular docking was performed to explore the binding mode.

Keywords diamides; cyclopropane; anti-cancer activities; target prediction; docking

1 Introduction

Cancer is a leading cause of death and one of the barriers threatening public health. Due to the aging of population and increasing prevalence of established risk factors such as smoking and physical inactivity, the burden of cancer incidence and mortality is growing rapidly around the world.^[1] According to the latest Global Cancer Statis-

tics, there were approximately 19.3 million new cancer cases and 10 million cancer deaths worldwide in 2020.^[2] Chemotherapy is recognized as an effective approach for cancer treatment. Since 1940s, over 50 licensed chemotherapeutic drugs have been introduced for clinical use.^[3] However, few drugs could satisfy the needs of clinicians and patients nowadays. The problems of drug, such as low effectiveness, weak selectivity, side effects and potential

* Corresponding author. E-mail: lizhong@ecust.edu.cn

Received June 29, 2021; revised August 2, 2021; published online September 2, 2021.

Project supported by the National Key Research Program of China (No. 2017YFD0200505).

国家重点研发计划(No. 2017YFD0200505)资助项目.

sequelae, are still the public concerns.^[4-5] Therefore, the search for lead-compounds with novel skeleton and development of drugs with new mode of action remain the goals in pharmaceutical industry.

The role of diamide derivatives has gained increasing prominence in the development of drugs. Each amide bond, which is highly polar and stable, provides two hydrogen bonding sites for diamides.^[6] And the planarity of amide bonds will install conformational rigidity on the molecules.^[7] The unique chemical properties of diamides contribute their broad biological potencies. Over the last decade, various diamide derivatives have been reported as anti-cancer agents (Figure 1).^[8-9] In addition, diamides are embedded in many bioactive compounds with analgesic, anti-inflammatory, antimicrobial, antiviral, and insecticidal activities.^[10-14]

At the same time, cyclopropane is a significant ring system in pharmaceutical chemistry. (A) Cyclopropane has strain energy in the tension structure. (B) The three C—C bonds of cyclopropane are in the same plane. (C) The vibrational frequencies of C—H bonds in cyclopropane are similar to those of olefins. (D) The six C—H bonds of cyclopropane are relatively rigid in orientation. Due to these four properties, cyclopropane is considered a privileged terminal substituent in drug discovery to improve the pharmaceutical properties of drug molecules, such as potency, selectivity and stability.^[15-16] For decades, cyclopropane fragments have been introduced in a few drugs or agrochemicals, exemplified by naltrexone, cabozantinib, prothioconazole and phenothrin (Figure 2A).^[17-20] Besides,

the characteristic structural rigidity of cyclopropane makes it an effective unit to control the conformation of a molecule. It can serve as a “tether” to restrict the orientation of substituents.^[21] In medicinal research, cyclopropyl rings are often introduced into drug molecules for further optimization because of the hypothesis that the conformation of resulting fixed molecule would mimic the bioactive conformation of the original one (Figure 2B).^[22-24] And as a result, the fixed molecules would have less conformational entropy upon binding to the target proteins, stronger potency and higher selectivity.

Previous reports showed that the relative spatial distance between the two terminal groups of diamides determines the activities of compounds.^[25] The diamides with similar terminal groups but different spatial conformations usually have different potency or target selectivity.^[26-28] Herein, in order to investigate potential anti-cancer agent with novel “chemical space”, 1,2-disubstituted cyclopropane was introduced into diamides as central component (Scheme 1). Consequently, substituents were restricted into *trans* or *cis*-conformations. Perfluoroisopropyl aniline was fixed in design progress since several reports demonstrated that this moiety could improve the potency or pharmacokinetic profile of anti-cancer agents including several diamides.^[29-31] A total of 31 diamide racemates were designed and synthesized. And their anti-cancer activities against four tumor cell lines, including MCF-7 (breast epithelial adenocarcinoma), BGC-823 (gastric cancer), HepG2 (hepatocellular liver carcinoma) and NCI-H460 (lung cancer), were evaluated. *cis*-Cyclopropane was then identified as the preferred

Figure 1 Selected diamides with biological activities

(A) Representative drugs and argichemicals containing cyclopropane

(B) Conformation restriction strategy through cyclopropane in drug research

Figure 2 Representative cases for introduction of cyclopropane in drugs and agrichemicals **Scheme 1** Molecular design of novel diamides conformationally restricted by cyclopropane

center for anti-cancer diamides. Furthermore, the potential target protein was identified by pharmacophore-mapping and assessed through molecular docking.

2 Results and discussion

2.1 Chemistry

As illustrated in Scheme 2, the title racemates were syn-

thesized following the reported procedures with some necessary modifications.^[32-34] The intermediate 2-methyl-4-(perfluoropropan-2-yl)aniline (**2**) was obtained by the alkylation of commercially available 2-methylaniline with 2-iodoperfluoropropane. And the *cis*- and *trans*-title racemates were then obtained by 2 different routes. The structures of all racemates were characterized by NMR and ESI analysis.

Scheme 2 General synthetic procedure for the title racemates

2.1.1 Synthetic route for title *cis*-cyclopropyl racemates **1-1** to **1-30**

Diethyl cyclopropane-1,2-dicarboxylate (**3**) was prepared from methyl 2-chloroacetate and ethyl acrylate in the presence of sodium hydride (NaH). **3** was hydrolyzed and then acidified to get the mixture of *cis*- and *trans*-cyclopropane-1,2-dicarboxylic acid **4**. Subsequently, the crude **4** was transformed to the key intermediate 1,2-cyclopropane dicarboxylic anhydride **5** using acetic anhydride (Ac_2O) at 200°C .

This anhydride had two advantages. First, it was unnecessary to separate the *cis*- and *trans*-acids because the *trans*- one would convert to its isomer at high temperature.^[35] Second, the anhydride was an activated intermediate for next amidation. The amidation of **2** was a challenge due to the electron-withdrawing effect of its *para*-fluoroisopropyl and the steric hindrance of its *ortho*-methyl. While traditional coupling reagents such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), 2-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) and benzotriazol-1-yl-oxy-tripyrrolidinophosphonium hexafluorophosphate (PyBOP), all failed for this amidation, **6** could be prepared in high yield from a mechanical-stirring heterogeneous reaction of **2** and **5**.

The reported procedures for amidation between anhydride and aniline were low in yield and inconvenient, because base was added as deacid reagent and the resulting mixture was then acidified to obtain the desired free acid.

After survey of various parameters including base, solvent and temperature, it was found that **5** could be converted to **6** in over 90% HPLC area percent without base in dichloromethane (DCM) after 5 h at 35°C . What is more, owing to the poor solubility in DCM of **6**, it could be easily obtained by filtration.

Finally, in the presence of HATU and *N,N*-diisopropylethylamine (DIPEA), **6** coupled with different amines to afford the title diamides **1-1** to **1-30** in about 70% yield.

2.1.2 Synthetic route for title *trans*-cyclopropyl racemate **1-31**

Starting from commercially available diethyl *trans*-cyclopropane-1,2-dicarboxylate **7**, **1-31** was obtained through two amidations. **7** was firstly hydrolyzed to get **8**. And then **8** coupled with **2** using the combination of *N*-methylimidazole (NMI) and *N,N,N',N'*-tetramethylchloroformamidinium hexafluorophosphate (TCFH).

NMI-TCFH is a mild and efficient method for the coupling between electron-deficient aniline and hindered carboxylic acid.^[36] At the same time, the stereogenic integrity would be maintained. NMI-TCFH could provide the intermediate **9** in about 40% isolated yield while other coupling agents still could not lead to satisfactory yields as mentioned above. In addition, running this reaction in reflux reduced the reaction time and slightly improved the yield.

After another hydrolysis, **9** transformed to acid **10**. Finally, the title *trans*-**1-31** was obtained through HATU-DIPEA in about 70% yield.

2.1.3 Conformational analysis of title racemates

The conformations of **1-2** (*cis*) and **1-31** (*trans*) were examined based on ^1H NMR and NOESY NMR experiments. The ^1H NMR spectrum revealed that the chemical shifts for H2 and H3 of **1-2** were respectively between δ 1.56~1.44 and 1.23~1.12. And the chemical shifts for H2 and H3 of **1-31** were similar in a range of δ 1.21~1.13. What's more, Nuclear overhauser effects (NOEs) were observed between H1-H2, H1-H4 and H2-H4 for **1-2** (Figure 3). Besides NOEs between H1-H2 and H3-H4 were observed for **1-31**. These facts indicated that **1-2** and **1-31** assumed the *cis*- and *trans*-conformation, respectively.

Figure 3 NOEs patterns (blue two-way arrows) for **1-2** and **1-31**

2.2 Biological activity

The anti-cancer activities of all racemates against MCF-7, BGC-823, HepG2 and NCI-H460 cell lines were evaluated *in vitro* by the standard 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl tetrazolium bromide (MTT) assay using 5-fluorouracil (5-Fu) as a positive control. The racemates whose inhibition rate was greater than 50% at $200\ \mu\text{g}\cdot\text{mL}^{-1}$, were chosen to study their IC_{50} values. The results were summarized in Tables 1~3.

Two racemates **1-2** and **1-31** were firstly designed and synthesized, which possessed the same terminal groups but different conformations. As shown in Table 1, no obvious anti-cancer activities were observed for *trans*-configurational **1-31**. To our delight, *cis*-**1-2** showed moderate inhibitory activities against all four cancer cell lines at the concentration of $200\ \mu\text{g}\cdot\text{mL}^{-1}$. In particular, **1-2** exhibited inhibition rates of more than 80% on both MCF-7 and NCI-H460 at this concentration. The results indicated that *cis*-cyclopropane was an acceptable central component of diamides for further study.

Four another diamides with different substituents R^1 (**1-1**: methyl; **1-3**: *tert*-butyl; **1-4**: benzyl; **1-30**: phenethyl) were then screened for their activities as the probes. As presented in Table 2, those racemates exhibited moderate

Table 1 Growth-inhibition against four cancer cell lines of diamides containing *trans* and *cis*-cyclopropanes

Compd.	Structure (Relative stereochemistry)	Growth-inhibition ($200\ \mu\text{g}\cdot\text{mL}^{-1}$)			
		MCF-7	BGC-823	HepG2	NCI-H460
1-2		83.46 ± 3.64	54.78 ± 1.16	48.13 ± 1.69	97.31 ± 0.18
1-31		19.42 ± 1.19	-28.92 ± 2.03	38.24 ± 5.37	14.20 ± 6.58

Table 2 Growth-inhibition against four cancer cell lines of some title diamides at $200\ \mu\text{g}\cdot\text{mL}^{-1}$

Compd.	R^1	Growth-inhibition ($200\ \mu\text{g}\cdot\text{mL}^{-1}$)			
		MCF-7	BGC-823	HepG2	NCI-H460
1-1	Methyl	37.23 ± 3.34	32.39 ± 1.17	23.47 ± 4.79	47.37 ± 0.50
1-2	Ethyl	83.46 ± 3.64	54.78 ± 1.16	48.13 ± 1.69	97.31 ± 0.18
1-3	<i>tert</i> -Butyl	85.73 ± 2.92	4.17 ± 0.52	57.57 ± 1.57	94.34 ± 2.25
1-4	Benzyl	96.83 ± 0.69	97.75 ± 0.08	99.14 ± 0.28	98.42 ± 0.16
1-30	Phenethyl	90.96 ± 0.28	91.19 ± 0.99	93.00 ± 8.44	97.46 ± 0.16

activity against four cancer cell lines at a concentration of $200\ \mu\text{g}\cdot\text{mL}^{-1}$. The primary bioassay results suggested that the introduction of an aromatic group at R^1 was positive because **1-4** (benzyl) and **1-30** (phenethyl) inhibited cell viability to a greater extent than the others. Since **1-4** was most active against all the cell lines with 96%~99% growth inhibition at $200\ \mu\text{g}\cdot\text{mL}^{-1}$, it was considered as the starting point for further optimization.

Then, the influence on anti-cancer activities of the position for substituents in R^1 was explored. Six diamides substituted by methyl (**1-5**, **1-6** and **1-7**) or chlorine (**1-8**, **1-9**, **1-10**) were designed and synthesized. As shown in Table 3, the introductions of methyl or chlorine led to increases in activities against 4 cell lines except that **1-10** (2-chloro) displayed a slight decreased activity against MCF-7. And a substituent on 4-position was satisfactory because both the potencies of **1-5** and **1-8** were respectively stronger than their isomers.

Aimed to further investigate the structure-activity relationship (SAR) of our diamides, different substituents were introduced to obtain the title diamides **1-11** to **1-22**. Most

of those racemates exhibited good anti-cancer activities against the cancer cell lines (Table 3). And several general trends emerged from the results. Most mono-substituted derivatives had stronger activities comparing with the non-(**1-4**) or di-substituted (**1-21** and **1-22**) racemates. And neither difluoromethoxyl (**1-15**) nor *tert*-butyl (**1-16**) substituents had good effects on the activity. While the introduction of fluorine (**1-11**) was dissatisfactory, brominated (**1-12**) and iodized (**1-13**) racemates both showed better activities than **1-4** against MCF-7, BGC-823 and NCI-H460.

On the potency against MCF-7, the electronic effect of substituents has little influence. Methyl (**1-5**), methoxyl (**1-14**), trifluoromethyl (**1-18**), phenoxyphenyl (**1-19**) and phenyl (**1-20**) all improved the anti-cancer activities whereas *tert*-butyl (**1-16**) showed less activities than **1-4**. It is assumed that spatial plane hydrophobic substituents were preferred. It was worth noting that **1-19** displayed the strongest activities against MCF-7 with an IC_{50} value of $8.38\ \mu\text{g}\cdot\text{mL}^{-1}$, which was comparable to that of 5-Fu.

Table 3 The IC_{50} against four cancer cell lines of diamides **1-1** to **1-30**

Compd.	R^1	$\text{IC}_{50}/(\mu\text{g}\cdot\text{mL}^{-1})$			
		MCF-7	BGC-823	HepG2	NCI-H460
1-1	Methyl	>200	>200	>200	>200
1-2	Ethyl	33.81 ± 2.06	>200	>200	29.80 ± 2.91
1-3	<i>tert</i> -Butyl	29.27 ± 5.83	>200	>200	38.38 ± 1.40
1-4	Benzyl	40.79 ± 2.18	50.90 ± 0.99	26.09 ± 2.97	26.78 ± 2.78
1-5	4-Methylbenzyl	32.29 ± 1.7	21.25 ± 3.03	10.44 ± 1.60	20.38 ± 1.21
1-6	3-Methylbenzyl	40.35 ± 5.19	35.50 ± 3.97	35.01 ± 1.54	30.59 ± 1.18
1-7	2-Methylbenzyl	33.80 ± 1.72	25.42 ± 2.29	23.20 ± 1.07	21.60 ± 1.60
1-8	4-Chlorobenzyl	36.89 ± 2.29	19.47 ± 5.37	17.65 ± 1.24	10.06 ± 1.82
1-9	3-Chlorobenzyl	36.80 ± 4.39	30.58 ± 2.65	28.62 ± 0.81	22.48 ± 0.71
1-10	2-Chlorobenzyl	80.47 ± 6.93	31.75 ± 4.74	22.75 ± 0.84	21.61 ± 1.30
1-11	4-Fluorobenzyl	67.45 ± 2.43	29.55 ± 2.70	27.07 ± 1.05	25.27 ± 2.85
1-12	4-Bromobenzyl	17.65 ± 0.99	28.29 ± 2.54	21.17 ± 0.81	21.01 ± 1.25
1-13	4-Iodobenzyl	27.57 ± 6.26	26.24 ± 2.70	26.29 ± 1.43	26.15 ± 1.36
1-14	4-Methoxybenzyl	20.78 ± 1.38	24.51 ± 2.57	24.14 ± 1.23	25.17 ± 1.84
1-15	4-Difluoromethoxybenzyl	75.61 ± 14.80	50.03 ± 2.97	39.52 ± 6.27	56.80 ± 10.58
1-16	4- <i>tert</i> -Butylbenzyl	>200	92.61 ± 16.39	>200	95.35 ± 17.62
1-17	4-Cyanobenzyl	47.75 ± 9.06	42.49 ± 4.12	37.92 ± 2.94	30.26 ± 3.15
1-18	4-Trifluoromethylbenzyl	25.66 ± 1.11	21.39 ± 1.11	21.23 ± 1.99	15.12 ± 1.43
1-19	(4-Phenoxyphenyl)methyl	8.38 ± 0.67	22.67 ± 0.80	19.30 ± 1.10	20.49 ± 1.32
1-20	4-Phenylbenzyl	25.88 ± 2.95	24.97 ± 1.27	27.06 ± 3.07	22.94 ± 1.15
1-21	2,4-Difluorobenzyl	40.84 ± 1.24	51.29 ± 1.09	23.52 ± 2.50	28.35 ± 2.28
1-22	2,6-Dichlorobenzyl	>200	>200	19.74 ± 0.79	39.89 ± 9.27
1-23	3,4,5-Trimethoxybenzyl	19.16 ± 3.01	34.39 ± 3.52	27.22 ± 2.61	20.39 ± 1.41
1-24	(Tetrahydrofuran-3-yl)methyl	153.9 ± 20.2	50.30 ± 6.65	46.88 ± 5.39	56.51 ± 11.21
1-25	(6-Chloropyridine-3-yl)methyl	27.12 ± 0.92	32.39 ± 3.46	34.15 ± 0.94	29.36 ± 1.79
1-26	(2-Chlorothiazole-5-yl)methyl	>200	>200	>200	>200
1-27	Furan-2-ylmethyl	25.8 ± 1.32	32.39 ± 4.40	15.42 ± 4.40	27.03 ± 2.49
1-28	Thiophene-3-ylmethyl	83.43 ± 4.37	>200	>200	>200
1-29	Naphthalene-1-ylmethyl	32.53 ± 2.82	22.91 ± 1.20	23.55 ± 1.97	20.52 ± 0.98
1-30	Phenethyl	59.05 ± 3.60	53.12 ± 1.16	26.76 ± 1.18	30.37 ± 2.36
5-Fu	—	8.23 ± 1.33	10.31 ± 1.40	7.09 ± 1.61	2.26 ± 0.80

For the potency against BGC-823, the replacement of hydrogen with only one halogen atom at any position of amino toluene was positive. The activities against BGC-823 of most racemates remained at the same level with IC_{50} values range from $20 \mu\text{g}\cdot\text{mL}^{-1}$ to $30 \mu\text{g}\cdot\text{mL}^{-1}$, so no clear SAR was observed. In particular, chlorinated **1-8** was most active with an IC_{50} value of $19.47 \mu\text{g}\cdot\text{mL}^{-1}$.

1-5 (4-Methyl) and **1-8** (4-chloro) were the two most active racemates against HepG2. And **1-5** has a comparable anti-cancer activity to 5-Fu with an IC_{50} value of $10.44 \mu\text{g}\cdot\text{mL}^{-1}$. While, **1-13** (4-iodobenzyl), **1-16** (4-*tert*-butylbenzyl) and **1-20** (4-phenylbenzyl) all displayed weaker potency than **1-4** against HepG2. It implied that a big steric hindrance would lead to a decreased activity.

Most of the substituents at 4-position increased the activity against NCI-H460. It was notable that **1-8** (4-chlorobenzyl) and **1-18** (4-trifluoromethyl benzyl), two racemates with electron-withdrawing substituents, both exhibited significant inhibitory activities against NCI-H460 with IC_{50} values of 10.06 and $15.12 \mu\text{g}\cdot\text{mL}^{-1}$, respectively.

To further evaluate the viability of amino toluene, several heterocycle-containing (**1-23** to **1-28**) and polycyclic-aromatic-containing (**1-29**) racemates were synthesized. The bioassay results are shown in Table 3. Changing the phenyl ring to tetrahydrofuran (**1-24**) led to the decrease of activities on all cell lines. It suggested once again that the aromaticity of R^1 was a critical structural element for our diamides to achieve high anti-cancer activities. Neither **1-26** [(2-chlorothiazole-5-yl)methyl] nor **1-28** (thiophene-3-ylmethyl) could inhibit the growth of those four cancer cells. It implied that the introduction of sulfur atom might had a negative effect. Fortunately, the racemates modified by pyridine (**1-25**) and furfuran (**1-27**) both exhibited better activities against MCF-7 and BGC-823 than **1-4**. The improvement might be a tribute to the addition of nitrogen and oxygen atoms which could usually increase the interaction of drug molecules with target proteins.

2.3 Target recognition

PharmMapper, a web portal, provides services to identify potential target candidates for small molecules through pharmacophore mapping approach.^[37-39] PharmMapper has a database with 23236 proteins and works by ‘probing’ given compounds into the database.

The promising anti-cancer activities of our diamides against MCF-7 encouraged us to further explore their potential action mechanism and target protein. The most active racemates **1-12** (4-bromobenzyl), **1-14** (4-methoxybenzyl), **1-19** [(4-phenoxyphenyl)methyl] and **1-23** (3,4,5-trimethoxybenzyl) were submitted to PharmMapper. And 300 putative targets for each racemate were reserved by PharmMapper. Those targets linked to human cancers were further manually identified according to the published reports.

After filtering by normalized fit score >0.9 , the details of racemates and their candidate target proteins were listed in Table 4. The results ‘fished’ by PharmMapper for our

four racemates were similar. As described in Figure 4, 12 proteins were identified as potential targets. All given racemates have high normalized fit score to five proteins including steryl-sulfatase (STS, PDB ID: 1p49), ephrin type-B receptor 4 (EphB4, PDB ID: 2vwu), kinesin-like protein KIF11 (KIF11, PDB ID: 2pg2), thymidylate synthase (TS, PDB ID: 1juj) and epidermal growth factor receptor (EGFR, PDB ID: 2ity). And it was worth noting that STS kept the highest score and EphB4 remained the sec-

Table 4 Top target proteins of active diamides and their scores^a

Compd.	Pharma model (PDB ID)	Normalized fit score
1-12	1p49	0.996
	2vwu	0.996
	2brb	0.988
	2pg2	0.980
	1juj	0.963
	2g01	0.956
	2ity	0.935
	2ovm	0.929
	1o4i	0.906
	1p49	0.996
1-14	2vwu	0.995
	2pg2	0.979
	1juj	0.977
	1ya8	0.957
	2ovm	0.944
	2ity	0.925
	1p49	0.996
	2vwu	0.994
	2pg2	0.980
	1juj	0.971
1-19	1ya8	0.971
	2g01	0.946
	2ity	0.928
	1o4i	0.903
	1p49	0.995
	2vwu	0.993
	2c3i	0.987
	2pg2	0.979
	1ya8	0.972
	1juj	0.967
1-23	2ocf	0.942
	2ity	0.930

^a 1p49: Steryl-sulfatase, linked to tumors such as breast or prostate cancers; 2vwu: Ephrin type-B receptor 4, linked to tumors such as colon and lung cancers; 2brb: Serine/threonine-protein kinase Chk1, linked to tumors such as Colon and lung cancers; 2pg2: Kinesin-like protein KIF11, linked to tumors such as breast and lung cancers; 1juj: Thymidylate synthase, linked to tumors such as brain and breast cancers; 2g01: Mitogen-activated protein kinase 8, linked to tumors such as gastric and prostate cancers; 2ity: Epidermal growth factor receptor, linked to tumors such as breast and liver cancers; 2ovm: Progesterone receptor, linked to tumors such as breast cancer; 1o4i: Proto-oncogene tyrosine-protein kinase Src, linked to tumors such as lung and pancreatic cancer; 1ya8: Liver carboxylesterase 1, linked to tumors such as liver and lung cancer; 2c3i: Proto-oncogene serine/threonine-protein kinase Pim-1, linked to tumors such as pancreatic and prostate cancer; 2ocf: Estrogen receptor, linked to tumors such as lung and prostate cancers.

and highest score for all the four racemates.

STS is the key enzyme responsible for the transform of dehydroepiandrosterone sulfate (DHEA-S) to DHEA.^[40-41] STS and estrogen sulfotransferase (ETS) together regulate the steroid homeostasis. Recently, STS has been identified as a promising pharmaceutical target for the treatment of hormones-sensitive diseases like breast and prostate cancer.^[42-43] And several STS inhibitors, such as estrone 3-*O*-sulfamate (EMATE) and 667 COUMATE, have been evaluated in clinical trials.^[44-45] Meanwhile, EphB4 plays

Figure 4 Racemate-target network

The network was constructed using Cytoscape. The targets in the outer ring are black; the racemate numbers in the inner ring are black; the racemate-target relationships are marked with grey lines.

an essential role in vascular development.^[46] However, the potential of EphB4 as an anti-cancer target remained to be evaluated because of its bidirectional signalling mechanisms. Both overexpression and silence of EphB4 have been linked to tumorigenicity.^[47]

The highest scores for all racemates and crucial role in breast cancer development of STS indicated that it might act as the potential anti-cancer target of our racemates.

2.4 Molecular docking

Molecular docking study was performed to further explore the potential binding mode of our racemates to human STS. **1-19**, the most active one, was selected as the ligand.

The STS catalytic site is a linear cavity surrounded by hydrophobic residues.^[48-49] As shown in Figure 5, both compounds of (1*S*,2*R*) and (1*R*,2*S*) conformation fit well with the pocket. In the binding mode of (1*S*,2*R*) **1-19**, each of the two amide moieties was involved in one hydrogen bonding interactions with Arg98 of STS. The characteristic perfluoroisopropyl moiety bounded in the hydrophobic groove at the entrance of the catalytic site. Resulted from the conformational restrict effect of cyclopropanyl core, the 4-phenoxyphenyl moiety extended into the deeper cave of the binding pocket. In the binding mode of (1*R*,2*S*) **1-19**, the perfluoroisopropyl and 4-phenoxyphenyl moieties laid in the similar position compare to that of (1*S*,2*R*) one. While, the amide moieties formed two hydrogen bonds with Gly181 and Thr484 of STS.

The binding mode analysis further demonstrated the rational selection of human STS target and significant anti-cancer activities of title diamides. And it laid a foundation for further optimization on chemical structure of those diamides with *cis*-conformational cyclopropane.

3 Conclusions

In summary, 31 diamide derivatives conformationally restricted by cyclopropane were designed and synthesized. The bioassay results showed that most of our racemates with *cis*-cyclopropane possessed certain anti-cancer activities on four tumor cell lines including MCF-7, BGC-823, HepG2 and NCI-H460. And racemate **1-19** exhibited the best inhibition on MCF-7 with an IC₅₀ value of (8.38 ± 0.67) mg·L⁻¹, which was comparable to that of 5-Fu. It suggested that **1-19** might be a lead for the discovery of novel anti-cancer agents. What's more, STS was identified as the potential target for our anti-cancer racemates aided by PharmMapper. As verification, molecular docking revealed the possible binding mode of **1-19** to STS. Further study of those diamides are in process in our laboratory.

4 Experimental section

4.1 Instruments and reagents

Melting points (m.p.) were recorded on a Büchi Melting Point B-540 melting point apparatus (Büchi, Flawil, Swit-

Figure 5 **1-19** binding in human STS

Overview of binding pocket on Human STS (A); 3D binding mode of (1*S*,2*R*) (B) and (1*R*,2*S*) (C) **1-19**.

zerlandand) without correction. ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra were recorded on a Bruker AM-400 spectrometer (Bruker, Fällanden, Switzerland) with $\text{DMSO}-d_6$ as solvent and TMS as internal standard. High-resolution electron mass spectra (ESI-TOF) were performed on a Waters Xevo G2 TOF spectrometer (Waters, Milford, America). Specific rotations were recorded on a Rudolph Analytical Autopol III polarimeter (Rudolph, Hackettstown, America). Analytical thin-layer chromatography (TLC) was performed on precoated plates (silica gel 60 F254), and spots were visualized with ultraviolet (UV) light (254 nm). All column chromatography was performed using 200~300 mesh silica gel.

Unless otherwise stated, all reactions were carried out under inert gas atmosphere. Unless otherwise noted, all reagents were obtained from commercial sources and used without further purification. Solvents were purchased from commercial sources, degassed by purging thoroughly with nitrogen and dried over activated molecular sieves.

4.2 Synthetic routes

The intermediates **2**, **3** and **5** were prepared according to the reported methods with necessary modifications and their characterization data was same as that in reference.^[29-31]

4.2.1 Synthesis of 2-methyl-4-(perfluoropropan-2-yl)-aniline (**2**)

To a mixture of *o*-toluidine (10.0 mmol) in methyl *tert*-butyl ether (MTBE, 150 mL) and water (150 mL) were added potassium carbonate (K_2CO_3 , 12.0 mmol), hexadecyl trimethyl ammonium bromide (CTMAB, 11.0 mmol) and sodium hydrosulfite ($\text{Na}_2\text{S}_2\text{O}_4$, 12.0 mmol). The resulting mixture was stirred at room temperature for 20 min and then treated with 1,1,1,2,3,3,3-heptafluoro-2-iodopropane (12 mmol). After completion, the two phases were separated and the water phase was extracted with ethyl acetate (EA, 80 mL) twice. The combined organic phases were dried over anhydrous sodium sulfate (Na_2SO_4) and concentrated *in vacuo* to give yellow oil 2-methyl-4-(perfluoropropan-2-yl)aniline (**2**), which could be used in the next step without further purification.

4.2.2 Synthesis of dimethyl cyclopropane-1,2-dicarboxylate (**3**)

To a suspension of sodium hydride (NaH, 60% dispersion in mineral oil, 24.0 mmol) in dry toluene (80 mL) cooled with an ice bath was added a mixture of methyl 2-chloroacetate (20.0 mmol) and methyl acrylate (20.0 mmol) dropwise. The resulting mixture was then warmed to room temperature and stirred for another 3 h. After completion, the reaction mixture was quenched with crushed ice (100 g) and then extracted with EA (50 mL) twice. The combined organic phases were dried over Na_2SO_4 and concentrated *in vacuo* to obtain yellow oil dimethyl cyclopropane-1,2-dicarboxylate (**3**), which could be directly used in the next step.

4.2.3 Synthesis of 3-oxabicyclo[3.1.0]hexane-2,4-dione (**5**)

To a solution of **3** (8.0 mmol) in methanol (50 mL) was added a solution of sodium hydroxide (NaOH, 16.8 mmol) in water (10 mL). The resulting mixture was stirred at reflux for 4 h. After completion, the reaction mixture was concentrated *in vacuo* and diluted with EA (50 mL). Then the pH was adjusted to 5~6 with dilute hydrochloric acid (HCl). The aqueous phase was extracted twice with EA (30 mL). The combined organic phases were dried over Na_2SO_4 and concentrated *in vacuo* to give crude cyclopropane-1,2-dicarboxylic acid (**4**, 7.4 mmol, 92.5%), which could be used in the next without further purification.

A mixture of crude **4** (5.0 mmol) and acetic anhydride (Ac_2O , 10.0 mmol) was stirred under 200 °C for 5 h. After completion, half of the Ac_2O was removed *in vacuo* and the resulting mixture was cooled to room temperature. The formed white crystal was collected by filtration, washed with dichloromethane (DCM, 5 mL) twice and dried to obtain white solid 3-oxabicyclo[3.1.0]hexane-2,4-dione (**5**), yield 76.4%, m.p. 58.7~59.8 °C (lit.^[31] b.p. 56~58 °C).

4.2.4 Synthesis of *cis*-2-((2-methyl-4-(perfluoropropan-2-yl)phenyl)carbamoyl)cyclopropane-1-carboxylic acid (**6**)

To a suspension of **5** (9.5 mmol) in DCM (30 mL) was added **2** dropwise (10.0 mmol). The mixture was mechanically stirred under 35 °C for 5 h. After completion, the white precipitate was collected by filtration, washed with dichloromethane (DCM, 5 mL) for three times and dried to obtain white solid **6**, yield 94.2%. m.p. 163.4~164.0 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 12.19 (s, 1H), 9.67 (s, 1H), 7.70 (d, $J=8.5$ Hz, 1H), 7.51~7.41 (m, 2H), 2.38~2.26 (m, 4H), 2.10~2.00 (m, 1H), 1.49~1.39 (m, 1H), 1.22~1.15 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 171.31, 167.58, 139.61, 132.24, 127.13~126.85 (m), 125.13~124.98 (m), 123.17~122.93 (m), 121.88~121.53 (m), 120.72 (d, $J=20.8$ Hz), 119.92~118.65 (m), 92.75~89.77 (m), 23.33, 21.61, 17.89, 10.90; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.10~-75.22 (m, 6F), -181.32~-181.47 (m, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{15}\text{H}_{13}\text{F}_7\text{NO}_3$ ($\text{M}+\text{H}$)⁺ 388.0784, found 388.0783.

4.2.5 Synthesis of *trans*-2-(ethoxycarbonyl)cyclopropane-1-carboxylic acid (**8**)

To a solution of **7** (10.0 mmol) in methanol (50 mL) was added a solution of NaOH (10.5 mmol) in water (8 mL). The resulting mixture was stirred at 50 °C for 2 h. After completion, the reaction mixture was concentrated *in vacuo* and diluted with EA (50 mL). Then the pH was adjusted to 5~6 with dilute HCl solution. The aqueous phase was extracted twice with EA (30 mL). The combined organic phases were dried over Na_2SO_4 and concentrated *in vacuo* to give white solid **8**, which could be used in the next step without further purification.

4.2.6 Synthesis of ethyl *trans*-2-((2-methyl-4-(perfluoropropan-2-yl)phenyl)carbamoyl)cyclopropane-1-carboxylate (**9**)

To a suspension of **8** (8.0 mmol) in acetonitrile (50 mL) was added **2** (6.4 mmol) and NMI (28 mmol) in order. Then TCFH (10 mmol) was added into the stirring reaction mixture in portions. The resulting mixture was stirred at reflux for 5 h. After completion, the reaction mixture was concentrated *in vacuo*, diluted with EA (50 mL) and washed with dilute HCl (0.2 mol/L, 60 mL) and saturated brine (60 mL). The organic phase was separated, dried over Na₂SO₄ and concentrated *in vacuo*. The crude product was purified by flash column chromatography to obtain white solid **9**, yield 38.8%. m.p. 85.1~85.9 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.90 (s, 1H), 7.85 (d, *J*=8.6 Hz, 1H), 7.53~7.42 (m, 2H), 4.12 (q, *J*=7.1 Hz, 2H), 2.63~2.55 (m, 1H), 2.34 (s, 3H), 2.05~1.06 (m, 1H), 1.40~1.30 (m, 2H), 1.22 (t, *J*=7.1 Hz, 3H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 171.60, 168.84, 139.26, 131.53, 127.24~127.01 (m), 124.40~124.29 (m), 123.31~123.10 (m), 121.86~121.74 (m), 120.72 (d, *J*=21.2 Hz, 1H), 119.62~119.51 (m), 95.15~93.50 (m), 60.62, 23.36, 21.24, 17.93, 14.39, 14.02; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: -75.15 (d, *J*=7.5 Hz, 6F), -181.41 (hept, 1F); HRMS (ESI-TOF) calcd for C₁₇H₁₇F₇NO₃ (M+H)⁺ 416.1098, found 416.1099.

4.2.7 Synthesis of *trans*-2-((2-methyl-4-(perfluoropropan-2-yl)phenyl)carbamoyl)cyclopropane-1-carboxylic acid (**10**)

Using a similar method as the preparation of **8**, **9** was hydrolyzed to prepare crude white solid **10** with yield 87.4%, which was directly used in the next step.

4.2.8 Synthesis of the title compositions **1-1** to **1-31**

A suspension of **6** (5.0 mmol) in DCM (30 mL) was treated with HATU (5.5 mmol) followed by DIPEA (20 mmol). The resulting mixture was stirred for 30 min. Then to it was added different amines R¹NH₂ dropwise. The reaction was stirred for another 5 h. After completion, the reaction mixture was treated with saturated brine (20 mL). The organic phase was washed with dilute hydrochloric acid (0.2 mol/L, 20 mL) and brine (20 mL), dried over Na₂SO₄ and then concentrated *in vacuo*. The crude product was purified by flash column chromatography to obtain the title compositions **1-1**~**1-30**.

1-31 was prepared from **10** by a similar method for the preparation of **1-1**~**1-30**.

cis-N¹-Methyl-N²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-1**): White solid, yield 61%. m.p. 159.1~160.7 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.61 (s, 1H), 8.10~7.99 (m, 1H), 7.78 (d, *J*=8.5 Hz, 1H), 7.51~7.39 (m, 2H), 2.57 (d, *J*=4.6 Hz, 3H), 2.31 (s, 3H), 2.19~2.08 (m, 1H), 2.04~1.93 (m, 1H), 1.53~1.43 (m, 1H), 1.22~1.13 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 169.46, 168.50, 139.77, 131.70, 127.03~126.82 (m), 124.53~124.39 (m), 123.22~122.99 (m), 121.95~121.32 (m), 120.32 (d, *J*=20.1 Hz),

119.30~118.77 (m), 95.43~94.62 (m), 25.76, 23.44, 23.21, 17.84, 10.93; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: -75.14 (d, *J*=7.5 Hz, 6F), -181.37 (hept, *J*=7.5 Hz, 1F); HRMS (ESI-TOF) calcd for C₁₆H₁₅F₇N₂NaO₂ (M+Na)⁺ 423.0920, found 423.0920.

cis-N¹-Ethyl-N²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-2**): White solid, yield 74%. m.p. 182.4~183.9 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.61 (s, 1H), 8.13 (t, *J*=5.2 Hz, 1H), 7.80 (d, *J*=8.5 Hz, 1H), 7.51~7.39 (m, 2H), 3.13~3.00 (m, 2H), 2.31 (s, 3H), 2.19~2.07 (m, 1H), 2.05~1.93 (m, 1H), 1.56~1.44 (m, 1H), 1.23~1.12 (m, 1H), 1.00 (t, *J*=7.2 Hz, 3H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 168.81, 168.49, 139.78, 131.63, 127.02~126.83 (m), 124.45~124.36 (m), 123.18~122.95 (m), 121.90~121.52 (m), 120.29 (d, *J*=20.5 Hz), 119.05~118.69 (m), 92.44~89.76 (m), 33.60, 23.68, 23.18, 17.84, 14.63, 10.96; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: -75.16 (d, *J*=7.5 Hz, 6F), -181.38 (hept, *J*=7.5 Hz, 1F); HRMS (ESI-TOF) calcd for C₁₇H₁₇F₇N₂NaO₂ (M+Na)⁺ 437.1076, found 437.1075.

cis-N¹-(*tert*-Butyl)-N²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-3**): White solid, yield 81%. m.p. 185.7~187.4 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.61 (s, 1H), 7.87~7.76 (m, 2H), 7.49~7.39 (m, 2H), 2.32 (s, 3H), 2.11~1.99 (m, 2H), 1.51~1.42 (m, 1H), 1.23 (s, 9H), 1.22~1.13 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 168.79, 168.65, 139.89, 131.44, 127.05~126.75 (m), 124.12~124.01 (m), 123.22~122.92 (m), 121.92~121.53 (m), 120.16 (d, *J*=20.5 Hz), 119.06~118.61 (m), 92.27~90.05 (m), 50.29, 28.44, 24.29, 23.65, 17.95, 11.11; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: -75.23 (d, *J*=7.5 Hz, 6F), -181.51 (hept, *J*=7.5 Hz, 1F); HRMS (ESI-TOF) calcd for C₁₉H₂₂F₇N₂O₂ (M+H)⁺ 443.1570, found 443.1571.

cis-N¹-Benzyl-N²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-4**): White solid, yield 88%. m.p. 109.9~110.5 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.56 (s, 1H), 8.62 (t, *J*=5.9 Hz, 1H), 7.80 (d, *J*=8.5 Hz, 1H), 7.49~7.39 (m, 2H), 7.28~7.10 (m, 5H), 4.38~4.13 (m, 2H), 2.28 (s, 3H), 2.26~2.16 (m, 4H), 2.14~2.04 (m, 1H), 1.59~1.51 (m, 1H), 1.24~1.15 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 168.97, 168.25, 139.78, 139.34, 131.53, 128.04, 127.14, 127.04~126.75 (m), 126.55, 124.40~124.29 (m), 123.20~122.91 (m), 121.93~121.51 (m), 120.22 (d, *J*=20.2 Hz), 119.19~118.92 (m), 92.52~91.04 (m), 42.24, 23.65, 23.20, 17.87, 10.73; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: -75.16 (d, *J*=7.5 Hz, 6F), -181.46 (hept, *J*=7.5 Hz, 1F); HRMS (ESI-TOF) calcd for C₂₂H₁₉F₇N₂O₂ (M+H)⁺ 499.1232, found 499.1231.

cis-N¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-N²-(4-methylbenzyl)cyclopropane-1,2-dicarboxamide (**1-5**): White solid, yield 83%. m.p. 105.1~106.0 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.56 (s, 1H), 8.58 (t, *J*=5.9 Hz, 1H), 7.79 (d, *J*=8.6 Hz, 1H), 7.51~7.39 (m, 2H), 7.10 (d, *J*=8.0 Hz, 2H), 6.99 (d, *J*=8.0 Hz, 1H), 4.33~4.09 (m,

2H), 2.28 (s, 3H), 2.25~2.14 (m, 4H), 2.13~2.03 (m, 1H), 1.58~1.50 (m, 1H), 1.22~1.14 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 168.89, 168.25, 139.77, 136.31, 135.56, 131.49, 128.59, 127.15, 127.04~126.72 (m), 124.36~124.28 (m), 123.19~122.86 (m), 120.80~120.55 (m), 120.18 (d, $J=19.9$ Hz), 119.76~119.59 (m), 92.60~91.00 (m), 41.98, 23.68, 23.21, 20.52, 17.87, 10.70; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.16 (d, $J=7.5$ Hz, 6F), -181.49 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{23}\text{H}_{22}\text{F}_7\text{N}_2\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 491.1570, found 491.1569.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(3-methylbenzyl)cyclopropane-1,2-dicarboxamide (**1-6**): White solid, yield 78%. m.p. 91.0~91.7 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.55 (s, 1H), 8.59 (t, $J=5.8$ Hz, 1H), 7.79 (d, $J=8.5$ Hz, 1H), 7.49~7.37 (m, 2H), 7.12~6.94 (m, 4H), 4.34~4.14 (m, 2H), 2.28 (s, 3H), 2.25~2.14 (m, 4H), 2.14~2.05 (m, 1H), 1.61~1.52 (m, 1H), 1.24~1.14 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 168.97, 168.29, 139.78, 139.22, 137.19, 131.55, 127.93, 127.78, 127.18, 127.55~126.73 (m), 124.44~124.28 (m), 124.27, 123.15~122.85 (m), 121.90~121.54 (m), 120.26 (d, $J=20.5$ Hz), 119.07~118.71 (m), 92.38~90.43 (m), 42.25, 23.69, 23.22, 20.71, 17.85, 10.76; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.18 (d, $J=7.5$ Hz, 6F), -181.42 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{23}\text{H}_{21}\text{F}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 513.1390, found 513.1390.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(2-methylbenzyl)cyclopropane-1,2-dicarboxamide (**1-7**): White solid, yield 81%. m.p. 126.4~127.8 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.56 (s, 1H), 8.49 (t, $J=5.6$ Hz, 1H), 7.80 (d, $J=8.5$ Hz, 1H), 7.50~7.39 (m, 2H), 7.21~7.16 (m, 1H), 7.14~7.05 (m, 2H), 7.04~6.96 (m, 1H), 4.33~4.14 (m, 2H), 2.29 (s, 3H), 2.25~2.16 (m, 4H), 2.15~2.07 (m, 1H), 1.60~1.51 (m, 1H), 1.24~1.16 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 169.03, 168.37, 139.63, 136.68, 135.44, 131.60, 129.73, 127.44, 127.03~126.75 (m), 126.69, 125.52, 124.48~124.36 (m), 123.16~122.87 (m), 121.28~121.05 (m), 120.39 (d, $J=20.5$ Hz), 119.53~119.30 (m), 92.27~90.46 (m), 40.45, 23.63, 23.13, 18.49, 17.80, 10.67; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.17 (d, $J=7.5$ Hz, 6F), -181.39 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{23}\text{H}_{22}\text{F}_7\text{N}_2\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 491.1570, found 491.1571.

*cis-N*¹-(4-Chlorobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-8**): White solid, yield 73%. m.p. 124.8~126.1 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.55 (s, 1H), 8.61 (t, $J=5.9$ Hz, 1H), 7.76 (d, $J=8.5$ Hz, 1H), 7.49~7.39 (m, 2H), 7.28~7.18 (m, 4H), 4.36~4.14 (m, 2H), 2.31~2.18 (m, 4H), 2.14~2.04 (m, 1H), 1.60~1.50 (m, 1H), 1.23~1.13 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 168.94, 168.20, 139.73, 138.51, 131.52, 131.11, 129.01, 127.92, 127.06~126.75 (m), 124.41~124.30 (m), 123.17~122.87 (m), 121.91~121.56 (m), 120.30 (d, $J=20.4$ Hz), 119.08~118.73 (m), 92.10~90.09 (m), 41.58, 23.45, 23.29, 17.86, 10.58; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.15 (d, $J=$

7.5 Hz, 6F), -181.41 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}^{35}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 533.0843, found 533.0842; HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}^{37}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 535.0814, found 535.0814.

*cis-N*¹-(3-Chlorobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-9**): White solid, yield 85%. m.p. 90.7~91.9 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.55 (s, 1H), 8.63 (t, $J=5.9$ Hz, 1H), 7.78 (d, $J=8.6$ Hz, 1H), 7.49~7.36 (m, 3H), 7.33 (s, 1H), 7.28~7.15 (m, 3H), 4.37~4.18 (m, 2H), 2.34~2.18 (m, 4H), 2.15~2.04 (m, 1H), 1.57~1.49 (m, 1H), 1.24~1.14 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 169.04, 168.23, 142.08, 139.74, 132.90, 129.84, 127.22, 126.97, 126.94, 126.93~126.73 (m), 126.49, 125.78, 124.43~126.38 (m), 123.05~122.89 (m), 121.90~121.51 (m), 120.26 (d, $J=20.2$ Hz), 119.05~118.65 (m), 92.87~89.79 (m), 41.73, 23.37, 23.30, 17.88, 10.69; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.16 (d, $J=7.5$ Hz, 6F), -181.39 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}^{35}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 533.0843, found 533.0842; HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}^{37}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 535.0814, found 535.0814.

*cis-N*¹-(2-Chlorobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-10**): White solid, yield 79%. m.p. 132.2~133.4 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.56 (s, 1H), 8.62 (t, $J=5.8$ Hz, 1H), 7.78 (d, $J=8.5$ Hz, 1H), 7.48~7.36 (m, 3H), 7.35~7.29 (m, 1H), 7.24~7.16 (m, 1H), 7.12~7.03 (m, 1H), 4.44~4.19 (m, 2H), 2.32~2.19 (m, 4H), 2.19~2.10 (m, 1H), 1.62~1.52 (m, 1H), 1.28~1.15 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 169.17, 168.22, 141.06, 139.74, 136.18, 131.78, 131.56, 128.88, 128.76, 128.32, 127.04~126.83 (m), 126.82, 124.45~124.34 (m), 123.12~122.95 (m), 121.30~120.94 (m), 120.29 (d, $J=19.6$ Hz), 119.70~119.30 (m), 92.10~89.83 (m), 39.97, 23.51, 23.24, 17.88, 10.67; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.15 (d, $J=7.5$ Hz, 6F), -181.39 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}^{35}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 533.0843, found 533.0842; HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}^{37}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 535.0814, found 535.0814.

*cis-N*¹-(4-Fluorobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-11**): White solid, yield 68%. m.p. 110.3~110.5 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.54 (s, 1H), 8.59 (t, $J=5.9$ Hz, 1H), 7.77 (d, $J=8.5$ Hz, 1H), 7.49~7.38 (m, 2H), 7.29~7.20 (m, 2H), 7.02~6.92 (m, 2H), 4.39~4.08 (m, 2H), 2.28 (s, 3H), 2.26~2.16 (m, 1H), 2.14~2.03 (m, 1H), 1.59~1.50 (m, 1H), 1.23~1.12 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) δ : 168.89, 168.21, 139.75, 135.57 (d, $J=3.0$ Hz), 131.51, 129.07 (d, $J=8.0$ Hz), 127.04~126.78 (m), 124.39~124.28 (m), 123.18~122.91 (m), 121.88~121.51 (m), 120.26 (d, $J=20.5$ Hz), 119.04~118.71 (m), 114.66 (d, $J=21.2$ Hz), 92.13~90.08 (m), 41.52, 23.52, 23.26, 17.85, 10.60; ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.17 (d, $J=7.5$ Hz, 6F), -116.55~-116.69 (m, 1F),

−181.42 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $C_{22}H_{19}F_8N_2O_2$ ($M+H$)⁺ 495.1319, found 495.1320.

*cis-N*¹-(4-Bromobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-12**): White solid, yield 74%. m.p. 144.5~146.2 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.55 (s, 1H), 8.61 (t, $J=5.9$ Hz, 1H), 7.76 (d, $J=8.5$ Hz, 1H), 7.49~7.40 (m, 2H), 7.36 (d, $J=8.4$ Hz, 2H), 7.19 (d, $J=8.4$ Hz, 2H), 4.35~4.12 (m, 2H), 2.32~2.18 (m, 4H), 2.13~2.04 (m, 1H), 1.60~1.50 (m, 1H), 1.24~1.14 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 168.96, 168.21, 139.71, 138.93, 131.55, 131.85, 129.41, 127.05~126.75 (m), 124.46~124.37 (m), 123.16~122.87 (m), 121.93~121.55 (m), 120.34 (d, $J=20.4$ Hz), 119.04~118.69 (m), 92.35~89.76 (m), 41.65, 23.44, 23.28, 17.86, 10.58; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ : −75.14 (d, $J=7.5$ Hz, 6F), −181.40 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $[C_{22}H_{18}^{79}BrF_7N_2O_2]$ ($M+H$)⁺ 555.0518, found 555.0518; HRMS (ESI-TOF) calcd for $C_{22}H_{18}^{81}BrF_7N_2O_2$ ($M+H$)⁺ 557.0504, found 557.0504.

*cis-N*¹-(4-Iodobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-13**): White solid, yield 69%. m.p. 154.4~155.3 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.55 (s, 1H), 8.60 (t, $J=5.9$ Hz, 1H), 7.75 (d, $J=8.5$ Hz, 1H), 7.54 (d, $J=8.3$ Hz, 2H), 7.49~7.38 (m, 2H), 7.04 (d, $J=8.3$ Hz, 2H), 4.32~4.10 (m, 2H), 2.32~2.16 (m, 4H), 2.13~2.03 (m, 1H), 1.60~1.49 (m, 1H), 1.23~1.13 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 168.95, 168.21, 139.73, 139.34, 136.76, 131.58, 129.60, 127.06~126.75 (m), 124.48~124.39 (m), 123.18~122.93 (m), 121.70~121.43 (m), 120.32 (d, $J=20.6$ Hz), 119.16~118.89 (m), 92.49~91.00 (m), 41.77, 23.46, 23.26, 17.87, 10.63; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ : −75.10 (d, $J=7.3$ Hz, 6F), −181.36 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $C_{22}H_{18}F_7N_2O_2I$ ($M+H$)⁺ 603.0379, found 603.0378.

*cis-N*¹-(4-Methoxybenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-14**): White solid, yield 86%. m.p. 81.5~82.6 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.57 (s, 1H), 8.55 (t, $J=5.7$ Hz, 1H), 7.77 (d, $J=8.6$ Hz, 1H), 7.46 (s, 1H), 7.42 (d, $J=8.8$ Hz, 1H), 7.16 (d, $J=8.5$ Hz, 2H), 6.78 (d, $J=8.6$ Hz, 2H), 4.28~4.12 (m, $J=14.8, 5.9$ Hz 2H), 3.69 (s, 3H), 2.29 (s, 3H), 2.23~2.15 (m, 1H), 2.12~2.02 (m, 1H), 1.59~1.49 (m, 1H), 1.22~1.15 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 168.91, 168.32, 158.11, 139.77, 131.55, 131.28, 128.54, 127.05~126.76 (m), 124.41~124.34 (m), 123.15~122.95 (m), 121.92~121.54 (m), 120.26 (d, $J=20.2$ Hz), 119.06~118.69 (m), 92.46~90.12 (m), 54.92, 41.75, 38.20, 23.72, 23.20, 17.87, 10.82; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ : −75.11~−75.23 (m, 6F), −183.30~−183.45 (m, 1F); HRMS (ESI-TOF) calcd for $C_{23}H_{22}F_7N_2O_3$ ($M+H$)⁺ 507.1520, found 507.1520.

*cis-N*¹-(4-(Difluoromethoxy)benzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-15**): White solid, yield 65%. m.p. 89.0~89.7 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.55 (s, 1H),

8.61 (t, $J=5.9$ Hz, 1H), 7.76 (d, $J=8.5$ Hz, 1H), 7.49~7.39 (m, 2H), 7.27 (d, $J=8.6$ Hz, 2H), 7.15 (t, $J=74.2$ Hz, 1H), 6.99 (d, $J=8.6$ Hz, 2H), 4.35~4.12 (m, 2H), 2.28 (s, 3H), 2.26~2.17 (m, 1H), 2.13~2.03 (m, 1H), 1.39~1.51 (m, 1H), 1.23~1.16 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 168.92, 168.24, 149.58, 139.73, 136.51, 128.76, 127.06~126.79 (m), 124.43~124.31 (m), 123.16~122.89 (m), 122.10~120.92 (m), 120.31 (d, $J=21.2$ Hz), 119.61~118.39 (m), 118.56, 116.29, 92.55~90.61 (m), 41.57, 23.49, 23.27, 17.86, 10.65; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ : −75.16 (d, $J=7.5$ Hz, 6F), −81.94 (d, $J=74.2$ Hz, 2F), −181.32 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $C_{23}H_{19}F_9N_2NaO_3$ ($M+Na$)⁺ 565.1151, found 565.1151.

*cis-N*¹-(4-(*tert*-Butyl)benzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-16**): White solid, yield 82%. m.p. 175.9~177.1 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.56 (s, 1H), 8.57 (t, $J=5.7$ Hz, 1H), 7.83 (t, $J=8.6$ Hz, 1H), 7.49~7.39 (m, 2H), 7.23 (d, $J=8.4$ Hz, 2H), 7.15 (d, $J=8.4$ Hz, 2H), 4.35~4.12 (m, 2H), 2.32~2.27 (m, 3H), 2.25~2.15 (m, 1H), 2.13~2.04 (m, 1H), 2.61~2.51 (m, 1H), 1.24~1.16 (m, 10H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 168.93, 168.34, 149.02, 139.77, 136.27, 131.47, 127.00, 126.99~126.73 (m), 124.77, 124.34~124.22 (m), 123.14~122.89 (m), 121.92~121.54 (m), 120.26 (d, $J=20.1$ Hz), 119.05~118.69 (m), 92.61~90.07 (m), 42.01, 33.99, 31.01, 23.72, 23.25, 17.87, 10.74; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ : −75.17 (d, $J=7.5$ Hz, 6F), −181.38 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $C_{26}H_{27}F_7N_2NaO_2$ ($M+Na$)⁺ 555.1859, found 555.1859.

*cis-N*¹-(4-Cyanobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-17**): White solid, yield 62%. m.p. 71.5~72.8 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.59 (s, 1H), 8.71 (d, $J=6.0$ Hz, 1H), 7.77 (d, $J=8.6$ Hz, 1H), 7.61 (d, $J=8.2$ Hz, 2H), 7.50~7.38 (m, 4H), 4.47~4.27 (m, 2H), 2.32~2.23 (m, 4H), 2.17~2.08 (m, 1H), 1.60~1.53 (m, 1H), 1.22~1.16 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 169.06, 168.19, 145.44, 139.69, 131.88, 131.57, 127.90, 127.04~126.74 (m), 124.49~124.38 (m), 123.16~122.90 (m), 121.92~121.54 (m), 120.36 (d, $J=20.6$ Hz), 119.04~118.74 (m), 118.72, 109.31, 92.46~90.15 (m), 41.97, 23.32, 23.27, 17.86, 10.49; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ : −75.22 (d, $J=7.5$ Hz, 6F), −181.55 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $C_{23}H_{18}F_7N_3NaO_2$ ($M+Na$)⁺ 524.1186, found 524.1186.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(4-(trifluoromethyl)benzyl)cyclopropane-1,2-dicarboxamide (**1-18**): White solid, yield 64%. m.p. 107.3~108.6 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.57 (s, 1H), 8.69 (m, 1H), 7.77 (d, $J=8.6$ Hz, 1H), 7.57~7.33 (m, 6H), 4.50~4.23 (m, 2H), 2.35~2.21 (m, 4H), 2.20~2.08 (m, 1H), 1.65~1.54 (m, 1H), 1.27~1.16 (m, 1H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 169.01, 168.21, 144.37, 139.69, 131.44, 127.74, 127.35 (q, $J=31.7$ Hz), 127.02~126.75 (m), 124.71 (q, $J=3.6$ Hz), 124.36~124.28 (m),

123.05~122.81 (m), 121.90~121.52 (m), 120.41 (d, $J=20.4$ Hz), 119.04~118.65 (m), 92.43~89.39 (m), 41.91, 38.12, 23.38, 17.81, 10.43; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -61.26 (m, 3F), -75.41 (d, $J=7.5$ Hz, 6F), -181.58 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{23}\text{H}_{18}\text{F}_{10}\text{N}_2\text{NaO}_2$ ($\text{M} + \text{Na}$) $^+$ 567.1105, found 567.1105.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(4-phenoxybenzyl)cyclopropane-1,2-dicarboxamide (**1-19**): White solid, yield 83%. m.p. 131.7~132.5 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.56 (s, 1H), 8.60 (t, $J=5.9$ Hz, 1H), 7.77 (d, $J=8.5$ Hz, 1H), 7.48~7.32 (m, 4H), 7.25 (d, $J=8.6$ Hz, 2H), 7.15~7.06 (m, 1H), 6.98~6.90 (m, 2H), 6.86 (d, $J=8.6$ Hz, 2H), 4.40~4.13 (m, 2H), 2.28 (s, 3H), 2.25~2.16 (m, 1H), 2.14~2.03 (m, 1H), 1.59~1.49 (m, 1H), 1.24~1.16 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 168.95, 168.27, 156.95, 155.18, 139.75, 134.64, 131.58, 129.90, 128.22, 127.05~126.76 (m), 124.43~124.32 (m), 123.13, 123.12~122.91 (m), 121.72~121.50 (m), 120.28 (d, $J=20.5$ Hz), 119.49~119.29 (m), 118.57, 118.12, 92.18~90.06 (m), 41.71, 23.60, 23.23, 17.89, 10.75; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.16 (d, $J=7.5$ Hz, 6F), -181.41 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{28}\text{H}_{23}\text{F}_7\text{N}_2\text{NaO}_3$ ($\text{M} + \text{Na}$) $^+$ 591.1496, found 591.1496.

*cis-N*¹-([1,1'-biphenyl]-4-ylmethyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-20**): White solid, yield 81%. m.p. 96.2~96.9 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.58 (s, 1H), 8.65 (t, $J=5.9$ Hz, 1H), 7.80 (d, $J=8.6$ Hz, 1H), 7.63~7.57 (m, 2H), 7.55~7.49 (m, 2H), 7.48~7.39 (m, 4H), 7.38~7.30 (m, 3H), 4.44~4.20 (m, 2H), 2.30 (s, 3H), 2.26~2.18 (m, 1H), 2.18~2.06 (m, 1H), 1.62~1.52 (m, 1H), 1.25~1.16 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.02, 168.28, 139.92, 139.77, 138.65, 138.61, 131.59, 128.81, 127.84, 127.25, 127.06~126.75 (m), 126.45, 126.39, 124.46~124.37 (m), 123.17~122.92 (m), 121.91~121.52 (m), 120.29 (d, $J=20.6$ Hz), 119.07~118.65 (m), 92.68~90.07 (m), 42.01, 23.65, 23.26, 17.89, 10.78; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.11~ -75.21 (m, 6F), -181.28~ -181.44 (m, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{28}\text{H}_{24}\text{F}_7\text{N}_2\text{NaO}_2$ ($\text{M} + \text{Na}$) $^+$ 553.1727, found 553.1727.

*cis-N*¹-(2,4-Difluorobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-21**): White solid, yield 71%. m.p. 131.8~133.1 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.55 (s, 1H), 8.60 (t, $J=5.7$ Hz, 1H), 7.76 (d, $J=8.5$ Hz, 1H), 7.49~7.39 (m, 2H), 7.38~7.28 (m, 1H), 7.19~7.08 (m, 1H), 6.82~6.72 (m, 1H), 4.26 (qd, $J=15.4$, 5.8 Hz, 2H), 2.28 (s, 3H), 2.26~2.19 (m, 1H), 2.15~2.06 (m, 1H), 1.60~1.51 (m, 1H), 1.23~1.15 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.08, 168.17, 160.55 (dd, $J=246.0$, 141.7 Hz), 160.43 (dd, $J=246.1$, 142.1 Hz), 139.71, 130.69 (dd, $J=9.7$, 6.0 Hz), 127.03~126.75 (m), 124.39~124.28 (m), 123.13~122.91 (m), 122.30 (dd, $J=15.0$, 3.6 Hz), 121.94~121.52 (m), 120.33 (d, $J=20.4$ Hz), 119.07~118.65 (m), 110.82

(dd, $J=20.9$, 3.5 Hz), 103.32 (t, $J=25.9$ Hz), 92.43~89.81 (m), 35.60 (d, $J=4.1$ Hz), 23.48, 23.19, 17.83, 10.58; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.16~ -75.30 (m, 6F), -112.58~ -112.73 (m, 1F), -114.79~ -114.91 (m, 1F), -181.35~181.54 (m, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{18}\text{F}_9\text{N}_2\text{O}_2$ ($\text{M} + \text{H}$) $^+$ 513.1225, found 513.1226.

*cis-N*¹-(2,6-Dichlorobenzyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-22**): White solid, yield 72%. m.p. 149.6~151.0 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.51 (s, 1H), 8.39 (t, $J=4.8$ Hz, 1H), 7.80 (d, $J=8.5$ Hz, 1H), 7.50~7.40 (m, 4H), 7.37~7.30 (m, 1H), 4.56~4.40 (m, 2H), 2.30 (s, 3H), 2.17~2.09 (m, 1H), 2.08~1.98 (m, 1H), 1.56~1.48 (m, 1H), 1.23~1.16 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.12, 168.19, 139.79, 135.54, 133.20, 130.17, 128.48, 127.02~126.71 (m), 124.38~124.25 (m), 123.16~122.91 (m), 121.92~121.61 (m), 120.32~120.20 (m), 119.16~118.95 (m), 92.39~90.04 (m), 39.99, 24.28, 22.53, 20.89, 17.87, 11.09; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.15 (d, $J=7.5$ Hz, 6F), -181.39 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{17}^{35}\text{Cl}_2\text{F}_7\text{N}_2\text{NaO}_2$ ($\text{M} + \text{Na}$) $^+$ 567.0453, found 567.0452; HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{17}^{35}\text{Cl}^{37}\text{ClF}_7\text{N}_2\text{NaO}_2$ ($\text{M} + \text{Na}$) $^+$ 569.0424, found 569.0424; HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{17}^{37}\text{Cl}_2\text{F}_7\text{N}_2\text{NaO}_2$ ($\text{M} + \text{Na}$) $^+$ 571.0394, found 571.0394.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(3,4,5-trimethoxybenzyl)cyclopropane-1,2-dicarboxamide (**1-23**): White solid, yield 86%. m.p. 137.4~138.1 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.62 (s, 1H), 8.55 (t, $J=5.7$ Hz, 1H), 7.73 (d, $J=8.5$ Hz, 1H), 7.49~7.38 (m, 2H), 6.58 (s, 2H), 4.30~4.14 (m, 2H), 3.64 (s, 6H), 3.59 (s, 3H), 2.30 (s, 3H), 2.25~2.17 (m, 1H), 2.14~2.06 (m, 1H), 1.61~1.53 (m, 1H), 1.24~1.16 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 168.97, 168.58, 152.66, 139.64, 136.16, 135.08, 131.92, 127.02~126.77 (m), 124.78~124.65 (m), 123.05~122.84 (m), 121.92~121.80 (m), 120.58 (d, $J=20.4$ Hz), 119.04~118.63 (m), 104.51, 92.59~89.80 (m), 59.80, 55.53, 42.53, 23.45, 23.13, 17.76, 10.80; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.30 (d, $J=7.5$ Hz, 6F), -181.52 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{25}\text{H}_{26}\text{F}_7\text{N}_2\text{NaO}_5$ ($\text{M} + \text{Na}$) $^+$ 567.1730, found 567.1729.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-((tetrahydrofuran-3-yl)methyl)cyclopropane-1,2-dicarboxamide (**1-24**): White solid, yield 73%. m.p. 139.6~141.2 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.56 (s, 0.5H, *rotamer* 1), 9.54 (s, 0.5H, *rotamer* 2), 8.20 (t, $J=5.6$ Hz, 1H), 7.82 (t, $J=8.4$ Hz, 0.5H, *rotamer* 1), 7.80 (t, $J=8.4$ Hz, 0.5H, *rotamer* 2), 7.50~7.39 (m, 2H), 3.82~3.66 (m, 2H), 3.60~3.50 (m, 1H), 3.20~3.02 (m, 2H), 2.31 (s, 3H), 2.21~2.00 (m, 2H), 1.87~1.62 (m, 3H), 1.54~1.39 (m, 2H), 1.24~1.14 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.60 (*rotamer* 1), 169.58 (*rotamer* 2), 168.59, 139.34, 132.06~131.76 (m), 127.07~126.82 (m), 124.80~124.51 (m), 123.15~122.92 (m), 121.97~

121.45 (m), 120.83 (d, $J=20.5$ Hz), 120.76 (d, $J=20.4$ Hz), 119.04~118.63 (m), 92.95~89.62 (m), 77.00 (rotamer 1), 76.88 (rotamer 2), 67.09, 42.83 (rotamer 1), 42.74 (rotamer 2), 28.27 (rotamer 1), 28.16 (rotamer 2), 24.88, 23.52~23.39 (m), 23.02, 17.66, 10.53~10.45 (m); ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.12~-75.26 (m, 6F), -181.40~-181.53 (m, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{20}\text{H}_{21}\text{F}_7\text{N}_2\text{NaO}_3$ ($\text{M}+\text{Na}$) $^+$ 493.1339, found 493.1339.

*cis-N*¹-((6-Chloropyridin-3-yl)methyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-25**): White solid, yield 66%. m.p. 145.1~146.0 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.56 (s, 1H), 8.65 (t, $J=5.8$ Hz, 1H), 8.31 (d, $J=1.8$ Hz, 1H), 7.72~7.65 (m, 2H), 7.49~7.39 (m, 2H), 7.30~7.24 (m, 1H), 4.35~4.19 (m, 2H), 2.31~2.16 (m, 4H), 2.12~2.03 (m, 1H), 1.57~1.48 (m, 1H), 1.31~1.13 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.01, 168.19, 148.85, 148.48, 139.67, 138.83, 134.67, 131.69, 127.07~126.82 (m), 124.59~124.50 (m), 123.60, 123.12~122.89 (m), 121.92~121.68 (m), 123.37 (d, $J=20.0$ Hz), 119.20~118.98 (m), 92.61~90.04 (m), 40.01, 23.31, 23.06, 17.86, 10.50; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.14 (d, $J=7.5$ Hz, 6F), -181.45 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{21}\text{H}_{17}^{35}\text{ClF}_7\text{N}_3\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 534.0795, found 534.0794; HRMS (ESI-TOF) calcd for $\text{C}_{21}\text{H}_{17}^{37}\text{ClF}_7\text{N}_3\text{NaO}_2$ ($\text{M}+\text{Na}$) $^+$ 536.0766, found 536.0766.

*cis-N*¹-((2-Chlorothiazol-5-yl)methyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-26**): White solid, yield 70%. m.p. 160.3~161.4 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.53 (s, 1H), 8.80 (t, $J=5.8$ Hz, 1H), 7.72 (d, $J=8.6$ Hz, 1H), 7.52 (s, 1H), 7.49~7.38 (m, 2H), 4.43~4.28 (m, 2H), 2.28 (s, 3H), 2.27~2.17 (m, 1H), 2.09~1.95 (m, 1H), 1.55~1.48 (m, 1H), 1.20~1.12 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.32, 167.87, 149.63, 139.87, 139.69, 139.20, 131.53, 127.09~126.85 (m), 124.43~124.36 (m), 123.10~122.84 (m), 121.93~121.71 (m), 120.29 (d, $J=20.5$ Hz), 119.06~118.66 (m), 92.28~89.94 (m), 35.35, 23.32, 22.94, 17.86, 10.43; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.14 (d, $J=7.5$ Hz, 6F), -181.47 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{19}\text{H}_{16}^{35}\text{ClF}_7\text{N}_3\text{O}_2\text{S}$ ($\text{M}+\text{H}$) $^+$ 518.0540, found 518.0540; HRMS (ESI-TOF) calcd for $\text{C}_{19}\text{H}_{16}^{37}\text{ClF}_7\text{N}_3\text{O}_2\text{S}$ ($\text{M}+\text{H}$) $^+$ 520.0510, found 520.0510.

*cis-N*¹-(Furan-2-ylmethyl)-*N*²-(2-methyl-4-(perfluoropropan-2-yl)phenyl)cyclopropane-1,2-dicarboxamide (**1-27**): White solid, yield 65%. m.p. 150.2~151.7 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.55 (s, 1H), 8.60 (t, $J=5.4$ Hz, 1H), 7.76 (d, $J=8.5$ Hz, 1H), 7.53 (s, 1H), 7.50~7.40 (m, 2H), 6.37~6.29 (m, 1H), 6.25~6.19 (m, 1H), 4.30~4.20 (m, 2H), 2.29 (s, 3H), 2.22~2.13 (m, 1H), 2.10~2.01 (m, 1H), 1.55~1.48 (m, 1H), 1.22~1.12 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 168.99, 168.16, 152.15, 141.93, 139.74, 131.78, 127.00~126.78 (m), 124.64~124.52 (m), 123.19~122.95 (m), 122.02~

121.63 (m), 120.31 (d, $J=20.5$ Hz), 119.16~118.79 (m), 110.32, 106.75, 92.60~90.44 (m), 40.94, 35.70, 22.86, 17.86, 10.86; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.15 (d, $J=7.5$ Hz, 6F), -181.46 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{20}\text{H}_{17}\text{F}_7\text{N}_2\text{NaO}_3$ ($\text{M}+\text{Na}$) $^+$ 489.1025, found 489.1024.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(thiophen-3-ylmethyl)cyclopropane-1,2-dicarboxamide (**1-28**): White solid, yield 79%. m.p. 143.5~144.2 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.58 (s, 1H), 8.57 (t, $J=5.6$ Hz, 1H), 7.79 (d, $J=8.5$ Hz, 1H), 7.50~7.35 (m, 3H), 7.29~7.21 (m, 1H), 7.04~6.95 (m, 1H), 4.34~4.19 (m, 2H), 2.30 (s, 3H), 2.25~2.15 (m, 1H), 2.13~2.03 (m, 1H), 1.60~1.50 (m, 1H), 1.24~1.15 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.02, 168.47, 140.05, 139.47, 131.89, 127.33, 127.11~126.83 (m), 125.92, 124.71~124.60 (m), 123.16~122.91 (m), 121.96~121.63 (m), 121.28, 120.66 (d, $J=20.5$ Hz), 119.88~119.52 (m), 92.76~90.00 (m), 37.95, 23.32, 23.19, 17.75, 10.47; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.14 (d, $J=7.5$ Hz, 6F), -181.43 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{20}\text{H}_{17}\text{F}_7\text{N}_2\text{NaO}_2\text{S}$ ($\text{M}+\text{Na}$) $^+$ 505.0797, found 505.0796.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-(naphthalen-1-ylmethyl)cyclopropane-1,2-dicarboxamide (**1-29**): White solid, yield 75%. m.p. 148.1~148.8 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.57 (s, 1H), 8.67 (t, $J=5.6$ Hz, 1H), 8.03 (d, $J=8.4$ Hz, 1H), 7.92 (d, $J=8.0$ Hz, 1H), 7.85~7.76 (m, 2H), 7.52~7.29 (m, 6H), 4.83~4.62 (m, 2H), 2.30 (s, 3H), 2.25~2.17 (m, 1H), 2.16~2.08 (m, 1H), 1.64~1.55 (m, 1H), 1.24~1.18 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 168.95, 168.30, 139.81, 134.36, 133.19, 131.48, 130.82, 128.38, 127.41, 127.05~126.79 (m), 126.04, 125.62, 125.53, 125.23, 124.32~124.21 (m), 123.47, 123.19~122.94 (m), 121.95~121.61 (m), 120.23 (d, $J=20.2$ Hz), 119.30~118.72 (m), 92.81~90.16 (m), 40.48, 23.81, 23.16, 17.88, 10.85; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.16 (d, $J=7.5$ Hz, 6F), -181.39 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{26}\text{H}_{22}\text{F}_7\text{N}_2\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 527.1570, found 527.1570.

*cis-N*¹-(2-Methyl-4-(perfluoropropan-2-yl)phenyl)-*N*²-phenethylcyclopropane-1,2-dicarboxamide (**1-30**): White solid, yield 81%. m.p. 119.4~120.6 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.59 (s, 1H), 8.26 (t, $J=5.4$ Hz, 1H), 7.83 (d, $J=8.5$ Hz, 1H), 7.50~7.39 (m, 2H), 7.28~7.11 (m, 5H), 3.33~3.20 (m, 2H), 2.75~2.63 (m, 2H), 2.31 (s, 3H), 2.20~2.10 (m, 1H), 2.06~1.96 (m, 1H), 1.54~1.46 (m, 1H), 1.24~1.15 (m, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ : 169.13, 168.51, 139.77, 139.39, 131.57, 128.51, 128.21, 127.05~126.79 (m), 125.96, 124.35~124.28 (m), 123.21~122.96 (m), 121.93~121.50 (m), 120.30 (d, $J=20.5$ Hz), 119.04~118.66 (m), 92.17~89.72 (m), 40.47, 38.18, 35.09, 23.19, 17.86, 10.99; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ : -75.19 (d, $J=7.5$ Hz, 6F), -181.41 (hept, $J=7.5$ Hz, 1F); HRMS (ESI-TOF) calcd for $\text{C}_{23}\text{H}_{22}\text{F}_7\text{N}_2\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 491.1570, found 491.1570.

trans-N¹-Ethyl-N²-(2-methyl-4-(perfluoropropan-2-yl)-phenyl)cyclopropane-1,2-dicarboxamide (**1-31**): White solid, yield 73.5%. m.p. 195.4~196.2 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.81 (s, 1H), 8.27 (t, *J*=5.4 Hz, 1H), 7.83 (d, *J*=8.6 Hz, 1H), 7.52~7.40 (m, 2H), 3.15~3.06 (m, 2H), 2.44~2.35 (m, 1H), 2.33 (s, 3H), 2.09~2.01 (m, 1H), 1.21~1.13 (m, 2H), 1.03 (t, *J*=7.2 Hz, 3H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ: 170.06, 169.65, 139.48, 131.47, 127.02~126.83 (m), 124.39~124.21 (m), 123.29~123.20 (m), 121.91~121.69 (m), 120.66~120.59 (m), 119.12~118.96 (m), 95.18~93.53 (m), 33.61, 22.78, 22.02, 17.96, 14.66, 13.12; ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: -75.13 (d, *J*=7.5 Hz, 6F), -181.36 (hept, *J*=7.5 Hz, 1F); HRMS (ESI-TOF) calcd for C₁₇H₁₈F₇N₂O₂ (M+H)⁺ 415.1256, found 415.1255.

4.3 General methods for anti-cancer activity evaluation

4.3.1 MTT assay

Cells in the logarithmic growth phase were seeded in 96-well plates at 10000 cells per milliliter. The cells were treated with different compounds (different gradients for different compounds: 0.781, 1.563, 3.125, 6.250 and 12.500 μg•mL⁻¹; 6.25, 12.50, 25.00, 50.00 and 100 μg•mL⁻¹; 10, 20, 40, 80, 160 μg•mL⁻¹) and cultured for a period of 48 h. 20 μL of MTT (Dojindo) was added to each well at a concentration of 5 mg•mL⁻¹. After incubated at 37 °C for 4 h, the medium was aspirated. 100 μL of DMSO was then added to each well to dissolve the formazan, and the plate was shaken on a shaker for 10 min. The optical density (OD) was measured by an enzyme-linked immunosorbent assay reader (Thermo Fisher Scientific, USA) at wavelengths of both 492 and 630 nm. Cell viability was calculated according to the following formula:

$$\text{Cell viability} = [\text{OD}_{\text{value of treatment group}} - \text{OD}_{\text{value of background}}] / [\text{OD}_{\text{value of control group}} - \text{OD}_{\text{value of background}}] \times 100\%$$

Results were Mean ± SD of three independent experiments.

4.4 General procedure of target prediction and molecule docking

4.4.1 Target prediction

The compositions in SDF format were submitted to PharmMapper (<http://59.78.96.61/pharmmapper/>; access date: March 30, 2021). The databases containing only human targets (the total number is 2241) were selected. The weight of hydrophobic, hydrogen bond receptor (HBA), hydrogen bond donor (HBD), positive, negative and aromatic scores were all set to 1. And the number of reserved matched targets remained at the default of 300. For all the other parameters not mentioned we used the default values. After completion, 300 drug targets results presented by PharmMapper according to the normalized fit scores. Through the investigation, targets unrelated to cancer were screened out manually. After filtering by normalized fit score > 0.9, the top targets and their relationships with our diamides were imported to Cytoscape for visual analysis.

4.4.2 Molecular docking

Molecular modeling was preformed according to the reported methods.^[48-49]

The racemates **1-19** exhibited best anti-cancer activity was selected for further molecular docking studies, getting insights into the putative binding mode with the screening target. The crystal structure of human STS (PDB code: 1P49) was acquired from the Protein Data Bank (<https://www.rcsb.org>) and was treated using Maestro 10.2 to remove the waters from the crystal structure. The 3D structure of composition **1-19** was firstly built in ChemBioDraw Ultra 14.0 and optimized by the Ligprep Module in Maestro 10.2. Both compounds of (1*S*,2*R*) and (1*R*,2*S*) conformation configurations were docked into the STS catalytic pocket simultaneously with Glide program with standard precision (SP) mode in default settings. 100 conformations were output in each ligand docking run. The superior poses were selected by Glide-Score for the further binding mode analysis.

Supporting Information (1) Supplementary anti-cancer activities data; (2) ¹H NMR, ¹³C NMR, ¹⁹F NMR and HRMS spectra; (3) target prediction of **1-12**, **1-14**, **1-19** and **1-23**; (4) 2D binding mode of **1-19**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Torre, L. A.; Bray, F.; Siegel, R. L.; Ferlay, J.; Lortet-Tieulent, J.; Jemal, A. *CA-Cancer J. Clin.* **2015**, *65*, 87.
- [2] Sung, H.; Ferlay, J.; Siegel, R. L.; Laversanne, M.; Soerjomataram, I.; Jemal, A.; Bray, F. *CA-Cancer J. Clin.* **2021**, *71*, 209.
- [3] Dickens, E.; Ahmed, S. *Surgery (Oxford)* **2018**, *36*, 134.
- [4] Jagoe, K. N. R. T.; Abalo, R. *Front. Pharmacol.* **2018**, *9*, 245.
- [5] Janganani, V.; Ponder, J.; Thakkar, S.; Thakkar, S.; Jordan, C. T.; Crooks, P. A. *Bioorg. Med. Chem.* **2017**, *25*, 3694.
- [6] Pattabiraman, V. R.; Bode, J. W. *Nature* **2011**, *480*, 471.
- [7] Kumari, S.; Carmona, A. V.; Tiwari, A. K.; Trippier, P. C. *J. Med. Chem.* **2020**, *63*, 12290.
- [8] Vlaar, C. P.; Castillo-Pichardo, L.; Medina, J. I.; Marrero-Serra, C. M.; Vélez, E.; Ramos, Z.; Hernández, E. *Bioorg. Med. Chem.* **2018**, *26*, 884.
- [9] Rai, U. S.; Isloor, A. M.; Shetty, P.; Pai, K. S. R.; Fun, H. K. *Ara-bian J. Chem.* **2015**, *8*, 317.
- [10] Shao, P. P.; Ok, D.; Fisher, M. H.; Garcia, M. L.; Kaczorowski, G. J.; Li, C.; Lyons, K. A.; Martin, W. J.; Meinke, P. T.; Priest, B. T.; Smith, M. M.; Wyvratt, M. J.; Ye, F.; Parsons, W. H. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1901.
- [11] Tarzia, G.; Shiatti, P.; Selva, D.; Favara, D.; Ceriani, S. *Eur. J. Med. Chem.* **1976**, *11*, 263.
- [12] Hickey, S. M.; Ashton, T. D.; Khosa, S. K.; Robson, R. N.; White, J. M.; Li, J.; Nation, R. L.; Yu, H. Y.; Elliott, A. G.; Butler, M. S.; Huang, J. X.; Cooper, M. A.; Pfeiffer, F. M. *Org. Biomol. Chem.* **2015**, *22*, 6255.
- [13] Saudi, M.; Zmurko, J.; Kaptein, S.; Rpzinski, J.; Gadakh, B.; Chaltin, P.; Marchand, A.; Neyts, J.; Aerschot, A. V. *Eur. J. Med. Chem.* **2016**, *12*, 158.
- [14] Tohnishi, M.; Nakao, H.; Furuya, T.; Seo, A.; Kodama, H.; Tsubata, K.; Fujioka, S.; Kodama, H.; Hirooka, T.; Nishimatsu, T. *J. Pestic. Sci.* **2005**, *30*, 354.
- [15] Galano, A.; Alvarez-Idaboy, J. R.; Vivier-Bunge, A. *Theor. Chem. Acc.* **2007**, *118*, 597.

- [16] Talele, T. T. *J. Med. Chem.* **2016**, *59*, 8712.
- [17] Julius, D. A. *Am. J. Psychiatry* **1979**, *136*, 782.
- [18] Yakes, F. M.; Chen, J.; Tan, J.; Yamaguchi, K.; Shi, Y.; Yu, P.; Qian, F.; Chu, F.; Bentzien, F.; Cancilla, B.; Orf, J.; You, A.; Laird, A. D.; Engst, S.; Lee, L.; Lesch, J.; Chou, Y.-C.; Joly, A. H. *J. Clin. Oncol.* **2017**, *35*, 591.
- [19] Chopra, N.; Nathan, P. D. *Expert Rev. Anticancer Ther.* **2015**, *15*, 749.
- [20] Scheciiter, M. S.; Sullivan, W. N.; Schoof, H. F.; Maddock, D. R.; Amyx, C. M.; Porter, J. E. *J. Med. Entomol.* **1974**, *11*, 231.
- [21] Matsui, K.; Kido, Y.; Watari, R.; Kashima, Y.; Yoshida, Y.; Shuto, S. *Chem. Eur. J.* **2017**, *23*, 3034.
- [22] Lavedan, C.; Forsberg, M.; Gentile, A. J. *Neuropharmacology* **2015**, *91*, 142.
- [23] Yamaguchi, K.; Kazuta, Y.; Hirano, K.; Yamada S.; Matsuda, A.; Shuto, S. *Bioorg. Med. Chem.* **2008**, *16*, 8875.
- [24] Martin, S. F.; Dorsey, G. O.; Gane, T.; Hillier, M. C.; Kessler, H.; Baur, M.; Mathä, B.; Erickson, J. W.; Bhat, T. N.; Munshi, S.; Gulnik, S. V.; Topol, I. A. *J. Med. Chem.* **1998**, *41*, 1581.
- [25] Chen, Y.; Le, V.; Xu, X.; Shao, X.; Liu, J.; Li, Z. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 3948.
- [26] Ozoe, Y.; Kita, T.; Ozoe, F.; Nakao, T.; Sato, K.; Hirase, K. *Pestic. Biochem. Physiol.* **2013**, *107*, 285.
- [27] Shang, J.; Liu, Q.; Wang, B.; Li, Z. *Chin. J. Org. Chem.* **2019**, *39*, 1489 (in Chinese).
(尚俊峰, 刘巧霞, 王宝雷, 李正名, 有机化学, **2019**, *39*, 1489.)
- [28] Chen, K.; Liu, Q.; Ni, J.; Zhu, H.; Li, Y.; Wang, Q. *Pest Manage. Sci.* **2014**, *71*, 1503.
- [29] Duan, J. J.-W.; Lu, Z.; Jiang, B.; Stachura, S.; Weigelt, C. A.; Sack, J. S.; Khan, J.; Ruzanov, M.; Galella, M. A.; Wu, D.-R.; Yarde, M.; Shen, D.-R.; Shuster, D. J.; Borowski, V.; Xie, J. H.; Zhang, L.; Vanteru, S.; Gupta, A. K.; Mathur, A.; Zhao, Q.; Foster, W.; Salter-Cid, L. M.; Carter, P. H.; Dhar, T. G. M. *ACS Med. Chem. Lett.* **2019**, *10*, 367.
- [30] Li, Z.; Xu, X.; Chen, Y.; Liu, J.; Li, W. *CN 103435562*, **2013** (in Chinese).
- [31] Li, Z.; Chen, Y.; Liu, A.; Li, Y.; Wang, B.; Pan, L.; Wan, Y.; Liu, J.; Chen, W. *CN 104402785*, **2015** (in Chinese).
- [32] Lepri, S.; Goracci, L.; Valeri, A.; Cruciani, G. *Eur. J. Med. Chem.* **2016**, *121*, 658.
- [33] Simpkins, L. M.; Bolton, S.; Pi, Z.; Sutton, J. C.; Kwon, C.; Zhao, G.; Magnin, D. R.; Augeri, D. J.; Gungor, T.; Rotella, D. P.; Sun, Z.; Liu, Y.; Slusarchyk, W. S.; Marcinkeviciene, J.; Robertson, J. G.; Wang, A.; Robl, J. A.; Atwal, K. S.; Zahler, R. L.; Parker, R. A.; Kirby, M. S.; Hamann, L. G. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 6476.
- [34] Wang, F.; Xu, X.; Wang, F.; Peng, L.; Zhang, Y.; Wang, L.; Wang, L. *Lett. Org. Chem.* **2015**, *12*, 741.
- [35] Milewska, M. J.; Gdaniec, M.; Poloński, T. *Tetrahedron: Asymmetry* **1996**, *7*, 3169.
- [36] Beutner, G. L.; Young, I. S.; Davies, M. L.; Hickey, M. R.; Park, H.; Stevens, J. M.; Ye, Q. *Org. Lett.* **2018**, *20*, 4218.
- [37] Liu, X.; Ouyang, S.; Yu, B.; Liu, Y.; Huang, K.; Gong, J.; Zheng, S.; Li, Z.; Li, H.; Jiang, H. *Nucleic Acids Res.* **2010**, *38*, 609.
- [38] Wang, X.; Pan, C.; Gong, J.; Liu, X.; Li, H. *J. Chem. Inf. Model.* **2016**, *56*, 1175.
- [39] Wang, X.; Shen, Y.; Wang, S.; Li, S.; Zhang, W.; Liu, X.; Lai, L.; Pei, J.; Li, H. *Nucleic Acids Res.* **2017**, *45*, 356.
- [40] Ferrante, P.; Messali, S.; Meroni, G.; Ballabio, A. *Eur. J. Hum. Genet.* **2002**, *10*, 813.
- [41] Hernandez-Guzman, F. G.; Higashiyama, T.; Pangborn, W.; Osawa, Y.; Ghosh, D. *J. Biol. Chem.* **2003**, *278*, 22989.
- [42] Geisler, J.; Sasano, H.; Chen, S.; Purohit, A. *J. Steroid Biochem. Mol. Biol.* **2011**, *125*, 39.
- [43] Purohit, A.; Foster, P. A. *J. Endocrinol.* **2012**, *212*, 99.
- [44] Leese, M. P.; Hejaz, H. A. M.; Mahon, M. F.; Newman, S. P.; Purohit, A.; Reed, M. J.; Potter, B. V. L. *J. Med. Chem.* **2005**, *48*, 5243.
- [45] A. Purohit, Woo, L. W. L.; Potter, B. V. L. *Cancer Res.* **2000**, *60*, 3394.
- [46] Kertesz, N.; Krasnoperov, V.; Reddy, R.; Leshanski, L.; Kumar, S. R.; Zozulya, S.; Gill, P. S. *Blood* **2006**, *107*, 2330.
- [47] Pasquale, E. B. *Nat. Rev. Cancer* **2010**, *10*, 165.
- [48] Dohle, W.; Jourdan, F. L.; Menchon, G.; Protta, A. E.; Foster, P. A.; Mannion, P.; Hamel, E.; Thomas, M. P.; Kasprzyk, P. G.; Ferrandis, E.; Steinmetz, M. O.; Leese, M. P.; Potter, B. V. L. *J. Med. Chem.* **2018**, *61*, 1031.
- [49] Maltais, R.; Djimeny, A. N.; Roy, J.; Barbeau, X.; Lambert, J.-P.; Poirier, D. *Bioorg. Med. Chem.* **2020**, *28*, 115368.

(Cheng, F.)