溶剂介导吡咯并[1,2-*a*]喹喔啉选择性 C—H 键碘代

刘亚丽 杨 振 李 阳 刘 岩* 刘 平*

(石河子大学化学化工学院 化工绿色过程兵团重点实验室 新疆石河子 832003)

摘要 开发了一种溶剂介导吡咯并[1,2-*a*]喹喔啉和 *N*-碘代丁二酰亚胺(NIS)的区域选择性 C—H 碘化, 分别以 CHCl_3 和 *N,N*-二甲基甲酰胺(DMF)为溶剂, 选择性地生成了 1-碘吡咯并[1,2-*a*]喹喔啉和 3-碘吡咯并[1,2-*a*]喹喔啉. 此外, 吡咯并[1,2-*a*]喹喔啉与 *N*-溴代丁二酰亚胺(NBS)的溴代反应得到主要产物为 1,3-二溴吡咯并喹喔啉. 该方法具有反应条件简单温和、区域选择性好、底物范围广、克级合成等特点. 此外, 还通过钯和碘催化的 C—X (X=C, S)键形成反应研究了卤化吡咯并[1,2-*a*]喹喔啉产物的进一步转化.

关键词 溶剂介导; 区域选择性; 吡咯并[1,2-*a*]喹喔啉; C—H 键碘代

Solvent Mediated Selective C—H Bond Iodination of Pyrrolo[1,2-*a*]quinoxaline

Liu, Yali Yang, Zhen Li, Yang Liu, Yan* Liu, Ping*

(Key Laboratory for Green Processing of Chemical Engineering of Xinjiang Bingtuan, School of Chemistry and Chemical Engineering, Shihezi University, Shihezi, Xinjiang 832003)

Abstract A solvent mediated regioselective C—H iodination of pyrrolo[1,2-*a*]quinoxaline with *N*-iodo-succinimide (NIS) has been developed. 1-Iodopyrrolo[1,2-*a*]quinoxalines and 3-iodopyrrolo[1,2-*a*]quinoxalines could be selectively synthesized by using CHCl_3 and *N,N*-dimethylformamide (DMF) as solvents, respectively. Furthermore, 1,3-dibromopyrroloquinoxalines were obtained as the main products by the bromination of pyrrolo[1,2-*a*]quinoxaline with *N*-bromo-succinimide (NBS). The method features simple and mild reaction conditions, good regioselectivity, broad substrate scope, and gram-scale synthesis. In addition, the further transformation of halogenated pyrrolo[1,2-*a*]quinoxaline products has also been investigated through palladium and iodine catalyzed C—X (X=C, S) bond formation reactions.

Keywords solvent-mediated; regioselectivity; pyrrolo[1,2-*a*]quinoxaline; C—H bond iodination

1 Introduction

Pyrrolo[1,2-*a*]quinoxalines function as antiparasitic agents,^[1] antifungal activity,^[2] 5-HT₃ receptors,^[3] inhibitors of human protein kinase Casein Kinase (2CK2) and protein kinase B (AKT),^[4] and so on^[5] in a wide range of biological processes.^[6] Although many approaches have been developed to synthesize pyrrolo[1,2-*a*]quinoxalines, usually 4-substituted pyrroloquinoxaline skeletons are constructed.^[7] This greatly limits the further development and application of pyrroloquinoxaline as a biologically active molecule in the field of medicine. In the past decade, the C—H bond functionalization as green and sustainable methodology has been achieved notable progress due to construct diverse organic molecules starting from simple

compounds.^[8] As we known, pyrrolo[1,2-*a*]quinoxaline has multiple reactive sites.^[9-12] Therefore, the direct C—H bond functionalization of pyrrolo[1,2-*a*]quinoxalines as the core to construct diverse pyrroloquinoxaline derivatives is an alternative protocol. In 2020, Nguyen *et al.* reported a C1—H chlorination of 4-aryl pyrrolo[1,2-*a*]quinoxalines with *N*-chloro-succinimide (NCS) in the presence of dimethyl sulfoxide (DMSO) as catalyst and CHCl_3 as solvent. Also, the bromination of pyrrolo[1,2-*a*]quinoxalines was achieved by using CuBr_2 as brominated reagent and $\text{K}_2\text{S}_2\text{O}_8$ as oxidant (Scheme 1a).^[11] However, the catalytic amount of DMSO and the stoichiometric amount of CuBr in the above reaction system are indispensable. Recently, a selective C3—H iodination of pyrrolo[1,2-*a*]quinoxalines with tetrabutylammonium iodide (TBAI) or I_2 was deve-

* Corresponding authors. E-mail: liuyan@shzu.edu.cn; liuping@shzu.edu.cn

Received July 15, 2021; revised August 20, 2021; published online September 2, 2021.

Project supported by the National Natural Science Foundation of China (Nos. 21563025, 22162022).

国家自然科学基金(Nos. 21563025, 22162022)资助项目.

Scheme 1 Selective halogenations of pyrrolo[1,2-*a*]quinoxalines

loped, and a series of novel 3-iodo-pyrrolo[1,2-*a*]-quinoxalines were obtained with excellent regioselectivity and broad substrate scope (Scheme 1b).^[12] On the basis of this work, herein we reported a solvent mediated regioselective C—H iodination of pyrrolo[1,2-*a*]quinoxaline and *N*-iodosuccinimide (NIS), and 1-iodopyrrolo[1,2-*a*]quinoxalines could be selectively synthesized by using CHCl₃ and *N,N*-dimethylformamide (DMF) as solvents, respectively (Scheme 1c). Compared with the previous work, the characteristics of this reaction are simple operation, mild reaction conditions and metal-free participation.

2 Results and discussion

To investigate the selective iodination conditions of pyrrolo[1,2-*a*]quinoxaline (Table 1), we first did the iodination of pyrrolo[1,2-*a*]quinoxaline and NIS by similar the brominated reaction conditions reported by Nguyen, unfortunately, only 17% yield of C1-iodinated product **2a** was obtained in the presence of 20 mol% of DMSO as catalyst and CHCl₃ as solvent at room temperature for 24 h (Entry 1). To our delights, the reaction could be achieved in the absence of DMSO, affording the desired product **2a** in 81% yield (Entry 2). Shortening the reaction time to 12 h, the reaction yield of **2a** was reduced to 76% (Entry 3). Switching NIS to I₂ or TBAI was detrimental to the reaction (Entries 4 and 5). Next, the solvent was screened. Interestingly, when the polar solvents such as DMF, DMSO, MeCN, EtOH, and MeOH were used as the reaction medium, selective C3—H iodinated product **3a** was obtained with 39%~72% yields of (Entries 6~10). Among them, DMF as the optimal solvent provided the product **3a** in 72% yield (Entry 6). Dichloromethane (DCM) and toluene were ineffective in this reaction (Entries 11 and 12). The above experimental results showed that the solvent played a vital role in the regioselectivity of the iodination reaction. Subsequently, I₂ as iodine source was tested again in DMF, but a lower yield was obtained (Entries 13). When 2 equiv. of

NIS was used, the yield was reduced to 40% (Entry 14). Prolonging the reaction time to 10 h resulted in no significant increase in the yield of **3a** (Entry 15). On the contrary, shortening the time to 3 h, the yield can be increased to 76% (Entry 16).

Table 1 Optimization of the reaction conditions

Entry	[I] (equiv.)	Solvent	<i>t</i> /h	Yield ^b /%	
				2a	3a
1 ^c	NIS (1.2)	CHCl ₃	24	17	—
2	NIS (1.2)	CHCl ₃	24	81	—
3	NIS (1.2)	CHCl ₃	12	76	—
4	I ₂ (1.2)	CHCl ₃	12	nr	—
5	TBAI (1.2)	CHCl ₃	12	nr	—
6	NIS (1)	DMF	6	—	72
7	NIS (1)	DMSO	6	—	56
8	NIS (1)	MeCN	6	—	52
9	NIS (1)	EtOH	6	—	51
10	NIS (1)	MeOH	6	—	39
11	NIS (1)	DCM	6	—	trace
12	NIS (1)	Toluene	6	—	nr
13	I ₂ (1)	DMF	6	—	21
14	NIS (2)	DMF	6	—	40
15	NIS (1)	DMF	10	—	73
16	NIS (1)	DMF	3	—	76

^a Reaction conditions: **1a** (0.2 mmol), solvent (3 mL), room temperature, in air.

^b Isolated yield. ^c 20 mol% of DMSO as oxidant was used under air.

With optimized conditions in hand, the scope of selective C—H iodination of pyrrolo[1,2-*a*]quinoxalines was evaluated (Table 2). Under the optimized reaction condition 1 (Table 1, Entry 2), the substituted pyrrolo[1,2-*a*]quinox-

line bearing 8-Me, and 8-F groups at the benzene ring position afforded the C1-iodinated products **2b** and **2c** in 63% and 67% yields, respectively. The iodination of 7-methylpyrrolo[1,2-*a*]quinoxaline could proceed smoothly to obtain the desired product **2d** in 67% yield. Subsequently, the iodinations of 4-aryl substituted pyrrolo[1,2-*a*]quinoxalines were investigated. The substrates with various substituents on *para*-positions of the benzene ring were well tolerated, providing the desired products **2e**~**2k** in 63%~89% yields with good regioselectivity. The results showed that the electronic effect of the substituents has no obvious influence on the reactivity. Under the optimized reaction condition 2 (Table 1, Entry 16), the regioselective C3-iodinated reactions of the 7-Me, 8-Me, and 8-F substituted pyrrolo[1,2-*a*]quinoxalines could proceed well, providing the corresponding products **3b**~**3d** in 52%~72% yields. However, 4-arylpyrrolo[1,2-*a*]quinoxalines were used as reactants under the same conditions to afford the C1-iodinated products **2e**, **2f**, **2i**, and **2k** in 45%~79% yields, and the C3-iodinated products **3e**, **3f**, **3i**, and **3k** were obtained in lower yields. This may be attributed to the electronic effect of the 4-aryl substituent on the reaction. Further studies showed that increasing the temperature was beneficial to the selectivity of C3-H iodination.

Next, the brominations of pyrrolo[1,2-*a*]quinoxaline derivatives were explored (Table 3). The bromination of pyrrolo[1,2-*a*]quinoxaline **1a** with NBS in the presence of DMF could proceed smoothly, providing the product 1,3-dibromopyrrolo[1,2-*a*]quinoxaline (**4a**) in 77% yield. The structure of **4a** was determined by single crystal X-ray diffraction (Figure 1). In addition, the reactions of 4-

arylpyrrolo[1,2-*a*]quinoxalines with NBS gave 1,3-dibrominated products **4b** and **4c** in 53% and 49% yields, respectively, and 1,2,3-tribrominated products **4b'** and **4c'** were obtained in 19% and 26% yields, respectively.

Figure 1 Crystal structure of **4a** (CCDC: 2073986)

Scheme 2 TBAI-promoted the sulfonylation of **4a**

Interestingly, when 1,3-dibromopyrrolo[1,2-*a*]quinoxaline was utilized as a substrate to react with TsNHNH₂ in the

Table 2 Scope of selective C—H iodination of pyrrolo[1,2-*a*]quinoxalines with NIS^a

^a Conditions 1: **1a** (0.2 mmol), NIS (1.2equiv.), CHCl₃ (3 mL), 24 h, r.t., in air, isolated yield; Condition 2: **1a** (0.2 mmol), NIS (1.0 equiv.), DMF (3 mL), 3 h, r.t., in air, isolated yield. ^b 100 °C, 6 h.

Table 3 Bromination of pyrrolo[1,2-*a*]quinoxaline with NBS^a

^a Reaction conditions: **1a** (0.2 mmol), DMF (3 mL), 3 h, r.t., under Ar atmosphere. ^b Isolated yield.

presence of TBAI, the sulfenylated product 1-bromo-3-tosylpyrrolo[1,2-*a*]quinoxaline (**5a**) was also obtained in 84% yield. The structure of **5a** was established by X-ray crystallographic analysis (Figure 2).

Figure 2 Crystal structure of **5a** (CCDC: 2062352)

The scale-up reaction was performed by using 6 mmol of pyrrolo[1,2-*a*]quinoxaline **1a** and NIS in the presence of 20 mL of CHCl₃, affording 1.063 g of **2a** in 60% yield (Scheme 3).

Scheme 3 Gram-scale synthesis of **2a**

To highlight the synthetic utility of 1-iodopyrrolo[1,2-*a*]quinoxaline, the further transformations of C1—I bond were investigated (Table 5). First, the Suzuki-Miyaura reactions of **2a** with arylboronic acids in the presence of 5 mol% Pd(PPh₃)₄ and Na₂CO₃ as base at 80 °C under Ar atmosphere were explored, and the desired products **6a**~**6f** were obtained in 49%~99% yields with good functional group tolerance. Next, the PdCl₂(PPh₃)₂/CuI-catalyzed Sonogashira reactions of **2a** with various terminal alkynes were performed by using Et₃N as base at 65 °C in the air (Table 5). The expected products **7a**~**7e** were obtained in 69%~79% yields. Also, 2-methylbut-3-yn-2-ol was suitable

substrate for this reaction, affording the product **7f** in 72% yield.

Table 4 Pd-catalyzed Suzuki-Miyaura reactions of **2a**^a

^a Reaction conditions: **2a** (0.2 mmol), arylboronic acid (2 equiv.), 1,4-dioxane/H₂O (2.5 mL, V:V = 4:1), 80 °C, 5 h, under Ar atmosphere. ^b Isolated yield; ^c 7 h.

Table 5 Pd-catalyzed Sonogashira reactions of **2a**^a

^a Reaction conditions: **2a** (0.2 mmol), terminal alkyne (1.5 equiv.), Et₃N (2 mL), 65 °C, 12 h, under air. ^b Isolated yield.

3 Conclusions

In summary, a regioselective C—H iodination of pyrrolo[1,2-*a*]quinoxaline and NIS has been developed. CHCl₃ as solvent resulted in C1—H iodination, so that 1-iodopyrrolo[1,2-*a*]quinoxalines could be formed. In the DMF solvent case, the C3—H iodination readily occurred to give 3-iodopyrrolo[1,2-*a*]quinoxalines. The present work represent novel strategies for selective C—H iodination of pyrrolo[1,2-*a*]quinoxaline. Furthermore, the bromination of pyrrolo[1,2-*a*]quinoxaline with NBS has also been investigated, and 1,3-dibromopyrroloquinoxalines were obtained as the main products. The successful constructions of C—C and C—S bonds by further derivatization of these products provided the possibility for the application of pyrroloquinoxaline derivatives in the fields of medicine and synthesis. The method features simple and mild reaction conditions, good regioselectivity, broad substrate scope, gram-

scale synthesis, and diverse transformations of the products. Further studies to elucidate the action of solvents on the regioselectivity of iodination reactions are ongoing.

4 Experimental section

4.1 General methods

Unless otherwise noted, all reactions were performed in Schlenk tubes. Melting points were determined with a fu-siometer and are not corrected. The materials obtained from commercial suppliers were used without further purification. For chromatography, Qingdao Ocean Chemical 200-300 mesh silica gel was employed. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker Avance III HD 400 MHz spectrometer in CDCl_3 solution and the chemical shifts were reported in δ relative to the internal standard tetramethylsilane (TMS, δ 0). High-resolution mass spectra (HRMS) were acquired in electrospray ionization (APCI) mode using a TOF mass analyzer.

4.2 General procedure for pyrrolo[1,2-*a*]quinoxalines with NIS in CHCl_3 (Conditions 1)

A mixture of pyrrolo[1,2-*a*]quinoxaline (0.2 mmol), NIS (0.24 mmol), and CHCl_3 (3 mL) was stirred at room temperature for 24 h. Upon completion, the solvent was removed under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the final products **2a**~**2k**.

4.3 General procedure for pyrrolo[1,2-*a*]quinoxalines with NIS in DMF (Conditions 2)

A mixture of pyrrolo[1,2-*a*]quinoxaline (0.2 mmol), NIS (0.2 mmol), and DMF (3 mL) was stirred at room temperature for 3 h. Upon completion, the reaction was quenched with the saturated solution of sodium thiosulfate (2 mL) and extracted with EtOAc (10 mL $\times$ 3). The combined organic layer was dried over anhydrous Na_2SO_4 , and the solvent was removed under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the final products **3a**~**3k**.

4.4 General procedure for bromination of pyrrolo[1,2-*a*]quinoxaline with NBS

A mixture of pyrrolo[1,2-*a*]quinoxaline (0.2 mmol), NBS (0.6 mmol), and DMF (3 mL) was stirred at room temperature under Ar atmosphere for 3 h. Upon completion, the solution was extracted with EtOAc (10 mL $\times$ 3). The combined organic layer was dried over anhydrous Na_2SO_4 , the solvent was removed under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the corresponding products **4a**~**4c**, **4b'**, and **4c'**.

4.5 General procedure for TBAI-promoted the sulfenylation of **4a**

A mixture of 1,3-dibromopyrrolo[1,2-*a*]quinoxaline (0.3 mmol), TsNHNH_2 (0.9 mmol), TBAI (0.6 mmol), TBHP (0.5 mL), and 1,4-dioxane (0.5 mL) was stirred at 80 $^\circ\text{C}$ for 12 h. Upon completion, the reaction was quenched the sat-

urated solution of sodium thiosulfate (2 mL) and extracted with dichloromethane (10 mL $\times$ 3). The combined organic layer was dried over anhydrous Na_2SO_4 , and the solvent was removed under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the corresponding products **5a** (84% yield).

4.6 General procedure for the gram-scale synthesis of **2a**

A mixture of **1a** (6 mmol), NIS (7.2 mmol), and CHCl_3 (20 mL) was stirred at room temperature for 36 h. Upon completion, the reaction was quenched the saturated solution of sodium thiosulfate (10 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layer was dried over anhydrous Na_2SO_4 , and the solvent was removed under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the final product **2a** (1.063 g, 60% yield).

4.7 Typical procedure for the Pd-catalyzed Suzuki-Miyaura reactions of **2a** with arylboronic acids

A mixture of **2a** (0.2 mmol), arylboronic acid (0.4 mmol), $\text{Pd}(\text{PPh}_3)_4$ (5 mol%), Na_2CO_3 (2 equiv.), and 1,4-dioxane/ H_2O ($V:V=4:1$, 2.5 mL) was stirred at 80 $^\circ\text{C}$ under Ar atmosphere for 5 h. Upon completion, the solution was washed with the saturated NaCl aqueous solution (10 mL) and extracted with dichloromethane (10 mL $\times$ 3). The organic phase was dried over Na_2SO_4 and concentrated under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the final product **6a**~**6f**.

4.8 Typical procedure for the Pd-catalyzed Sonogashira reactions of **2a** with terminal alkynes

A mixture of **2a** (0.2 mmol), terminal alkyne (0.3 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (5 mol%), CuI (20 mol%), and Et_3N (2.0 mL) was stirred at 65 $^\circ\text{C}$ for 12 h under air atmosphere. Upon completion, an appropriate amount of silica gel was added to the solution. The mixture was concentrated under reduced pressure. The crude was purified by flash chromatography on silica gel (petroleum ether/EtOAc) to give the final product **7a**~**7f**.

4.9 Characterization data of the products

1-Iodopyrrolo[1,2-*a*]quinoxaline (**2a**):^[13] 47.6 mg, 81% yield. Brown solid, m.p. 164~165 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 9.47 (d, $J=8.3$ Hz, 1H), 8.67 (s, 1H), 8.01~7.91 (m, 1H), 7.61~7.40 (m, 2H), 7.05 (d, 1H), 6.86 (d, $J=4.1$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.0, 137.4, 131.0, 130.7, 130.1, 127.6, 126.8, 126.2, 115.2, 110.2, 61.3; HRMS (APCI) calcd for $\text{C}_{11}\text{H}_7\text{IN}_2$ [$\text{M}+\text{H}$] $^+$ 294.9727, found 294.9723.

1-Iodo-8-methylpyrrolo[1,2-*a*]quinoxaline (**2b**): 38.8 mg, 63% yield. Light yellow solid, m.p. 185~186 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 9.29 (s, 1H), 8.62 (s, 1H), 7.83 (d, $J=8.2$ Hz, 1H), 7.35~7.22 (m, 1H), 7.04 (d, $J=4.1$ Hz, 1H), 6.84 (d, $J=4.1$ Hz, 1H), 2.55 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 143.6, 137.1, 135.0, 130.7, 130.0, 129.4, 127.2, 127.0, 114.8, 109.5, 60.2, 22.1; HRMS (APCI) calcd

for $C_{12}H_9IN_2 [M+H]^+$ 308.9883, found 308.9879.

8-Fluoro-1-iodopyrrolo[1,2-*a*]quinoxaline (**2c**): 41.8 mg, 67% yield. Light yellow solid, m.p. 182~183 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.22 (dd, $J=11.1, 2.6$ Hz, 1H), 8.63 (s, 1H), 7.99~7.86 (m, 1H), 7.23~7.18 (m, 1H), 7.07 (d, $J=4.1$ Hz, 1H), 6.87 (d, $J=4.1$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 143.8, 131.8, 131.8, 130.3, 127.6, 113.6, 113.4, 109.9, 102.0, 101.7, 77.4, 77.0, 76.7, 61.1; ^{19}F NMR (376 MHz, $CDCl_3$) δ : 109.6; HRMS (APCI) calcd for $C_{11}H_6FIN_2 [M+H]^+$ 312.9632, found 312.9627.

1-Iodo-7-methylpyrrolo[1,2-*a*]quinoxaline (**2d**): 41.3 mg, 67% yield. Light yellow solid, m.p. 179~180 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.35 (d, $J=8.7$ Hz, 1H), 8.65 (s, 1H), 7.75 (s, 1H), 7.34 (d, $J=10.5$ Hz, 1H), 7.03 (d, $J=4.1$ Hz, 1H), 6.85 (d, $J=4.1$ Hz, 1H), 2.49 (s, 4H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 144.5, 136.8, 135.7, 130.5, 130.0, 127.54, 127.52, 126.9, 114.5, 109.7, 60.7, 20.9; HRMS (APCI) calcd for $C_{12}H_9IN_2 [M+H]^+$ 308.9883, found 308.9877.

1-Iodo-4-(4-methoxyphenyl)pyrrolo[1,2-*a*]quinoxaline (**2e**): Conditions 1: 66.4 mg, 83% yield; Conditions 2: 36 mg, 45% yield. Light yellow solid, m.p. 146~147 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.49 (d, $J=9.7$ Hz, 1H), 8.03 (d, $J=9.6$ Hz, 1H), 7.88 (d, $J=8.7$ Hz, 2H), 7.56~7.45 (m, 2H), 7.13~6.99 (m, 3H), 6.94 (d, $J=4.2$ Hz, 1H), 3.89 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 161.0, 153.0, 137.4, 130.4, 130.2, 130.1, 129.8, 128.6, 127.4, 125.8, 125.8, 114.9, 114.0, 111.2, 77.4, 77.0, 76.7, 60.9, 55.5; HRMS (APCI) calcd for $C_{18}H_{13}IN_2O [M+H]^+$ 401.0145, found 401.0139.

4-(4-Bromophenyl)-1-iodopyrrolo[1,2-*a*]quinoxaline (**2f**): Conditions 1: 75.4 mg, 84% yield; Conditions 2: 66.5 mg, 74% yield. White solid, m.p. 172~173 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.52 (d, $J=8.7$ Hz, 1H), 8.04 (d, $J=9.1$ Hz, 1H), 7.79 (d, $J=8.4$ Hz, 2H), 7.67 (d, $J=8.4$ Hz, 3H), 7.60~7.47 (m, 2H), 7.10 (d, $J=4.2$ Hz, 1H), 6.89 (d, $J=4.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 152.3, 137.2, 136.7, 131.8, 131.8, 130.4, 130.3, 129.4, 128.7, 127.6, 126.3, 126.0, 124.3, 115.0, 111.0; HRMS (APCI) calcd for $C_{17}H_{10}BrIN_2 [M+H]^+$ 448.9145, found 448.9138.

1-Iodo-4-(4-nitrophenyl)pyrrolo[1,2-*a*]quinoxaline (**2g**): 52.3 mg, 63% yield. Yellow solid, m.p. 234~235 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.60~9.49 (m, 1H), 8.39 (d, $J=8.8$ Hz, 2H), 8.10 (d, $J=8.8$ Hz, 2H), 8.05 (dd, $J=7.8, 1.7$ Hz, 1H), 7.63~7.57 (m, 1H), 7.56~7.51 (m, 1H), 7.14 (d, $J=4.2$ Hz, 1H), 6.89 (d, $J=4.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 150.0, 147.6, 142.8, 136.0, 129.6, 129.4, 128.7, 128.1, 127.8, 126.8, 125.9, 125.2, 123.3, 122.8, 114.0, 109.8, 61.0; HRMS (APCI) calcd for $C_{17}H_{10}IN_3O_2 [M+H]^+$ 415.9890, found 415.9883.

1-Iodo-4-phenylpyrrolo[1,2-*a*]quinoxaline (**2h**): 65.9 mg, 89% yield. Light yellow solid, m.p. 148~149 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.52 (d, $J=9.7$ Hz, 1H), 8.06 (d, $J=7.5$ Hz, 1H), 7.99~7.83 (m, 2H), 7.53 (d, $J=4.2$ Hz, 4H), 7.52~7.47 (m, 1H), 7.09 (d, $J=4.2$ Hz, 1H), 6.93 (d, $J=4.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 153.5,

137.9, 137.3, 130.4, 129.8, 129.8, 129.8, 128.8, 128.7, 128.6, 127.5, 126.1, 125.9, 114.9, 111.3, 60.9; HRMS (APCI) calcd for $C_{17}H_{10}IN_3O_2 [M+H]^+$ 371.0040, found 371.0033.

1-Iodo-4-(*p*-tolyl)pyrrolo[1,2-*a*]quinoxaline (**2i**): Conditions 1: 62.2 mg, 81% yield; Conditions 2: 60.7 mg, 79% yield. Light yellow solid, m.p. 141~142 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.43 (d, $J=9.7$ Hz, 1H), 8.10~7.92 (m, 2H), 7.74 (d, $J=8.1$ Hz, 3H), 7.48~7.39 (m, 3H), 7.27 (d, $J=7.9$ Hz, 4H), 7.00 (d, $J=4.2$ Hz, 2H), 6.87 (d, $J=4.2$ Hz, 2H), 2.39 (s, 6H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 153.3, 139.7, 137.2, 134.8, 130.1, 129.6, 129.1, 128.5, 128.4, 127.2, 125.7, 125.6, 114.6, 111.0, 60.7, 21.3; HRMS (APCI) calcd for $C_{18}H_{13}IN_2 [M+H]^+$ 385.0196, found 385.0194.

4-(4-Chlorophenyl)-1-iodopyrrolo[1,2-*a*]quinoxaline (**2j**): 59.1 mg, 73% yield. White solid, m.p. 172~173 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.52 (d, $J=8.1$ Hz, 1H), 8.03 (d, $J=7.5$ Hz, 1H), 7.86 (d, $J=8.0$ Hz, 2H), 7.52 (t, $J=9.4$ Hz, 5H), 7.10 (d, $J=3.8$ Hz, 1H), 6.89 (d, $J=3.8$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 152.0, 137.0, 136.0, 135.8, 130.2, 129.8, 129.3, 128.7, 128.6, 127.4, 126.1, 125.8, 114.7, 110.9, 61.3; HRMS (APCI) m/z calcd for $C_{17}H_{10}ClIN_2 [M+H]^+$ 404.9650, found 404.9645.

4-(4-Fluorophenyl)-1-iodopyrrolo[1,2-*a*]quinoxaline (**2k**): Conditions 1: 60.6 mg, 78% yield; Conditions 2: 54.3 mg, 70% yield. White solid, m.p. 172~173 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.48 (d, $J=9.6$ Hz, 1H), 8.02 (dd, $J=7.6, 1.9$ Hz, 2H), 7.90 (dd, $J=8.6, 5.5$ Hz, 3H), 7.50 (p, $J=7.3$ Hz, 3H), 7.22 (t, $J=8.6$ Hz, 3H), 7.06 (d, $J=4.2$ Hz, 1H), 6.87 (d, $J=2.6$ Hz, 0H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 164.8, 162.3, 152.1, 137.0, 133.7, 130.4, 130.1, 129.3, 128.4, 127.3, 125.9, 125.7, 115.5, 115.3, 114.6, 110.8, 61.1; ^{19}F NMR (376 MHz, $CDCl_3$) δ : 110.8; HRMS (APCI) calcd for $C_{17}H_{10}FIN_2 [M+H]^+$ 388.9945, found 388.9938.

3-Iodopyrrolo[1,2-*a*]quinoxaline (**3a**)^[12]: 44.7 mg, 76% yield. Light yellow solid, m.p. 177~178 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.70 (s, 1H), 7.98 (dd, $J=8.0, 1.5$ Hz, 1H), 7.88 (d, $J=2.8$ Hz, 1H), 7.82 (dd, $J=8.2, 1.5$ Hz, 1H), 7.53 (td, $J=8.2, 7.8, 1.5$ Hz, 1H), 7.46 (td, $J=7.8, 1.4$ Hz, 1H), 6.98 (d, $J=2.8$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.3, 135.7, 130.2, 128.0, 127.2, 126.4, 125.6, 120.9, 115.4, 113.0, 59.9; HRMS (APCI) calcd for $C_{11}H_7IN_2 [M+H]^+$ 294.9727, found 294.9723.

4-Iodo-8-methylpyrrolo[1,2-*a*]quinoxaline (**3b**)^[12]: 44.4 mg, 72% yield. Light yellow solid, m.p. 175~176 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.61 (s, 1H), 7.80 (d, $J=2.7$ Hz, 2H), 7.56 (s, 1H), 7.28~7.21 (m, 1H), 6.94 (d, $J=2.8$ Hz, 1H), 2.52 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 144.5, 138.8, 133.9, 130.0, 127.1, 127.0, 126.7, 121.0, 115.2, 113.2, 59.6, 21.9; HRMS (APCI) calcd for $C_{12}H_9IN_2 [M+H]^+$ 308.9883, found 308.9876.

8-Fluoro-3-iodopyrrolo[1,2-*a*]quinoxaline (**3c**)^[12]: 32.5 mg, 52% yield. Light yellow solid, m.p. 180~181 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.67 (s, 1H), 7.98 (dd, $J=9.0, 5.8$ Hz, 1H), 7.82~7.76 (m, 2H), 7.50 (dd, $J=9.0, 2.7$ Hz, 1H), 7.20 (ddd, $J=8.9, 8.2, 2.7$ Hz, 1H), 7.02 (d, $J=2.9$ Hz,

1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 144.7, 132.3, 132.2, 121.6, 115.7, 113.9, 113.7, 100.3, 100.1, 60.3; ^{19}F NMR (376 MHz, CDCl_3) δ : 109.4; HRMS (APCI) calcd for $\text{C}_{11}\text{H}_6\text{FIN}_2$ $[\text{M} + \text{H}]^+$ 312.9632, found 312.9624.

3-Iodo-7-methylpyrrolo[1,2-*a*]quinoxaline (**3d**)^[12]: 39.4 mg, 64% yield. Light yellow solid, m.p. 79~80 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.64 (s, 1H), 7.80 (d, $J=2.7$ Hz, 1H), 7.74 (s, 1H), 7.65 (d, $J=8.4$ Hz, 1H), 7.30 (dd, $J=8.4$, 1.7 Hz, 1H), 6.93 (d, $J=2.8$ Hz, 1H), 2.48 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.3, 135.8, 135.6, 130.0, 129.3, 126.5, 125.2, 120.8, 115.4, 112.9, 59.7, 21.1; HRMS (APCI) calcd for $\text{C}_{12}\text{H}_9\text{IN}_2$ $[\text{M} + \text{H}]^+$ 308.9883, found 308.9879.

3-Iodo-4-(4-methoxyphenyl)pyrrolo[1,2-*a*]quinoxaline (**3e**)^[12]: 13.6 mg, 17% yield. Light yellow solid, m.p. 176~177 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.99 (dd, $J=7.9$, 1.6 Hz, 1H), 7.93 (d, $J=2.9$ Hz, 1H), 7.80 (dd, $J=8.1$, 1.4 Hz, 1H), 7.58~7.40 (m, 4H), 7.09~6.98 (m, 3H), 3.89 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 160.7, 155.1, 135.6, 131.4, 130.1, 129.7, 127.7, 127.2, 125.7, 124.3, 124.0, 116.3, 113.6, 113.2, 60.6, 55.4; HRMS (APCI) calcd for $\text{C}_{18}\text{H}_{13}\text{IN}_2\text{O}$ $[\text{M} + \text{H}]^+$ 401.0145, found 401.0142.

2-Iodo-4-(*p*-tolyl)pyrrolo[1,2-*a*]quinoxaline (**3i**)^[12]: 4.6 mg, 6% yield. Light yellow solid, m.p. 152~153 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.99 (dd, $J=7.9$, 1.5 Hz, 1H), 7.92 (d, $J=2.9$ Hz, 1H), 7.79 (dd, $J=8.1$, 1.3 Hz, 1H), 7.51~7.40 (m, 4H), 7.32 (d, $J=7.8$ Hz, 2H), 7.01 (d, $J=2.9$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 155.5, 139.3, 135.6, 134.3, 130.2, 129.8, 128.8, 127.7, 127.3, 125.7, 124.2, 124.0, 116.2, 113.2, 60.6, 21.6; HRMS (APCI) calcd for $\text{C}_{18}\text{H}_{13}\text{IN}_2$ $[\text{M} + \text{H}]^+$ 385.0196, found 385.0188.

4-(4-Fluorophenyl)-3-iodopyrrolo[1,2-*a*]quinoxaline (**3k**)^[12]: Conditions 1: 10.9 mg, 14% yield; Conditions 2: 25.6 mg, 33% yield. Light yellow solid, m.p. 140~141 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.98 (dd, $J=7.9$, 1.4 Hz, 1H), 7.94 (d, $J=2.9$ Hz, 1H), 7.80 (dd, $J=8.2$, 1.1 Hz, 1H), 7.60~7.42 (m, 4H), 7.20 (d, $J=8.7$ Hz, 2H), 7.02 (d, $J=2.9$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 154.3, 135.4, 133.2, 133.2, 131.9, 131.8, 130.2, 128.0, 127.3, 125.8, 124.1, 124.0, 116.4, 115.3, 115.1, 113.2, 60.5; ^{19}F NMR (376 MHz, CDCl_3) δ : 112.0; HRMS (APCI) calcd for $\text{C}_{17}\text{H}_{10}\text{FIN}_2$ $[\text{M} + \text{H}]^+$ 388.9945, found 388.9937.

1,3-Dibromopyrrolo[1,2-*a*]quinoxaline (**4a**)^[13]: 50.2 mg, 77% yield. White solid, m.p. 162~163 °C; ^1H NMR (400 MHz, CDCl_3) δ : 9.25 (d, $J=9.8$ Hz, 1H), 8.02~7.95 (m, 2H), 7.54~7.49 (m, 7H), 7.47 (d, $J=8.3$ Hz, 3H), 6.93 (s, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 143.1, 136.9, 130.2, 128.5, 127.0, 126.1, 125.6, 120.4, 115.0, 98.8, 95.6, 77.2, 76.8, 76.5; HRMS (APCI) calcd for $\text{C}_{11}\text{H}_6\text{Br}_2\text{N}_2$ $(\text{M} + \text{H})^+$: 324.8970, found 324.8968.

1,3-Dibromo-4-(4-chlorophenyl)pyrrolo[1,2-*a*]quinoxaline (**4b**): 46.3 mg, 53% yield. Light yellow solid, m.p. 202~203 °C; ^1H NMR (400 MHz, CDCl_3) δ : 9.25 (d, $J=9.8$ Hz, 1H), 8.02~7.95 (m, 2H), 7.54~7.49 (m, 7H), 7.47 (d, $J=8.3$ Hz, 3H), 6.93 (s, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.7, 136.4, 135.5, 135.3, 130.6, 130.1, 128.1,

126.8, 126.2, 123.8, 122.7, 115.0, 99.7, 96.2; HRMS (APCI) calcd for $\text{C}_{17}\text{H}_9\text{Br}_2\text{ClN}_2$ $[\text{M} + \text{H}]^+$ 434.8894, found 434.8888.

1,2,3-Tribromo-4-(4-chlorophenyl)pyrrolo[1,2-*a*]quinoxaline (**4b'**): 19.6 mg, 19% yield. White solid, m.p. 241~242 °C; ^1H NMR (400 MHz, CDCl_3) δ : 9.29 (d, $J=9.5$ Hz, 1H), 8.04~7.97 (m, 1H), 7.55 (t, $J=6.3$ Hz, 2H), 7.53~7.46 (m, 5H); ^{13}C NMR (101 MHz, CDCl_3) δ : 152.1, 136.1, 135.5, 135.3, 130.6, 130.4, 128.2, 127.4, 127.2, 126.6, 123.8, 115.2, 114.0, 101.7, 99.0; HRMS (APCI) calcd for $\text{C}_{17}\text{H}_8\text{Br}_3\text{ClN}_2$ $[\text{M} + \text{H}]^+$ 512.7999, found 512.7990.

1,3-Dibromo-4-(4-fluorophenyl)pyrrolo[1,2-*a*]quinoxaline (**4c**): 41.2 mg, 49% yield. Light yellow solid, m.p. 167~168 °C; ^1H NMR (400 MHz, CDCl_3) δ : 9.30~9.20 (m, 1H), 8.03~7.96 (m, 1H), 7.55 (dd, $J=8.7$, 5.4 Hz, 3H), 7.53~7.47 (m, 3H), 7.18 (t, $J=8.7$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 164.9, 162.4, 153.2, 136.6, 133.3 (d, $J=3.3$ Hz), 131.4, 131.3, 130.3, 128.3, 127.0, 126.4, 124.2, 123.0, 115.2 (d, $J=4.5$ Hz), 115.0, 99.9, 96.5; ^{19}F NMR (376 MHz, CDCl_3) δ : 111.8; HRMS (APCI) calcd for $\text{C}_{17}\text{H}_9\text{Br}_2\text{FN}_2$ $[\text{M} + \text{H}]^+$ 418.9189, found 418.9181.

1,2,3-Tribromo-4-(4-fluorophenyl)pyrrolo[1,2-*a*]quinoxaline (**4c'**): 26.0 mg, 26% yield. White solid, m.p. 233~234 °C; ^1H NMR (400 MHz, CDCl_3) δ : 9.29 (d, $J=9.8$ Hz, 1H), 8.01 (d, $J=9.6$ Hz, 1H), 7.59~7.51 (m, 4H), 7.20 (t, $J=8.7$ Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 164.7, 162.3, 152.2, 136.1, 133.0, 131.2, 131.1, 130.3, 127.4, 127.1, 126.6, 123.9, 115.20, 115.15, 114.98, 101.7, 99.1; ^{19}F NMR (376 MHz, CDCl_3) δ : 111.4; HRMS (APCI) calcd for $\text{C}_{17}\text{H}_8\text{Br}_3\text{FN}_2$ (M): 495.8225, found 495.8222.

1-Bromo-3-tosylpyrrolo[1,2-*a*]quinoxaline (**5a**): 101.1 mg, 84% yield. Light yellow solid, m.p. 187~188 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.33 (dd, $J=8.3$, 1.4 Hz, 1H), 7.26 (s, 1H), 6.36 (dd, $J=7.7$, 1.9 Hz, 1H), 6.20 (d, $J=8.4$ Hz, 2H), 5.92 (qd, $J=7.2$, 1.7 Hz, 3H), 5.69 (d, $J=8.3$ Hz, 2H), 0.77 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.4, 143.8, 137.3, 137.0, 131.4, 130.6, 130.4, 130.2, 128.9, 128.2, 128.0, 127.3, 127.1, 125.2, 118.6, 21.7; HRMS (APCI) calcd for $\text{C}_{18}\text{H}_{13}\text{BrN}_2\text{O}_2\text{S}$ $[\text{M} + \text{H}]^+$ 400.9954, found 400.9947.

1-Phenylpyrrolo[1,2-*a*]quinoxaline (**6a**)^[14]: 34.2 mg, 70% yield. Brown oily liquid; ^1H NMR (400 MHz, CDCl_3) δ : 9.52 (dd, $J=8.2$, 1.5 Hz, 1H), 8.06 (dd, $J=7.7$, 1.9 Hz, 1H), 7.90 (dd, $J=6.7$, 3.0 Hz, 3H), 7.53 (dd, $J=6.3$, 3.6 Hz, 5H), 7.49 (dd, $J=7.2$, 1.5 Hz, 1H), 7.09 (d, $J=4.2$ Hz, 1H), 6.93 (d, $J=4.2$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 153.3, 137.7, 137.1, 130.2, 129.6, 129.6, 129.6, 128.6, 128.5, 128.4, 127.3, 125.9, 125.7, 114.7, 111.1, 60.7; HRMS (APCI) calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2$ $[\text{M} + \text{H}]^+$ 245.1073, found 245.1071.

1-(*p*-Tolyl)pyrrolo[1,2-*a*]quinoxaline (**6b**): 40.8 mg, 79% yield. Yellow solid, m.p. 122~123 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.80 (s, 1H), 7.93 (d, $J=8.0$ Hz, 1H), 7.47 (d, $J=8.5$ Hz, 1H), 7.41 (d, $J=8.0$ Hz, 2H), 7.37~7.29 (m, 3H), 7.14 (t, $J=7.9$ Hz, 1H), 6.95 (d, $J=4.0$ Hz, 1H), 6.74 (d, $J=4.0$ Hz, 1H), 2.48 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 145.9, 138.4, 137.0, 132.4, 130.9, 129.8, 129.4,

129.2, 128.9, 127.4, 126.2, 124.7, 116.5, 116.4, 107.3, 21.2; HRMS (APCI): m/z calcd for $C_{18}H_{14}N_2$ $[M+H]^+$ 259.1230, found 259.1226.

1-(4-Methoxyphenyl)pyrrolo[1,2-*a*]quinoxaline (**6c**): 47.2 mg, 86% yield. Yellow solid, m.p. 142~143 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.63 (s, 1H), 7.77 (d, $J=8.0$ Hz, 1H), 7.29 (dd, $J=11.4, 8.6$ Hz, 3H), 7.18 (t, $J=7.6$ Hz, 1H), 6.98 (t, $J=7.9$ Hz, 1H), 6.87 (d, $J=8.8$ Hz, 2H), 6.78 (d, $J=4.0$ Hz, 1H), 6.57 (d, $J=4.0$ Hz, 1H), 3.75 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 159.7, 145.8, 137.0, 132.1, 130.8, 129.7, 129.0, 127.3, 126.2, 126.1, 124.6, 116.4, 116.2, 113.9, 107.2, 55.2; HRMS (APCI) calcd for $C_{18}H_{14}N_2O$ $[M+H]^+$ 275.1179, found 275.1177.

1-(4-Vinylphenyl)pyrrolo[1,2-*a*]quinoxaline (**6d**): 26.5 mg, 49% yield. Yellow solid, m.p. 113~114 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.81 (s, 1H), 7.94 (d, $J=8.0$ Hz, 1H), 7.56~7.47 (m, 3H), 7.36 (t, $J=7.6$ Hz, 1H), 7.16 (t, $J=7.9$ Hz, 1H), 6.97 (d, $J=4.0$ Hz, 1H), 6.87~6.76 (m, 2H), 5.88 (d, $J=17.6$ Hz, 1H), 5.37 (d, $J=10.9$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.8, 137.6, 137.0, 136.0, 133.1, 132.0, 129.8, 129.6, 128.8, 127.7, 126.4, 126.3, 124.8, 116.8, 116.5, 114.7, 107.5; HRMS (APCI) calcd for $C_{19}H_{14}N_2$ $[M+H]^+$ 271.1230, found 271.1227.

1-(Naphthalen-2-yl)pyrrolo[1,2-*a*]quinoxaline (**6e**): 50.0 mg, 85% yield. Yellow solid, m.p. 122~123 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.85 (s, 1H), 8.05 (s, 1H), 7.97 (d, $J=9.6$ Hz, 3H), 7.93 (s, 1H), 7.90 (d, $J=9.4$ Hz, 1H), 7.59 (d, $J=3.5$ Hz, 2H), 7.57 (d, $J=1.7$ Hz, 1H), 7.48 (d, $J=8.5$ Hz, 1H), 7.35 (t, $J=7.6$ Hz, 1H), 7.10~7.04 (m, 1H), 7.01 (d, $J=4.0$ Hz, 1H), 6.86 (d, $J=4.0$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.8, 137.0, 133.0, 132.8, 132.1, 131.1, 129.7, 128.8, 128.3, 128.2, 128.0, 127.7, 127.6, 127.0, 126.7, 126.5, 126.3, 124.8, 117.0, 116.5; HRMS (APCI) calcd for $C_{21}H_{14}N_2$ $[M+H]^+$ 295.1230, found 295.1229.

1-(Thiophen-3-yl)pyrrolo[1,2-*a*]quinoxaline (**6f**): 49.6 mg, 99% yield. Yellow liquid. 1H NMR (400 MHz, $CDCl_3$) δ : 8.80 (s, 1H), 7.94 (d, $J=7.9$ Hz, 1H), 7.54 (d, $J=6.0$ Hz, 1H), 7.50~7.33 (m, 3H), 7.22 (s, 1H), 7.20 (s, 1H), 6.90 (dd, $J=22.6, 3.8$ Hz, 2H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.6, 145.5, 136.9, 134.2, 129.9, 129.3, 127.8, 127.3, 126.7, 124.9, 123.5, 118.5, 115.9, 107.1, 106.9; HRMS (APCI) calcd for $C_{15}H_{10}N_2S$ $[M+H]^+$ 251.0637, found 251.0635.

1-(*p*-Tolylethynyl)pyrrolo[1,2-*a*]quinoxaline (**7a**): 40.1 mg, 71% yield. Yellow solid, m.p. 100~101 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.38 (d, $J=7.9$ Hz, 1H), 8.75 (s, 2H), 7.96 (d, $J=7.5$ Hz, 1H), 7.57~7.43 (m, 6H), 7.21 (s, 2H), 7.11 (d, $J=4.2$ Hz, 2H), 6.86 (d, $J=4.2$ Hz, 2H), 2.41 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.0, 138.9, 136.8, 130.8, 129.8, 129.7, 129.1, 127.0, 126.9, 125.4, 121.1, 119.4, 115.4, 112.5, 107.2, 96.8, 82.1, 21.4; HRMS (APCI) calcd for $C_{20}H_{14}N_2$ $(M+H)^+$: 283.1230, found 283.1228.

1-(Phenylethynyl)pyrrolo[1,2-*a*]quinoxaline (**7b**): 39.7 mg, 74% yield. Yellow solid, m.p. 108~109 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.38 (d, $J=7.9$ Hz, 1H), 8.76 (s, 2H), 7.97 (d, $J=7.7$ Hz, 1H), 7.63 (d, $J=7.9$ Hz, 3H), 7.55~

7.47 (m, 3H), 7.47~7.39 (m, 4H), 7.13 (d, $J=4.2$ Hz, 2H), 6.87 (d, $J=4.2$ Hz, 2H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.0, 136.8, 130.9, 129.7, 128.6, 128.4, 127.1, 127.0, 125.4, 122.5, 121.3, 115.4, 112.3, 107.3, 96.6, 82.7; HRMS (APCI) calcd for $C_{19}H_{12}N_2$ $[M+H]^+$ 269.1073, found 269.1068.

1-([1,1'-Biphenyl]-4-ylethynyl)pyrrolo[1,2-*a*]quinoxaline (**7c**): 52.4 mg, 76% yield. Yellow solid, m.p. 152~153 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.42 (d, $J=9.2$ Hz, 1H), 8.79 (s, 1H), 8.00 (d, $J=7.8$ Hz, 1H), 7.73~7.61 (m, 7H), 7.59~7.45 (m, 4H), 7.39 (t, $J=7.3$ Hz, 1H), 7.16 (d, $J=4.2$ Hz, 1H), 6.90 (d, $J=4.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 145.0, 141.4, 140.0, 136.9, 131.4, 129.8, 129.8, 128.8, 127.7, 127.2, 127.1, 126.9, 125.5, 121.5, 121.4, 115.5, 112.4, 107.4, 96.7, 83.4; HRMS (APCI) calcd for $C_{25}H_{16}N_2$ $[M+H]^+$ 345.1386, found 345.1384.

1-((4-Methoxyphenyl)ethynyl)pyrrolo[1,2-*a*]quinoxaline (**7d**): 41.2 mg, 69% yield. Yellow solid, m.p. 111~112 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.38 (d, $J=8.1$ Hz, 1H), 8.75 (s, 1H), 7.96 (d, $J=7.6$ Hz, 1H), 7.59~7.43 (m, 4H), 7.10 (d, $J=3.8$ Hz, 1H), 6.95 (d, $J=8.0$ Hz, 2H), 6.86 (d, $J=4.2$ Hz, 1H), 3.86 (s, 2H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 159.9, 145.0, 136.8, 132.5, 129.8, 129.7, 127.0, 126.9, 125.4, 121.0, 115.4, 114.6, 114.1, 112.7, 107.3, 96.7, 81.4, 55.2. HRMS (APCI) calcd for $C_{20}H_{14}N_2O$ $[M+H]^+$ 299.1179, found 299.1177.

1-((4-Chlorophenyl)ethynyl)pyrrolo[1,2-*a*]quinoxaline (**7e**): 47.8 mg, 79% yield. Yellow solid, m.p. 149~150 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.26 (d, $J=9.7$ Hz, 1H), 8.74 (s, 1H), 7.95 (d, $J=9.6$ Hz, 1H), 7.51 (s, 1H), 7.50~7.45 (m, 3H), 7.38 (s, 2H), 7.09 (d, $J=4.2$ Hz, 1H), 6.84 (d, $J=4.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 144.9, 136.8, 134.6, 131.9, 129.8, 129.6, 128.7, 127.2, 127.0, 125.4, 121.5, 120.9, 115.2, 111.8, 107.3, 95.5, 83.7; HRMS (APCI) calcd for $C_{19}H_{11}ClN_2$ $[M+H]^+$ 303.0684, found 303.0681.

2-Methyl-4-(pyrrolo[1,2-*a*]quinoxalin-1-yl)but-3-yn-2-ol (**7f**): 36.0 mg, 72% yield. Yellow solid, m.p. 154~155 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 9.00 (d, $J=8.2$ Hz, 1H), 8.67 (d, $J=3.6$ Hz, 1H), 7.90 (d, $J=9.1$ Hz, 1H), 7.22 (d, $J=7.6$ Hz, 2H), 6.89 (t, $J=4.8$ Hz, 1H), 6.70~6.64 (m, 1H), 1.75 (s, 7H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 144.7, 136.0, 129.3, 129.2, 126.7, 126.4, 125.1, 121.0, 114.9, 112.0, 107.2, 101.7, 75.5, 65.3, 30.8; HRMS (APCI) calcd for $C_{16}H_{14}N_2O$ $[M+H]^+$ 251.1179, found 251.1177.

Supporting Information Single crystal data of compounds **4a** and **5a**; 1H NMR, ^{13}C NMR and ^{19}F NMR spectra. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Ronga, L.; Del Favero, M.; Cohen, A.; Soum C.; Le Pape P.; Savrimoutou S.; Pinaud, N.; Mullie C.; Daulouede, S.; Vincendeau, P.; Farvacques, N.; Agnamey, P.; Pagniez, F.; Hutter, S.; Azas, N.; Sonnet, P.; Guillon, J. *Eur. J. Med. Chem.* **2014**, *81*, 378.
(b) van Heerden, L.; Cloete, T. T.; Breytenbach, J. W.; de Kock, C.;

- Smith, P.; Breytenbach, J. C.; N'Da; D. D. *Eur. J. Med. Chem.* **2012**, *55*, 335.
- (c) Guillon, J.; Mouray, E.; Moreau, S.; Mullie, C.; Forfar, I.; Desplat, V.; Belisle-Fabre, S.; Pinaud, N.; Ravanella, F.; Le-Naour, A.; Le-Naour, Leger, J.-M.; Gosmann, G.; Jarry, C.; Deleris, G.; Sonnet, P.; Grellier, P. *Eur. J. Med. Chem.* **2011**, *46*, 2310.
- (d) Jonet, A.; Guillon, J.; Mullie, C.; Cohen, A.; Bentzinger, G.; Schneider, J.; Taudon, N.; Hutter, S.; Azas, N.; Moreau, S.; Savrimoutou, S.; Agnamey, P.; Dassonville-Klimpt, A.; Sonnet, P. *Med. Chem.* **2018**, *14*, 293.
- (e) Guillon, J.; Cohen, A.; Gueddouda, N. M.; Das, R. N.; Moreau, S.; Ronga, L.; Savrimoutou, S.; Basmaciyan, L.; Monnier, A.; Monget, M.; Rubio, S.; Garnerin, T.; Azas, N.; Mergny, J.-L.; Mullie, C.; Sonnet, P. *J. Enzyme Inhib. Med. Chem.* **2017**, *32*, 547.
- [2] Xu, H.; Fan, L. *Eur. J. Med. Chem.*, **2011**, *46*, 1919.
- [3] Morelli, E.; Gemma, S.; Budriesi, R.; Campiani, G.; Novellino, E.; Fattorusso, C.; Catalanotti, B.; Coccone, S. S.; Ros, S.; Borrelli, G.; Kumar, V.; Persico, M.; Fiorini, I.; Nacci, V.; Ioan, P.; Chiarini, A.; Hamon, M.; Cagnotto, A.; Mennini, T.; Fracasso, C.; Colovic, M.; Caccia, S.; Butini, S. *J. Med. Chem.* **2009**, *52*, 3548.
- [4] (a) Guillon, J.; Le Borgne, M.; Rimbault, C.; Moreau, S.; Savrimoutou, S.; Pinaud, N.; Baratin, S.; Marchivie, M.; Roche, S.; Bollacke, A.; Pecci, A.; Alvarez, L.; Despla, V.; Jose, J. *Eur. J. Med. Chem.* **2013**, *65*, 205.
- (b) Desplat, V.; Moreau, S.; Gay, A.; Fabre, S. B.; Thiolat, D.; Massip, S.; Macky, G.; Godde, F.; Mossalayi, D.; Jarry, C.; Guillon, J. *J. Enzyme Inhib. Med. Chem.* **2010**, *25*, 204.
- [5] (a) Brindisi, M.; Brogi, S.; Maramai, S.; Grillo, A.; Borrelli, G.; Butini, S.; Novellino, E.; Allara, M.; Ligresti, A.; Campiani, G.; Di Marzo, V.; Gemma, S. *RSC Adv.* **2016**, *6*, 64651.
- (b) Desplat, V.; Vincenzi, M.; Lucas, R.; Moreau, S.; Savrimoutou, S.; Pinaud, N.; Lesbordes, J.; Peyrilles, E.; Marchivie, M.; Routier, S.; Sonnet, P.; Rossi, F.; Ronga, L.; Guillon, J. *Eur. J. Med. Chem.* **2016**, *113*, 214.
- (c) Desplat, V.; Vincenzi, M.; Lucas, R.; Moreau, S.; Savrimoutou, S.; Rubio, S.; Pinaud, N.; Bigat, D.; Enriquez, E.; Marchivie, M.; Routier, S.; Sonnet, P.; Rossi, F.; Ronga, L.; Guillon, J. *Chem-MedChem* **2017**, *12*, 940.
- (d) Wang, T.; Tang, Y.; Yang, Y.; An, Q.; Sang, Z.; Yang, T.; Liu, P.; Zhang, T.; Deng, Y.; Luo, Y.; *Bioorg. Med. Chem. Lett.* **2018**, *28*, 2084.
- (e) Gemma, S.; Colombo, L.; Forloni, G.; Savini, L.; Fracasso, C.; Caccia, S.; Salmona, M.; Brindisi, M.; Joshi, B. P.; Tripaldi, P.; Giorgi, O.; Tagliatela-Scafati, E.; Novellino, I.; Fiorini, G.; Campiani, S.; Butini, S. *Org. Biomol. Chem.* **2011**, *9*, 5137.
- [6] (a) Huang, A.; Ma, C. *Mini-Rev. Med. Chem.* **2013**, *13*, 607.
- (b) Campiani, G.; Aiello, F.; Fabbrini, M.; Morelli, E.; Ramunno, A.; Armoroli, S.; Nacci, V.; Garofalo, A.; Greco, G.; Novellino, E.; Maga, G.; Spadari, S.; Bergamini, A.; Ventura, L.; Bongiovanni, B.; Capozzi, M.; Bolacchi, F.; Marini, S.; Coletta, M.; Guiso, G.; Caccia, S. *J. Med. Chem.* **2001**, *44*, 305.
- (c) Glennon, R. A.; Daoud, M. K.; Dukat, M.; Teitler, M.; Herick-Davis, K.; Purohit, A.; Syed, H. *Bioorg. Med. Chem.* **2003**, *11*, 4449.
- (d) Carta, A.; Loriga, M.; Paglietti, G.; Mattana, A.; Fiori, P. L.; Mollicotti, P.; Sechi, L.; Zanetti, S. *Eur. J. Med. Chem.* **2004**, *39*, 195.
- (e) Guillon, J.; Forfar, I.; Mamani-Matsuda, M.; Desplat, V.; Saliège, M.; Thiolat, D.; Massip, S.; Tabourier, A.; Léger J.-M.; Dufaure, B.; Haumont, G.; Jarry, C.; Mossalayi, D. *Bioorg. Med. Chem.* **2007**, *15*, 194.
- (f) Moarbess, G.; Deleuze-Masquefa, C.; Bonnard, V.; Gayraud-Paniagua, S.; Vidal, J. R.; Bressolle, F.; Pinguet, P.; Bonnet, P. A. *Bioorg. Med. Chem.* **2008**, *16*, 6601.
- [7] (a) Kalinin, A. A.; Islamova, L. N.; Fazleeva, G. M. *Chem. Heterocycl. Compd.* **2019**, *55*, 584.
- (b) Cong, W.; Wang, L.; Yu, F.; Li, J. *Chin. J. Org. Chem.* **2018**, *38*, 2866.
- [8] (a) Qin, Y.; Zhu, L.; Luo, S. *Chem. Rev.* **2017**, *117*, 9433.
- (b) Ping, L.; Chung, D. S.; Bouffard, J.; Lee, S. G. *Chem. Soc. Rev.* **2017**, *46*, 4299.
- (c) Liu, C. X.; Gu, Q.; You, S. L. *Trends Chem.* **2020**, *2*, 737.
- (d) Hartwig, J. F.; Larsen, M. A. *ACS Cent. Sci.* **2017**, *46*, 4299.
- (e) Zheng, Q.; Jiao, N. *Chem. Soc. Rev.* **2016**, *45*, 4590.
- (f) Sun, K.; Xiao, F.; Yu, B.; He, W. *Chin. J. Catal.* **2021**, *13*, 63850.
- (g) Xie, L.; Peng, S.; Yang, L.; Peng, C.; Lin, Y.; Yu, X.; Cao, Z.; Peng, Y.; He, W. *Green Chem.* **2021**, *23*, 374.
- (h) Chen, J.; Wu, H.; Gui, Q.; Yan, S.; Deng, J.; Lin, Y.; Cao, Z.; He, W. *Chin. J. Catal.* **2021**, *42*, 1445.
- (i) Chen, J.; Zhong, C.; Gui, Q.; Zhou, Y.; Fang, Y.; Liu, K.; Lin, Y.; Cao, Z.; He, W. *Chin. Chem. Lett.* **2021**, *32*, 475.
- (j) Zhu, X.; Li, X.; Li, X. *Org. Chem. Front.* **2021**, *8*, 3128.
- (k) Chen, Q.; Yang, Y.; Wang, X. *Chin. J. Org. Chem.* **2020**, *40*, 454 (in Chinese).
- (陈倩雯, 杨耀成, 王霞, 有机化学, **2020**, *40*, 454.)
- (l) Hao, W.; Wang, Y.; Yang, G.; Liu, Y. *Chin. J. Org. Chem.* **2017**, *37*, 2678 (in Chinese).
- (郝文燕, 王昱赞, 杨国敏, 刘云云, 有机化学, **2017**, *37*, 2678.)
- (m) Gan, Z.; Li, G.; Yang, X. *Sci. China. Chem.* **2020**, *63*, 1652.
- (n) Sun, K.; Lv, Y.; Wang, J. *Org. Lett.* **2015**, *17*, 4408.
- (o) Luo, J.; Xu, X.; Zhao, Y. *Chin. J. Org. Chem.* **2017**, *37*, 2873 (in Chinese).
- (骆钧飞, 徐星, 赵延超, 梁洪泽, 有机化学, **2017**, *37*, 2873.)
- [9] Yang, Z.; He, J.; Wei, Y.; Li, W.; Liu, P.; *Org. Biomol. Chem.* **2020**, *18*, 3360.
- [10] Yang, Z.; He, J.; Wei, Y.; Li, W.; Liu, P.; Zhao, J.; Wei, Y. *Org. Biomol. Chem.* **2020**, *18*, 9088.
- [11] Le, H.; Hoang, T.; Tran, T.; Nguyen, C.; Chiem, L.; Phan, N.; Nguyen, T. *Tetrahedron Lett.* **2021**, *67*, 152879.
- [12] Liu, Y.; Wei, Y.; Yang, Z.; Li, Y.; Liu, Y.; Liu, P. *Org. Biomol. Chem.* **2021**, *19*, 5191.
- [13] Cheeseman, G. W. H.; Roy, P. D. *J. Chem. Soc. C*, **1968**, 2848.
- [14] Reeves, J. T.; Fandrick, D. R.; Tan, Z.; Song, J. J.; Lee, H.; Yee, N. K.; Senanayake, C. H. *J. Org. Chem.* **2010**, *75*, 992.

(Fan, Y.)