电化学条件下喹喔啉-2(1*H*)-酮 C(3)—H 功能化的研究进展李丹丹* 王晓辰 李闪闪 付晨雨
李倩倩 许东涛 马莹莹

(许昌学院化工与材料学院 表面微纳米材料研究所 河南省储能与转化微纳米材料重点实验室 河南许昌 461000)

摘要 喹喔啉酮及其衍生物是一类重要的氮杂环化合物,广泛存在于具有各种生物活性的天然产物、药物及功能材料中,因此,喹喔啉酮的 C—H 功能化引起了化学工作者的广泛关注. 电化学合成无需外加氧化还原试剂,直接利用电子这一“清洁试剂”参与氧化还原反应,具有反应条件温和及原子经济性好等特点,十分符合绿色化学符合绿色化学和可持续发展的要求. 随着对电化学合成反应机理的深入研究以及反应设备的标准化,该方法已经成为喹喔啉酮类化合物功能化的强有力工具. 综述了有机电化学在喹喔啉-2(1*H*)-酮 C(3)—H 功能化领域所取得的研究进展,对反应的条件和机理进行了系统的总结,并对该领域所面临的挑战及发展方向进行了展望与探讨.

关键词 电化学; 喹喔啉-2(1*H*)-酮; C(3)—H 功能化

Recent Advances in Electrochemical C(3)—H Functionalization of Quinoxalin-2(1*H*)-onesLi, Dandan* Wang, Xiaochen Li, Shanshan Fu, Chenyu
Li, Qianqian Xu, Dongtao Ma, Yingying

(Key Laboratory of Micro-Nano Materials for Energy Storage and Conversion of Henan Province, Institute of Surface Micro and Nano Materials, College of Chemical and Materials Engineering, Xuchang University, Xuchang, Henan 461000)

Abstract Quinoxalinone and its derivatives are a class of important nitrogen heterocyclic compounds, which widely exist in natural products, drugs and functional materials. Therefore, the C—H functionalization of quinoxalinone has attracted extensive attention of chemical workers. In recent years, electrochemical synthesis directly uses electron as a “clean reagent” to participate in redox reaction without additional redox reagent, which has the characteristics of mild reaction conditions and good atom economy. This method meets the requirements of green chemistry and sustainable development. With the in-depth study on the mechanism of electrochemical synthesis and the standardization of reaction equipment, this method has become a powerful tool for functionalization of quinoxalineones. The recent advances in the electrochemical C(3)—H functionalization of quinoxalin-2(1*H*)-one are summarized. The reaction transformation conditions and mechanisms are systematically discussed, and the challenges and future directions of this field are included.

Keywords electrochemical synthesis; quinoxalin-2(1*H*)-one; C(3)—H functionalization

1 Introduction

In recent years, the direct C—H functionalization has received continuous attention because of its high efficiency and atomic economy characteristics. Quinoxalin-2(1*H*)-ones, as a privileged heterocyclic unit, can be widely found in bioactive natural products, drugs and functional materi-

als.^[1] In particular, the C(3) functionalized quinoxalin-2(1*H*)-ones exhibit a variety of biological and pharmacological activities, such as anticancer,^[1e,1e,1j,2] anti-inflammatory,^[1e,1e,3] and drug resistance antagonism.^[1f,1n] Besides, polymers derived from these compounds can be used as semiconductors, which also establish their importance in

* Corresponding author. E-mail: dandanli2015@outlook.com

Received July 18, 2021; revised September 4, 2021; published online September 14, 2021.

Project supported by the National Natural Science Foundation of China (No. 21901219), the Key Project of Education Department of Henan Province (No. 20B150024), the Young Key Teacher Training Plan of Henan Provincial Higher Education in 2021 (No. 2021GGJS146), the Open Subject Research Fund of the Key Laboratory of Drug Quality Control and Evaluation in Henan Province and the Young Backbone Teachers Project of Xuchang University.

国家自然科学基金(No. 21901219)、河南教育厅重点项目(No. 20B150024)、2021 年度河南省高等教育青年骨干教师培养计划(No. 2021GGJS146)、河南省药品质量控制与评价重点实验室(郑州大学)开放课题科研基金和许昌学院杰出青年骨干资助项目.

the field of materials science.^[4] Therefore, continuous endeavors have been focused on developing efficient methods for the C(3)—H functionalization of quinoxalin-2-ones.^[5] With the development of green and sustainable chemistry, electrosynthesis has attracted considerable attention and has emerged as a new tool for organic synthesis.^[6] Compared with traditional methods for C—H functionalization, electrosynthesis eliminates the generation of waste by using electrons as mass-free reagent instead of chemical redox reagents. In particular, electrochemical C(3)—H functionalization of quinoxalin-2(1*H*)-ones has been established as one of the most fundamental tools for the economical, efficient and green synthesis of 3-substituted quinoxalinone derivatives. According to existing reports, the reaction pathways can be roughly classified into three categories: (1) Quinoxalin-2(1*H*)-one undergoes anodic oxidation, a nucleophilic addition to R²H, followed by anodic oxidation and deprotonation to afford the final product (Scheme 1, a). (2) Quinoxalin-2(1*H*)-one undergoes protonation first, followed by a radical addition to R²•, and then further anodic oxidation and deprotonation lead to the desired product (Scheme 1, b). (3) The transformation that include radical addition, anodic oxidation and deprotonation (Scheme 1, c). Although many reviews have summarized on the C(3)—H functionalization of quinoxalin-2(1*H*)-ones,^[5e,5j-5m,7] there is no systematic summary of the electrochemical C(3)—H

functionalization of quinoxalin-2(1*H*)-ones. Herein, we summarize the recent advances in the electrochemical C(3)—H functionalization of quinoxalin-2-ones as well as the discussion of their mechanisms.

2 Electrochemical C—H alkylation of quinoxalin-2(1*H*)-ones

2.1 *N*-Hydroxyphthalimide esters as alkylation reagents

The *N*-hydroxyphthalimide (NHP) esters, easily prepared from carboxylic acids, have recently been extensively involved in alkylation reaction as a free radical precursor. In 2019, Xu, Zeng and coworkers^[8] demonstrated the first NiCl₂-catalyzed electro-reductive decarboxylative coupling of quinoxalinones with *N*-hydroxyphthalimide esters (Scheme 2). By utilizing NiCl₂ as catalyst, graphite felt and nickel foam as the electrode materials, different substitutions on the phenyl of quinoxalin-2(1*H*)-ones could give good yields, and it was also found that *N*-protected quinoxalin-2(1*H*)-ones could be compatible with the reaction. At the same time, a variety of NHP esters involving primary, secondary, tertiary carboxylic acids and also amino acids, all could be transformed into the corresponding products in satisfactory yields.

Scheme 1 Reaction pathways for electrochemical C(3)—H functionalization of quinoxalin-2-ones

Scheme 2 Nickel-catalyzed electrochemical reductive decarboxylative coupling of *N*-hydroxyphthalimide esters with quinoxalinones

Control experiments indicated that the cyclohexyl radical might be the reaction intermediate, which would go through a nucleophilic addition reaction with quinoxalinones complex **B**, leading to the radical cation intermediate **C**, which loses a proton to afford intermediate **D**, followed by single-electron oxidation and remove of Ni(II) species to form target products (Scheme 3).

Subsequently, a method without the metal-catalysis for the electrochemical decarboxylative C(3) alkylation of quinoxalin-2(1*H*)-ones was reported by Wang group (Scheme 4).^[9] In this work, a series of *N*-hydroxyphthalimide esters containing chain, cyclic, primary, secondary and tertiary carboxylic acids, were all well tolerated in this electrochemical oxidative alkylation and furnished the alkylated quinoxalinones. Furthermore, various substituted quinoxalinones were suitable for this procedure. It is worth mentioning that sensitive allyl and alkyne groups on the quinoxalinones are also retained under the electrochemical

conditions.

Mechanism research indicated that *N*-phenylmethacrylamide obtains an electron from cathode to give the intermediate **A**. Subsequently, radical anion **A** undergoes N—O bond cleavage and decarboxylation to generate alkyl radical **B**, which undergoes oxidative addition with quinoxalin-2(1*H*)-ones to yield the radical intermediate **2**, followed by 1,2-H shift, single-electron oxidation, and deprotonation process to access the final product (Scheme 3).

2.2 Alkyl boronic compounds as alkylation reagents

Organic boron compounds have a prominent role in chemical synthesis due to their high stability, low toxicity and extensive reactivity.^[10] In 2021, Wang group^[11] discovered an oxidative C—H alkylation of quinoxalin-2(1*H*)-ones under electrochemical conditions with organoboron compounds as alkylation reagent (Scheme 6).

Scheme 3 Possible mechanism

Scheme 4 Electrochemical decarboxylative C(3) alkylation of quinoxalin-2(1*H*)-ones

Scheme 5 Possible mechanism

Without redox mediator or catalyst, a wide range of quinoxalin-2(1*H*)-ones reacted well with *n*-Bu₄NPF₆ as the electrolyte, graphite plates as electrodes, trifluoroacetic acid as an additive, assembled in an undivided cell. Moreover, the method could be applied not only for primary, secondary and tertiary alkylboronic acids, but also for organotrifluoroborates and boronic esters (Scheme 6).

Scheme 6 Electrochemical oxidative C—H alkylation of quinoxalin-2(1*H*)-ones with organoboron compounds

As shown in Scheme 7, a possible mechanism has been proposed for electro-oxidative C—H alkylation of quinoxalin-2(1*H*)-ones. Initially, alkylboronic acid is oxidized at the anode via a single electron transfer (SET) process to form an alkyl radical. Meanwhile, the alkyl radical is intercepted by the protonated quinoxaline-2(1*H*)-one to pro-

duce the radical cation **A**. Subsequently, intermediate **A** undergoes deprotonation to generate C-radical intermediate **B**. Further anodic oxidation and deprotonation of intermediate **B** afford the C(3)-alkylated quinoxalin-2(1*H*)-ones.

Scheme 7 Possible mechanism

2.3 Alkyl carbazates as alkylation reagents

In 2020, Wang and co-workers^[12] revealed a deoxygenative C(3)-alkylation of quinoxalin-2(1*H*)-ones by using carbazates as a new type of electrochemically activated alkylating agent (Scheme 8). With a carbon anode and a platinum plate cathode, various substituted quinoxalin-2(1*H*)-ones were suitable for this procedure. It is noteworthy that the primary, secondary and tertiary alkyl radicals obtained through anodic oxidation can react smoothly in this transformation and afford the corresponding products with a satisfactory yield.

According to the mechanistic investigation, a stepwise dehydrazination-decarboxylation sequence was involved in the C(3)-alkylation of quinoxalin-2(1*H*)-ones (Scheme 9). Initially, alkyl carbazate is oxidized at the anode and deprotonated to afford hydrazinecarboxylate radical **A** and diazenecarboxylate **B**. Then, the diazenecarboxylate is

Scheme 8 Electrochemical deoxygenative alkylation of quinoxalin-2(1*H*)-ones with alkyl carbazates

further oxidized and deprotonated to form diazene **C**, which undergoes the cleavage of C—N bond to generate acyl radical **D** and releases molecular nitrogen. The subsequent decarboxylation affords the alkyl radical. The subsequent decarboxylation affords the alkyl radical, followed by the reaction with quinoxalin-2(1*H*)-ones, oxidation and deprotonation to produce the desired product.

Scheme 9 Possible mechanism

Scheme 10 Electrochemical cross-coupling of C(sp²)-H with aryldiazonium salts via a paired electrolysis

3 Electrochemical C—H arylation of quinoxalin-2(1*H*)-ones

3.1 Aryldiazonium salts as arylation reagents

As a common aryl radical precursor, aryldiazonium salts are widely used in the construction of aryl C—C bonds.^[13] In 2019, Zeng group^[14] revealed an efficient electrochemical C(3) arylation of quinoxalin-2(1*H*)-ones with aryldiazonium salts under paired electrolysis conditions (Scheme 10). In a simple undivided cell, the electrolysis was conducted at low controlled potentials without external supporting electrolyte. A variety of quinoxalin-2(1*H*)-ones were suitable for this C(sp²)-H arylation reaction and gave

the corresponding products in satisfactory yields. Meanwhile, aryldiazonium salts could tolerate different substituents on the aromatic ring and afforded the desired products. It was found that the electronic effect had an obvious effect on the transformation. And the substrates with electron-donating groups gave higher yield than that of electron-withdrawing substituted aryldiazonium salts. Notably, aryldiazonium salts substituted 2-I or 2-Br couldn't provide the corresponding products owing to the possible reduction reaction of aryl bromides or aryl iodides in the electro-

chemical condition. In addition, the authors also attempted the heterocyclic aryldiazonium salt as the aryl radical precursor.

Control experiments indicated that cathodic reduction led to the formation of the aryl radical. According to the mechanism study and research reports, the possible reaction mechanism is outlined in Scheme 11. Firstly, aryldiazonium salts could be reduced at the surface of the cathode, followed by loss of molecular N_2 to afford aryl radical. Then, the aryl radical undergoes a radical addition to quinoxalin-2(1*H*)-one, generating the intermediate **A**. Finally, the intermediate **A** is oxidized at the anode to finish the desired arylation product. Nevertheless, the pathway which intermediate **A** reacts with aryldiazonium salts followed by deprotonation to form the product, along with the release of aryl radical can't be excluded.

Scheme 11 Possible mechanism

3.2 Pyrroles as arylation reagents

In 2019, Wang *et al.*^[15] reported an electrochemical heteroarylation of quinoxalin-2(1*H*)-ones with pyrroles via cross-dehydrogenation coupling (Scheme 12). The reaction could be compatible with different substituted quinoxalin-2(1*H*)-ones and gave the corresponding products. Unfortunately, the author did not make a further study on this reaction.

3.3 Cyanopyridines as arylation reagents

Recently, the electrochemical reduction of cyanopyridines has arisen increasing attention.^[16] Wen, Sun, Yang and coworkers^[17] reported the electroreductive-induced C(3) pyridylation of quinoxalin-2(1*H*)-ones with cyanopyridines (Scheme 13). The reaction was performed in an undivided cell under a constant current of 15 mA with CH_2Cl_2 /hexafluoroisopropanol (HFIP) as the solvent, carbon felt as anode and Pt as counter electrode. A broad range of functional groups bearing quinoxalin-2(1*H*)-ones were well tolerated and provided good to excellent yields. It is worth noting that some C(2) or C(4) replaced cyanopyridines were not compatible in this transformation (Scheme 13). It might

Scheme 12 Electrochemical heteroarylation of quinoxalin-2(1*H*)-ones with pyrroles

Scheme 13 Electroreductive-induced C(3) pyridylation of quinoxalin-2(1*H*)-ones with cyanopyridines

be due to the position of the substituent and the electronic effect.

This proposed mechanism is shown in Scheme 14. Firstly, cyanopyridine undergoes proton exchange with HFIP to generate intermediate **A**, which obtains an electron at the cathode producing the intermediate **B**. Then, intermediate **B** reacts with quinoxalin-2(1*H*)-one to provide the intermediate **C**, which is subjected to oxidation and decyanation to furnish the free radical **D**. Finally, radical **D** is converted into the desired product via the dehydrogenation and isomerization.

Scheme 14 Possible mechanism

4 Electrochemical C—H trifluoromethylation of quinoxalin-2(1*H*)-ones

Trifluoromethyl, as an important chemical group, has been widely used in organic synthesis and drug molecules because of its high lipid solubility, good metabolic stability, high electronegativity and bioavailability.^[18] The group of Zeng^[19] reported their efforts on bromide ions-mediated electrosynthesis of Minisci-type trifluoromethylation of

quinoxaline-2(1*H*)-ones in an undivided cell (Scheme 15). Due to the sulfonyl hypobromite intermediates generated *in situ*, the CF₃ radical could be regulated and controlled at a low concentration, thus improving the efficiency of indirect electrolysis. Utilizing graphite plate as the anode and Pt net as the cathode, a variety of quinoxaline-2(1*H*)-ones could be transformed into the corresponding products.

Scheme 15 Bromide ions-mediated electrosynthesis of Minisci-type trifluoromethylation of quinoxaline-2(1*H*)-ones

Mechanistic elucidation shows that Br₂, which is generated from the anodic oxidation of bromide, reacts with Zn(SO₂CF₃)₂ to give sulfonyl hypobromite **A** or sulfonyl bromide **B**. Subsequently, sulfonyl bromide decomposition yields the CF₃ free radical via a cathodic reduction or a homolytic cleavage process, along with the release of one molecule of SO₂. Then, the CF₃ free radical adds to the quinoxalin-2(1*H*)-one to afford the radical intermediate **C**, which is successively oxidized and deprotonated to deliver the desired product (Scheme 16).

Scheme 16 Possible mechanism

5 Electrochemical C—H cyanation of quinoxalin-2(1*H*)-ones

Nitrile is an important structural part of natural products and drugs.^[20] Organic nitriles are versatile intermediates for the production of pharmaceuticals, pesticides, dyes, pigments, polyamides and various fine chemicals.^[20c,21] In addition, nitriles can be used as a variety of intermediates and widely converted to other functional groups, such as amines, acids, aldehydes, amides, isocyanates, ketones, esters and heterocycles.^[22] Therefore, the facile introduction of a cyano group to organic compounds is very attractive for organic chemists. Very recently, Sun group^[23] developed an electro-oxidative method for efficient C—H cyanation of quinoxalin-2(1*H*)-ones with TMSCN as a nucleophilic cyano source in a mixed solvent of CH₃CN and KH₂PO₄/K₂HPO₄ (pH=9) buffer solution (Scheme 17). The protocol features oxidant free, ambient condition, a broad substrate scope, excellent functional group tolerance and good to excellent yield of the desired product. Notably, quinoxalin-2(1*H*)-ones bearing *N*-substituents (chain alkyl, cycloalkyl, cyanomethyl, ester group, phenacyl, 3-bromo-propyl allyl, propargyl, benzyl, *ect.*) successfully transformed the C(3) alkylated derivative in moderate to excellent yields. In addition, quinoxalin-2(1*H*)-ones possessing both electron-donating groups and electron-withdrawing groups could afford the desired products under this electrochemical oxidative conditions.

Scheme 17 Electrooxidative C(3) cyanation of quinoxalin-2(1*H*)-ones with TMSCN

A plausible mechanism of this electrochemical oxidative C—H cyanation of quinoxalin-2(1*H*)-ones is outlined in Scheme 18. Initially, quinoxalin-2(1*H*)-one is oxidized at the anode via a single-electron transfer (SET) pathway to afford the intermediate **A**, which then undergoes nucleophilic attack of TMSCN to produce intermediate **B**. Finally, the intermediate **B** is further oxidized and deprotonated to furnish the final product.

Scheme 18 Possible mechanism

6 Electrochemical C—H amination of quinoxalin-2(1*H*)-ones

The construction of C—N bond plays an important role in pharmaceutical chemistry, materials science and organic synthesis.^[24] The direct C—H amination has attracted much attention due to its high efficiency and atomic economy. In 2019, Zeng group^[25] developed an efficient protocol for the electro-oxidative C(3)—H amination of quinoxalin-2(1*H*)-ones, which allowed aliphatic amines and azoles as the nitrogen reagents (Scheme 19). Utilizing graphite rod as the anode and platinum plate as the cathode, a variety of 3-aminoquinoxalin-2(1*H*)-ones were obtained in moderate to excellent yields under electrochemical oxidation conditions. The reaction mechanism was initiated from the nucleophilic addition of the amine to the protonated quinoxaline-2(1*H*)-one.

Based on the cyclic voltammetry and control experiments, the authors proposed two reaction mechanism pathways and also suggested that the pathway involving a radical process is most likely (Scheme 20). In pathway 1, amine is oxidized in the anode to form an *N*-centered radical **A**, which undergoes a radical addition to yield the radical cation **C**. Then, the intermediate **C** is converted into the corresponding product via a single-electron transfer (SET) oxidation processes and deprotonation. In pathway 2, amine undergoes nucleophilic addition with protonated quinoxaline-2(1*H*)-one to produce the intermediate **B**, followed by

Scheme 19 Electro-oxidative C(3)—H amination of quinoxalin-2(1*H*)-ones

further oxidation and decarboxylation to afford the final

product.

7 Electrochemical C—H alkoxylation of quinoxalin-2(1*H*)-ones

Generally, alcohols participate in the formation of C—O bonds as nucleophiles. In 2020, Li, Yu and co-workers^[26] exploited an efficient electrochemical protocol for the synthesis of 3-alkoxylated quinoxalin-2(1*H*)-ones through dehydrogenative C—H/O—H cross-coupling (Scheme 21). With graphite rod as anode and Pt plate as cathode, different quinoxalin-2(1*H*)-ones bearing electron-rich and electron-poor substituents reacted smoothly, affording the corresponding products in moderate to good yields under standard conditions. Meanwhile, primary alcohols were well tolerated in this protocol. For the secondary alcohols, the yields decreased slightly owing to the steric hindrance. Unfortunately, for the tertiary alcohols and phenols, the transformation was not realized.

Mechanism studies revealed that a radical process might be involved in the reaction (Scheme 22). First, single-electron-transfer (SET) oxidation of the quinoxalin-2(1*H*)-one by the anode leads to a radical-cation intermediate A. Subsequent nucleophilic addition of ROH to intermediate

Scheme 20 Possible mechanism

Scheme 21 Electrochemical dehydrogenative C—H/O—H cross-coupling for the synthesis of 3-alkoxylated quinoxalin-2(1*H*)-ones

A generates a radical cation **B**. Further anodic oxidation and deprotonation lead to the desired product.

Scheme 22 Possible mechanism

8 Electrochemical C—H thioetherification of quinoxalin-2(1H)-ones

Over the past decade, oxidative C—H/S—H cross-coupling has received continuous attention because of the highly atom-economical and environmental friendly

characteristics of the approach.^[27] In 2020, the group of Li^[28] demonstrated an electrooxidative cross-coupling between thiols and quinoxalin-2(1H)-ones for the synthesis of 3-thioquinoxalinones with platinum electrodes as both the anode and the cathode (Scheme 23). Both electron-donating and electron-withdrawing groups on the phenyl ring of quinoxalin-2(1H)-ones were tolerated under the reaction conditions to provide the desired products in acceptable yields. Different *N*-substituted quinoxalin-2(1H)-ones also tolerated the reaction conditions, but it showed less responsive than the unprotected quinoxalin-2(1H)-one. In addition, aliphatic thiols worked well and provided moderate to good yields, while the thiophenols showed poor reactivity in the transformation.

Control experiments revealed that the sulfur radical was likely to be a reaction intermediate, which would add to protonated quinoxaline-2(1H)-one **A**, delivering **C**-radical cation species **B**. Oxidation of intermediate **B** at the anode, followed by deprotonation finished the 3-thioquinoxalinones (Scheme 24).

9 Electrochemical C—H phosphorylation of quinoxalin-2(1H)-ones

Organophosphorus compounds widely exist in pesticides, organic ligands, materials and bioactive compounds.^[29] It is of great significance to effectively construct different C—P

Scheme 23 Electrooxidative cross-coupling for the synthesis of 3-thioquinoxalinones

Scheme 24 Possible mechanism

bonds. In 2019, the group of Zeng^[30] developed an efficient electrochemical phosphonation of quinoxalin-2(1*H*)-ones with H-phosphonates or H-phosphine oxides via C—H/P—H dehydrogenative cross-coupling (Scheme 25). This protocol provided a concise and practical route to 3-phosphonated quinoxalin-2(1*H*)-ones with excellent functional group tolerance and broad substrate scope.

The mechanistic studies revealed that the reaction proceeded through a radical process (Scheme 26). Initially, quinoxalin-2(1*H*)-one is oxidized at the anodic, lead to the formation of immediate **A**, which may undergo homogeneous electron transfer with H-phosphonates or H-phos-

phine oxides to generate the phosphorus-centered radical **B**, along with the regeneration of quinoxalin-2(1*H*)-one. Subsequently, radical **C** undergoes radical addition to the protonated quinoxalin-2(1*H*)-one with the formation of a carbon radical intermediate **D**. Finally, the intermediate **D** is converted into the corresponding product via single-electron transfer (SET) oxidation and deprotonation processes.

Subsequently, Wang and coworkers^[15] achieved the coupling of quinoxalin-2(1*H*)-ones with diarylphosphine oxides using similar electrochemical oxidative methods (Scheme 27). This reaction exhibited good functional

Scheme 25 Electrochemical phosphonation of quinoxalin-2(1*H*)-ones via C—H/P—H dehydrogenative cross-coupling

Scheme 26 Possible mechanism

Scheme 27 Electrochemical oxidative coupling of quinoxalin-2(1*H*)-ones with diarylphosphine oxides

group tolerance and a wide range of substrate scope and gave moderate to high yields. The reaction mechanism proposed by the author is similar to the Zeng group.

10 Conclusions

In this review, we have summarized recent developments in electrochemical C(3)—H functionalization of quinoxalin-2-ones, including the formation of C—C, C—O, C—N, C—S and C—P bonds. Compared with the traditional C—H functionalization reaction, the electrosynthesis C—H functionalization has a high atom-economy, because it does not need to sacrifice oxidants. In the case of adjustable operating current and potential, some sensitive functional groups could be well tolerated. Mechanistic insights could help us to understand the nature of these reactions. However, there are still some major challenges to be addressed in this field. (1) The scope of electrochemical C(3)—H functionalization of quinoxaline-2-one may need to be further expanded, such as acylation, amidation and halogenation. (2) The functionalized products of quinoxaline-2-one still have shortcomings in drug design and development and lead compound modification, thus we should pay more attention to the development of pharmaceutical chemistry, and further deepened the related research. Overall, electrochemical C(3)—H functionalization offer a new opportunity for green synthesis of various C(3)-substituted quinoxalin-2(1H)-ones in the areas of pharmaceutical chemistry and functional materials. Importantly, with the development of sustainable chemistry and green chemistry, electrochemical synthesis, as a safe and more economical alternative to traditional synthesis methods, will attract more attention. We believe that this review will provide a useful reference and important guidance for the further development of electrochemical synthesis and its application for the functionalization of quinoxalin-2-ones.

References

- [1] (a) Waring, M. J.; Ben-Hadda, T.; Kotchevar, A. T.; Ramdani, A.; Touzani, R.; Elkadiri, S.; Hakkou, A.; Bouakka, M.; Ellis, T. *Molecules* **2002**, *7*, 641.
 (b) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893.
 (c) Refaat, H. M.; Moneer, A. A.; Khalil, O. M. *Arch. Pharm. Res.* **2004**, *27*, 1093.
 (d) Carta, A.; Piras, S.; Loriga, G.; Paglietti, G. *Mini-Rev. Med. Chem.* **2006**, *6*, 1179.
 (e) El-Hawash, S. A. M.; Habib, N. S.; Kassem, M. A. *Arch. Pharm.* **2006**, *339*, 564.
 (f) Meyer, E.; Joussef, A. C.; de Souza, L. D. B. P. *Synth. Commun.* **2006**, *36*, 729.
 (g) Moarbess, G.; Deleuze-Masquefa, C.; Bonnard, V.; Gayraud-Paniagua, S.; Vidal, J.-R.; Bressolle, F.; Pinguet, F.; Bonnet, P.-A. *Bioorg. Med. Chem.* **2008**, *16*, 6601.
 (h) Ramli, Y.; Benzeid, H.; Bouhfid, R.; Rodi, Y. K.; Ferfra, S.; Essassi, E. M. *Stud. Cercet. Stiint.: Chim. Ing. Chim., Biotechnol., Ind. Aliment.* **2010**, *11*, 67.
 (i) Liu, R.; Huang, Z.; Murray, M. G.; Guo, X.; Liu, G. *J. Med. Chem.* **2011**, *54*, 5747.
 (j) Galal, S. A.; Khairat, S. H. M.; Ragab, F. A. F.; Abdelsamie, A. S.; Ali, M. M.; Soliman, S. M.; Mortier, J.; Wolber, G.; El Diwani, H. I. *Eur. J. Med. Chem.* **2014**, *86*, 122.
 (k) Hussain, S.; Parveen, S.; Hao, X.; Zhang, S.; Wang, W.; Qin, X.; Yang, Y.; Chen, X.; Zhu, S.; Zhu, C.; Ma, B. *Eur. J. Med. Chem.* **2014**, *80*, 383.
 (l) Khattab, S. N.; Abdel Moneim, S. A. H.; Bekhit, A. A.; El Massry, A. M.; Hassan, S. Y.; El-Faham, A.; Ali Ahmed, H. E.; Amer, A. *Eur. J. Med. Chem.* **2015**, *93*, 308.
 (m) Qin, X.; Hao, X.; Han, H.; Zhu, S.; Yang, Y.; Wu, B.; Hussain, S.; Parveen, S.; Jing, C.; Ma, B.; Zhu, C. *J. Med. Chem.* **2015**, *58*, 1254.
 (n) Cil, O.; Phuan, P.-W.; Lee, S.; Tan, J.; Haggie, P. M.; Levin, M. H.; Sun, L.; Thiagarajah, J. R.; Ma, T.; Verkman, A. S. *Cell Mol. Gastroenterol. Hepatol.* **2016**, *2*, 317.
 (o) Shi, L.; Li, X.; Hu, W.; Wu, J.; Zhou, H.; Zhou, H. *Mini-Rev. Med. Chem.* **2018**, *18*, 392.
 [2] Weiwer, M.; Spoonamore, J.; Wei, J.; Guichard, B.; Ross, N. T.; Masson, K.; Silkworth, W.; Dandapani, S.; Palmer, M.; Scherer, C. A.; Stern, A. M.; Schreiber, S. L.; Munoz, B. *ACS Med. Chem. Lett.* **2012**, *3*, 1034.
 [3] (a) Smits, R. A.; Lim, H. D.; Hanzer, A.; Zuiderveld, O. P.; Guaita, E.; Adami, M.; Coruzzi, G.; Leurs, R.; de Esch, I. J. P. *J. Med. Chem.* **2008**, *51*, 2457.
 (b) Abu-Hashem, A. A.; Gouda, M. A.; Badria, F. A. *Eur. J. Med. Chem.* **2010**, *45*, 1976.
 (c) Mangold, S. L.; Prost, L. R.; Kiessling, L. L. *Chem. Sci.* **2012**, *3*, 772.
 [4] Quinn, J.; Guo, C.; Ko, L.; Sun, B.; He, Y.; Li, Y. *RSC Adv.* **2016**, *6*, 22043.
 [5] (a) Xie, L.-Y.; Bai, Y.-S.; Xu, X.-Q.; Peng, X.; Tang, H.-S.; Huang, Y.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Green Chem.* **2020**, *22*, 1720.
 (b) Wang, L.; Bao, P.; Liu, W.; Liu, S.; Hu, C.; Yue, H.; Yang, D.; Wei, W. *Chin. J. Org. Chem.* **2018**, *38*, 3189 (in Chinese).
 (王雷雷, 鲍鹏丽, 刘伟伟, 刘思彤, 胡昌松, 岳会兰, 杨道山, 魏伟, 有机化学, **2018**, *38*, 3189.)
 (c) Wu, Y.; Chen, J.-Y.; Ning, J.; Jiang, X.; Deng, J.; Deng, Y.; Xu, R.; He, W.-M. *Green Chem.* **2021**, *23*, 3950.
 (d) Shi, J.; Wei, W. *Chin. J. Org. Chem.* **2020**, *40*, 2170 (in Chinese).
 (时建伟, 魏伟, 有机化学, **2020**, *40*, 2170.)
 (e) Mao P.; Zhu, J.; Yuan, J.; Yang, L.; Xiao, Y.; Zhang, C. *Chin. J. Org. Chem.* **2019**, *39*, 1529 (in Chinese).
 (毛璞, 朱军亮, 袁金伟, 杨亮茹, 肖咏梅, 张长森, 有机化学, **2019**, *39*, 1529.)
 (f) Xie, L.-Y.; Peng, S.; Yang, L.-H.; Peng, C.; Lin, Y.-W.; Yu, X.; Cao, Z.; Peng, Y.-Y.; He, W.-M. *Green Chem.* **2021**, *23*, 374.
 (g) Yi, R.; He, W. *Chin. J. Org. Chem.* **2021**, *41*, 1267 (in Chinese).
 (易荣楠, 何卫民, 有机化学, **2021**, *41*, 1267.)
 (h) Xie, L.-Y.; Liu, Y.-S.; Ding, H.-R.; Gong, S.-F.; Tan, J.-X.; He, J.-Y.; Cao, Z.; He, W.-M. *Chin. J. Catal.* **2020**, *41*, 1168.
 (i) Chen, J.-Y.; Wu, H.-Y.; Gui, Q.-W.; Yan, S.-S.; Deng, J.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Chin. J. Catal.* **2021**, *42*, 1445.
 (j) Ke, Q.; Yan, G.; Yu, J.; Wu, X. *Org. Biomol. Chem.* **2019**, *17*, 5863.
 (k) Rostoll-Berenguer, J.; Blay, G.; Pedro, J. R.; Vila, C. *Eur. J. Org. Chem.* **2020**, *2020*, 6148.
 (l) Monika, M.; Selvakumar, S. *Synthesis* **2019**, *51*, 4113.
 (m) Ghosh, P.; Das, S. *Synth. Commun.* **2020**, *50*, 2266.
 [6] (a) Frontana-Urbe, B. A.; Little, R. D.; Ibanez, J. G.; Palma, A.; Vasquez-Medrano, R. *Green Chem.* **2010**, *12*, 2099.
 (b) Jiang, Y.; Xu, K.; Zeng, C. *Chem. Rev.* **2018**, *118*, 4485.
 (c) Jiao, K.-J.; Xing, Y.-K.; Yang, Q.-L.; Qiu, H.; Mei, T.-S. *Acc. Chem. Res.* **2020**, *53*, 300.
 (d) Karkas, M. D. *Chem. Soc. Rev.* **2018**, *47*, 5786.
 (e) Ma, C.; Fang, P.; Mei, T.-S. *ACS Catal.* **2018**, *8*, 7179.
 (f) Maeda, H.; Ohmori, H. *Acc. Chem. Res.* **1999**, *32*, 72.
 (g) Minter, S. D.; Baran, P. *Acc. Chem. Res.* **2020**, *53*, 545.
 (h) Möhle, S.; Zirbes, M.; Rodrigo, E.; Gieshoff, T.; Wiebe, A.; Waldvogel, S. R. *Angew. Chem., Int. Ed.* **2018**, *57*, 6018.

- (i) Novaes, L. F. T.; Liu, J.; Shen, Y.; Lu, L.; Meinhardt, J. M.; Lin, S. *Chem. Soc. Rev.* **2021**, *50*, 7941.
- (j) Röckl, J. L.; Pollok, D.; Franke, R.; Waldvogel, S. R. *Acc. Chem. Res.* **2020**, *53*, 45.
- (k) Sauermann, N.; Meyer, T. H.; Qiu, Y.; Ackermann, L. *ACS Catal.* **2018**, *8*, 7086.
- (l) Waldvogel, S. R.; Lips, S.; Selt, M.; Riehl, B.; Kampf, C. J. *Chem. Rev.* **2018**, *118*, 6706.
- (m) Wang, H.; Gao, X.; Lv, Z.; Abdelilah, T.; Lei, A. *Chem. Rev.* **2019**, *119*, 6769.
- (n) Yamamoto, K.; Kuriyama, M.; Onomura, O. *Acc. Chem. Res.* **2020**, *53*, 105.
- (o) Yan, M.; Kawamata, Y.; Baran, P. S. *Chem. Rev.* **2017**, *117*, 13230.
- (p) Yi, H.; Zhang, G.; Wang, H.; Huang, Z.; Wang, J.; Singh, A. K.; Lei, A. *Chem. Rev.* **2017**, *117*, 9016.
- (q) Yuan, Y.; Lei, A. *Acc. Chem. Res.* **2019**, *52*, 3309.
- [7] Tan, Y.; Wang, J.; Zhang, H.-Y.; Zhang, Y.; Zhao, J. *Front. Chem.* **2020**, *8*, 582.
- [8] Lian, F.; Xu, K.; Meng, W.; Zhang, H.; Tan, Z.; Zeng, C. *Chem. Commun.* **2019**, *55*, 14685.
- [9] Niu, K.; Song, L.; Hao, Y.; Liu, Y.; Wang, Q. *Chem. Commun.* **2020**, *56*, 11673.
- [10] Traister, K. M.; Molander, G. A. Synthesis and Application of Organoboron Compounds, In *Topics in Organometallic Chemistry*, Vol. 49, Eds.: Fernández, E.; Whiting, A., Springer, Switzerland, **2015**, pp. 117~151.
- [11] Niu, K.; Hao, Y.; Song, L.; Liu, Y.; Wang, Q. *Green Chem.* **2021**, *23*, 302.
- [12] Gao, Y.; Wu, Z.; Yu, L.; Wang, Y.; Pan, Y. *Angew. Chem., Int. Ed.* **2020**, *59*, 10859.
- [13] Mo, F.; Qiu, D.; Zhang, L.; Wang, J. *Chem. Rev.* **2021**, *121*, 5741.
- [14] Jiang, Y.-Y.; Dou, G.-Y.; Zhang, L.-S.; Xu, K.; Little, R. D.; Zeng, C.-C. *Adv. Synth. Catal.* **2019**, *361*, 5170.
- [15] Hu, C.; Hong, G.; Zhou, C.; Tang, Z.-C.; Han, J.-W.; Wang, L.-M. *Asian J. Org. Chem.* **2019**, *8*, 2092.
- [16] (a) Lehnher, D.; Lam, Y.-H.; Nicastrì, M. C.; Liu, J.; Newman, J. A.; Regalado, E. L.; DiRocco, D. A.; Rovis, T. *J. Am. Chem. Soc.* **2020**, *142*, 468.
- (b) Nicastrì, M. C.; Lehnher, D.; Lam, Y.-H.; DiRocco, D. A.; Rovis, T. *J. Am. Chem. Soc.* **2020**, *142*, 987.
- (c) Zhang, S.; Li, L.; Li, X.; Zhang, J.; Xu, K.; Li, G.; Findlater, M. *Org. Lett.* **2020**, *22*, 3570.
- (d) Zhang, X.; Yang, C.; Gao, H.; Wang, L.; Guo, L.; Xia, W. *Org. Lett.* **2021**, *23*, 3472.
- [17] Wen, J.; Yang, X.; Yan, K.; Qin, H.; Ma, J.; Sun, X.; Yang, J.; Wang, H. *Org. Lett.* **2021**, *23*, 1081.
- [18] (a) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320.
- (b) Nie, J.; Guo, H.-C.; Cahard, D.; Ma, J.-A. *Chem. Rev.* **2011**, *111*, 455.
- (c) Tomashenko, O. A.; Grushin, V. V. *Chem. Rev.* **2011**, *111*, 4475.
- (d) Liu, T.; Shen, Q. *Eur. J. Org. Chem.* **2012**, *2012*, 6679.
- [19] Dou, G.-Y.; Jiang, Y.-Y.; Xu, K.; Zeng, C.-C. *Org. Chem. Front.* **2019**, *6*, 2392.
- [20] (a) Scheuer, P. J. *Acc. Chem. Res.* **1992**, *25*, 433.
- (b) Fleming, F. *Nat. Prod. Rep.* **1999**, *16*, 597.
- (c) Fleming, F. F.; Yao, L.; Ravikumar, P. C.; Funk, L.; Shook, B. C. *J. Med. Chem.* **2010**, *53*, 7902.
- [21] (a) Larock, R. C. *Comprehensive Organic Transformations*, Wiley-VCH, New York, **1989**, p. 819.
- (b) Caboni, P.; Sammelson, R. E.; Casida, J. E. *J. Agric. Food Chem.* **2003**, *51*, 7055.
- (c) Kuhn, P.; Thomas, A.; Antonietti, M. *Macromolecules* **2009**, *42*, 319.
- [22] (a) Liskey, C. W.; Liao, X.; Hartwig, J. F. *J. Am. Chem. Soc.* **2010**, *132*, 11389.
- (b) Wade, J. R. *Organic Chemistry*, 8th ed., Pearson Education Inc, Glenview, Illinois, USA, **2013**.
- (c) Zhang, S.; del Pozo, J.; Romiti, F.; Mu, Y.; Torker, S.; Hoveyda, A. H. *Science* **2019**, *364*, 45.
- [23] Zhan, Y.; Li, Y.; Tong, J.; Liu, P.; Sun, P. *Eur. J. Org. Chem.* **2021**, *2021*, 2193.
- [24] (a) Vitaku, E.; Smith, D. T.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 10257.
- (b) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. *Chem. Soc. Rev.* **2016**, *45*, 546.
- [25] Li, K.-J.; Xu, K.; Liu, Y.-G.; Zeng, C.-C.; Sun, B.-G. *Adv. Synth. Catal.* **2019**, *361*, 1033.
- [26] Jiang, X.; Yang, L.; Ye, Z.; Du, X.; Fang, L.; Zhu, Y.; Chen, K.; Li, J.; Yu, C. *Eur. J. Org. Chem.* **2020**, *2020*, 1687.
- [27] Song, C.; Liu, K.; Dong, X.; Chiang, C.-W.; Lei, A. *Synlett* **2019**, *30*, 1149.
- [28] Zhou, J.; Li, Z.; Sun, Z.; Ren, Q.; Zhang, Q.; Li, H.; Li, J. *J. Org. Chem.* **2020**, *85*, 4365.
- [29] (a) Demmer, C. S.; Krosgaard-Larsen, N.; Bunch, L. *Chem. Rev.* **2011**, *111*, 7981.
- (b) Montchamp, J.-L. *Acc. Chem. Res.* **2014**, *47*, 77.
- (c) Mady, M. F.; Kelland, M. A. *Energy Fuels* **2017**, *31*, 4603.
- (d) Ntatsopoulos, V.; Maccioniuk, K.; Mucha, A.; Vassiliou, S.; Berlicki, L. *Eur. J. Med. Chem.* **2018**, *159*, 307.
- [30] Li, K.-J.; Jiang, Y.-Y.; Xu, K.; Zeng, C.-C.; Sun, B.-G. *Green Chem.* **2019**, *21*, 4412.

(Zhao, C.)