Rh(III)催化偶氮苯与碳酸亚乙烯基酯的环化反应

张文杰^a 李超^a 王博超^a 高慧^{*,a,b} 李洪基^{*,a}^(a) 淮北师范大学化学与材料科学学院 教育部绿色和精准合成化学及应用重点实验室 安徽淮北 235000)^(b) 广西师范大学 省部共建药用资源化学与药物分子工程国家重点实验室 广西桂林 541004)

摘要 报道了利用导向基策略实现 Rh(III)催化芳基偶氮苯与碳酸亚乙烯基酯的环化反应。该反应体系可以一步构建吡唑类化合物骨架。值得指出的是碳酸亚乙烯基酯中的一个碳原子参与吡唑环的构建,另一个碳原子用于醛的生成。该反应能够高效合成一系列吡唑衍生物,产物收率中等到良好。

关键词 Rh 催化; 偶氮苯; 碳酸亚乙烯基酯; 环化反应

Rh(III)-Catalyzed Annulation of Azobenzenes with Vinylene Carbonate

Zhang, Wenjie^a Li, Chao^a Wang, Bochao^a Gao, Hui^{*,a,b} Li, Hongji^{*,a}^(a) Key Laboratory of Green and Precise Synthetic Chemistry and Applications, Ministry of Education, School of Chemistry and Materials Science, Huaibei Normal University, Huaibei, Anhui 235000)^(b) State Key Laboratory for Chemistry and Molecular Engineering of Medicinal Resources, Guangxi Normal University, Guilin, Guangxi 541004)

Abstract A mild Rh(III)-catalyzed annulation of azobenzenes with vinylene carbonate via inert C—H bond activation has been developed. In this reaction system, vinylene carbonate provides one vinyl carbon for the construction of indazole backbone, and leaves the other one for the formation of aldehyde. This annulation enables the direct synthesis of 2*H*-indazole-3-carbaldehyde with excellent regioselectivity in moderate to good yields.

Keywords Rh catalysis; azobenzene; vinylene carbonate; annulation

1 Introduction

The indazole belongs to an important class of *N*-containing heterocycles, which features a benzene ring fused to a pyrazole molecule.^[1] Particularly, 2*H*-indazole-containing molecules having various functionalities could well control its pharmacological activities, which have been found in several best-selling drugs.^[2] In this regard, increasing attention has been focused on the construction of the indazole skeleton by using diverse strategies, including metal-catalyzed coupling,^[3a] cycloaddition of arynes with sydnone,^[3b] organophotocatalytic functionalization and some other methods.^[3] Among these works, one of the most effective method for the indazole synthesis is the transition metal-catalyzed C—H bond activation and functionaliza-

tion starting from the azobenzenes and aldehydes.^[4-6] For example, in 2013, we^[4a] have established a Pd-catalyzed *ortho*-acylation of azobenzene with aldehydes, followed by a reductive cyclization to form 2*H*-indazoles. Simultaneously, Ellman and coworkers^[4b] reported an efficient Rh-catalyzed cyclization of azobenzenes with aldehydes to construct indazole backbone in one-step. In 2014, Ellman's group^[4c] found that use of cheap Cobalt catalysis could enable the cyclization of azobenzenes with aldehydes. Additionally, Glorius *et al.*^[5] reported an oxidative annulation of arylimidates with organo azides by Rh catalysis to construct the 1*H*-indazoles skeleton.

Then, the significant advance of methods for the construction of 2*H*-indazoles using C—H bond activation has been achieved.^[6] However, these reactions generally require

* Corresponding authors. E-mail: gaohui200032@163.com; hongjili@chnu.edu.cn

Received July 18, 2021; revised September 16, 2021; published online September 26, 2021.

Project supported by the National Natural Science Foundation of China (No. 21772061), the Research Funding for Outstanding Young Talents in Colleges and Universities of Anhui Province (No. gxyq2020015), and the State Key Laboratory for Chemistry and Molecular Engineering of Medicinal Resources (Guangxi Normal University) (No. CMEMR2021-B14).

国家自然科学基金(No. 21772061)、安徽省高校优秀人才(No. gxyq2020015)、省部共建药用资源化学与药物分子工程国家重点实验室(广西师范大学)(No. CMEMR2021-B14)资助项目。

Previous work: The roles of vinylene carbonate

Vinylene carbonate serving as a C2 Synthon

Vinylene carbonate serving as a C1 Synthon

This work: Rh(III)-catalyzed annulation of azobenzenes with vinylene carbonate

Scheme 1 Roles of vinylene carbonate in the inert C—H bond activation

harsh reaction conditions and the difficult late-stage transformation of products extremely limited their further applications in synthetic chemistry.

Recently, the commercially available vinylene carbonate has received increasing attention for its unique roles in the formation of new chemical bonds through transition metal catalysis.^[7] For instance, Tanaka *et al.*^[8] in early 2009 found that the vinylene carbonate could act as an “ethynol” surrogate (C2 synthon) for the phenol synthesis. In 2019, Miura’s group^[9a] disclosed the critical role of vinylene carbonate as an “acetylene” substitute (C2 synthon) in Rh-catalyzed C—H bond activation for the construction of heterocycles. Recently, some other groups such as those of Zhang,^[9b] and Zhou^[9c] have independently reported the excellent works involving the vinylene carbonate, thus providing various synthetic methods for the construction of *N*-heterocyclic compounds (Schemes 1a~1b). The past years have witnessed the powerful application of vinylene carbonate (C2 synthon) in transition-metal-catalyzed C—H bond activation reactions,^[9] but vinylene carbonate acting as a C1 synthon for this purpose tend to be rarely reported.

Very recently, Ma’s group and Cheng’s group separately reported the construction of heterocycles including 4-methylpyrrolo[1,2-*a*]quinoxalines and 4-methylquinazolines using the strategy of C—H bond activation. Remarkably, one of the vinyl carbon within vinylene carbonate was used for forming heterocycle backbone, and the other one left for methyl group (Schemes 1c~1d).^[10] In fact, there are only two examples with respect to the vinylene carbonate as a C1 synthon for heterocycle synthesis. Therefore, exploring the versatile application of vinylene carbonate in organic synthesis is highly appealing.

We reasoned that the works listed above, along with our recent interests in the functionalization of azobenzene,^[11] would serve as a basis for establishing a mild protocol to realize the *N*-heterocycle synthesis starting from azobenzenes and commercially available vinylene carbonate (Scheme 1e). In this work, we will disclose an efficient Rh(III)-catalyzed annulation of vinylene carbonate with azobenzenes, which can afford a series of 2*H*-indazole-3-carbaldehydes that cannot be achieved through traditional synthetic method.^[12] To the best of our knowledge, the new reaction model of vinylene carbonate has never been reported, in which vinylene carbonate provides one vinyl carbon for indazole backbone and leaves the other one for the formation of aldehyde (Scheme 1e).

2 Results and discussion

Initially, azobenzene **1a** and vinylene carbonate **2** were chosen as the model substrates to examine the feasibility of the Rh(III)-catalyzed annulation reaction and the results are listed in Table 1. With [Cp*RhCl₂]₂ as a catalyst and 2,2,2-trifluoroethanol (TFE) as a solvent, it is found that the annulation reaction did not proceed without a suitable additive (Entry 1). Fortunately, the annulation reaction occurred with 5 mol% [Cp*RhCl₂]₂ and 80 mol% of AgSbF₆ affording the desired 2-phenyl-2*H*-indazole-3-carbaldehyde (**3a**) in 60% isolated yield (Entry 2). Replacing the AgSbF₆ with AgTFA led to the formation of **3a** in decreasing yield (Entry 3). It was next found that the use of AgOAc as an additive could promote this annulation of azobenzene with vinylene carbonate under the above conditions (Entry 4). Subsequently, several common catalysts were evaluated on the annulation of **2** with **1a** in TFE. The result showed that only [Cp*Rh(MeCN)₃](SbF₆)₂ gave **3a** in 67% yield, and [Cp*CoCl₂]₂ or [Cp*IrCl₂]₂ did not work under the similar conditions (Entries 5~7). Further, It was found that the use of 80 mol% AgOAc under the above reaction conditions could generate **3a** in 71% yield (Entries 8~9). Subsequently, the reaction parameters, including the reaction time and reaction temperature, were carefully examined (Entries 10~13). The result showed that high temperature or longer reaction time led to diminished yields of **3a**. For this observation, it was found that the aldehyde group was prone to partial oxidation leading to diminished yield of product **3a**. Finally, the solvent effect was discussed (see Table S1 in supporting information). It was found that the reaction was

moderately successful in 1,2-dichloroethane (DCE) with **3a** isolated in 67% yield. Some other solvents failed to improve the annulation reaction of **1a** with vinylene carbonate **2**.

Table 1 Optimization of reaction conditions^{a,b}

Entry	Catalyst	Additive	Yield ^b /%
1	[Cp*RhCl ₂] ₂	—	n.r.
2	[Cp*RhCl ₂] ₂	AgSbF ₆	60
3	[Cp*RhCl ₂] ₂	AgTFA	40
4	[Cp*RhCl ₂] ₂	AgOAc	72
5	[Cp*Rh(MeCN) ₃](SbF ₆) ₂	AgOAc	67
6	[Cp*CoCl ₂] ₂	AgOAc	n.r.
7	[Cp*IrCl ₂] ₂	AgOAc	n.r.
8	[Cp*RhCl ₂] ₂	AgOAc	66 ^c
9	[Cp*RhCl ₂] ₂	AgOAc	71 ^d
10	[Cp*RhCl ₂] ₂	AgOAc	68 ^e
11	[Cp*RhCl ₂] ₂	AgOAc	53 ^f
12	[Cp*RhCl ₂] ₂	AgOAc	56 ^g
13	[Cp*RhCl ₂] ₂	AgOAc	62 ^h

^a Reaction conditions: azobenzene (**1a**, 0.20 mmol), vinylene carbonate (**2**, 0.40 mmol), catalyst (5 mol%), additive (80 mol%), TFE (2.0 mL) at 60 °C in air for 8 h. ^b Isolated yield. ^c AgOAc (50 mol%). ^d AgOAc (100 mol%). ^e 40 °C. ^f 80 °C. ^g 5 h. ^h 10 h. Cp* = pentamethylcyclopentadienyl. n.r. = no reaction.

With the optimized conditions in hand, the generality of the Rh(III)-catalyzed annulation reaction with a number of diversely substituted azobenzenes was then tested. The result is listed in Table 2. Firstly, some symmetric azobenzenes bearing *para*-substituted electron-donating groups (Me, Et, *i*-Pr and *t*-Bu) delivered the desired 2*H*-indazole products in moderate yields (**3b** ~ **3e**). Introduction of electron-withdrawing groups, such as F and ester, generated the regarding products in improved yields (**3f** ~ **3g**). The reactions of *meta*-substituted azobenzenes with vinylene carbonate afforded the desired products in 70% ~ 79% yields (**3h** ~ **3j**). Unsurprisingly, the sterically hindered

(*E*)-1,2-di-*o*-tolylidiazene reacted with vinylene carbonate smoothly to form the **3k** in 45% yield. The azobenzenes with disubstitution also allowed for the formation of indazoles in moderate yields (**3l** ~ **3m**). Moreover, some unsymmetrical azobenzenes were specifically synthesized and used to test their reactions under standard conditions, in which high regioselectivity was observed. The use of azobenzenes containing Cl and Br delivered the according products in 74% (**3n**) and 80% (**3o**) yields, respectively. However, 4-iodo substituted azo substrate generated **3p** as a single product in decreasing yield. Instead of halogens with phenyl group led to the formation of **3q** in good isolated yield. The more electron-withdrawing groups, such as ester and nitrile, were also well tolerated under the optimal conditions, generating the products **3r** ~ **3s** in 76 ~ 79% yield. Introduction of electron-rich group into azobenzene, such as CF₃O and MeO, produced the products **3t** and **3u** in acceptable yields. Next, the annulation of 3-bromo substituted phenyl azo with vinylene carbonate under above conditions were investigated, which gave **3v** in 69% yield. The replacement of Br with ester group delivered the target product **3w** in comparable yield. Finally, disubstituted aryl azo compounds allowed direct access to 2*H*-indazoles, furnishing the products (**3x** ~ **3y**) in acceptable yields.

The gram-scale cyclization of azobenzene **1a** with **2** can be performed under the above conditions, except for prolonging reaction time to 13 h, generating the desired **3a** in 62% yield (Scheme 2a). The late-stage transformations of **3a** were also achieved to afford the unsaturated alkenes by Wittig reaction and condensation of aldehyde with methyl 2-cyanoacetate, respectively (Schemes 2b ~ 2c).

Subsequently, several experiments were conducted to understand the reaction pathway of this Rh(III)-catalyzed annulation (Scheme 3). In absence of vinylene carbonate **2**, D₂O was added into the reaction system, and the result showed that 25% D was incorporated into the phenyl ring, which supported that the step of H/D scrambling was reversible (Scheme 3a). Furthermore, intramolecular and intermolecular kinetic isotope effect (KIE) experiment with **1a**/[D₁₀]-**1a** showed that the similar KIE value was determined by using ¹H NMR analysis, indicating that C—H bond activation maybe not involved in the rate-determining step (Schemes 3b ~ 3c).

(a) Gram-scale synthesis of **3a**

Typical transformations of **3a**

Scheme 2 Gram-scale synthesis of **3a** and typical transformations

Table 2 Scope of azobenzenes^{a,b}

^a Reaction conditions: azobenzene (**1**, 0.20 mmol), vinylene carbonate (**2**, 0.40 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (5 mol%), AgOAc (80 mol%), TFE (2.0 mL) at 60 °C in air for 8 h. ^b Isolated yield.

Based on the previous work and the above experiments,^[4,10–11] a possible reaction mechanism was proposed and listed in Scheme 4. Firstly, rhodacycle complex **I** was obtained through the coordination of Rh(III) center and azobenzene **1a**. Secondly, the insertion of the Rh—C bond within **I** into C=C double bond resulted in the formation of rhodacycle **II** or **II'**. It should be noted that the formation of intermediate **II'** would lead to a methylation process,^[10b] but which can be ruled out according to the result from Table 1. Subsequently, rhodacycle **II** underwent an intramolecular cyclization with the assistance of HX to generate the product **3a** and Rh(III) catalyst, along with releasing the CO₂ gas.

3 Conclusions

In summary, we have demonstrated that the annulation of azobenzenes with vinylene carbonate can be realized by

using Rh catalysis for the efficient construction of 2H-indazole backbone. The reaction is tolerant of a wide range of substituents and affords 2H-indazole-3-carbaldehydes in good yields. Further investigations into the role of vinylene carbonate in the Rh-catalyzed annulation, and application of the vinylene carbonate for more challenging transformations are currently underway in our laboratory.

4 Experimental section

4.1 General information

¹H NMR and ¹³C NMR spectra were recorded on 400, 600 MHz Bruker FT-NMR spectrometers (400 MHz and 100 MHz, 600 MHz and 150 MHz, respectively). All chemical shifts are given with reference to tetramethylsilane (TMS) as an internal standard. High resolution mass spectroscopy data of the products were collected on a Thermo Fisher Scientific LTQ FTICR-MS instrument.

Scheme 3 Mechanistic studies

Scheme 4 Possible reaction mechanism

The starting materials, such as symmetric azobenzenes and unsymmetric azobenzenes, are prepared according to the reported methods.^[14] Vinylene carbonate was purchased from commercial supplier Energy Chemical Company in China. Products were purified by flash chromatography on 200~300 mesh silica gels, eluting with petroleum ether/ethyl acetate ($V:V=5:1$ to $20:1$).

4.2 Synthesis of compounds **3a**~**3y**

General procedure for the gram-scale synthesis of **3a**: A

100 mL round flask equipped with a magnetic stirrer bar was charged with azobenzene (**1a**, 910.0 mg, 5 mmol), $[Cp^*RhCl_2]_2$ (154.5 mg, 0.25 mmol), AgOAc (667.6 mg, 4 mmol), and vinylene carbonate (**2a**, 860.5 mg, 10 mmol), then 2,2,2-trifluoroethanol (TFE, 40 mL) was injected into the tube. The reaction tube was placed in an oil bath and stirred at 60 °C for 12 h. After that, the mixture in reaction tube was monitored by thin-layer chromatography (TLC). The crude product was purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=20:1$), affording the product **3a** as a white solid (0.668 g, 62% yield).

The synthetic procedures of compounds **3b**~**3y** are the same as that of **3a**.

2-Phenyl-2H-indazole-3-carbaldehyde (3a):^[17] White solid, 32.0 mg, 72% yield. m.p. 73.1~74.6 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 10.12 (s, 1H), 8.32 (d, $J=8.4$ Hz, 1H), 7.92 (d, $J=8.4$ Hz, 1H), 7.67 (d, $J=7.0$ Hz, 2H), 7.64~7.59 (m, 3H), 7.51~7.47 (m, 1H), 7.45 (t, $J=7.5$ Hz, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 180.0, 148.6, 138.8, 132.2, 130.1, 129.6, 127.8, 127.1, 126.5, 123.4, 121.1, 118.7. HRMS (ESI) calcd for $C_{14}H_{11}N_2O$ $[M+H]^+$ 223.0866; found 223.0868.

5-Methyl-2-(p-tolyl)-2H-indazole-3-carbaldehyde (3b): White solid; 31.5 mg, 63% yield. m.p. 134.2~135.1 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 10.07 (s, 1H), 8.08 (s, 1H), 7.80 (d, $J=9.0$ Hz, 1H), 7.53 (d, $J=7.8$ Hz, 2H), 7.39 (d, $J=7.8$ Hz, 2H), 7.31 (d, $J=8.4$ Hz, 1H), 2.52 (s, 3H), 2.49 (s, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 180.3, 147.4, 140.3, 137.4, 136.4, 131.6, 130.5, 130.1, 126.2, 123.8, 119.4, 118.3, 22.1, 21.3. HRMS (ESI) calcd for $C_{16}H_{15}N_2O$ $[M+H]^+$ 251.1179; found 251.1175.

5-Ethyl-2-(4-ethylphenyl)-2H-indazole-3-carbaldehyde

(**3c**): White solid, 36.2 mg, 65% yield. m.p. 127.3 ~ 128.3 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.09 (s, 1H), 8.10 (s, 1H), 7.82 (d, *J*=9.0 Hz, 1H), 7.56 (d, *J*=8.4 Hz, 2H), 7.42 (d, *J*=7.8 Hz, 2H), 7.36~7.34 (m, 1H), 2.82 (q, *J*=7.4 Hz, 2H), 2.78 (q, *J*=7.4 Hz, 2H), 1.33 (t, *J*=7.8 Hz, 3H), 1.32 (t, *J*=7.8 Hz, 3H); ¹³C NMR (151 MHz, CDCl₃) δ: 180.3, 147.5, 146.5, 143.8, 136.5, 131.8, 129.6, 129.0, 126.3, 123.8, 118.4, 118.2, 29.5, 28.6, 15.6, 15.5. HRMS (ESI) calcd for C₁₈H₁₉N₂O [M + H]⁺ 279.1492; found 279.1490.

5-Isopropyl-2-(4-isopropylphenyl)-2*H*-indazole-3-carbaldehyde (**3d**): White solid, 41.6 mg, 68% yield. m.p. 113.1 ~ 114.2 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.10 (s, 1H), 8.12 (s, 1H), 7.84 (d, *J*=9.0 Hz, 1H), 7.57~7.55 (m, 2H), 7.45~7.43 (m, 2H), 7.40~7.39 (m, 1H), 3.11~3.02 (m, 2H), 1.35 (s, 3H), 1.34 (s, 3H), 1.33 (s, 3H), 1.32 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 180.4, 151.1, 148.3, 147.7, 136.6, 131.9, 128.3, 127.6, 126.3, 123.7, 118.5, 116.7, 34.7, 34.0, 23.91, 23.90. HRMS (ESI) calcd for C₂₀H₂₃N₂O [M + H]⁺ 307.1805; found 307.1804.

5-(*tert*-Butyl)-2-(4-(*tert*-butyl)phenyl)-2*H*-indazole-3-carbaldehyde (**3e**): White solid, 44.8 mg, 67% yield. m.p. 153.3 ~ 154.4 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.11 (s, 1H), 8.25 (s, 1H), 7.84 (d, *J*=9.0 Hz, 1H), 7.60 (t, *J*=7.2 Hz, 3H), 7.56 (d, *J*=7.8 Hz, 2H), 1.42 (s, 9H), 1.40 (s, 9H). ¹³C NMR (150 MHz, CDCl₃) δ: 180.5, 153.4, 150.5, 147.2, 136.3, 132.1, 127.3, 126.5, 126.0, 123.5, 118.1, 115.5, 35.4, 35.0, 31.3, 31.2. HRMS (ESI) calcd for C₂₂H₂₇N₂O [M + H]⁺ 335.2118; found 335.2118.

5-Fluoro-2-(4-fluorophenyl)-2*H*-indazole-3-carbaldehyde (**3f**): White solid, 36.1 mg, 70% yield. m.p. 83.1 ~ 84.7 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.06 (s, 1H), 7.90~7.88 (m, 2H), 7.66~7.64 (m, 2H), 7.32 (t, *J*=8.4 Hz, 2H), 7.30~7.27 (m, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 179.27, 163.3 (d, *J*=250.5), 161.8 (d, *J*=246.0), 145.8, 134.7, 132.7, 128.1 (d, *J*=9.3), 123.7 (d, *J*=12.9), 120.9 (d, *J*=9.9), 119.5 (d, *J*=28.5), 116.8 (d, *J*=23.3), 104.3 (d, *J*=25.4); ¹⁹F NMR (565 MHz, CDCl₃) δ: -109.41, -111.54. HRMS (ESI) calcd for C₁₄H₉F₂N₂O [M + H]⁺ 259.0677; found 259.0681.

Ethyl-2-(4-(ethoxycarbonyl)phenyl)-3-formyl-2*H*-indazole-5-carboxylate (**3g**): White solid, 59.3 mg, 81% yield. m.p. 191.1 ~ 192.3 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.18 (s, 1H), 9.04 (s, 1H), 8.31 (d, *J*=8.4 Hz, 2H), 8.12 (d, *J*=9.0 Hz, 1H), 7.93 (d, *J*=9.0 Hz, 1H), 7.78 (d, *J*=8.4 Hz, 2H), 4.45 (q, *J*=3.6 Hz, 4H), 1.45 (t, *J*=7.2 Hz, 3H), 1.44 (t, *J*=7.2 Hz, 3H); ¹³C NMR (151 MHz, CDCl₃) δ: 179.1, 166.2, 165.2, 149.9, 141.8, 133.3, 132.3, 131.0, 129.4, 128.1, 126.3, 124.6, 123.1, 118.7, 61.7, 61.4, 14.4, 14.3. HRMS (ESI) calcd for C₂₀H₁₉N₂O₅ [M + H]⁺ 367.1288; found 367.1289.

6-Methyl-2-(*m*-tolyl)-2*H*-indazole-3-carbaldehyde (**3h**): White solid, 39.5 mg, 79% yield. m.p. 149.8 ~ 150.1 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.07 (s, 1H), 8.17 (d, *J*=8.4 Hz, 1H), 7.64 (d, *J*=1.2 Hz, 1H), 7.48~7.45 (m, 2H), 7.42 (d, *J*=7.8 Hz, 1H), 7.37 (d, *J*=7.8 Hz, 1H), 7.27~7.26 (m, 1H), 2.51 (s, 3H), 2.48 (s, 3H); ¹³C NMR (150 MHz,

CDCl₃) δ: 180.2, 149.1, 139.9, 138.7, 137.8, 132.1, 130.7, 129.9, 129.2, 126.9, 123.6, 121.7, 120.6, 117.0, 22.1, 21.3. HRMS (ESI) calcd for C₁₆H₁₅N₂O [M + H]⁺ 251.1179; found 251.1174.

6-Ethyl-2-(3-ethylphenyl)-2*H*-indazole-3-carbaldehyde (**3i**): White solid, 38.9 mg, 70% yield. m.p. 128.3 ~ 128.9 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.07 (s, 1H), 8.20 (d, *J*=8.4 Hz, 1H), 7.67 (s, 1H), 7.50~7.48 (m, 2H), 7.44 (d, *J*=8.4 Hz, 1H), 7.40 (d, *J*=7.2 Hz, 1H), 7.30 (d, *J*=9.0 Hz, 1H), 2.81 (q, *J*=7.4 Hz, 2H), 2.77 (q, *J*=7.4 Hz, 2H), 1.33 (t, *J*=7.2 Hz, 3H), 1.30 (t, *J*=7.2 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 180.3, 149.1, 146.2, 144.3, 138.8, 132.1, 129.6, 129.4, 129.0, 125.8, 123.8, 121.9, 120.7, 115.7, 29.4, 28.7, 15.3. HRMS (ESI) calcd for C₁₈H₁₉N₂O [M + H]⁺ 279.1492; found 279.1489.

6-Chloro-2-(3-chlorophenyl)-2*H*-indazole-3-carbaldehyde (**3j**): White solid, 44.7 mg, 77% yield. m.p. 93.1 ~ 94.7 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.11 (s, 1H), 8.24 (d, *J*=9.0 Hz, 1H), 7.89 (s, 1H), 7.71 (s, 1H), 7.61~7.54 (m, 3H), 7.39 (d, *J*=8.4 Hz, 1H), 5.30 (s, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 179.4, 148.8, 139.4, 135.7, 134.2, 132.3, 130.6, 130.6, 128.8, 126.6, 124.5, 122.3, 121.9, 117.7. HRMS (ESI) calcd for C₁₄H₉N₂Cl₂O [M + H]⁺ 291.0086; found 291.0084.

7-Methyl-2-(*o*-tolyl)-2*H*-indazole-3-carbaldehyde (**3k**): yellow oil; 22.5 mg, 45% yield; ¹H NMR (600 MHz, CDCl₃) δ: 9.74 (s, 1H), 8.05 (d, *J*=8.4 Hz, 1H), 7.43 (t, *J*=7.8 Hz, 1H), 7.38~7.31 (m, 3H), 7.28 (t, *J*=7.8 Hz, 1H), 7.18 (d, *J*=9.0 Hz, 1H), 2.64 (s, 3H), 2.04 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 178.8, 147.8, 136.8, 134.5, 132.1, 130.3, 129.5, 128.1, 126.8, 126.4, 125.8, 125.8, 121.3, 117.2, 16.2, 15.9. HRMS (ESI) calcd for C₁₆H₁₅N₂O [M + H]⁺ 251.1179; found 251.1176.

2-(3,4-Dimethylphenyl)-5,6-dimethyl-2*H*-indazole-3-carbaldehyde (**3l**): White solid, 41.7 mg, 75% yield. m.p. 151.3 ~ 152.7 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.04 (s, 1H), 8.05 (s, 1H), 7.63 (s, 1H), 7.43 (d, *J*=1.2 Hz, 1H), 7.35~7.31 (m, 2H), 2.43 (s, 6H), 2.37 (d, *J*=3.0 Hz, 6H); ¹³C NMR (150 MHz, CDCl₃) δ: 180.4, 148.2, 138.7, 138.3, 138.0, 137.7, 136.6, 131.5, 130.4, 127.2, 123.7, 122.4, 119.7, 117.4, 21.0, 20.7, 19.8, 19.6. HRMS (ESI) calcd for C₁₈H₁₉N₂O [M + H]⁺ 279.1492; found 279.1497.

6-Bromo-2-(3-bromo-4-methylphenyl)-5-methyl-2*H*-indazole-3-carbaldehyde (**3m**): White solid, 59.2 mg, 73% yield. m.p. 173.5 ~ 174.3 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.07 (s, 1H), 8.17 (s, 1H), 8.14 (s, 1H), 7.87 (d, *J*=2.4 Hz, 1H), 7.50~7.48 (m, 1H), 7.46 (d, *J*=8.1 Hz, 1H), 2.57 (s, 3H), 2.53 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 179.6, 148.2, 140.6, 137.4, 137.1, 131.6, 131.3, 129.9, 126.2, 125.4, 125.0, 122.9, 121.7, 120.9, 23.9, 22.9. HRMS (ESI) calcd for C₁₆H₁₃Br₂N₂O [M + H]⁺ 406.9389; found 406.9393.

2-(4-Chlorophenyl)-2*H*-indazole-3-carbaldehyde (**3n**):^[18] White solid, 37.9 mg, 74% yield. m.p. 112.3 ~ 113.8 °C; ¹H NMR (600 MHz, CDCl₃) δ: 10.12 (s, 1H), 8.29~8.27 (m, 1H), 7.931~7.89 (m, 1H), 7.63~7.58 (m, 4H), 7.50~7.48 (m, 1H), 7.46~7.43 (m, 1H); ¹³C NMR

(150 MHz, CDCl_3) δ : 179.3, 148.7, 137.4, 136.3, 132.0, 129.8, 128.0, 127.6, 127.3, 123.8, 120.8, 118.7. HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{10}\text{ClN}_2\text{O}$ $[\text{M} + \text{H}]^+$ 257.0476; found 257.0478.

2-(4-Bromophenyl)-2*H*-indazole-3-carbaldehyde (**3o**): White solid, 48.0 mg, 80% yield. m.p. 123.6~124.1 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.12 (s, 1H), 8.28 (d, $J=7.8$ Hz, 1H), 7.90 (d, $J=8.4$ Hz, 1H), 7.75 (d, $J=9.0$ Hz, 2H), 7.55 (d, $J=8.4$ Hz, 2H), 7.50~7.43 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 179.2, 148.7, 137.9, 132.8, 132.0, 128.0, 127.8, 127.4, 124.3, 123.8, 120.8, 118.7. HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{10}\text{BrN}_2\text{O}$ $[\text{M} + \text{H}]^+$ 300.9971; found 300.9972.

2-(4-Iodophenyl)-2*H*-indazole-3-carbaldehyde (**3p**): White solid, 44.5 mg, 64% yield. m.p. 102.4~103.6 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.13 (s, 1H), 8.28 (d, $J=8.4$ Hz, 1H), 7.95 (d, $J=8.4$ Hz, 2H), 7.90 (d, $J=8.4$ Hz, 1H), 7.50~7.43 (m, 2H), 7.42 (d, $J=8.4$ Hz, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 179.3, 148.7, 138.8, 138.5, 131.9, 128.1, 127.9, 127.4, 123.8, 120.9, 118.7, 95.9. HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{10}\text{IN}_2\text{O}$ $[\text{M} + \text{H}]^+$ 348.9832; found 348.9830.

2-([1,1'-Biphenyl]-4-yl)-2*H*-indazole-3-carbaldehyde (**3q**): White solid, 49.5 mg, 83% yield. m.p. 146.1~147.6 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.20 (s, 1H), 8.33 (d, $J=8.3$ Hz, 1H), 7.93 (d, $J=8.4$ Hz, 1H), 7.84~7.82 (m, 2H), 7.75 (d, $J=8.4$ Hz, 2H), 7.69~7.66 (m, 2H), 7.53~7.49 (m, 3H), 7.47~7.42 (m, 2H); ^{13}C NMR (151 MHz, CDCl_3) δ : 179.96, 148.62, 143.18, 139.49, 137.83, 132.20, 129.08, 128.24, 128.21, 127.84, 127.29, 127.15, 126.72, 123.59, 121.06, 118.70. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{15}\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 299.1179; found 299.1180.

Ethyl 4-(3-formyl-2*H*-indazol-2-yl)benzoate (**3r**): White solid, 44.7 mg, 76% yield. m.p. 93.4~94.3 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.15 (s, 1H), 8.29 (d, $J=8.4$ Hz, 3H), 7.91 (d, $J=8.5$ Hz, 1H), 7.76 (d, $J=8.4$ Hz, 2H), 7.50~7.48 (m, 1H), 7.46~7.43 (m, 1H), 4.59 (q, $J=7.2$ Hz, 2H), 1.44 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 179.3, 165.3, 148.8, 142.1, 132.1, 131.9, 130.9, 128.2, 127.5, 126.3, 123.9, 121.0, 118.8, 61.7, 14.3. HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$ 295.1077; found 295.1075.

4-(3-Formyl-2*H*-indazol-2-yl)benzonitrile (**3s**): White solid, 38.0 mg, 77% yield. m.p. 100.6~101.3 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.20 (s, 1H), 8.29~8.22 (m, 1H), 7.96~7.88 (m, 3H), 7.87~7.80 (m, 2H), 7.56~7.44 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 178.3, 149.1, 142.4, 133.4, 131.7, 128.5, 127.8, 127.0, 124.7, 120.5, 118.9, 117.6, 113.9. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{10}\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 248.0818; found 248.0822.

2-(4-(Trifluoromethoxy)phenyl)-2*H*-indazole-3-carbaldehyde (**3t**): White solid, 37.3 mg, 61% yield. m.p. 126.6~127.9 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.15 (s, 1H), 8.29 (d, $J=8.4$ Hz, 1H), 7.91 (d, $J=9.0$ Hz, 1H), 7.74~7.72 (m, 2H), 7.50~7.46 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 179.1, 150.1, 148.7, 137.2, 132.1, 128.1, 127.9, 127.4, 123.8, 121.9, 120.8, 120.4 (d, $J=257.6$ Hz), 118.7. ^{19}F NMR (565 MHz, CDCl_3) δ : -57.85. HRMS (ESI)

calcd for $\text{C}_{15}\text{H}_{10}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 307.0689; found 307.0692.

2-(4-Methoxyphenyl)-2*H*-indazole-3-carbaldehyde (**3u**): White solid, 23.2 mg, 43% yield. m.p. 142.3~143.4 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.08 (s, 1H), 8.30 (d, $J=7.8$ Hz, 1H), 7.90 (d, $J=8.4$ Hz, 1H), 7.58 (d, $J=9.0$ Hz, 2H), 7.48 (t, $J=7.2$ Hz, 1H), 7.43 (d, $J=7.2$ Hz, 1H), 7.10 (d, $J=9.0$ Hz, 2H), 3.92 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 180.18, 160.79, 148.32, 132.23, 131.68, 127.64, 127.61, 126.95, 123.25, 120.98, 118.56, 114.67, 55.73. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{13}\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 253.0972; found 253.0974.

2-(3-Bromophenyl)-2*H*-indazole-3-carbaldehyde (**3v**): White solid, 41.4 mg, 69% yield. m.p. 103.5~105.1 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.14 (s, 1H), 8.29 (d, $J=8.4$ Hz, 1H), 7.91 (d, $J=9.0$ Hz, 1H), 7.88 (s, 1H), 7.74 (d, $J=8.4$ Hz, 1H), 7.61 (d, $J=7.8$ Hz, 1H), 7.53~7.44 (m, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 179.4, 148.7, 139.9, 133.2, 132.1, 130.7, 129.5, 128.1, 127.5, 125.1, 123.7, 123.2, 120.9, 118.7. HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{10}\text{BrN}_2\text{O}$ $[\text{M} + \text{H}]^+$ 300.9971; found 300.9969.

Methyl 3-(3-formyl-2*H*-indazol-2-yl)benzoate (**3w**): White solid, 39.8 mg, 71% yield. m.p. 91.3~92.6 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.14 (s, 1H), 8.34 (t, $J=1.8$ Hz, 1H), 8.31 (d, $J=8.4$ Hz, 1H), 8.27 (d, $J=7.8$ Hz, 1H), 7.92 (d, $J=8.4$ Hz, 1H), 7.89~7.88 (m, 1H), 7.72 (t, $J=7.8$ Hz, 1H), 7.52~7.45 (m, 2H), 3.97 (s, 3H); ^{13}C NMR (151 MHz, CDCl_3) δ : 179.3, 165.5, 148.7, 139.1, 132.1, 131.8, 131.0, 130.5, 129.8, 128.1, 127.4, 123.8, 120.9, 118.8, 52.7. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{13}\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$ 281.0921; found 281.0921.

2-(3,5-Dimethylphenyl)-2*H*-indazole-3-carbaldehyde (**3x**): White solid, 30.0 mg, 60% yield. m.p. 165.4~166.5 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.11 (s, 1H), 8.31 (d, $J=8.4$ Hz, 1H), 7.91 (d, $J=8.4$ Hz, 1H), 7.49~7.42 (m, 2H), 7.28 (s, 2H), 7.22 (s, 1H), 2.44 (s, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 180.5, 148.4, 139.7, 138.5, 132.2, 131.7, 127.7, 127.0, 124.2, 123.2, 121.2, 118.6, 21.3. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 251.1179; found 251.1183.

2-(3,5-Dichlorophenyl)-2*H*-indazole-3-carbaldehyde (**3y**): White solid, 43.5 mg, 75% yield. m.p. 82.1~83.5 °C; ^1H NMR (600 MHz, CDCl_3) δ : 10.18 (s, 1H), 8.26 (d, $J=8.4$ Hz, 1H), 7.89 (d, $J=8.4$ Hz, 1H), 7.61 (s, 2H), 7.59 (s, 1H), 7.50 (t, $J=7.8$ Hz, 1H), 7.46 (t, $J=7.8$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 178.7, 148.8, 140.4, 136.0, 131.9, 130.2, 128.4, 127.7, 125.0, 124.1, 120.7, 118.8. HRMS (ESI) calcd for $\text{C}_{14}\text{H}_9\text{Cl}_2\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 291.0086; found 291.0090.

4.3 General procedure for the transformation of **3a** into **4**

Compound **3a** (44.4 mg, 0.20 mmol) was dissolved in 1,4-dioxane (2 mL) in a 10 mL reaction tube with a magnetic stirrer bar. Then methyltriphenylphosphonium bromide (71.4 mg, 0.2 mmol) and NaOH (0.3 g) were added the mixture in an oil bath and stirred at 70 °C for 5 h.^[15] After

cooling to room temperature, the reaction mixture was extracted with CH_2Cl_2 repeatedly. The organic phases were combined. The solvent was evaporated and the residue was purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=20:1$), and afforded 2-phenyl-3-vinyl-2*H*-indazole (**4**).^[19] Yellow oil, 37.9 mg, 86% yield. ^1H NMR (600 MHz, CDCl_3) δ : 7.92 (d, $J=8.4$ Hz, 1H), 7.78 (d, $J=9.0$ Hz, 1H), 7.59 (d, $J=7.8$ Hz, 2H), 7.55 (t, $J=7.8$ Hz, 2H), 7.50 (d, $J=7.2$ Hz, 1H), 7.39~7.34 (m, 1H), 7.19 (t, $J=7.8$ Hz, 1H), 6.84~6.79 (m, 1H), 6.03 (d, $J=18.0$ Hz, 1H), 5.53 (d, $J=11.4$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 148.9, 139.7, 133.2, 129.1, 128.9, 126.8, 126.3, 125.0, 122.8, 120.6, 120.4, 118.0, 117.7.

4.4 General procedure for the transformation of **3a** into **5**

The compound **3a** (39.4 mg, 0.20 mmol), methyl 2-cyanoacetate (19.8 mg, 0.20 mol), NaOH (9.6 mg, 0.24 mol) and MeOH (1 mL) were added into a reaction tube (10 mL) at room temperature.^[16] The resulted mixture was stirred for 3 h. After that, the mixture was quenched with water and extracted with CH_2Cl_2 repeatedly. The solvent was evaporated and the residue was further purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=5:1$), affording methyl 2-cyano-3-(2-phenyl-2*H*-indazol-3-yl)acrylate (**5**). Yellow oil, 54.6 mg, 90% yield. ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ : 8.18~8.16 (m, 2H), 7.95 (d, $J=8.4$ Hz, 1H), 7.74 (d, $J=7.2$ Hz, 2H), 7.69~7.67 (m, 3H), 7.54 (t, $J=7.2$ Hz, 1H), 7.46 (t, $J=7.8$ Hz, 1H), 3.85 (s, 3H); ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$) δ : 162.7, 149.0, 141.7, 139.1, 130.6, 130.2, 128.3, 128.0, 126.9, 125.6, 122.3, 122.2, 119.3, 116.3, 103.1, 54.0. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{N}_3\text{O}_2$ $[\text{M}+\text{H}]^+$ 304.1081; found 304.1086.

4.5 General procedure for the H/D exchange experiment

A reaction tube (10 mL) equipped with a magnetic stirrer bar was charged with azobenzene (**1a**, 36.4 mg, 0.20 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (6.2 mg, 0.01 mmol), AgOAc (26.7 mg, 0.16 mmol), and D_2O (0.1 mL), then freshly distilled TFE (2 mL) was injected into the reaction tube. The reaction tube was placed in an oil bath and stirred at 60 °C for 1 h. After that, the mixture was quenched with water and extracted with CH_2Cl_2 repeatedly. The obtained solution was concentrated under vacuum to yield the crude product, which was further purified by flash chromatography (silica gel; petroleum ether/ethyl acetate, $V:V=30:1$), affording a mixture of **1a**/ $[\text{D}_2]$ -**1a** ($\text{H/D}=3:1$). Then, 25% of $[\text{D}_2]$ -**1a** in the mixture was determined by ^1H NMR analysis. (Figure S1)

4.6 General procedure for the KIE experiments

(a) To an oven-dried reaction tube (10 mL) equipped with a magnetic stirrer bar was charged with deuterated azobenzene ($[\text{D}_5]$ -**1a**, 37.4 mg, 0.20 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (6.2 mg, 0.01 mmol), AgOAc (26.7 mg, 0.16 mmol), and vinylene carbonate (**2a**, 34.4 mg, 0.40 mmol), then 2,2,2-trifluoroethanol (TFE, 2 mL) was injected into the

tube. The reaction tube was placed in an oil bath and stirred at 60 °C for 4 h. After that, the mixture in reaction tube was detected by TLC. The crude product was purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=20:1$), affording the products $[\text{D}_4]$ -**3a** and $[\text{D}_5]$ -**3a** (15.8 mg, ca. 35% yield). The KIE value ($\text{KIE}=1.63$) was determined by ^1H NMR analysis (Figure S2).

(b) To an oven-dried reaction tube (10 mL) equipped with a magnetic stirrer bar was charged with deuterated azobenzene ($[\text{D}_{10}]$ -**1a**, 19.2 mg, 0.10 mmol), azobenzene (**1a**, 18.2 mg, 0.10 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (6.2 mg, 0.01 mmol), AgOAc (26.7 mg, 0.16 mmol), and vinylene carbonate (**2a**, 34.4 mg, 0.40 mmol), then 2,2,2-trifluoroethanol (TFE, 2 mL) was injected into the tube. The reaction tube was placed in an oil bath and stirred at 60 °C for 4 h. After that, the mixture in reaction tube was detected by TLC. The crude product was purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=20:1$), affording the products **3a** and $[\text{D}_9]$ -**3a** (18.6 mg, ca. 41% yield). The KIE value ($\text{KIE}=1.56$) was determined by ^1H NMR analysis (Figure S3).

Supporting Information ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra of new compounds **3**~**5**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References and note

- [1] (a) Dong, J.; Zhang, Q.; Wang, Z.; Huang, G.; Li, S. *ChemMedChem* **2018**, *13*, 1490.
(b) Ghosh, S.; Mondal, S.; Hajra, A. *Adv. Synth. Catal.* **2020**, *21*, 37684.
- [2] (a) Angelis, M. D.; Stossi, F.; Carlson, K. A.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. *J. Med. Chem.* **2005**, *48*, 1132.
(b) Aman, W.; Lee, J.; Kim, M.; Yang, S.; Jung, H.; Hah, J.-M. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 1188.
- [3] (a) Yong, W.-S.; Park, S.; Yun, H.; Lee, P. H. *Adv. Synth. Catal.* **2016**, *358*, 1958.
(b) Wu, C.; Fang, Y.; Larock, R. C.; Shi, F. *Org. Lett.* **2010**, *12*, 2234.
(c) Vidyacharan, S.; Ramanjaneyulu, B. T.; Jang, S.; Kim, D.-P. *ChemSusChem* **2019**, *12*, 2581.
(d) Kumar, M. R.; Park, A.; Park, N.; Lee, S. W. *Org. Lett.* **2011**, *13*, 3542.
(e) Wang, C.-D.; Liu, R.-S. *Org. Biomol. Chem.* **2012**, *10*, 8948.
(f) Geng, X.; Wang, C. *Org. Lett.* **2015**, *17*, 2434.
(g) Long, Z.; Wang, Z.; Zhou, D.; Wan, D.; You, J. *Org. Lett.* **2017**, *19*, 2777.
(h) Li, X.; Ye, X.; Wei, C.; Shan, C.; Wojtas, L.; Wang, Q.; Shi, X. *Org. Lett.* **2020**, *22*, 4151.
- [4] (a) Li, H.; Li, P.; Wang, L. *Org. Lett.* **2013**, *15*, 620.
(b) Lian, Y.; Bergman, R. G.; Lavis, L. D.; Ellman, J. A. *J. Am. Chem. Soc.* **2013**, *135*, 7122.
(c) Hummel, J. R.; Ellman, J. A. *J. Am. Chem. Soc.* **2015**, *137*, 490.
(d) Chen, X.; Wang, Y.; Wang, S.; Kong, D.; Wen, L.; Zhai, R.; Zhao, K.; Bai, L.; Li, Y. *Chin. J. Org. Chem.* **2020**, *40*, 688 (in Chinese).
(陈训, 王颖, 王烁今, 孔杜林, 文丽君, 翟锐锐, 赵珂, 白丽丽, 李友宾, 有机化学, **2020**, *40*, 688.)
- [5] (a) Zhao, D.; Vásquez-Céspedes, S.; Glorius, F. *Angew. Chem., Int. Ed.* **2015**, *54*, 1657.
(b) Wang, L.; Shao, Y.; Chen, F.; Qian, P.-C.; Cheng, J. *Chin. J.*

- Org. Chem.* **2021**, doi: 10.6023/cjoc202106023.
- [6] (a) Jeong, T.; Han, S. H.; Han, S.; Sharma, S.; Park, J.; Lee, J. S.; Kwak, J. H.; Jung, Y. H.; Kim, I. S. *Org. Lett.* **2016**, *18*, 232.
 (b) Long, Z.; Yang, Y.; You, J. *Org. Lett.* **2017**, *19*, 2781.
 (c) Cai, S.; Lin, S.; Yi, X.; Xi, C. *J. Org. Chem.* **2017**, *82*, 512.
 (d) Wang, N.; Liu, L.; Xu, W.; Zhang, M.; Huang, Z.; Shi, D.; Zhao, Y. *Org. Lett.* **2019**, *21*, 365.
 (e) Jiang, C.; Chen, S.; Gong, J.; Yang Z. *Acta Chim. Sinica* **2020**, *78*, 928 (in Chinese).
 (江崇国, 陈斯嘉, 龚建贤, 杨震, 化学学报, **2020**, *78*, 928.)
- [7] (a) Hilf, S.; Grubbs, R. H.; Kilbinger, A. F. M. *J. Am. Chem. Soc.* **2008**, *130*, 11040.
 (b) Kim, K. H.; Park, B. R.; Lim, J. W.; Kim, J. N. *Tetrahedron Lett.* **2011**, *52*, 3463.
 (c) Wang, Z.; Xue, F.; Hayashi, T. *Angew. Chem., Int. Ed.* **2019**, *58*, 11054.
 [8] Hara, H.; Hirano, M.; Tanaka, K. *Org. Lett.* **2009**, *11*, 1337.
 [9] (a) Ghosh, K.; Nishii, Y.; Miura, M. *ACS Catal.* **2019**, *9*, 11455.
 (b) Li, X.; Huang, T.; Song, Y.; Qi, Y.; Li, L.; Li, Y.; Xiao, Q.; Zhang, Y. *Org. Lett.* **2020**, *22*, 5925.
 (c) Wang, Z.-H.; Wang, H. Li, L.; Zhou, M.-D. *Org. Lett.* **2021**, *23*, 995.
 (d) Ghosh, K.; Nishii, Y.; Miura, M. *Org. Lett.* **2020**, *22*, 3547.
 (e) Mihara, G.; Ghosh, K.; Nishii, Y.; Miura, M. *Org. Lett.* **2020**, *22*, 5706.
 [10] (a) Nan, J.; Ma, Q.; Yin, J.; Liang, C.; Tian, L.; Ma, Y. *Org. Chem. Front.* **2021**, *8*, 1764.
 (b) Wang, C.; Fan, X.; Chen, F.; Qian, P.-C.; Cheng, J. *Chem. Commun.* **2021**, *57*, 3929.
 [11] (a) Li, H.; Li, P.; Zhao, Q.; Wang, L. *Chem. Commun.* **2013**, *49*, 9170.
 (b) Li, H.; Li, P.; Tan, H.; Wang L. *Chem.-Eur. J.* **2013**, *19*, 14432.
 (c) Li, H.; Xie, X.; Wang L. *Chem. Commun.*, **2014**, *50*, 4218.
 (d) Deng, H.; Li, H.; Wang, L. *Org. Lett.* **2015**, *17*, 2450.
 (e) Tan, H.; Li, H.; Ji, W.; Wang, L. *Angew. Chem., Int. Ed.*, **2015**, *54*, 8374.
 (f) Yang, W.; Li, P.; Wang, L. *Chem. Commun.*, **2015**, *51*, 7520.
 (g) Deng, H.; Li, H.; Wang, L. *Org. Lett.* **2016**, *18*, 3110.
 (h) Zhang, W.; Deng, H.; Li, H. *Org. Chem. Front.*, **2017**, *4*, 2202.
 (i) Zhang, W.; Bu, J.; Wang, L.; Li, P.; Li H. *Org. Chem. Front.* **2021**, *8*, 5054.
 [12] Shirtcliff, L. D.; Weakley, T. J. R.; Haley, M. M.; Köhler, F.; Herges, R. *J. Org. Chem.* **2004**, *69*, 6979.
 [13] During our preparation of this manuscript, Kim's group reported a Rh(III)-catalyzed [4+1]/[4+2] annulation of azobenzenes with vinyl carbonate in DCE, which affords a series of indazoles and dihydrocinnolinones in good yields. In this work, only the azobenzene bearing electro-donating groups can produce the 2*H*-indazole bearing a hydro methyl group. However, our manuscript first discloses an aldehyde formed from the vinyl carbonate, in which TFE plays a critical role for this chemical event. Particularly, the aldehyde can be easily transformed into other useful compounds. Therefore, a slightly different mechanism for this annulation was proposed herein. For the work reported by Kim, see: Park, M. S.; Moon, K.; Oh, H.; Lee, J. Y.; Ghosh, P.; Kang, J. Y.; Park, J. S.; Mishra, N. K.; Kim, I. S. *Org. Lett.* **2021**, *23*, 5081.
 [14] (a) Zhang, C.; Jiao, N. *Angew. Chem., Int. Ed.* **2010**, *49*, 6174.
 (b) Wang, Z.-C.; Yin, Z.-P.; Zhu, F.-X.; Li, Y.-H.; Wu, X.-F. *ChemCatChem* **2017**, *9*, 3637.
 [15] Delmas, M.; Bigot, Y. L.; Gaset, A.; Gorrichon, J. P. *Synth. Commun.* **1981**, *11*, 125.
 [16] Ranjbar-Karimi, R.; Beiki-Shoraki, K.; Amiri, A. *Monatsh. Chem.* **2010**, *141*, 1101.
 [17] Shirtcliff, L. D.; Weakley, T. J. R.; Haley, M. M.; Köhler, F.; Herges, R. *J. Org. Chem.* **2004**, *69*, 6979.
 [18] (a) Guo, S.; Sun, L.; Li, X.; Zhang, X.; Fan, X. *Adv. Synth. Catal.* **2020**, *362*, 913.
 (b) Bhattacharjee, S.; Laru, S.; Ghosh, P.; Hajra, A. *J. Org. Chem.* **2021**, *86*, 10866.
 [19] Wu, C.; Fang, Y.; Larock, R. C.; Shi, F. *Org. Lett.* **2010**, *12*, 2234.

(Fan, Y.)