有机催化邻羟基官能化的对甲亚基苯醌和 β -氧代酰基吡唑的不对称 [4+2] 环化合成光学活性 3,4-二氢香豆素衍生物

赵敏^a 李霏^a 程益政^b 王有名^a 周正洪^{*,a}

(^a 南开大学化学学院 元素有机化学国家重点实验室 天津 300071)

(^b 南开大学化学学院化学系 天津 300071)

摘要 发展了一种有机催化的邻羟基苯基取代对甲亚基苯醌和 β -氧代酰基吡唑之间的不对称 [4+2] 环化反应, 为光学活性 3,4-二氢香豆素衍生物的获取提供了有效途径。在衍生自奎宁的手性双功能方酰胺催化下, 一系列邻羟基苯基取代对甲亚基苯醌和 β -氧代酰基吡唑可以在温和的条件下顺利地发生反应, 以高产率和优秀的非对映及对映选择性得到相应的多取代 3,4-二氢香豆素衍生物。

关键词 不对称合成; 3,4-二氢香豆素; 邻羟基苯基取代对甲亚基苯醌; β -氧代酰基吡唑; 有机催化

Optically Active 3,4-Dihydrocoumarins via Organocatalyzed Asymmetric [4+2] Annulation of *ortho*-Hydroxyl Functionalized *p*-Quinone Methides with β -Keto Acylpyrazoles

Zhao, Min^a Li, Fei^a Cheng, Yizheng^b Wang, Youming^a Zhou, Zhenghong^{*,a}

(^a Institute and State Key Laboratory of Elemento-organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071)

(^b Department of Chemistry, College of Chemistry, Nankai University, Tianjin 300071)

Abstract An organocatalyzed asymmetric formal [4+2] annulation of *ortho*-hydroxyl functionalized *p*-quinone methides with β -keto acylpyrazoles has been developed, which provides an efficient approach to access optically active 3,4-dihydrocoumarins. Under the catalysis of a quinine-derived bifunctional squaramide, the reactions of a wide range of *ortho*-hydroxyphenyl substituted *p*-quinone methides and β -keto acylpyrazoles took place smoothly to generate the corresponding polysubstituted 3,4-dihydrocoumarins in good yields with high levels of diastereo- and enantio-selectivities (up to 96% yield, >19/1 *dr* and 98% *ee*) under mild reaction conditions.

Keywords asymmetric synthesis; 3,4-dihydrocoumarin; *ortho*-hydroxyphenyl-substituted *para*-quinone methide; β -keto acylpyrazole; organocatalysis

1 Introduction

3,4-Dihydrocoumarins, representing an important class of oxygen-containing heterocycles, are privileged structural motifs commonly present in a broad array of natural products and synthetic bioactive substances.^[1] Compounds incorporating a dihydrocoumarin core structure not only exhibit a wide range of biological and pharmacological activities,^[2] but also serve as useful building blocks in the synthesis of some other important compounds.^[3] Consequently, the development of effective protocols to con-

struct these architectures in a catalytic asymmetric fashion has attracted growing interest for synthetic chemists, and many synthetic approaches have been disclosed over the years.^[4] Among the diverse approaches, a very viable and reliable practice for the synthesis of optically active dihydrocoumarins is the organocatalyzed [4+2] annulation of *o*-quinone methides (*o*-QMs) with different two-atom synthons.^[5]

Due to the zwitterionic resonance structure and the possibility of aromatization upon conjugate addition, *para*-quinone methides (*p*-QMs) have actively participated as

* Corresponding author. E-mail: z.h.zhou@nankai.edu.cn

Received May 31, 2021; revised June 22, 2021; published online July 5, 2021.

Dedicated to the 100th anniversary of Chemistry at Nankai University.

1,6-acceptors in enantioselective synthesis.^[6] Moreover, it is noteworthy that the installation of an *ortho* hydroxyl group on the phenyl ring converts *p*-QMs into a novel type of donor-Michael acceptor substrates, which could be applied in organocatalyzed asymmetric cascade cyclizations for the construction of diverse oxygen-containing heterocyclic skeletons. In 2016, Enders described a pioneering work of asymmetric [4+2] cycloaddition of *ortho*-hydroxyphenyl-substituted *p*-QMs with isatin-derived enates to access chiral chroman derivatives.^[7] Thereafter, *ortho*-hydroxyphenyl substituted *p*-QMs have been extensively studied because of their high reactivity profile as four-atom partner for the catalytic asymmetric cycloaddition reactions, including [4+1]^[8], [4+2]^[7,9] and [4+3] cyclizations.^[10] Despite the impressive achievements in organocatalyzed asymmetric [4+2] annulation base on *ortho*-hydroxyphenyl substituted *p*-QMs, surprisingly, applications of *ortho*-hydroxyphenyl substituted *p*-QMs in stereo-controlled construction of 3,4-dihydrocoumarin scaffold have been rarely explored.^[9b,9d,9e] Therefore, the development of a new strategy for the tandem reaction of *p*-QMs access to enantiomerically enriched 3,4-dihydrocoumarins is still a challenging and meaningful work.

Recently, we^[5] have demonstrated β -keto acylpyrazoles as an efficient two atom building block for the construction of dihydrocoumarin framework upon treatment with *o*-QMs. However, the use of stabilized *o*-QM substrate to ensure satisfactory enantioselectivity makes it impossible to introduction of different substituents on the phenyl ring. We envisioned that this disadvantage could be overcome if we change the reaction partner from *o*-QM to *ortho*-hydroxyphenyl substituted *p*-QMs. The envisaged process has a specific challenge that need to be overcome, namely, control over enantioselectivity at a remote δ -position which is far from the catalyst binding site. As a part of our ongoing research on the development of approaches to biorelevant heterocycles,^[11] we herein report the organocatalyzed [4+2] cyclization of *ortho*-hydroxyphenyl substituted *p*-QMs and β -keto acylpyrazoles, which provide a new access to 3,4-dihydrocoumarins in high levels of diastereo- and enantio-selectivity.

2 Results and discussion

To test the feasibility of our design, *p*-quinone methide (**1a**) and 1-phenyl-3-(1*H*-pyrazol-1-yl)propane-1,3-dione (**2a**) were initially selected as the model substrates to evaluate the reactivity of this transformation (Table 1). The model reaction in dichloromethane at 25 °C proceeded smoothly by using 10 mol% thiourea **I** (Figure 1) as a chiral bifunctional Lewis base catalyst, providing the desired [4+2] annulation product (3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3aa**) in 81% yield with >19/1 *dr* and 77% *ee* (Entry 1). Other thioureas, such as (1*R*,2*R*)-1,2-diphenylethane-1,2-diamine derived thioureas **IIa**, **IIb** and cinchona alkaloid-based thiourea **III** (Figure 1) also proved to be efficient for this reac-

tion, delivering the desired cyclization product **3aa** with comparable or better stereochemical outcomes as thiourea **I** (Entries 2~4 vs. Entry 1). To further improve the enantioselectivity of this reaction, squaramide catalysts **III**~**VII** bearing different chiral diamine backbone and *N*-substituent were evaluated. Generally, slightly better performances were observed for squaramide catalysts (Entries 5~10). Among the evaluated squaramide catalyst, compound **VII** gave the best result (99% yield, >19/1 *dr*, 94% *ee*, Entry 10).

Table 1 Catalyst screening^a

Entry	Catalyst	Time/h	Yield ^b /%	<i>Dr</i> ^c	<i>ee</i> ^d /%
1	I	8	81	>19/1	77
2	IIa	12	89	>19/1	67
3	IIb	12	85	>19/1	70
4	III	8	90	>19/1	−89
5	IV	8	80	>19/1	80
6	Va	12	84	>19/1	78
7	Vb	12	83	>19/1	76
8	VIa	8	89	>19/1	−77
9	VIb	8	86	>19/1	−79
10	VIc	8	79	>19/1	−83
11	VII	8	90	>19/1	−94

^a Unless otherwise specified, all reactions were carried out with **1a** (0.20 mmol), **2a** (0.10 mmol) in the presence of 10 mol% catalyst in solvent (1 mL) at 25 °C. ^b Isolated yield. ^c Determined by ¹H NMR analysis. ^d Determined by HPLC analysis using a chiral stationary phase.

Having identified chiral bifunctional squaramide **VII** as the optimum catalyst for the reaction, other factors, such as solvent, catalyst loading, reaction temperature, as well as substrate concentration, influencing the reaction were thoroughly investigated employing the reaction between 1-phenyl-3-(1*H*-pyrazol-1-yl)propane-1,3-dione (**1a**) with *o*-quinone methide (**2a**) as the model. The results are listed in Table 2.

Solvent evaluation revealed that the solvent has a dramatic effect on the outcome of the reaction. Generally, better enantioselectivity was observed in chloroalkane solvents such as dichloromethane (Entry 1, 94% *ee*), chloroform (Entry 2, 88% *ee*), 1,2-dichloroethane (Entry 3,

Figure 1 Catalyst candidates

Table 2 Optimization of the reaction conditions^a

Entry	Solvent	Time/h	Yield ^b /%	<i>dr</i> ^c	<i>ee</i> ^d /%
1	CH ₂ Cl ₂	8	84	>19/1	94
2	CHCl ₃	8	80	>19/1	88
3	Cl(CH ₂) ₂ Cl	7	78	>19/1	90
4	PhCl	6	79	>19/1	84
5	PhF	6	75	>19/1	80
6	PhCH ₃	6	76	>19/1	76
7	EtOAc	48	45	>19/1	70
8	CH ₃ CN	48	43	>19/1	81
9 ^e	CH ₂ Cl ₂	36	80	>19/1	93
10 ^f	CH ₂ Cl ₂	8	85	>19/1	51
11 ^g	CH ₂ Cl ₂	5	80	>19/1	88
12 ^h	CH ₂ Cl ₂	12	83	>19/1	85

^a Unless otherwise specified, all reactions were carried out with **1a** (0.20 mmol), **2a** (0.10 mmol) in the presence of 10 mol% catalyst **VII** in solvent (1 mL) at 25 °C. ^b Isolated yield. ^c Determined by ¹H NMR analysis. ^d Determined by HPLC analysis using a chiral stationary phase. ^e The reaction was performed at 0 °C. ^f The reaction was carried out at 25 °C with a catalyst loading of 5 mol%. ^g The reaction was conducted with a substrate concentration of 0.2 mol/L. ^h The reaction took place with a substrate concentration of 0.05 mol/L.

90% *ee*) than other commonly used solvent, and dichloromethane was the best of choice for the reaction. It was found that there was a negligible temperature effect on the stereo-control of the reaction. When the reaction was car-

ried out at 0 °C, the corresponding annulation product **3aa** was obtained with almost the same *ee* value at the expense of reaction time (Entry 1 vs. Entry 9). When adjusting the loading of the catalyst **VII** from 10 mol% to 5 mol%, the enantioselectivity of the reaction decreased sharply (Entry 1 vs. Entry 10). Further studies revealed that substrate concentration had an important effect on the enantioselectivity. The best result was obtained when the reaction was carried out with a β -keto acylpyrazoles concentration of 1 mol/L in dichloromethane (Entry 1, 94% *ee*). An increase or decrease in the substrate concentration led to slightly lowered *ee* values (Entries 11 and 12, 88% *ee* and 85% *ee*, respectively).

With the optimum reaction conditions in hand (10 mol% catalyst **VII** in dichloromethane with a substrate concentration of 0.1 mol/L at 25 °C), we sought to investigate the generality of this asymmetric [4+2] annulation process. The results are collected in Table 3.

Firstly, the *ortho*-hydroxyphenyl substituted *para*-quinone methide (*p*-QM) component was evaluated. It was found that a broad range of *ortho*-hydroxyphenyl substituted *p*-QMs (**1**) could readily participate in this formal [4+2] annulation reaction. Generally, the reaction proceeded smoothly, and the desired annulation products **3** were obtained as single diastereomers in good yields with excellent enantioselectivities (86~98% *ee*). Both electron-withdrawing (Entries 7~11) and electron-donating substituents (Entries 2~6) on the phenyl group of **1** were readily tolerated. In terms of reactivity, *p*-QMs bearing electron-donating substituent were generally less reactive than those containing electron-neutral or electron-withdrawing substituent (Entries 2~4, 6 vs. Entries 1, 7~11). Moreover, the substituent position has a considerable effect on the reaction of substrates bearing chloro substituent. In contrast to 4-chloro-substituted *p*-QM (**1h**), for which excellent enantioselectivity was obtained, a remarkably decreased *ee* value was observed for 5-chloro-substituted one (**1i**) (Entry 8, 97% *ee* vs. Entry 9, 86% *ee*). Secondly, the generality of β -keto acylpyrazoles **2** was investigated by changing the substituents on the phenyl group (Entries 1 and 12~20). β -keto acylpyrazoles bea-

Table 3 Substrate scope of the organocatalyzed formal [4+2] annulations^a

Entry	3 (R, Ar)	Time/h	Yield ^b /%	dr ^c	ee ^d /%
1	3aa (H, Ph)	8	84	>19/1	94
2	3ba (4-Me, Ph)	48	80	>19/1	92
3	3ca (5-Me, Ph)	48	81	>19/1	97
4	3da (3-MeO, Ph)	32	80	>19/1	96
5	3ea (4-MeO, Ph)	8	85	>19/1	93
6	3fa (5-MeO, Ph)	32	84	>19/1	93
7	3ga (5-F, Ph)	5	86	>19/1	95
8	3ha (4-Cl, Ph)	8	82	>19/1	97
9	3ia (5-Cl, Ph)	8	82	>19/1	86
10	3ja (4-Br, Ph)	8	77	>19/1	98
11	3ka (5-Br, Ph)	8	68	>19/1	95
12	3ab (H, 4-MeC ₆ H ₄)	8	79	>19/1	92
13	3ac (H, 3-MeC ₆ H ₄)	8	78	>19/1	93
14	3ad (H, 2-MeC ₆ H ₄)	8	85	>19/1	90
15	3ae (H, 4-MeOC ₆ H ₄)	8	85	>19/1	91
16	3af (H, 3,4-(MeO) ₂ C ₆ H ₃)	8	66	>19/1	86
17	3ag (H, 4-FC ₆ H ₄)	8	83	>19/1	93
18	3ah (H, 3-ClC ₆ H ₄)	8	90	>19/1	96
19	3ai (H, 4-BrC ₆ H ₄)	8	81	>19/1	94
20	3aj (H, 3-BrC ₆ H ₄)	8	80	>19/1	95
21	3ak (H, 2-Naphthyl)	8	89	>19/1	87
22	3al (H, 2-Thienyl)	8	90	>19/1	93

^a Unless otherwise specified, all reactions were carried out with **1** (0.20 mmol), **2** (0.10 mmol) in the presence of 10 mol% catalyst **VII** in dichloromethane (1 mL) at room temperature. ^b Isolated yield. ^c Determined by ¹H NMR analysis. ^d Determined by HPLC analysis using a chiral stationary phase.

ring electron-neutral (Entry 1), electron-rich (Entries 12~16), or electron-poor substituents (Entries 17~20) could well participated in the asymmetric [4+2] cyclization to yield dihydrocoumarins **3** with overall high stereoselectivities (>19/1 *dr*, 86~96% *ee*) and generally good yields (66~90%) within 8 h. In detail, the substitution pattern clearly had some effect on the stereochemical outcome of the reaction. Compared with monomethoxy substituted substrate **2e**, a lower *ee* value was observed for dimethoxy substituted one (**2f**) (Entry 15 vs. Entry 16). In addition, by replacing the phenyl group of the β -keto acylpyrazole **2a** with 2-naphthalenyl, the reaction also ran smoothly to afford the annulation product **3ak** in good yield albeit with a somewhat decreased enantioselectivity (Entry 21). Electron-rich 2-thienyl substituted β -keto acylpyrazole **2l** was also a good reaction partner, delivering the desired annulation product **3al** in 90% yield with 93% *ee* (Entry 22).

The relative and absolute configuration of the product **3ai** is unequivocally established by X-ray analysis (Figure

2), and the remaining configurations are assumed by analogy.^[12]

Figure 2 X-ray crystal structure of (3*S*,4*S*)-**3ai**. Most of the hydrogen atoms have been omitted for clarity.

In order to demonstrate the practical utility of this newly developed transformation, we have performed a gram-scale cascade conjugate addition/lactonization reaction between **1a** and **2a** (Eq. 1). The desired 3,4-dihydrocoumarin **3aa** was obtained in 80% yield with a slightly improved stereochemical outcome albeit with a prolonged reaction time. Moreover, squaramide catalyst **VII** could be recovered with a recovery rate of 90% and reused without any loss of catalytic activity, the cyclization product **3aa** was obtained in 80% yield with >19/1 *dr* and 94% *ee*.

The 3,4-dihydrocoumarin products obtained in these formal [4+2] annulations would find more application in organic synthesis. For example, basic hydrolytic decarboxylation of 3,4-dihydrocoumarin **3aa** and subsequent methylation of the phenolic hydroxyl group afforded chiral β,β -diaryl substituted ketone **4**, an important type of intermediates for the synthesis of many pharmaceuticals and natural products, without appreciable loss in optical purity (Eq. 2). Moreover, according to a literature procedure,^[30] basic hydrolytic decarboxylation of 3,4-dihydrocoumarin

3aa and subsequent trifluoroborane promoted triethylsilane reduction delivered *cis*-2,4-disubstituted chroman **5** with a slightly erosion in optical purity (Eq. 3). Methylation of **3aa** with 1.5 equiv. of iodomethane preferentially occurred at the C(3) position to provide the corresponding C-methylation product **6** as a single diastereomer with maintaining the optical purity (Eq. 4).

At first glance, this cascade reaction seems to be triggered by the asymmetric 1,6-conjugate addition of β -keto acylpyrazoles to *p*-QMs. In this situation, it is difficult to realize remote stereocontrol since the binding and nucleophilic attack site is far apart. However, high levels of enantioselectivity were observed for all cases. To gain insight into the reaction mechanism, some control experiments were carried out. The masking of phenolic hydroxyl group in *p*-QM **1a** by methyl ether formation, replacement of hydroxyl group with a methyl group or changing the position of hydroxyl group from *ortho*- to *meta*-position all led

to no conversion of substrates, which indicate that triggering the reaction by 1,6-conjugate addition could be ruled out. Under the basic reaction condition, the acidic phenolic hydroxyl group could be deprotonated to generate a phenoxide ion intermediate with two resonance structures, **1a⁻** and **1a'⁻**. Protonation of the phenoxide ion **1a⁻** and **1a'⁻** resulted in the building of the tautomerization equilibrium between *p*-QM **1a** and *o*-QM **1a'** (Scheme 1). Theoretical calculations manifest that the energy barrier of tautomerization **1a** to **1a'** is 28.1 kJ/mol, indicating that the transformation of *p*-QM to *o*-QM was not difficult.^[9d] Therefore, *o*-QM **1a'** *in situ* generated by *p*-QM **1a** was the most possibly reaction partner of β -keto acylpyrazole **2a**, and the reaction was triggered by the Michael addition of β -keto acylpyrazole **2a** to *o*-QM **1a'**. According to this mechanism, the acidity strength of the phenolic hydroxyl group will be a key factor to influence the reactivity of *p*-QM substrates **1**, which means more acidic, faster tautomerization, thus more reactive. The observed phenomenon that *p*-QMs bearing electron-donating substituent were generally less reactive than those containing electron-neutral or electron-withdrawing substituent provides a solid support for this mechanism.

The proposed catalytic cycle is depicted in Figure 3. Firstly, *o*-QM **1a'** was formed via base-catalyzed tautomerization of *p*-QM **1a**, which is the rate-determining step of the whole transformation. Then, the deprotonation of 1-phenyl-3-(1*H*-pyrazol-1-yl)propane-1,3-dione (**2a**) forming a hydrogen-bond-paired enolate **A**, which nucleophilic attacks the *o*-QM (**1a'**) with the restoration of aromaticity as the driving force to generate a phenoxide ion intermediate **B**. Intramolecular lactonization furnishes the final [4+2] annulation product **3aa** with the 3*S*,4*S* absolute configuration as the predominant enantiomer and accompanied with the release of the pyrazole moiety.

3 Conclusions

In summary, we have disclosed a highly enantioselective formal [4+2] annulation of *ortho*-hydroxyphenyl-substituted *para*-quinone methides with β -keto acylpyrazoles by employing a quinine-based bifunctional squaramide as a chiral Lewis base catalyst. A series of optically pure and highly functionalized 3,4-dihydrocoumarin derivatives bearing two contiguous stereocenters were readily obtained with good isolated yields, excellent diastereo- and enantioselectivities under mild reaction conditions. The reaction is believed to be triggered by the Michael addition of the *in*

Scheme 1 Base catalyzed tautomerization between **1a** and **1a'**

Figure 3 Proposed catalytic cycle

situ formed *o*-QMs via the base-catalyzed tautomerization of the corresponding *p*-QMs. Moreover, the practicality and utility of this protocol were demonstrated by the scale-up synthesis and facile derivatization of the product.

4 Experimental section

4.1 Instruments and reagents

NMR spectra were obtained with a 400 MHz spectrometer (^1H 400 MHz, ^{13}C 100.6 MHz) in CDCl_3 . The chemical shifts are reported as δ values relative to tetramethylsilane. HRMS spectra were recorded with a Q-TOF mass spectrometer, equipped with an ESI source. Optical rotation values were measured with instruments operating at $\lambda=589$ nm, corresponding to the sodium D line at 20 °C. Enantiomeric excesses were determined by HPLC analysis with a chiral stationary phase. Materials were obtained from commercial suppliers and were used without further purification.

4.2 General procedure for squaramide VII-catalyzed asymmetric [4+2] annulation of *ortho*-hydroxyphenyl-substituted *para*-quinone methides and β -keto acylpyrazoles

A mixture of *ortho*-hydroxyphenyl-substituted *para*-quinone methide (**1**, 0.20 mmol), β -keto acylpyrazole (**2**, 0.10 mmol) and squaramide catalyst VII (6.3 mg, 0.010 mmol) in dichloromethane (1 mL) was stirred at 25 °C until the reaction reached completion (monitored by TLC). After the removal of the solvent under reduced pressure, the crude product was purified by chromatography on a silica gel column (200~300 mesh, PE/EtOAc, *V*:*V*=10:1~5:1) to afford the desired 3,4-dihydrocoumarin **3**. The title compounds were fully characterized by ^1H

NMR and ^{13}C NMR spectra, HRMS, and specific rotation data.

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3aa**): White solid, 38.3 mg, 84% yield. m.p. 118~121 °C; $[\alpha]_{\text{D}}^{20} +84.0$ (*c* 1.0, CH_3OH); >19/1 *dr*, 94% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (d, *J*=7.6 Hz, 2H), 7.58 (t, *J*=7.6 Hz, 1H), 7.45 (t, *J*=7.6 Hz, 2H), 7.31 (t, *J*=7.6 Hz, 1H), 7.18 (d, *J*=8.0 Hz, 1H), 7.09 (t, *J*=7.6 Hz, 1H), 6.99 (d, *J*=7.2 Hz, 1H), 6.93 (s, 2H), 5.16 (s, 1H), 4.97 (d, *J*=6.4 Hz, 1H), 4.65 (d, *J*=6.4 Hz, 1H), 1.34 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 194.0, 165.5, 153.2, 150.9, 136.5, 135.6, 133.8, 129.6, 128.8 (2C), 128.7, 128.6, 124.9, 124.4, 124.2, 116.8, 55.3, 44.8, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{32}\text{NaO}_4$ [*M*+*Na*] $^+$ 479.2193, found 479.2193. HPLC analysis [Chiralpak AD-H column, *V*(Hexane):*V*(2-propanol)=95:5, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=11.00 (major) and 14.67 min (minor).

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-7-methylchroman-2-one (**3ba**): White solid, 37.6 mg, 80% yield. m.p. 171~174 °C; $[\alpha]_{\text{D}}^{20} +66.1$ (*c* 1.3, CH_3OH); >19/1 *dr*, 92% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, *J*=7.2 Hz, 2H), 7.58 (t, *J*=7.2 Hz, 1H), 7.45 (t, *J*=7.6 Hz, 2H), 6.99 (s, 1H), 6.93 (s, 2H), 6.89 (d, *J*=8.4 Hz, 1H), 6.86 (d, *J*=8.0 Hz, 1H), 5.14 (s, 1H), 4.94 (d, *J*=6.4 Hz, 1H), 4.59 (d, *J*=6.4 Hz, 1H), 2.36 (s, 3H), 1.34 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 194.1, 165.8, 153.2, 150.8, 139.1, 136.5, 135.5, 133.7, 130.0, 128.8, 128.6, 128.4, 125.7, 124.1, 121.1, 117.3, 55.6, 44.6, 34.3, 30.1, 21.1; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_4$ [*M*+*Na*] $^+$ 493.2349, found 493.2348. HPLC analysis [Chiralpak IC column, *V*(Hexane):*V*(2-propanol)=90:10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=17.55 (major) and 25.52 min (minor).

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-6-methylchroman-2-one (**3ca**): White solid, 38.1 mg, 81% yield. m.p. 128~131 °C; $[\alpha]_{\text{D}}^{20} +53.3$ (*c* 1.8, CH_3OH); >19/1 *dr*, 97% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.88 (d, *J*=7.6 Hz, 2H), 7.59 (t, *J*=7.6 Hz, 1H), 7.46 (t, *J*=7.6 Hz, 2H), 7.09 (d, *J*=8.4 Hz, 1H), 7.06 (d, *J*=8.0 Hz, 1H), 6.93 (s, 2H), 6.81 (s, 1H), 5.16 (s, 1H), 4.92 (d, *J*=5.6 Hz, 1H), 4.56 (d, *J*=5.2 Hz, 1H), 2.24 (s, 3H), 1.36 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.9, 165.6, 153.2, 148.9, 136.6, 135.3, 134.5, 133.8, 130.2, 129.4, 129.1, 128.8, 128.7, 124.0, 123.6, 116.6, 56.0, 45.0, 34.3, 30.1, 20.8; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_5$ [*M*+*Na*] $^+$ 509.2298, found 509.2295. HPLC analysis [Chiralpak IC column, *V*(Hexane):*V*(2-propanol)=90:10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=18.79 (major) and 23.20 min (minor).

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-8-methoxychroman-2-one (**3da**): White solid, 39.0 mg, 80% yield. m.p. 120~123 °C; $[\alpha]_{\text{D}}^{20} +104.0$ (*c* 1.0, CH_3OH); >19/1 *dr*, 96% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, *J*=7.6 Hz, 2H), 7.57 (t, *J*=7.2 Hz, 1H), 7.43 (t, *J*=7.6 Hz, 2H), 7.02 (t, *J*=8.0 Hz, 1H), 6.93 (s, 2H), 6.91 (d, *J*=8.0 Hz, 1H), 6.57 (d, *J*=7.6 Hz, 1H), 5.14 (br s,

1H), 4.96 (d, $J=6.8$ Hz, 1H), 4.66 (d, $J=6.8$ Hz, 1H), 3.93 (s, 3H), 1.34 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.9, 164.9, 153.2, 147.6, 140.3, 136.5, 135.8, 133.7, 129.5, 128.7, 128.6, 125.8, 124.7, 124.3, 120.1, 120.0, 111.5, 56.2, 55.0, 44.8, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_5$ $[\text{M}+\text{Na}]^+$ 509.2298, found 509.2293. HPLC analysis [Chiralpak AD-H column, $V(\text{Hexane})$: $V(2\text{-propanol})=90:10$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=10.80$ (major) and 20.95 min (minor).

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-7-methoxychroman-2-one (**3ea**): White solid, 41.4 mg, 85% yield. m.p. 140~142 °C; $[\alpha]_D^{20}+76.3$ (c 1.3, CH_3OH); >19/1 *dr*, 93% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, $J=7.6$ Hz, 2H), 7.58 (t, $J=7.6$ Hz, 1H), 7.45 (t, $J=7.6$ Hz, 2H), 6.93 (s, 2H), 6.88 (d, $J=8.4$ Hz, 1H), 6.73 (d, $J=2.0$ Hz, 1H), 6.64 (dd, $J=8.4, 2.0$ Hz, 1H), 5.15 (s, 1H), 4.93 (d, $J=6.0$ Hz, 1H), 4.57 (d, $J=6.0$ Hz, 1H), 3.81 (s, 3H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 194.1, 165.6, 160.0, 153.1, 151.7, 136.5, 135.4, 133.8, 130.2, 129.3, 128.8, 128.6, 124.0, 116.0, 111.1, 102.2, 55.8, 55.5, 44.4, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_5$ $[\text{M}+\text{Na}]^+$ 509.2298, found 509.2295. HPLC analysis [Chiralpak OD-H column, $V(\text{Hexane})$: $V(2\text{-propanol})=90:10$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=16.90$ (major) and 22.35 min (minor).

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-6-methoxychroman-2-one (**3fa**): White solid, 40.8 mg, 84% yield. m.p. 148~151 °C; $[\alpha]_D^{20}+81.3$ (c 1.2, CH_3OH); >19/1 *dr*, 93% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.87 (d, $J=7.6$ Hz, 2H), 7.58 (t, $J=7.2$ Hz, 1H), 7.45 (t, $J=7.6$ Hz, 2H), 7.11 (d, $J=8.8$ Hz, 1H), 6.93 (s, 2H), 6.82 (dd, $J=8.8, 2.8$ Hz, 1H), 6.50 (d, $J=2.8$ Hz, 1H), 5.15 (s, 1H), 4.92 (d, $J=6.4$ Hz, 1H), 4.59 (d, $J=6.8$ Hz, 1H), 3.70 (s, 3H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 194.0, 165.6, 156.5, 153.2, 144.9, 136.6, 135.6, 133.7, 129.6, 128.8, 128.7, 125.4, 124.2, 117.6, 113.9, 113.7, 55.6, 55.4, 45.0, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_5$ $[\text{M}+\text{Na}]^+$ 509.2298, found 509.2295. HPLC analysis [Chiralpak AD-H column, $V(\text{Hexane})$: $V(2\text{-propanol})=90:10$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=11.31$ (major) and 13.10 min (minor).

(3*S*,4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-6-fluorochroman-2-one (**3ga**): White solid, 40.8 mg, 86% yield. m.p. 162~165 °C; $[\alpha]_D^{20}+64.3$ (c 1.1, CH_3OH); >19/1 *dr*, 95% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (d, $J=7.2$ Hz, 2H), 7.59 (t, $J=7.2$ Hz, 1H), 7.46 (t, $J=7.2$ Hz, 2H), 7.14 (dd, $J=8.4, 4.4$ Hz, 1H), 7.00 (t, $J=7.6$ Hz, 1H), 6.92 (s, 2H), 6.71 (d, $J=6.8$ Hz, 1H), 5.20 (s, 1H), 4.96 (d, $J=6.8$ Hz, 1H), 4.63 (d, $J=6.8$ Hz, 1H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.7, 165.1, 159.3 (d, $J=243.9$ Hz), 153.4, 146.9 (d, $J=1.9$ Hz), 136.8, 135.6, 133.9, 128.9, 128.8, 128.6, 126.5 (d, $J=7.7$ Hz), 124.1, 118.2 (d, $J=8.5$ Hz), 115.5 (d, $J=23.7$ Hz), 115.2 (d, $J=24.5$ Hz), 54.8, 44.8, 34.3, 30.1; HRMS (ESI) calcd

for $\text{C}_{30}\text{H}_{31}\text{FNaO}_4$ $[\text{M}+\text{Na}]^+$ 497.2097, found 497.2099. HPLC analysis [Chiralpak AD-H column, $V(\text{Hexane})$: $V(2\text{-propanol})=95:5$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=11.05$ (major) and 16.56 min (minor).

(3*S*,4*S*)-3-Benzoyl-7-chloro-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ha**): White solid, 40.2 mg, 82% yield. m.p. 125~128 °C; $[\alpha]_D^{20}+72.3$ (c 1.3, CH_3OH); >19/1 *dr*, 97% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, $J=7.6$ Hz, 2H), 7.59 (t, $J=7.6$ Hz, 1H), 7.46 (t, $J=7.6$ Hz, 2H), 7.19 (d, $J=1.6$ Hz, 1H), 7.07 (dd, $J=8.0, 2.0$ Hz, 1H), 6.94 (d, $J=8.4$ Hz, 1H), 6.91 (s, 2H), 5.18 (s, 1H), 4.95 (d, $J=6.4$ Hz, 1H), 4.60 (d, $J=6.4$ Hz, 1H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.6, 164.7, 153.4, 151.3, 136.7, 135.3, 134.0 (2 C), 129.7, 129.3, 128.9, 128.7, 125.1, 124.0, 123.0, 117.3, 55.2, 44.5, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{31}\text{ClNaO}_4$ $[\text{M}+\text{Na}]^+$ 513.1803, found 513.1801. HPLC analysis [Chiralpak IC column, $V(\text{Hexane})$: $V(2\text{-propanol})=90:10$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=14.51$ (major) and 19.77 min (minor).

(3*S*,4*S*)-3-Benzoyl-6-chloro-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ia**): White solid, 40.2 mg, 82% yield. m.p. 151~154 °C; $[\alpha]_D^{20}+61.4$ (c 1.1, CH_3OH); >19/1 *dr*, 86% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.88 (d, $J=7.2$ Hz, 2H), 7.60 (t, $J=6.8$ Hz, 1H), 7.47 (t, $J=7.2$ Hz, 2H), 7.27 (d, $J=7.2$ Hz, 1H), 7.12 (d, $J=8.4$ Hz, 1H), 7.01 (s, 1H), 6.91 (s, 2H), 5.21 (s, 1H), 4.95 (d, $J=4.8$ Hz, 1H), 4.58 (d, $J=4.4$ Hz, 1H), 1.36 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.5, 164.8, 153.4, 149.4, 136.8, 135.1, 134.0, 130.0, 129.2, 128.9, 128.7, 128.6, 126.0, 124.0, 118.2, 55.2, 44.8, 34.3, 30.0; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{31}\text{ClNaO}_4$ $[\text{M}+\text{Na}]^+$ 513.1803, found 513.1795. HPLC analysis [Chiralpak IC column, $V(\text{Hexane})$: $V(2\text{-propanol})=90:10$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=10.89$ (major) and 17.32 min (minor).

(3*S*,4*S*)-3-Benzoyl-7-bromo-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ja**): White solid, 41.1 mg, 77% yield. m.p. 135~138 °C; $[\alpha]_D^{20}+56.8$ (c 1.0, CH_3OH); >19/1 *dr*, 97% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, $J=7.6$ Hz, 2H), 7.59 (t, $J=7.6$ Hz, 1H), 7.46 (t, $J=7.6$ Hz, 2H), 7.34 (s, 1H), 7.22 (d, $J=8.0$ Hz, 1H), 6.90 (s, 2H), 6.88 (d, $J=8.0$ Hz, 1H), 5.17 (s, 1H), 4.94 (d, $J=6.0$ Hz, 1H), 4.58 (d, $J=6.0$ Hz, 1H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.6, 164.7, 153.4, 151.4, 136.8, 135.4, 134.0, 130.0, 129.2, 128.9, 128.7, 128.0, 124.1, 123.6, 121.6, 120.2, 55.1, 44.6, 34.4, 30.1; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{31}\text{BrNaO}_4$ $[\text{M}+\text{Na}]^+$ 557.1298, found 557.1296. HPLC analysis [Chiralpak IC column, $V(\text{Hexane})$: $V(2\text{-propanol})=90:10$, flow rate=1.0 mL/min, wavelength=254 nm]: $t_R=15.37$ (major) and 22.27 min (minor).

(3*S*,4*S*)-3-Benzoyl-6-bromo-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ka**): White solid, 36.3 mg, 68% yield. m.p. 155~158 °C; $[\alpha]_D^{20}+50.6$ (c 1.5, CH_3OH); >19/1 *dr*, 95% *ee*. ^1H NMR (400 MHz, CDCl_3)

δ : 7.87 (d, $J=7.6$ Hz, 2H), 7.61 (t, $J=7.6$ Hz, 1H), 7.47 (t, $J=7.6$ Hz, 2H), 7.42 (dd, $J=8.4, 1.6$ Hz, 1H), 7.16 (d, $J=1.6$ Hz, 1H), 7.07 (d, $J=8.4$ Hz, 1H), 6.91 (s, 2H), 5.20 (s, 1H), 4.93 (d, $J=5.6$ Hz, 1H), 4.58 (d, $J=5.6$ Hz, 1H), 1.36 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.4, 164.6, 153.5, 150.0, 136.8, 135.1, 134.0, 131.9, 131.5, 129.3, 128.9, 128.7, 126.4, 124.0, 118.6, 117.5, 55.4, 44.8, 34.4, 30.0; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{31}\text{BrNaO}_4$ $[\text{M}+\text{Na}]^+$ 557.1298, found 557.1298. HPLC analysis [Chiralpak IC column, $V(\text{Hexane}) : V(2\text{-propanol})=90 : 10$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=10.96$ (major) and 17.75 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(4-methylbenzoyl)chroman-2-one (**3ab**): White solid, 37.1 mg, 79% yield. m.p. 120~123 °C; $[\alpha]_D^{20} +47.7$ (c 1.3, CH_3OH); >19/1 *dr*, 92% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.78 (d, $J=8.4$ Hz, 2H), 7.30 (t, $J=8.4$ Hz, 1H), 7.25 (d, $J=8.4$ Hz, 2H), 7.17 (d, $J=8.0$ Hz, 1H), 7.08 (t, $J=7.6$ Hz, 1H), 6.99 (d, $J=7.2$ Hz, 1H), 6.93 (s, 2H), 5.15 (s, 1H), 4.94 (d, $J=6.0$ Hz, 1H), 4.62 (d, $J=6.0$ Hz, 1H), 2.41 (s, 3H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 193.4, 165.6, 153.2, 150.9, 144.9, 136.5, 132.9, 129.9, 129.5, 128.8, 128.7, 124.9, 124.3, 124.1, 116.8, 55.4, 44.9, 34.3, 30.1, 21.7; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_4$ $[\text{M}+\text{Na}]^+$ 493.2349, found 493.2348. HPLC analysis [Chiralpak IC column, $V(\text{Hexane}) : V(2\text{-propanol})=90 : 10$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=24.63$ (major) and 27.19 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(3-methylbenzoyl)chroman-2-one (**3ac**): White solid, 36.7 mg, 78% yield. m.p. 138~141 °C; $[\alpha]_D^{20} +58.4$ (c 1.3, CH_3OH); >19/1 *dr*, 93% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.67 (s, 1H), 7.66 (d, $J=6.8$ Hz, 1H), 7.40 (d, $J=7.2$ Hz, 1H), 7.34 (t, $J=7.2$ Hz, 1H), 7.31 (t, $J=7.2$ Hz, 1H), 7.18 (d, $J=7.6$ Hz, 1H), 7.08 (dt, $J=7.6, 0.8$ Hz, 1H), 6.98 (d, $J=7.2$ Hz, 1H), 6.93 (s, 2H), 5.16 (s, 1H), 4.96 (d, $J=6.4$ Hz, 1H), 4.63 (d, $J=6.4$ Hz, 1H), 2.39 (s, 3H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 194.2, 165.6, 153.2, 151.0, 138.7, 136.5, 135.5, 134.6, 129.8, 129.1, 128.8, 128.7 (2 C), 125.8, 124.9, 124.3, 124.1, 116.8, 55.5, 44.9, 34.3, 30.1, 21.3; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_4$ $[\text{M}+\text{Na}]^+$ 493.2349, found 493.2348. HPLC analysis [Chiralpak IC column, $V(\text{Hexane}) : V(2\text{-propanol})=90 : 10$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=16.81$ (major) and 19.38 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(2-methylbenzoyl)chroman-2-one (**3ad**): White solid, 40.0 mg, 85% yield. m.p. 126~129 °C; $[\alpha]_D^{20} +74.3$ (c 1.6, CH_3OH); >19/1 *dr*, 90% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.52 (d, $J=7.2$ Hz, 1H), 7.21~7.35 (m, 4H), 7.14 (t, $J=7.2$ Hz, 2H), 7.06 (t, $J=7.2$ Hz, 1H), 6.87 (s, 2H), 6.86 (d, $J=7.2$ Hz, 1H), 5.19 (s, 1H), 4.80 (d, $J=9.2$ Hz, 1H), 4.63 (d, $J=9.2$ Hz, 1H), 1.97 (s, 3H), 1.31 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 197.8, 166.1, 153.2, 150.9, 138.6, 137.3, 136.4, 131.7, 131.6, 128.7, 128.3, 128.1, 127.9, 125.6, 125.5, 124.8, 124.6, 116.7, 57.5, 44.5, 34.2, 30.0, 19.8; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{34}\text{NaO}_4$ $[\text{M}+\text{Na}]^+$

493.2349, found 493.2349. HPLC analysis [Chiralpak IC column, $V(\text{Hexane}) : V(2\text{-propanol})=90 : 10$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=16.61$ (major) and 23.56 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(4-methoxybenzoyl)chroman-2-one (**3ae**): White solid, 41.3 mg, 85% yield. m.p. 112~115 °C; $[\alpha]_D^{20} +50.0$ (c 1.4, CH_3OH); >19/1 *dr*, 91% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.88 (d, $J=8.4$ Hz, 2H), 7.30 (t, $J=7.6$ Hz, 1H), 7.17 (d, $J=8.0$ Hz, 1H), 7.08 (t, $J=7.6$ Hz, 1H), 7.02 (d, $J=7.6$ Hz, 1H), 6.93 (s, 2H), 6.92 (d, $J=8.8$ Hz, 2H), 5.15 (s, 1H), 4.92 (d, $J=6.0$ Hz, 1H), 4.62 (d, $J=5.6$ Hz, 1H), 3.87 (s, 3H), 1.35 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 192.1, 165.6, 164.1, 153.2, 151.0, 136.5, 131.1, 130.1, 128.7, 128.4, 124.8, 124.4, 124.1, 116.8, 114.0, 55.5, 55.2, 44.9, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{34}\text{NaO}_5$ $[\text{M}+\text{Na}]^+$ 509.2298, found 509.2299. HPLC analysis [Chiralpak AD-H column, $V(\text{Hexane}) : V(2\text{-propanol})=90 : 10$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=17.06$ (major) and 25.49 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(3,4-dimethoxybenzoyl)chroman-2-one (**3af**): White solid, 34.1 mg, 66% yield. m.p. 125~128 °C; $[\alpha]_D^{20} +29.2$ (c 2.6, CH_3OH); >19/1 *dr*, 86% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.57 (dd, $J=8.4, 2.0$ Hz, 1H), 7.43 (d, $J=2.0$ Hz, 1H), 7.31 (dt, $J=8.4, 1.6$ Hz, 1H), 7.17 (d, $J=7.2$ Hz, 1H), 7.09 (dt, $J=7.6, 1.2$ Hz, 1H), 7.02 (dd, $J=7.6, 0.8$ Hz, 1H), 6.94 (s, 2H), 6.88 (d, $J=8.4$ Hz, 1H), 5.16 (s, 1H), 4.94 (d, $J=6.0$ Hz, 1H), 4.64 (d, $J=6.0$ Hz, 1H), 3.95 (s, 3H), 3.88 (s, 3H), 1.36 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 192.1, 165.7, 154.0, 153.2, 151.0, 149.3, 147.1, 136.5, 130.1, 128.8, 128.5, 124.9, 124.4, 124.0, 123.6, 116.8, 110.7, 110.0, 56.2, 56.0, 55.1, 45.0, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{32}\text{H}_{36}\text{NaO}_6$ $[\text{M}+\text{Na}]^+$ 539.2404, found 539.2399. HPLC analysis [Chiralpak IC column, $V(\text{Hexane}) : V(2\text{-propanol})=75 : 25$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=23.39$ (major) and 29.58 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(4-fluorobenzoyl)chroman-2-one (**3ag**): White solid, 39.4 mg, 83% yield. m.p. 122~125 °C; $[\alpha]_D^{20} +50.9$ (c 1.1, CH_3OH); >19/1 *dr*, 93% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.89 (dd, $J=8.4, 5.6$ Hz, 2H), 7.31 (t, $J=7.6$ Hz, 1H), 7.08~7.18 (m, 4H), 7.00 (d, $J=7.6$ Hz, 1H), 6.92 (s, 2H), 5.17 (s, 1H), 4.91 (d, $J=7.2$ Hz, 1H), 4.65 (d, $J=6.8$ Hz, 1H), 1.34 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 192.4, 167.3, 165.3, 164.7, 152.0 (d, $J=242.9$ Hz), 136.6, 132.2, 131.3 (d, $J=9.3$ Hz), 129.4, 128.8, 128.6, 124.9, 124.6, 124.2, 116.8, 115.9 (d, $J=21.9$ Hz), 55.1, 44.7, 34.3, 30.1; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{31}\text{FNaO}_4$ $[\text{M}+\text{Na}]^+$ 497.2099, found 497.2112. HPLC analysis [Chiralpak IC column, $V(\text{Hexane}) : V(2\text{-propanol})=90 : 10$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R=12.83$ (major) and 15.29 min (minor).

(3*S*,4*S*)-3-(3-Chlorobenzoyl)-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ah**): White solid, 44.1 mg, 90% yield. m.p. 133~136 °C; $[\alpha]_D^{20} +58.6$ (c 1.4,

CH₃OH); >19/1 *dr*, 96% *ee*. ¹H NMR (400 MHz, CDCl₃) δ : 7.78 (s, 1H), 7.72 (d, *J*=8.0 Hz, 1H), 7.54 (d, *J*=8.0 Hz, 1H), 7.39 (t, *J*=8.0 Hz, 1H), 7.32 (t, *J*=7.2 Hz, 1H), 7.18 (d, *J*=8.0 Hz, 1H), 7.10 (t, *J*=7.6 Hz, 1H), 6.97 (d, *J*=7.6 Hz, 1H), 6.92 (s, 2H), 5.17 (s, 1H), 4.88 (d, *J*=7.6 Hz, 1H), 4.65 (d, *J*=7.6 Hz, 1H), 1.35 (s, 18H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 193.1, 165.2, 153.3, 150.9, 137.4, 136.7, 135.1, 133.6, 130.1, 129.1, 128.9, 128.7, 128.6, 126.6, 125.0, 124.5, 124.3, 116.9, 55.4, 44.8, 34.3, 30.1; HRMS (ESI) calcd for C₃₀H₃₁ClNaO₄ [M + Na]⁺ 513.1803, found 518.1800. HPLC analysis [Chiralpak IC column, *V*(Hexane) : *V*(2-propanol)=90 : 10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=10.38 (major) and 12.52 min (minor).

(3*S*,4*S*)-3-(4-Bromobenzoyl)-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ai**): White solid, 43.2 mg, 81% yield. m.p. 151~154 °C; [α]_D²⁰ +71.6 (*c* 1.2, CH₃OH); >19/1 *dr*, 94% *ee*. ¹H NMR (400 MHz, CDCl₃) δ : 7.71 (d, *J*=8.4 Hz, 2H), 7.58 (d, *J*=8.4 Hz, 2H), 7.31 (t, *J*=7.6 Hz, 1H), 7.16 (d, *J*=8.0 Hz, 1H), 7.10 (t, *J*=7.6 Hz, 1H), 6.99 (d, *J*=7.2 Hz, 1H), 6.92 (s, 2H), 5.17 (br. s, 1H), 4.89 (d, *J*=7.2 Hz, 1H), 4.64 (d, *J*=7.2 Hz, 1H), 1.35 (s, 18H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 193.1, 165.2, 153.3, 150.8, 136.6, 134.5, 130.0, 129.3, 129.1, 128.9, 128.6, 125.0, 124.5, 124.2, 116.8, 55.2, 44.7, 34.3, 30.1; HRMS (ESI) calcd for C₃₀H₃₁BrNaO₄ [M + Na]⁺ 557.1298, found 557.1296. HPLC analysis [Chiralpak IC column, *V*(Hexane) : *V*(2-propanol)=90 : 10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=13.54 (major) and 17.42 min (minor).

(3*S*,4*S*)-3-(3-Bromobenzoyl)-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3aj**): White solid, 42.7 mg, 80% yield. m.p. 112~115 °C; [α]_D²⁰ +103.4 (*c* 1.2, CH₃OH); >19/1 *dr*, 95% *ee*. ¹H NMR (400 MHz, CDCl₃) δ : 7.94 (s, 1H), 7.76 (d, *J*=8.0 Hz, 1H), 7.69 (d, *J*=8.4 Hz, 1H), 7.32 (dt, *J*=8.4, 2.8 Hz, 1H), 7.18 (d, *J*=8.4 Hz, 1H), 7.10 (t, *J*=7.6 Hz, 1H), 6.97 (d, *J*=7.6 Hz, 1H), 6.92 (s, 2H), 5.17 (s, 1H), 4.88 (d, *J*=7.6 Hz, 1H), 4.65 (d, *J*=7.2 Hz, 1H), 1.35 (s, 18H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 193.0, 165.2, 153.3, 150.9, 137.6, 136.7, 136.5, 131.5, 130.3, 129.1, 128.9, 128.7, 127.0, 125.0, 124.4, 124.3, 123.1, 116.9, 55.3, 44.8, 34.3, 30.1; HRMS (ESI) calcd for C₃₀H₃₁BrNaO₄ [M + Na]⁺ 557.1298, found 557.1296. HPLC analysis [Chiralpak IC column, *V*(Hexane) : *V*(2-propanol)=90 : 10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=10.81 (major) and 13.24 min (minor).

(3*S*,4*S*)-3-(2-Naphthoyl)-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)chroman-2-one (**3ak**): White solid, 45.1 mg, 89% yield. m.p. 121~124 °C; [α]_D²⁰ +73.8 (*c* 1.6, CH₃OH); >19/1 *dr*, 87% *ee*. ¹H NMR (400 MHz, CDCl₃) δ : 8.45 (s, 1H), 7.89~7.96 (m, 4H), 7.58~7.67 (m, 2H), 7.33 (t, *J*=8.8 Hz, 1H), 7.21 (d, *J*=8.4 Hz, 1H), 7.09 (t, *J*=7.6 Hz, 1H), 6.99 (d, *J*=8.0 Hz, 1H), 6.98 (s, 2H), 5.14 (s, 1H), 5.13 (d, *J*=4.8 Hz, 1H), 4.71 (d, *J*=4.8 Hz, 1H), 1.34 (s, 18H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 193.9, 165.6, 153.2, 151.0, 136.6, 135.8, 132.78, 132.4, 130.7, 129.9,

129.6, 129.0, 128.9, 128.8, 127.8, 127.1, 124.9, 124.3, 124.2, 123.9, 116.9, 55.7, 45.1, 34.3, 30.1; HRMS (ESI) calcd for C₃₄H₃₄NaO₄ [M + Na]⁺ 529.2349, found 529.2347. HPLC analysis [Chiralpak IC column, *V*(Hexane) : *V*(2-propanol)=90 : 10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=22.30 (major) and 26.05 min (minor).

(3*S*,4*S*)-4-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-3-(thiophene-2-carbonyl)chroman-2-one (**3al**): White solid, 41.7 mg, 90% yield. m.p. 161~163 °C; [α]_D²⁰ +54.0 (*c* 1.4, CH₃OH); >19/1 *dr*, 93% *ee*. ¹H NMR (400 MHz, CDCl₃) δ : 7.74 (dd, *J*=7.6, 0.4 Hz, 1H), 7.67 (dd, *J*=8.0, 0.8 Hz, 1H), 7.31 (dt, *J*=7.6, 1.2 Hz, 1H), 7.09~7.13 (m, 2H), 7.04 (d, *J*=7.2 Hz, 1H), 6.95 (s, 2H), 5.17 (s, 1H), 4.76 (d, *J*=5.4 Hz, 1H), 4.72 (d, *J*=5.4 Hz, 1H), 1.36 (s, 18H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 185.9, 165.0, 153.2, 150.9, 142.6, 136.5, 135.3, 133.4, 129.4, 128.8, 128.7, 128.3, 124.9, 124.5, 124.3, 116.7, 56.7, 44.9, 34.3, 30.1; HRMS (ESI) calcd for C₂₈H₃₀NaO₄S [M + Na]⁺ 485.1757, found 485.1752. HPLC analysis [Chiralpak AD-H column, *V*(Hexane) : *V*(2-propanol)=90 : 10, flow rate=1.0 mL/min, wavelength=254 nm]: *t*_R=15.82 (minor) and 26.05/18.74 min (major).

4.3 Transformation of 3,4-dihydrocoumarin **3aa** to the corresponding β,β -diaryl ketone **4**

To a solution of **3aa** (137 mg, 0.3 mmol) in tetrahydrofuran (3 mL) was added a solution of sodium hydroxide (32 mg, 0.8 mmol) in water (3 mL), and the resulting mixture was stirred at 40 °C for 1.5 h. After cooling to room temperature, the reaction mixture was extracted with ethyl acetate (5 mL $\times$ 3). The combined organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The crude product was used directly in the following reaction without purification.

To a solution of the aforementioned crude product in acetone (2 mL) were added potassium carbonate (249 mg, 1.8 mmol) and methyl iodide (298 mg, 2.1 mmol). After stirring at room temperature for 24 h, the resulting mixture was diluted with water (10 mL) and extracted with ethyl acetate (5 mL $\times$ 3). The combined organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel (100~200 mesh, *V*(petroleum ether) : *V*(ethyl acetate)=15 : 1) to afford the desired β,β -diaryl ketone **4**.

(*S*)-3-(3,5-Di-*tert*-butyl-4-methoxyphenyl)-3-(2-methoxyphenyl)-1-phenylpropan-1-one (**4**): Colorless oil, 59.2 mg, 43% yield. [α]_D²⁰ -25.8 (*c* 0.81, CH₃OH); >19/1 *dr*, 95% *ee*. ¹H NMR (400 MHz, CDCl₃) δ : 7.91 (d, *J*=7.2 Hz, 2H), 7.52 (t, *J*=7.2 Hz, 1H), 7.42 (t, *J*=7.2 Hz, 2H), 7.16 (dt, *J*=7.6, 1.6 Hz, 1H), 7.13 (dd, *J*=7.2, 1.6 Hz, 1H), 7.11 (s, 2H), 6.88 (t, *J*=7.6 Hz, 1H), 6.84 (d, *J*=8.4 Hz, 1H), 5.06 (t, *J*=7.2 Hz, 1H), 3.77 (s, 3H), 3.66 (dd, *J*=7.6, 2.0 Hz, 2H), 1.34 (s, 18H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 199.2, 157.5, 156.9, 142.8, 137.5, 136.9, 133.0, 132.7, 128.4, 128.1 (2C), 127.2, 126.3, 120.5, 110.7, 64.0,

55.3, 44.0, 40.3, 35.7, 32.1; HRMS (ESI) calcd for $C_{31}H_{38}NaO_3$ $[M+Na]^+$ 481.2713, found 481.2718. HPLC analysis [Chiralpak OD-H column, $V(\text{Hexane}) : V(2\text{-propanol}) = 95 : 5$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R = 5.75$ (minor) and 8.12 min (major).

4.4 Transformation of 3,4-dihydrocoumarin **3aa** to the corresponding *cis*-chromane **5**

To a solution of **3aa** (137 mg, 0.3 mmol) in tetrahydrofuran (3 mL) was added a solution of sodium hydroxide (32 mg, 0.8 mmol) in water (3 mL), and the resulting mixture was stirred at 40 °C for 1.5 h. After cooling to room temperature, the reaction mixture was extracted with ethyl acetate (5 mL $\times$ 3). The combined organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The crude product was used directly in the following reaction without purification.

To a solution of the aforementioned crude product in dichloromethane (2 mL) were added successively triethylsilane (523 mg, 4.5 mmol) and boron trifluoride diethyl etherate (170 mg, 1.2 mmol) at 0 °C. The resulting mixture was stirred at the same temperature for 3 h and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel [100~200 mesh, $V(\text{petroleum ether}) : V(\text{ethyl acetate}) = 15 : 1$] to give the corresponding *cis*-chromane **5**.

2,6-Di-*tert*-butyl-4-((2*R*,4*S*)-2-phenylchroman-4-yl)-phenol (**5**): White solid, 56.0 mg, 45% yield. m.p. 182~185 °C; $[\alpha]_D^{20} + 32.4$ (c 0.95, CH_3OH); >19/1 *dr*, 87% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.50 (d, $J = 7.2$ Hz, 2H), 7.40 (t, $J = 7.2$ Hz, 2H), 7.32 (t, $J = 7.2$ Hz, 1H), 7.09~7.19 (m, 2H), 6.98 (s, 2H), 6.94 (d, $J = 8.0$ Hz, 1H), 6.80 (t, $J = 7.6$ Hz, 1H), 5.24 (d, $J = 10.0$ Hz, 1H), 5.10 (s, 1H), 4.27 (dd, $J = 12.4, 5.6$ Hz, 1H), 2.44 (ddd, $J = 13.6, 5.6, 2.0$ Hz, 1H), 2.20~2.29 (m, 1H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 155.4, 152.4, 141.4, 135.9, 134.6, 129.7, 128.5, 127.9, 127.5, 126.3, 126.0, 125.0, 120.3, 116.7, 78.3, 43.56, 40.6, 34.35, 30.4; HRMS (ESI) calcd for $C_{29}H_{34}NaO_2$ $[M+Na]^+$ 437.2451, found 437.2455. HPLC analysis [Chiralpak OD-H column, $V(\text{Hexane}) : V(2\text{-propanol}) = 98 : 2$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R = 4.84$ (major) and 5.60 min (minor).

4.5 Transformation of 3,4-dihydrocoumarin **3aa** to the corresponding 3-methylated 3,4-dihydrocoumarin **6**

To a solution of **3aa** (45.6 mg, 0.1 mmol) in acetone (2 mL) were added potassium carbonate (20.7 mg, 0.15 mmol) and iodomethane (21.3 mg, 0.15 mmol). The resulting mixture was stirred at room temperature for 24 h. Water (20 mL) was added and the reaction mixture was extracted with dichloromethane (10 mL $\times$ 3). The combined organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel [100~200 mesh, $V(\text{petroleum ether}) : V(\text{ethyl acetate}) = 15 : 1$] to give the corresponding 3-methylated 3,4-dihydrocoumarin **6**.

(4*S*)-3-Benzoyl-4-(3,5-di-*tert*-butyl-4-hydroxyphenyl)-3-methylchroman-2-one (**6**): White solid, m.p. 144~148 °C; 42.3 mg, 90% yield. $[\alpha]_D^{20} + 98.6$ (c 0.82, CH_3OH); >19/1 *dr*, 98% *ee*. ^1H NMR (400 MHz, CDCl_3) δ : 7.41 (t, $J = 7.2$ Hz, 1H), 7.28 (t, $J = 7.2$ Hz, 2H), 7.27 (d, $J = 7.2$ Hz, 1H), 7.08 (t, $J = 8.0$ Hz, 2H), 7.22 (d, $J = 8.0$ Hz, 1H), 6.74 (s, 2H), 5.08 (s, 1H), 4.42 (s, 1H), 1.75 (s, 3H), 1.20 (s, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 199.6, 169.9, 153.3, 151.0, 136.6, 136.2, 131.4, 129.4, 128.9, 128.7, 128.3, 126.9, 125.7, 125.3, 124.4, 116.4, 57.7, 53.3, 34.0, 29.8, 24.8; HRMS (ESI) calcd for $C_{31}H_{34}NaO_4$ $[M+Na]^+$ 493.2349, found 493.2352. HPLC analysis [Chiralpak OD-H column, $V(\text{Hexane}) : V(2\text{-propanol}) = 99.5 : 0.5$, flow rate = 1.0 mL/min, wavelength = 254 nm]: $t_R = 8.75$ (major) and 12.06 min (minor).

Supporting Information Crystallography data, copies of NMR and HRMS spectra as well as chiral HPLC chromatograms of the prepared optically active 3,4-dihydrocoumarins **3aa**~**3ka**, **3ab**~**3al** and compounds **4**~**6** derived from **3aa**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Asai, F.; Iinuma, M.; Tanaka, T.; Mizuno, M. *Phytochemistry* **1991**, *30*, 3091.
(b) Asai, F.; Iinuma, M.; Tanaka, T.; Mizuno, M. *Heterocycles* **1992**, *33*, 229.
(c) Asai, F.; Iinuma, M.; Tanaka, T.; Takenaka, M.; Mizuno, M. *Phytochemistry* **1992**, *31*, 2487.
(d) Ngadjui, B. T.; Kapche, G. W. F.; Tamboue, H.; Abegaz, B. M.; Connolly, J. D. *Phytochemistry* **1999**, *51*, 119.
(e) Seo, E.-K.; Wani, M. C.; Wall, M. E.; Navarro, H.; Mukherjee, R.; Farnsworth, N. R.; Kinghorn, A. D. *Phytochemistry* **2000**, *55*, 35.
(f) Wu, P.-L.; Hsu, Y.-L.; Zao, C.-W.; Damu, A. G.; Wu, T.-S. *J. Nat. Prod.* **2005**, *68*, 1180.
(g) Wungwintaweeikul, B.; Umehara, K.; Miyase, T.; Noguchi, H. *Phytochemistry* **2011**, *72*, 495.
(h) Ee, G. C. L.; Mah, S. H.; The, S. S.; Rahmani, M.; Go, R.; Taufiq-Yap, Y. H. *Molecules* **2011**, *16*, 9721.
- [2] For selected examples, see: (a) Bedalov, A.; Gattbonton, T.; Irvine, W. P.; Gottschling, D. E.; Simon, J. A. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, *98*, 15113.
(b) Posakony, J.; Hirao, M.; Stevens, S.; Simon, J. A.; Bedalov, A. *J. Med. Chem.* **2004**, *47*, 2635.
(c) Roelens, F.; Huvaere, K.; Dhooge, W.; Van Cleemput, M.; Comhaire, F.; de Keukeleire, D. *Eur. J. Med. Chem.* **2005**, *40*, 1042.
(d) Zhang, X.; Wang, H.; Song, Y.; Nie, L.; Wang, L.; Liu, B.; Shen, P.; Liu, Y. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 949.
(e) Modranka, J.; Albrecht, A.; Jakubowski, R.; Krawczyk, H.; Roóżalski, M.; Krajewska, U.; Janecka, A.; Wyrębska, A.; Różańska, B.; Janecki, T. *Bioorg. Med. Chem.* **2012**, *20*, 5017.
(f) Sashidhara, K. V.; Singh, S. P.; Singh, S. V.; Srivastava, R. K.; Srivastava, K.; Saxena, J. K.; Puri, S. K. *Eur. J. Med. Chem.* **2013**, *60*, 497.
(g) Zhang, K.; Ding, W.; Sun, J.; Zhang, B.; Lu, F.; Lai, R.; Zou, Y.; Yedid, G. *Biochimie* **2014**, *107*, 203.
(h) Deredas, D.; Huben, K.; Janecka, A.; Długosz, A.; Pomorska, D. K.; Mirowski, M.; Krajewska, U.; Janecka, T.; Krawczyk, H. *MedChemComm* **2016**, *7*, 1745.
- [3] (a) Song, F.; Lu, S.; Gunnet, J.; Xu, J. Z.; Wines, P.; Proost, J.; Liang, Y.; Baumann, C.; Lenhard, J.; Murray, W. V.; Demarest, K.

- T.; Kuo, G.-H. *J. Med. Chem.* **2007**, *50*, 2807.
- (b) Ballerini, E.; Minuti, L.; Piermatti, O.; Pizzo, F. *J. Org. Chem.* **2009**, *74*, 4311.
- [4] For a review on the asymmetric construction of 3,4-dihydrocoumarins, see: Leitis, Z. *Chem. Heterocycl. Compd.* **2016**, *52*, 527.
- [5] (a) Alden-Danforth, E.; Scerba, M. T.; Lectka, T. *Org. Lett.* **2008**, *10*, 4951.
- (b) Lü, H.; You, L.; Ye, S. *Adv. Synth. Catal.* **2009**, *351*, 2822.
- (c) Izquierdo, J.; Orue, A.; Scheidt, K. A. *J. Am. Chem. Soc.* **2013**, *135*, 10634.
- (d) Lee, A.; Scheidt, K. A. *Chem. Commun.* **2015**, *51*, 3407.
- (e) Caruana, L.; Mondatori, M.; Corti, V.; Morales, S.; Mazzanti, A.; Fochi, M.; Bernardi, L. *Chem.-Eur. J.* **2015**, *21*, 6037.
- (f) Zhou, D.; Mao, K.; Zhang, J.; Yan, B.; Wang, W.; Xie, H. *Tetrahedron Lett.* **2016**, *57*, 5649.
- (g) Yu, X.-Y.; Chen, J.-R.; Wei, Q.; Cheng, H.-G.; Liu, Z.-C.; Xiao, W.-J. *Chem.-Eur. J.* **2016**, *22*, 6774.
- (h) Zhang, Y.-C.; Zhu, Q.-N.; Yang, X.; Zhou, L.-J.; Shi, F. *J. Org. Chem.* **2016**, *81*, 1681.
- (i) Jin, J.-H.; Li, X.-Y.; Luo, X.; Fossey, J. S.; Deng, W.-P. *J. Org. Chem.* **2017**, *82*, 5424.
- (j) Wu, B.; Yu, Z.; Gao, X.; Lan, Y.; Zhou, Y.-G. *Angew. Chem., Int. Ed.* **2017**, *56*, 4006.
- (k) Chen, X.; Song, R.; Liu, Y.; Ooi, C. Y.; Jin, Z.; Zhu, T.; Wang, H.; Hao, L.; Chi, Y. R. *Org. Lett.* **2017**, *19*, 5892.
- (l) Zhang, T.; Ma, C.; Zhou, J.-Y.; Mei, G.-J.; Shi, F. *Adv. Synth. Catal.* **2018**, *360*, 1128.
- (m) Alden-Danforth, E.; Scerba, M. T.; Lectka, T. *Org. Lett.* **2018**, *10*, 4951.
- (n) Cui, L.; Lv, D.; Wang, Y.; Fan, Z.; Li, Z.; Zhou, Z. *J. Org. Chem.* **2018**, *82*, 4221.
- (o) Lv, D.; Zhao, M.; Wang, Y.; Zhou, Z. *Org. Biomol. Chem.* **2019**, *17*, 9636.
- (p) Kim, K. S.; Jang, J.; Kim, D. Y. *ChemistrySelect* **2020**, *5*, 13259.
- [6] For selected reviews on *p*-QMs, see: (a) Caruana, L.; Fochi, M.; Bernardi, L. *Molecules* **2015**, *20*, 11733.
- (b) Parra, A.; Tortosa, M. *ChemCatChem* **2015**, *7*, 1524.
- (c) Chauhan, P.; Kaya, U.; Enders, D. *Adv. Synth. Catal.* **2017**, *359*, 888.
- (d) Lima, C. G. S.; Pauli, F. P.; Costa, D. C. S.; de Souza, A. S.; Ferreira, V. F.; de Carvalho da Silva, F. *Eur. J. Org. Chem.* **2020**, 2650.
- (e) Wang, J.-Y.; Hao, W.-J.; Tu, S.-J.; Jiang, B. *Org. Chem. Front.* **2020**, *7*, 1743.
- [7] Zhao, K.; Zhi, Y.; Shu, T.; Valkonen, A.; Rissanen, K.; Enders, D. *Angew. Chem., Int. Ed.* **2016**, *55*, 12104.
- [8] (a) Liu, L.; Yuan, Z.; Pan, R.; Zeng, Y.; Lin, A.; Yao, H.; Huang, Y. *Org. Chem. Front.* **2018**, *5*, 623.
- (b) Zielke, K.; Kováč, O.; Winter, M.; Pospíšil, J.; Waser, M. *Chem.-Eur. J.* **2019**, *25*, 8163.
- (c) Tan, J.-P.; Yu, P.; Wu, J.-H.; Chen, Y.; Pan, J.; Jiang, C.; Ren, X.; Zhang, H.-S.; Wang, T. *Org. Lett.* **2019**, *21*, 7298.
- (d) Lu, H.; Zhang, H.-X.; Tan, C.-Y.; Liu, J.-Y.; Wei, H.; Xu, P.-F. *J. Org. Chem.* **2019**, *84*, 10292.
- [9] (a) Zhang, L.; Zhou, X.; Li, P.; Liu, Z.; Liu, Y.; Sun, Y.; Li, W. *RSC Adv.* **2017**, *7*, 39216.
- (b) Zhang, L.; Liu, Y.; Liu, K.; Liu, Z.; He, N.; Li, W. *Org. Biomol. Chem.* **2017**, *15*, 8743.
- (c) Duan, C.; Ye, L.; Xu, W.; Li, X.; Chen, F.; Zhao, Z.; Li, X. *Chin. Chem. Lett.* **2018**, *29*, 1273.
- (d) Zhang, Z.-P.; Xie, K.-X.; Yang, C.; Li, M.; Li, X. *J. Org. Chem.* **2018**, *83*, 364.
- (e) Zhang, Z.-P.; Chen, L.; Li, X.; Cheng, J.-P. *J. Org. Chem.* **2018**, *83*, 2714.
- (f) Jiang, X.-L.; Wu, S.-F.; Wang, J.-R.; Mei, G.-J.; Shi, F. *Adv. Synth. Catal.* **2018**, *360*, 4225.
- (g) Yang, G.-H.; Zhao, Q.; Zhang, Z.-P.; Zheng, H.-L.; Chen, L.; Li, X. *J. Org. Chem.* **2019**, *84*, 7883.
- (h) You, Y.; Li, T.-T.; Yuan, S.-P.; Xie, K.-X.; Wang, Z.-H.; Zhao, J.-Q.; Zhou, M.-Q.; Yuan, W.-C. *Chem. Commun.* **2020**, *56*, 439.
- (i) Tan, J.-P.; Zhang, H.; Jiang, Z.; Chen, Y.; Ren, X.; Jiang, C.; Wang, T. *Adv. Synth. Catal.* **2020**, *362*, 1058.
- (j) Zhao, M.-X.; Xiang, J.; Zhao, Z.-Q.; Zhao, X.-L.; Shi, M. *Org. Biomol. Chem.* **2020**, *18*, 1637.
- (k) Roy, S.; Pradhan, S.; Kumar, K.; Chatterjee, I. *Org. Chem. Front.* **2020**, *7*, 1388.
- (l) Wu, S.-F.; Tu, M.-S.; Hang, Q.-Q.; Zhang, S.; Ding, H.; Zhang, Y.-C.; Shi, F. *Org. Biomol. Chem.* **2020**, *18*, 5388.
- (m) Xiang, M.; Li, C.-Y.; Song, X.-J.; Zou, Y.; Huang, Z.-C.; Li, X.; Tian, F.; Wang, L.-X. *Chem. Commun.* **2020**, *56*, 14825.
- [10] (a) Li, W.; Yuan, H.; Liu, Z.; Zhang, Z.; Cheng, Y.; Li, P. *Adv. Synth. Catal.* **2018**, *360*, 2460.
- (b) Liu, Q.; Li, S.; Chen, X.-Y.; Rissanen, K.; Enders, D. *Org. Lett.* **2018**, *20*, 3622.
- (c) Jiang, F.; Yuan, F.-R.; Jin, L.-W.; Mei, G.-J.; Shi, F. *ACS Catal.* **2018**, *8*, 10234.
- (d) Sun, M.; Ma, C.; Zhou, S.-J.; Lou, S.-F.; Xiao, J.; Jiao, Y.; Shi, F. *Angew. Chem., Int. Ed.* **2019**, *58*, 8703.
- [11] (a) Xiao, Y.; Wang, Y.; Zhou, Z. *Chin. J. Org. Chem.* **2019**, *39*, 2203 (in Chinese).
- (肖园园, 王有名, 周正洪. 有机化学, **2019**, *39*, 2203.)
- (b) Xiao, Y.; Jiang, R.; Wang, Y.; Zhou, Z. *Adv. Synth. Catal.* **2018**, *360*, 1961.
- (c) Zhang, H.; Wang, Y.; Zhou, Z. *Tetrahedron* **2018**, *74*, 6071.
- (d) Chen, S.; Wang, Y.; Zhou, Z. *J. Org. Chem.* **2016**, *81*, 11432.
- (e) Zheng, Y.; Cui, L.; Wang, Y.; Zhou, Z. *J. Org. Chem.* **2016**, *81*, 4340.
- (f) Wang, Y.; Pan, J.; Jiang, R.; Wang, Y.; Zhou, Z. *Adv. Synth. Catal.* **2016**, *358*, 195.
- (g) Zheng, Y.; Wang, Y.; Zhou, Z. *Chem. Commun.* **2015**, *51*, 11652.
- (h) Wu, L.; Wang, Y.; Song, H.; Tang, L.; Zhou, Z.; Tang, C. *ChemCatChem* **2014**, *6*, 649.
- (i) Wu, L.; Wang, Y.; Song, H.; Tang, L.; Zhou, Z.; Tang, C. *Chem. Asian J.* **2013**, *8*, 2204.
- (j) Wu, L.; Wang, Y.; Song, H.; Tang, L.; Zhou, Z.; Tang, C. *Adv. Synth. Catal.* **2013**, *355*, 1053.
- (k) Lu, A.; Hu, K.; Wang, Y.; Song, H.; Zhou, Z.; Fang, J.; Tang, C. *J. Org. Chem.* **2012**, *77*, 6208.
- [12] CCDC-2086168 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

(Cheng, F.)