串联氧化碳-氢键氧官能化和噁唑环化构建 2-酰基苯并噁唑

涂 志 赵保丽 万结平 王超莉* 刘云云*

(江西师范大学化学化工学院 南昌 330022)

摘要 通过铜催化 *N*-(邻羟基苯基)苯乙酰胺的一锅串联亚甲基碳-氢键氧官能化以及噁唑环化, 实现了有用的 2-酰基苯并噁唑类化合物的合成. 除了具有基于一步多键转化的独特步骤经济性, 以同样原料选择性可控的 2-亚甲基苯并噁唑合成以及在合成苯并噁唑方面的扩展应用也是本工作的重要优势.

关键词 串联; 碳-氢键氧官能化; 环化; 2-酰基苯并噁唑; 选择性

Aerobic Construction of 2-Acyl Benzoxazole by Tandem C—H Oxygenation and Oxazole Ring Formation

Tu, Zhi Zhao, Baoli Wan, Jieping Wang, Chaoli* Liu, Yunyun*

(College of Chemistry and Chemical Engineering, Jiangxi Normal University, Nanchang 330022)

Abstract The synthesis of useful 2-acyl benzoxazoles has been realized via the tandem methylene C—H oxygenation and oxazole annulation of *N*-(*o*-hydroxyphenyl)phenylacetamides via copper catalysis in one-pot fashion. Besides featuring the incomparable step-economy resulting from the multi-step transformation, the tunable synthesis of diverse 2-methylene functionalized benzoxazoles using identical starting materials as well as the expanded synthesis of corresponding benzothiazoles is also attractive advantage of the work.

Keywords tandem; C—H oxygenation; annulation; 2-acyl benzoxazole; selectivity

1 Introduction

As fundamental heterocyclic moiety, benzoxazole has been observed as the core structure in a vast number of functional organic molecules, including medicines, agrochemicals, materials as well as biologically relevant candidates.^[1] Particularly, as typical functionalized benzoxazoles, the 2-acyl benzoxazoles have been identified to possess enriched and special biological profiles, such as the inhibitory activity against fatty acid amide hydrolase,^[2] *sn*-1-diacylglycerol lipase α ,^[3] human cysteine proteases,^[4] channel-activating protease prostasin,^[5] as well as anti-bacteria activity.^[6] In addition, the carbonyl fragment in C2-site of the 2-acyl benzoxazoles endows them with additional application in the synthesis of diverse benzoxazole derivatives with various substructures.^[7] Accordingly, the synthesis of 2-acyl benzoxazoles has attracted significant attention, and a plethora of different tactics have been consequently developed targeting on the synthesis of 2-acyl benzoxazoles.^[8] Specifically, the C2-coupling car-

bonylation of benzoxazoles,^[9] the 2-acylation reactions of benzoxazoles with different acylating reagents,^[10] the benzoxazole annulation of proper substrates containing the prior installed acyl structure,^[11] the oxygenation of the 2-methylene/methyl benzoxazoles,^[12] the tandem benzoxazole annulation and unsaturated carbon-carbon bond oxygenation,^[13] and the coupling reactions using 2-metalated or trichloro-functionalized benzoxazoles^[14] constitute the major methods towards the synthesis of 2-acyl benzoxazoles.

By surveying these presently known works on 2-acyl benzoxazole synthesis, it can be found that one or more of the operations, including the use of prior prepared benzoxazole as substrate, the reliance on prior installed acyl structure and/or the transformation of an active functional groups for the acyl C=O generation, are yet the mandatory conditions. The additional works required in preparing such functionalized starting materials have thus undermined the step economy and the efficient application of these methods. In this context, developing practical approaches fea-

* Corresponding authors. E-mail: chemliuyunyun@jxnu.edu.cn; wangchaoli123@163.com

Received September 18, 2021; revised October 14, 2021; published online October 20, 2021.

Project supported by the Natural Science Foundation of Jiangxi Province (No. 20202BABL203008).

江西省自然科学基金(No. 20202BABL203008)资助项目.

turing simple and readily available starting materials by means of tandem transformations of hetero ring annulation and the acyl generation is urgently desired.

With the inspiration of our investigation on the simple *o*-aminophenol participated organic reactions and related works from other groups,^[15] it is envisaged that devising a method for the straightforward synthesis via one-pot C—H oxygenation and benzoxazole annulation is possible by using easily available *N*-(*o*-hydroxyphenyl)phenylacetamides as substrates. Herein, we report a new method toward the synthesis of 2-acyl benzoxazoles via step efficient one-pot C—H bond oxygenation^[16] and heterocycle formation under aerobic atmosphere.^[17] In addition, the tunable synthesis of various 2-methylene functionalized benzoxazoles has also been realized by simply modifying the operation to metal-free reaction conditions.

2 Results and discussion

To start the research work, the phenylacetamide **1a** was selected for the transformation under different reaction conditions. Then a series of different solvents were independently employed as the reaction medium, and the 2-acyl benzoxazoles **2a** and **3a** were both observed in the entry using AcOH (Entry 1, Table 1). Other solvents such as dimethyl sulfoxide (DMSO), *N,N*-dimethylformamide (DMF), EtOH or toluene were not practical for such reactions (Entries 2~5, Table 1). Delightfully, evidently improved yield of product **2a** was observed when CuSO₄ was used in the reaction (Entry 6, Table 1). Encouraged by the catalytic activity of copper reagent, an array of different copper salts, including Cu(OAc)₂, CuI and CuBr were then screened, respectively. From these results, CuBr was identified as the ideal copper species for the selective formation of **2a** (Entries 7~9, Table 1). The subsequent examination on the effect of the CuBr loading led to the observation that 30 mol% CuBr was proper for the synthesis of 2-acyl benzoxazole (Entries 10~12). On the other hand, varying the reaction temperature was not helpful for the synthesis of **2a** (Entries 13~14, Table 1). Following the selective conditions for **2a** formation, we then made efforts in finding conditions for the synthesis of 2-methylene benzoxazole (**3a**). Interestingly, simply employing non-metal acid *p*-toluenesulfonic acid (*p*-TSA) as catalyst enabled the selective synthesis of **3a** with excellent yield at the loading of 40 mol% (Entries 15~18, Table 1). Finally, conducting the reaction under nitrogen atmosphere with the catalysis of CuBr gave product **2a** with only 11% yield possibly resulting from the small amount of oxygen resolved in the solvent, and **3a** was simultaneous given as the major product (Entry 19, Table 1). This result suggested that air was the most possible oxygen source in the newly generated acyl group in **2a**. The reaction under oxygen was also examined, but no improved yield of **2a** was provided (Entry 20, Table 1).

To verify the application scope of the catalytic transformations, the reactions targeting on the synthesis of 2-acyl

Table 1 Optimization on the conditions for 2-acyl benzoxazole synthesis^a

Entry	Catalyst	Solvent	Yield/% of 2a	Yield/% of 3a
1	—	AcOH	27	49
2	—	DMSO	0	Trace
3	—	DMF	0	Trace
4	—	EtOH	0	15
5	—	Toluene	0	0
6	CuSO ₄	AcOH	45	37
7	Cu(OAc) ₂	AcOH	74	11
8	CuI	AcOH	77	Trace
9	CuBr	AcOH	80	Trace
10 ^c	CuBr	AcOH	69	18
11 ^d	CuBr	AcOH	78	Trace
12 ^e	CuBr	AcOH	78	Trace
13 ^{d,f}	CuBr	AcOH	56	17
14 ^{d,g}	CuBr	AcOH	75	Trace
15	<i>p</i> -TSA	AcOH	Trace	74
16 ^h	<i>p</i> -TSA	AcOH	Trace	76
17 ⁱ	<i>p</i> -TSA	AcOH	Trace	80
18 ^j	<i>p</i> -TSA	AcOH	Trace	79
19 ^{d,k}	CuBr	AcOH	11	59
20 ^{d,l}	CuBr	AcOH	77	Trace

^a General conditions: **1a** (0.2 mmol), catalyst (0.2 mmol), solvent (2 mL), stirred at 120 °C for 12 h in sealed tube. ^b Isolated yield. ^c CuBr (0.04 mmol). ^d CuBr (0.06 mmol). ^e CuBr (0.08 mmol). ^f Reaction at 100 °C. ^g Reaction at 140 °C. ^h *p*-TSA (0.06 mmol). ⁱ *p*-TSA (0.08 mmol). ^j *p*-TSA (0.09 mmol). ^k Reaction under N₂. ^l Reaction under O₂.

benzoxazoles **2** were conducted by using substrates **1** containing various substituents in different sites. When Ar was fixed as phenyl group, a series of different functional groups such as methyl, F, Cl and Br, different positions of *o*-aminophenol fragment could be tolerated to afford the titled products (**2a**~**2g**, Table 2). The present data indicated that the electron withdrawing halogen substitution in *o*-aminophenol backbone led to the synthesis of corresponding product with slightly lower yields (**2e**~**2g**, Table 2). On the other hand, more diverse products were synthesized by varying the functional group in the phenyl ring of the phenylacetamide section. It was interesting that the electron withdrawing group (EWG) in phenyl ring enabled the product formation with better yields than those ones containing electron donating group (EDG) in this substructure (**2h**~**2i**, **2k**~**2l**, **2n**~**2o** vs **2j**, **2m** and **2p**, Table 2). Notably, as examples of heteroaryl-based acetamides, the 2-(thiophen-2-yl)acetamide derived substrates were also successfully transformed into corresponding thiophen-2-yl functionalized 2-acyl benzoxazoles with moderate yields (**2a**~**2s**, Table 2).

Based on the previously acquired data on the selective formation of 2-methylene functionalized benzoxazole via the catalysis of *p*-TSA, the scope on the synthesis of such

Table 2 Scope on the selective synthesis of 2-acyl benzoxazoles^{a,b}

^a General conditions: **1** (0.2 mmol), CuBr (0.06 mmol) in AcOH (2 mL), stirred at 120 °C for 12 h. ^b Isolated yield. ^c Yield from 2 mmol scale reaction under open air.

benzoxazoles was also investigated by employing starting

materials **1** containing diverse substructures. Delightfully, satisfactory results on the synthesis of 2-methylene benzoxazoles **3** were obtained from this section. First, the aryl acetamides substrates bearing enriched structural diversity in both the phenyl ring of the benzoxazole skeleton (**3a**~**3g**, Table 3), the C2-methylene site (**3h**~**3j**, Table 3) as well as productions containing functionalized groups in both phenyls (**3k**~**3p**, Table 3) could be accessed with moderate to excellent yields. In addition, the 2-(thiophen-2-ylmethyl)benzoxazoles could also be practically synthesized via corresponding heteroaryl-based substrates. More notably, this selective synthesis was found to tolerate the synthesis of 3-alkyl functionalized 2-methylene benzoxazoles using corresponding alkyl amides as substrates (**3q**~**3r**, Table 3).

Interestingly, when the acrylamide **4** was used as substrate in AcOH, the product **6a** functionalized with an ester structure in the C2-side chain of the benzoxazole was provided wherein an oxa-Michael addition of the acetic acid to the C=C double bond was involved. Later, such unprecedented benzoxazole derivatives **6** were synthesized by employing different linear carboxylic acids via this interesting tandem protocol (Scheme 1). However, analogous reaction was not observed when benzoic acid was used to react with **4** in DMSO at 120 °C.

Scheme 1 Selective synthesis of ester-functionalized 2-methylene benzoxazoles

Encouraged by the results in the selective synthesis of products **2** and **3**, the reactions of *N*-(2-mercaptophenyl)-2-phenylacetamide (**7**) were then carried out under the reaction conditions employed in Tables 2 and 3, respectively. To our delight, the 2-acyl benzothiazole **8** and 2-benzyl benzothiazole **9** were both smoothly afforded (Scheme 2), indicating the expanded scope of the present methods in the synthesis of corresponding benzothiazole derivatives.

Scheme 2 Expanded synthesis of 2-acyl and 2-methylene benzothiazole

Subsequently, for the sake of probing the possible reaction mechanism of providing 2-acyl benzoxazoles, a series of control experiments were carried out. First, subjecting

Table 3 Scope on the selective synthesis of 2-methylene benzoxazoles^{a,b}

^a General conditions: **1** (0.2 mmol), CuBr (0.06 mmol) in AcOH (2 mL), stirred at 120 °C for 12 h. ^b Isolated yield.

methylene benzoxazole **3a** to the standard conditions of

CuBr catalysis provided 2-benzoyl benzoxazole (**2a**) with excellent yield (Eq. 1), supporting that the C—H oxygenation might possibly take place after the benzoxazole ring formation. In addition, employing 2,2,6,6-tetramethylpiperidinoxy (TEMPO) under the standard conditions for the synthesis of **2a** and **3a** was also performed, respectively. While the synthesis of **2a** was clearly suppressed (Eq. 2), no evident inhibition of TEMPO to the formation of **3a** was observed (Eq. 3). The results indicated that single electron transfer (SET) process might be involved in the synthesis of products **2**, and not during the formation of products **3**.

According to the results given by the control experiment in Eqs. 1~3 and Table 1 (Entry 19). The possible mechanism for the reactions providing benzoxazoles **2** and **3** is proposed (Scheme 3). Initially, in the presence of proton or Lewis acid, the carbonyl group in the amide substrate is activated to enable the intramolecular addition of the hy-

Scheme 3 Possible reaction mechanisms for 2-methylene and 2-acyl benzoxazole formation

droxyl, which provides intermediate **A**. The simple dehydration from **A** leads to the production of product **3a**. The generation of water in this step may bring slight negative effect to the subsequent production of 2-acylbenzoxazole. Subsequently, in the presence of Cu(I) catalyst, the single electron transfer (SET) from Cu(I) to oxygen affords oxygen radical anion and Cu(II). The oxygen free radical can couple the methylene C—H bond in **3a** to give anion intermediate **B** and the hydrogen free radical. The reaction between the hydrogen free radical and Cu(II) provides proton and regenerates Cu(I) catalyst. The incorporation of **B** with the proton then gives intermediate **C**. The elimination of water initiated by the O—O bond cleavage in **C** yields the 2-benzoyl benzoxazole **2a**.

3 Conclusions

In conclusion, by employing simple and readily available *N*-(*o*-hydroxyphenyl) phenylacetamides as starting materials, the synthesis of 2-acyl benzoxazoles via Cu(I)-catalyzed tandem benzoxazole annulation and C(sp³)—H oxygenation has been accomplished for the first time. In addition, simply modifying the reaction conditions to non-metal operation or using acrylamide-based substrates enables the selective synthesis of 2-methylene and 2-ester elaborated alkyl benzoxazoles, respectively. Besides the selective reaction pathways, the present method for the synthesis of 2-acyl benzoxazole synthesis is of special advantages in using air as oxygen atom source, low cost copper catalyst as well as the efficient transformation on the stable C(sp³)—H bonds, which make the synthetic approach attractive for the preparation of these useful heterocyclic scaffolds with environmentally benign operations.

4 Experimental section

4.1 General information

All reactions were performed under air atmosphere in sealed tube. Amides **1**, **4** and **7** were prepared following literature procedures.^[18] All other chemicals and solvents used in the experiments were obtained from commercial sources and used directly without further treatment. ¹H NMR and ¹³C NMR were recorded in 400 MHz apparatus. The frequency for ¹H NMR and ¹³C NMR test were 400 and 100 MHz, respectively. The chemical shifts were reported using TMS as internal standard. HRMS results were acquired under ESI model in a spectrometer equipped with TOF analyzer. Melting points were rested in the X-4A apparatus without prior temperature correction.

4.2 General procedure for the synthesis of 2-acyl benzoxazoles **2** and 2-benzoyl benzothiazole **8**

In a 25 mL reaction tube equipped with stirring bar were charged with amide **1** or **7** (0.2 mmol), CuBr (0.06 mmol) and AcOH (2 mL). The tube was then sealed with Teflon cap and subjected to oil bath heating at 120 °C and stirring for 12 h. The vessel was then allowed to cool down to room temperature, and 10 mL of water was added. The hetero-

geneous mixture was extracted with ethyl acetate (10 mL × 3). The combined organic phase was dried over Na₂SO₄. After filtration, the acquired solution was collected and the solvent was removed at reduced pressure. The residue was subjected to silica gel column chromatography to give pure products by using mixed petroleum ether and ethyl acetate (*V* : *V* = 100 : 1) as eluent.

4.3 Procedure for the synthesis of **2a** at 2 mmol scale reaction

In a 50 mL reaction tube equipped with stirring bar were charged with amide **1** (2 mmol), CuBr (0.6 mmol) and AcOH (10 mL). The tube was then sealed with Teflon cap and subjected to oil bath heating at 120 °C and stirring for 12 h. The vessel was then allowed to cool down to room temperature, and 30 mL water was added. The heterogeneous mixture was extracted with ethyl acetate (30 mL × 3). The combined organic phase was dried over Na₂SO₄. After filtration, the acquired solution was collected and the solvent was removed at reduced pressure. The residue was subjected to silica gel column chromatography to give pure products by using mixed petroleum ether and ethyl acetate (*V* : *V* = 100 : 1) as eluent.

Benzo[*d*]oxazol-2-yl(phenyl)methanone (**2a**): Yield 78%, 35 mg. Yellow solid, m.p. 59~60 °C (lit.^[18] 60~61 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.55 (d, *J* = 7.6 Hz, 2H), 7.94 (d, *J* = 7.8 Hz, 1H), 7.73~7.65 (m, 2H), 7.55 (q, *J* = 7.6 Hz, 3H), 7.46 (t, *J* = 7.4 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 180.5, 157.1, 150.4, 140.8, 135.0, 134.3, 131.0, 128.6, 128.4, 125.7, 122.4, 111.9.

(7-Methylbenzo[*d*]oxazol-2-yl)(phenyl)methanone (**2b**): Yield 53%, 25 mg. Yellow solid, m.p. 82~83 °C (lit.^[11b] 86~88 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.6~8.56 (m, 2H), 7.68 (t, *J* = 7.4 Hz, 1H), 7.56 (t, *J* = 7.6 Hz, 2H), 7.50 (d, *J* = 8.2 Hz, 1H), 7.41 (t, *J* = 7.8 Hz, 1H), 7.24 (d, *J* = 7.4 Hz, 1H), 2.71 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 180.5, 156.5, 150.3, 140.3, 135.1, 134.2, 133.2, 131.1, 128.6, 128.2, 125.9, 109.0, 16.5.

(6-Methylbenzo[*d*]oxazol-2-yl)(phenyl)methanone (**2c**):^[10b] Yield 66%, 31 mg. Yellow solid, m.p. 76~77 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.54 (m, 2H), 7.81~7.78 (m, 1H), 7.69~7.65 (m, 1H), 7.55 (t, *J* = 7.8 Hz, 2H), 7.48 (s, 1H), 7.29 (d, *J* = 1.4 Hz, 1H), 2.54 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 180.5, 156.9, 150.8, 139.5, 138.7, 135.2, 134.1, 131.0, 128.6, 127.4, 121.7, 111.7, 22.1.

(5-Methylbenzo[*d*]oxazol-2-yl)(phenyl)methanone (**2d**): Yield 62%, 29 mg. Yellow solid, m.p. 91~92 °C (lit.^[11b] 91~93 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.53 (d, *J* = 7.2 Hz, 2H), 7.72~7.65 (m, 2H), 7.56 (t, *J* = 7.8 Hz, 3H), 7.35 (d, *J* = 8.4 Hz, 1H), 2.51 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 180.6, 157.3, 148.7, 141.0, 135.8, 135.1, 134.2, 131.0, 129.9, 128.6, 121.9, 111.2, 21.5.

(5-Fluorobenzo[*d*]oxazol-2-yl)(phenyl)methanone (**2e**): Yield 40%, 19 mg. Yellow solid, m.p. 86~88 °C (lit.^[11b] 88~90 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.54 (d, *J* = 7.4 Hz, 2H), 7.73~7.65 (m, 2H), 7.64~7.55 (m, 3H), 7.30 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 180.2, 160.6

(d, $^1J_{\text{CF}}=242\text{Hz}$), 158.6, 146.8, 141.4, 134.8, 134.5, 131.0, 128.7, 116.7 (d, $^2J_{\text{CF}}=27\text{Hz}$), 112.4 (d, $^3J_{\text{CF}}=10\text{Hz}$), 108.2 (d, $^2J_{\text{CF}}=25\text{Hz}$).

(5-Chlorobenzo[d]oxazol-2-yl)(phenyl)methanone (**2f**): Yield 53%, 27 mg. Yellow solid, m.p. 99~100 °C (lit.^[11b] 98~100 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.56~8.48 (m, 2H), 7.94~7.88 (m, 1H), 7.69 (t, $J=7.8\text{ Hz}$, 1H), 7.63 (d, $J=8.8\text{ Hz}$, 1H), 7.57 (t, $J=7.6\text{ Hz}$, 2H), 7.51 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 180.1, 158.1, 148.9, 141.7, 134.7, 134.5, 131.3, 131.0, 128.9, 128.7, 122.1, 112.7.

(5-Bromobenzo[d]oxazol-2-yl)(phenyl)methanone (**2g**): Yield 56%, 34 mg. Yellow solid, m.p. 110~111 °C (lit.^[11b] 112~114 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.57~8.50 (m, 2H), 8.09 (d, $J=1.6\text{ Hz}$, 1H), 7.73~7.65 (m, 2H), 7.62~7.56 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 180.1, 157.9, 149.4, 142.2, 134.7, 134.5, 131.5, 131.0, 128.7, 125.2, 118.5, 113.1.

Benzo[d]oxazol-2-yl(4-fluorophenyl)methanone (**2h**):^[11b] Yield 83%, 40 mg. Yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.66 (m, 2H), 7.93 (d, $J=8.4\text{ Hz}$, 1H), 7.70 (d, $J=8.2\text{ Hz}$, 1H), 7.55 (t, $J=7.8\text{ Hz}$, 1H), 7.47 (t, $J=7.6\text{ Hz}$, 1H), 7.23 (t, $J=8.6\text{ Hz}$, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 178.7, 166.6 (d, $^1J_{\text{CF}}=256\text{Hz}$), 156.9, 150.4, 140.7, 133.9 (d, $^3J_{\text{CF}}=10\text{Hz}$), 131.4 (d, $^4J_{\text{CF}}=3\text{Hz}$), 128.5, 125.8, 122.4, 115.9 (d, $^2J_{\text{CF}}=22\text{Hz}$), 111.9.

Benzo[d]oxazol-2-yl(4-chlorophenyl)methanone (**2i**): Yield 84%, 43 mg. White solid, m.p. 97~98 °C (lit.^[11b] 100~101 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.55 (d, $J=8.6\text{ Hz}$, 2H), 7.93 (d, $J=8.0\text{ Hz}$, 1H), 7.70 (d, $J=8.2\text{ Hz}$, 1H), 7.55 (m, 3H), 7.47 (t, $J=7.4\text{ Hz}$, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 179.0, 156.8, 150.4, 141.0, 140.7, 133.3, 132.4, 129.0, 128.6, 125.8, 122.4, 111.9.

Benzo[d]oxazol-2-yl(4-methoxyphenyl)methanone (**2j**): Yield 45%, 23 mg. Yellow solid, m.p. 92~93 °C (lit.^[11b] 89~90 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.40 (d, $J=9.0\text{ Hz}$, 2H), 7.81 (m, 1H), 7.50~7.45 (m, 1H), 7.39~7.35 (m, 1H), 7.31 (d, $J=8.2\text{ Hz}$, 1H), 7.26 (s, 1H), 7.00 (d, $J=9.0\text{ Hz}$, 2H), 3.89 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 152.5, 149.9, 146.3, 131.8, 131.4, 130.4, 129.1, 126.8, 125.5, 116.1, 113.8, 55.4.

(4-Fluorophenyl)(6-methylbenzo[d]oxazol-2-yl)methanone (**2k**): Yield 51%, 26 mg. Yellow solid, m.p. 138~139 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.68~8.63 (m, 2H), 7.79 (d, $J=8.2\text{ Hz}$, 1H), 7.49 (s, 1H), 7.29 (d, $J=8.2\text{ Hz}$, 1H), 7.23 (t, $J=8.6\text{ Hz}$, 2H), 2.55 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 178.7, 166.5 (d, $^1J_{\text{CF}}=256\text{Hz}$), 156.7, 150.8, 139.7, 138.6, 133.9 (d, $^3J_{\text{CF}}=9\text{Hz}$), 131.5 (d, $^4J_{\text{CF}}=3\text{Hz}$), 127.4, 121.7, 115.8 (d, $^2J_{\text{CF}}=22\text{Hz}$), 111.7, 22.1. ESI-HRMS calcd for $\text{C}_{15}\text{H}_{11}\text{FNO}_2$ $[\text{M}+\text{H}]^+$ 256.0768, found 256.0777.

(4-Chlorophenyl)(6-methylbenzo[d]oxazol-2-yl)methanone (**2l**): Yield 61%, 33 mg. Yellow solid, m.p. 118~119 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.53 (d, $J=8.4\text{ Hz}$, 2H), 7.78 (d, $J=8.2\text{ Hz}$, 1H), 7.52 (d, $J=8.4\text{ Hz}$, 2H), 7.47 (s, 1H), 7.28 (d, $J=8.4\text{ Hz}$, 1H), 2.54 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 179.0, 156.6, 150.8, 140.9, 139.8,

138.6, 133.4, 132.4, 128.9, 127.5, 121.7, 111.7, 22.1; ESI-HRMS Calcd for $\text{C}_{15}\text{H}_{11}\text{ClNO}_2$ $[\text{M}+\text{H}]^+$ 272.0473, found 272.0486.

(4-Methoxyphenyl)(6-methylbenzo[d]oxazol-2-yl)methanone (**2m**): Yield 54%, 29 mg. Yellow solid, m.p. 134~135 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.60 (d, $J=9.0\text{ Hz}$, 2H), 7.78 (d, $J=8.2\text{ Hz}$, 1H), 7.48 (s, 1H), 7.28~7.25 (m, 1H), 7.03 (d, $J=9.0\text{ Hz}$, 2H), 3.91 (s, 3H), 2.54 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 178.8, 164.6, 157.1, 150.7, 139.1, 138.7, 133.5, 128.1, 127.2, 121.5, 113.9, 111.7, 55.6, 22.1. ESI-HRMS calcd for $\text{C}_{16}\text{H}_{14}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 268.0968, found 268.0981.

(5-Chlorobenzo[d]oxazol-2-yl)(4-fluorophenyl)methanone (**2n**): Yield 61%, 34 mg. Yellow solid, m.p. 155~156 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.65 (m, 2H), 7.92 (d, $J=1.8\text{ Hz}$, 1H), 7.65 (d, $J=8.8\text{ Hz}$, 1H), 7.53 (m, 1H), 7.28~7.23 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 178.3, 166.7 (d, $^1J_{\text{CF}}=257\text{Hz}$), 157.9, 148.9, 141.6, 134.0 (d, $^3J_{\text{CF}}=9\text{Hz}$), 131.4, 131.0 (d, $^3J_{\text{CF}}=8\text{Hz}$), 129.0, 122.1, 116.1 (d, $^2J_{\text{CF}}=22\text{Hz}$), 112.7. ESI-HRMS calcd for $\text{C}_{14}\text{H}_8\text{ClFNO}_2$ $[\text{M}+\text{H}]^+$ 276.0222, found 276.0233.

(5-Chlorobenzo[d]oxazol-2-yl)(4-chlorophenyl)methanone (**2o**): Yield 63%, 37 mg. Yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.53 (d, $J=7.0\text{ Hz}$, 2H), 7.91 (d, $J=1.8\text{ Hz}$, 1H), 7.64 (d, $J=8.8\text{ Hz}$, 1H), 7.56~7.50 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 178.6, 157.8, 148.9, 141.6, 141.3, 133.0, 132.4, 131.4, 129.1, 129.1, 122.1, 112.7. ESI-HRMS calcd for $\text{C}_{14}\text{H}_8\text{Cl}_2\text{NO}_2$ $[\text{M}+\text{H}]^+$ 291.9927, found 291.9946.

(5-Chlorobenzo[d]oxazol-2-yl)(4-methoxyphenyl)methanone (**2p**): Yield 37%, 21 mg. Yellow solid, m.p. 157~158 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.34 (d, $J=8.8\text{ Hz}$, 2H), 7.73 (d, $J=2.2\text{ Hz}$, 1H), 7.35 (m, 1H), 7.19 (s, 1H), 6.93 (d, $J=8.8\text{ Hz}$, 2H), 3.83 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.8, 150.7, 144.8, 132.3, 131.6, 130.6, 130.2, 128.5, 126.4, 117.1, 113.9, 55.5. ESI-HRMS calcd for $\text{C}_{15}\text{H}_{11}\text{ClNO}_3$ $[\text{M}+\text{H}]^+$ 288.0422, found 288.0433.

Benzo[d]oxazol-2-yl(thiophen-2-yl)methanone (**2q**):^[11a] Yield 51%, 23 mg. Yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.74 (dd, $J=3.8, 1.2\text{ Hz}$, 1H), 7.96~7.92 (m, 1H), 7.85 (dd, $J=4.8, 1.2\text{ Hz}$, 1H), 7.73~7.68 (m, 1H), 7.54 (m, 1H), 7.47 (m, 1H), 7.28~7.25 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.1, 156.8, 150.6, 140.7, 140.6, 137.6, 137.0, 128.7, 128.4, 125.8, 122.3, 111.9.

(6-Methylbenzo[d]oxazol-2-yl)(thiophen-2-yl)methanone (**2r**): Yield 49%, 24 mg. Yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.74 (d, $J=3.6\text{ Hz}$, 1H), 7.83 (m, 2H), 7.50 (s, 1H), 7.31~7.26 (m, 2H), 2.55 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.2, 151.0, 140.7, 139.5, 138.6, 137.4, 136.7, 128.6, 127.4, 121.6, 111.8, 22.1. ESI-HRMS calcd for $\text{C}_{13}\text{H}_{10}\text{Cl}_2\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$ 244.0427, found 244.0441.

(5-Chlorobenzo[d]oxazol-2-yl)(thiophen-2-yl)methanone (**2s**): Yield 47%, 25 mg. Yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.72 (dd, $J=3.8, 1.0\text{ Hz}$, 1H), 7.92 (d, $J=1.8\text{ Hz}$, 1H), 7.87 (dd, $J=4.8, 1.0\text{ Hz}$, 1H), 7.64 (d, $J=8.8\text{ Hz}$, 1H), 7.51 (dd, $J=8.8, 2.0\text{ Hz}$, 1H), 7.27 (dd, $J=3.2, 1.6\text{ Hz}$, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.7, 157.8,

149.1, 141.7, 140.4, 137.7, 137.3, 131.3, 128.8, 128.8, 122.0, 112.7. ESI-HRMS calcd for $C_{12}H_7ClNO_2S [M+H]^+$ 263.9881, found 263.9891.

Benzo[d]thiazol-2-yl(phenyl)methanone (**8**): Yield 42%, 20 mg. Yellow solid, m.p. 97~98 °C (lit.^[11b] 98~99 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 8.61~8.53 (m, 2H), 8.27~8.22 (m, 1H), 8.02 (m, 1H), 7.70~7.65 (m, 1H), 7.60~7.52 (m, 4H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 185.4, 167.1, 153.9, 137.0, 135.0, 133.9, 131.3, 128.5, 127.6, 126.9, 125.7, 122.2.

4.4 General procedure for the synthesis of 2-methylene benzoxazoles **3**, ester functionalized benzoxazoles **6** and 2-benzothiazole **9**

In a 25 mL reaction tube equipped with stirring bar were charged with amide **1**, **4** or **7** (0.2 mmol), *p*-TSA (0.08 mmol for the synthesis of **3** and **9** only) and RCO_2H (2 mL, AcOH in the synthesis of **3** and **9**). The tube was then sealed with Teflon cap and subjected to oil bath heating at 120 °C with stirring. After the reaction time of 12 h, the vessel was allowed to cool down to room temperature, and 10 mL of water was added. The heterogeneous mixture was extracted with ethyl acetate (10 mL $\times$ 3). The combined organic phase was dried over Na_2SO_4 . After filtration, the acquired solution was collected and the solvent was removed at reduced pressure. The acquired residue was subjected to silica gel column chromatography to give pure products by using mixed petroleum ether and ethyl acetate ($V:V=80:1$) as eluent.

2-Benzylbenzo[d]oxazole (**3a**):^[19a] Yield 80%, 33 mg. Yellow liquid. 1H NMR (400 MHz, $CDCl_3$) δ : 7.71~7.66 (m, 1H), 7.47~7.42 (m, 1H), 7.35 (m, 4H), 7.30~7.23 (m, 3H), 4.26 (s, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 165.2, 151.1, 141.3, 134.8, 129.0, 128.9, 127.3, 124.7, 124.2, 119.8, 110.5, 35.3.

2-Benzyl-7-methylbenzo[d]oxazole (**3b**): Yield 71%, 32 mg. White liquid. 1H NMR (400 MHz, $CDCl_3$) δ : 7.37 (d, $J=7.0$ Hz, 2H), 7.31 (t, $J=7.4$ Hz, 2H), 7.25 (d, $J=7.4$ Hz, 2H), 7.15 (t, $J=7.8$ Hz, 1H), 7.07 (d, $J=7.4$ Hz, 1H), 4.27 (s, 2H), 2.61 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 164.3, 150.3, 140.5, 135.1, 130.2, 129.0, 128.8, 127.2, 124.8, 124.4, 107.8, 35.3, 16.6. ESI-HRMS calcd for $C_{15}H_{14}NO [M+H]^+$ 224.1071, found 224.1084.

2-Benzyl-6-methylbenzo[d]oxazole (**3c**): Yield 67%, 30 mg. Yellow solid, m.p. 38~39 °C (lit.^[19a] 35~36 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 7.55 (d, $J=8.0$ Hz, 1H), 7.39~7.30 (m, 4H), 7.29~7.25 (m, 2H), 7.10 (d, $J=8.0$ Hz, 1H), 4.25 (s, 2H), 2.45 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 139.0, 134.9, 129.0, 128.8, 127.3, 125.4, 119.1, 110.7, 35.3, 21.7.

2-Benzyl-5-methylbenzo[d]oxazole (**3d**): Yield 63%, 28 mg. White solid, m.p. 49~50 °C (lit.^[19b] 49~51 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 7.47 (s, 1H), 7.34 (m, 5H), 7.28~7.24 (m, 1H), 7.08 (d, $J=8.0$ Hz, 1H), 4.25 (s, 2H), 2.44 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 165.3, 141.5, 134.9, 134.0, 129.0, 128.8, 127.3, 125.8, 119.7, 109.8, 35.3, 21.4.

2-Benzyl-5-fluorobenzo[d]oxazole (**3e**): Yield 53%, 24 mg. Yellow solid, m.p. 46~47 °C (lit.^[19a] 48~49 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 7.35 (m, 6H), 7.30~7.25 (m, 1H), 7.00 (m, 1H), 4.24 (s, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 167.1, 159.9 (d, $^1J_{CF}=239$ Hz), 147.4, 142.1 (d, $^3J_{CF}=13$ Hz), 134.5, 128.9 (d, $^3J_{CF}=13$ Hz), 127.4, 112.3 (d, $^2J_{CF}=26$ Hz), 110.7 (d, $^3J_{CF}=10$ Hz), 106.3 (d, $^2J_{CF}=25$ Hz), 35.3.

2-Benzyl-5-chlorobenzo[d]oxazole (**3f**):^[19b] Yield 63%, 31 mg. Yellow solid, m.p. 60~61 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.67~7.64 (m, 1H), 7.37~7.32 (m, 5H), 7.31~7.27 (m, 1H), 7.26~7.23 (m, 1H), 4.25 (s, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 166.7, 142.5, 134.4, 129.7, 129.0, 128.9, 127.5, 125.0, 119.9, 111.2, 35.3.

2-Benzyl-5-bromobenzo[d]oxazole (**3g**):^[19b] Yield 58%, 33 mg. Yellow solid, m.p. 52~53 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.84~7.80 (m, 1H), 7.40 (m, 1H), 7.37~7.31 (m, 5H), 7.31~7.26 (m, 1H), 4.26 (s, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 143.0, 134.4, 129.0, 128.9, 127.8, 127.5, 122.9, 117.0, 111.7, 35.3.

2-(4-Fluorobenzyl)benzo[d]oxazole (**3h**): Yield 78%, 35 mg. Yellow solid, m.p. 49~50 °C (lit.^[19a] 49~51 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 7.71~7.66 (m, 1H), 7.45 (m, 1H), 7.34 (m, 2H), 7.31~7.26 (m, 2H), 7.02 (t, $J=8.6$ Hz, 2H), 4.23 (s, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 164.9, 162.1 (d, $^1J_{CF}=244$ Hz), 151.0, 141.3, 130.6 (d, $^3J_{CF}=8$ Hz), 130.5 (d, $^4J_{CF}=3$ Hz), 124.8, 124.3, 119.9, 115.7 (d, $^2J_{CF}=22$ Hz), 110.5, 34.4.

2-(4-Chlorobenzyl)benzo[d]oxazole (**3i**): Yield 86%, 42 mg. White solid, m.p. 73~74 °C (lit.^[19b] 78~80 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 7.68 (m, 1H), 7.46 (m, 1H), 7.31 (d, $J=3.8$ Hz, 5H), 7.29 (d, $J=4.2$ Hz, 1H), 4.23 (s, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 164.6, 141.2, 133.3, 133.2, 130.4, 129.0, 124.9, 124.3, 119.9, 110.5, 34.6.

2-(4-Methoxybenzyl)benzo[d]oxazole (**3j**): Yield 57%, 27 mg. Yellow solid, m.p. 40~41 °C (lit.^[20] 43~45 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 7.70~7.66 (m, 1H), 7.47~7.43 (m, 1H), 7.32~7.26 (m, 4H), 6.87 (d, $J=8.6$ Hz, 2H), 4.20 (s, 2H), 3.77 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 165.6, 158.9, 130.1, 126.8, 124.7, 124.2, 119.8, 114.3, 110.5, 55.3, 34.5.

2-(4-Fluorobenzyl)-6-methylbenzo[d]oxazole (**3k**): Yield 59%, 28 mg. Yellow solid, m.p. 50~51 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.54 (d, $J=8.0$ Hz, 1H), 7.36~7.30 (m, 2H), 7.25 (s, 1H), 7.11 (d, $J=8.0$ Hz, 1H), 7.05~6.98 (m, 2H), 4.20 (s, 2H), 2.44 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 164.4, 162.1 (d, $^1J_{CF}=244$ Hz), 151.3, 139.0, 135.2, 130.6 (d, $^3J_{CF}=8$ Hz), 125.5, 119.2, 115.7 (d, $^2J_{CF}=21$ Hz), 110.6, 34.4, 21.7. ESI-HRMS calcd for $C_{15}H_{13}FNO [M+H]^+$ 242.0976, found 242.0987.

2-(4-Chlorobenzyl)-6-methylbenzo[d]oxazole (**3l**): Yield 68%, 35 mg. White solid, m.p. 53~54 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.54 (d, $J=8.0$ Hz, 1H), 7.28 (s, 4H), 7.24 (s, 1H), 7.10 (d, $J=8.0$ Hz, 1H), 4.19 (s, 2H), 2.44 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 164.1, 151.3, 139.0, 135.3, 133.4, 133.3, 130.4, 129.0, 125.5, 119.2, 110.7, 34.6, 21.7. ESI-HRMS calcd for $C_{15}H_{13}ClNO [M+H]^+$ 258.0680,

found 258.0696.

2-(4-Methoxybenzyl)-6-methylbenzo[d]oxazole (**3m**): Yield 62%, 31 mg. Yellow solid, m.p. 144~145 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.54 (d, *J*=8.0 Hz, 1H), 7.30~7.24 (m, 3H), 7.09 (m, 1H), 6.87 (d, *J*=8.6 Hz, 2H), 4.18 (s, 2H), 3.77 (s, 3H), 2.44 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 165.0, 158.8, 151.3, 135.0, 139.1, 130.0, 126.9, 125.3, 119.1, 114.2, 110.6, 55.3, 34.4, 21.7. ESI-HRMS calcd for C₁₆H₁₆NO₂ [M+H]⁺ 254.1176, found 254.1190.

5-Chloro-2-(4-fluorobenzyl)benzo[d]oxazole (**3n**): Yield 70%, 37 mg. Yellow solid, m.p. 61~62 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.66 (d, *J*=1.8 Hz, 1H), 7.38 (d, *J*=8.6 Hz, 1H), 7.33 (m, 2H), 7.27 (m, 1H), 7.04 (t, *J*=8.6 Hz, 2H), 4.23 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 166.4, 162.2 (d, ¹*J*_{CF}=244 Hz), 149.6, 142.4, 130.6 (d, ³*J*_{CF}=8 Hz), 130.3 (d, ⁴*J*_{CF}=3 Hz), 129.8, 125.1, 119.9, 115.8 (d, ³*J*_{CF}=22 Hz), 111.2, 34.4. ESI-HRMS calcd for C₁₄H₁₀Cl-FNO [M+H]⁺ 262.0429, found 262.0444.

5-Chloro-2-(4-chlorobenzyl)benzo[d]oxazole (**3o**): Yield 69%, 38 mg. Yellow liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.66 (d, *J*=1.6 Hz, 1H), 7.38 (d, *J*=8.6 Hz, 1H), 7.31 (d, *J*=1.6 Hz, 4H), 7.27 (d, *J*=7.6 Hz, 1H), 4.23 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 166.1, 149.6, 142.4, 133.5, 132.8, 130.4, 129.8, 129.1, 125.2, 119.9, 111.2, 34.6. ESI-HRMS calcd for C₁₄H₁₀Cl₂NO [M+H]⁺ 278.0134, found 278.0146.

5-Chloro-2-(4-methoxybenzyl)benzo[d]oxazole (**3p**): Yield 42%, 23 mg. Yellow liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.65 (d, *J*=1.8 Hz, 1H), 7.36 (d, *J*=8.6 Hz, 1H), 7.28 (d, *J*=8.6 Hz, 2H), 7.24 (m, 1H), 6.88 (d, *J*=8.6 Hz, 2H), 4.19 (s, 2H), 3.78 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.1, 159.0, 149.6, 142.5, 130.1, 129.6, 126.4, 124.9, 119.8, 114.3, 111.2, 55.3, 34.4. ESI-HRMS calcd for C₁₅H₁₃ClNO₂ [M+H]⁺ 274.0629, found 274.0641.

2-(Thiophen-2-ylmethyl)benzo[d]oxazole (**3q**):^[19a] Yield 62%, 27 mg. Yellow liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.70 (m, 1H), 7.49 (m, 1H), 7.31 (m, 2H), 7.23 (d, *J*=5.0 Hz, 1H), 7.05 (d, *J*=3.2 Hz, 1H), 6.98 (dd, *J*=5.0, 3.6 Hz, 1H), 4.48 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 164.1, 151.0, 141.2, 135.9, 127.1, 126.9, 125.2, 124.9, 124.3, 120.0, 110.6, 29.5.

6-Methyl-2-(thiophen-2-ylmethyl)benzo[d]oxazole (**3r**): Yield 52%, 24 mg. Yellow liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.56 (d, *J*=8.0 Hz, 1H), 7.28 (s, 1H), 7.24~7.20 (m, 1H), 7.12 (d, *J*=8.0 Hz, 1H), 7.03 (d, *J*=3.0 Hz, 1H), 6.97 (dd, *J*=5.0, 3.4 Hz, 1H), 4.45 (s, 2H), 2.46 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.6, 151.3, 138.9, 136.1, 135.3, 127.1, 126.8, 125.5, 125.1, 119.3, 110.7, 29.5, 21.7. ESI-HRMS calcd for C₁₃H₁₂NOS [M+H]⁺ 230.0634, found 230.0644.

5-Chloro-2-(thiophen-2-ylmethyl)benzo[d]oxazole (**3s**): Yield 49%, 24 mg. Yellow liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.67 (d, *J*=2.0 Hz, 1H), 7.39 (d, *J*=8.6 Hz, 1H), 7.27 (dd, *J*=8.6, 2.0 Hz, 1H), 7.23 (dd, *J*=5.0, 1.0 Hz, 1H), 7.04 (d, *J*=2.8 Hz, 1H), 6.97 (dd, *J*=5.0, 3.4 Hz, 1H), 4.46 (d, *J*=0.8 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 165.5,

149.8, 142.4, 135.5, 129.8, 127.2, 127.1, 125.4, 125.2, 120.0, 111.3, 29.5. ESI-HRMS calcd for C₁₂H₉ClNOS [M+H]⁺ 250.0088, found 250.0094.

2-Propylbenzo[d]oxazole (**3t**):^[21] Yield 72%, 23 mg. Yellow solid, m.p. 170~171 °C (lit.^[21] 178~179 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.70~7.64 (m, 1H), 7.48 (m, 1H), 7.32~7.25 (m, 2H), 2.91 (t, *J*=7.4 Hz, 2H), 1.93 (m, 2H), 1.05 (t, *J*=7.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 167.2, 150.8, 141.3, 124.4, 124.0, 119.5, 110.2, 30.5, 20.2, 13.7.

2-Isobutylbenzo[d]oxazole (**3u**): Yield 68%, 24 mg. Yellow solid, m.p. 236~238 °C (lit.^[22] 240~241 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.70~7.66 (m, 1H), 7.50~7.46 (m, 1H), 7.31~7.27 (m, 2H), 2.81 (d, *J*=7.2 Hz, 2H), 2.31 (m, 1H), 1.04 (d, *J*=6.6 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 166.6, 150.8, 141.3, 124.4, 124.0, 119.5, 110.3, 37.6, 27.5, 22.4.

2-(Benzo[d]oxazol-2-yl)ethyl acetate (**6a**): Yield 31%, 13 mg. White liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.69 (m, 1H), 7.50 (m, 1H), 7.32 (m, 2H), 4.59 (t, *J*=6.4 Hz, 2H), 3.28 (t, *J*=6.4 Hz, 2H), 2.04 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.7, 163.5, 150.8, 141.2, 124.8, 124.3, 119.8, 110.4, 60.6, 28.6, 20.8. ESI-HRMS calcd for C₁₁H₁₂-NO₃ [M+H]⁺ 206.0812, found 206.0824.

2-(Benzo[d]oxazol-2-yl)ethyl propionate (**6b**): Yield 27%, 12 mg. White liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.69 (dd, *J*=6.0, 3.0 Hz, 1H), 7.50 (dd, *J*=6.2, 3.0 Hz, 1H), 7.32 (dd, *J*=6.0, 3.2 Hz, 2H), 4.59 (t, *J*=6.4 Hz, 2H), 3.29 (t, *J*=6.4 Hz, 2H), 2.32 (q, *J*=7.6 Hz, 2H), 1.11 (t, *J*=7.6 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 174.2, 163.6, 150.8, 141.1, 124.8, 124.3, 119.8, 110.4, 60.5, 28.6, 27.42, 9.0. ESI-HRMS calcd for C₁₂H₁₄NO₃ [M+H]⁺ 220.0968, found 220.0981.

2-(Benzo[d]oxazol-2-yl)ethyl butyrate (**6c**): Yield 26%, 12 mg. White liquid. ¹H NMR (400 MHz, CDCl₃) δ: 7.70 (m, 1H), 7.50 (m, 1H), 7.33 (m, 2H), 4.59 (t, *J*=6.4 Hz, 2H), 3.29 (t, *J*=6.4 Hz, 2H), 2.28 (t, *J*=7.4 Hz, 2H), 1.61 (m, 2H), 0.90 (t, *J*=7.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 173.4, 163.6, 150.8, 141.1, 124.8, 124.3, 119.7, 110.4, 60.4, 36.0, 28.6, 18.3, 13.6. ESI-HRMS calcd for C₁₃H₁₆NO₃ [M+H]⁺ 234.1125, found 234.1137.

2-Benzylbenzo[d]thiazole (**9**): Yield 49%, 22 mg. Yellow solid, m.p. 102~103 °C (lit.^[19a] 105~106 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.00 (d, *J*=8.0 Hz, 1H), 7.78 (d, *J*=7.8 Hz, 1H), 7.45 (t, *J*=7.4 Hz, 1H), 7.36 (d, *J*=6.6 Hz, 4H), 7.33~7.27 (m, 2H), 4.45 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 171.3, 153.1, 137.2, 135.6, 129.2, 128.9, 127.4, 126.0, 124.9, 122.8, 121.54, 40.7.

Supporting Information ¹H NMR and ¹³C NMR spectra of all products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] For selected reviews, see:
 - (a) Ghoshal, T.; Patel, T. M. *Future J. Pharm. Sci.* **2020**, 6, no. 94.
 - (b) Sattar, R.; Mukhtar, R.; Atif, M.; Hasnain, M.; Irfan, A. J. *Het-*

- erocycl. Chem.* **2020**, *57*, 2079.
- (c) Singh, S.; Veeraswamy, G.; Bhattarai, D.; Goo, J.-I.; Lee, K.; Choi, Y. *Asian J. Org. Chem.* **2015**, *4*, 1338.
- (c) Sun, Z.; Xu, M.; Wang, Y.; Hu, X. *Chin. J. Org. Chem.* **2020**, *40*, 4203 (in Chinese).
- (孙泽众, 徐苗, 王云侠, 胡向东, 有机化学, **2020**, *40*, 4203.)
- [2] (a) Myllymäki, M. J.; Saario, S. M.; Kataja, A. O.; Castillo-Melendez, J. A.; Nevalainen, T.; Juvonen, R. O.; Jävinen, T.; Koshinen, A. M. *J. Med. Chem.* **2007**, *50*, 4236.
- (b) Boger, D. L.; Sato, H.; Lerner, A. E.; Hedrick, M. P.; Fecik, R. A.; Miyauchi, H.; Wikie, G. D.; Austin, B. J.; Patricelli, M. P.; Cravatt, B. F. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97*, 5044.
- [3] Janssen, F. J.; Bagge, M. P.; Hummel, J. J. A.; Overkleeft, H. S.; Cravatt, B. F.; Boger, D. L.; van der Stelt, M. *J. Med. Chem.* **2015**, *58*, 9742.
- [4] McGrath, M. E.; Sprengeler, P. A.; Hill, C. M.; Martichonok, V.; Cheung, H.; Somoza, J. R.; Palmer, J. T.; Janc, J. W. *Biochemistry* **2003**, *42*, 15018.
- [5] Tully, D. C.; Vidal, A.; Chatterjee, A. K.; Williams, J. A.; Roberts, M. J.; Petrassi, H. M.; Sraggon, G.; Bursulaya, B.; Pacoma, R.; Shipway, A.; Schumacher, A. M.; Danahay, H.; Harris, J. L. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 5895.
- [6] Wolfart, K.; Molar, A.; Kawase, M.; Motohashi, N.; Molnar, J. *Biol. Pharm. Bull.* **2004**, *27*, 1462.
- [7] (a) Huang, J.; Zhou, Z.; Zhou, M.; Miao, M.; Li, H.; Hu, Q. *Bioorg. Med. Chem.* **2018**, *26*, 1653.
- (b) Tobisu, M.; Chatani, N.; Asaumi, T.; Amako, K.; Ie, Y.; Fukumoto, Y.; Murai, S. *J. Am. Chem. Soc.* **2000**, *122*, 12663.
- (c) Komanduri, V.; Krische, M. J. *J. Am. Chem. Soc.* **2006**, *128*, 16448.
- [8] For a recent review, see: Aljaar, N.; Gujjarappa, R.; Al-Refai, M.; Shtaiwi, M.; Malakar, C. C. *J. Heterocycl. Chem.* **2019**, *56*, 2730.
- [9] Wu, X.-F.; Anbarasan, P.; Neumann, H.; Beller, M. *Angew. Chem., Int. Ed.* **2010**, *49*, 7316.
- [10] (a) Karthik, S.; Gandhi, T. *Org. Lett.* **2017**, *19*, 5486.
- (b) Yang, K.; Chen, X.; Wang, Y.; Li, W.; Kadi, A. A.; Fun, H.-K.; Sun, H.; Zhang, Y.; Li, G.; Lu, H. *J. Org. Chem.* **2015**, *80*, 11065.
- (c) Yang, K.; Zhang, C.; Wang, P.; Zhang, Y.; Ge, H. *Chem.-Eur. J.* **2014**, *20*, 7241.
- (d) Sharma, U. K.; Sharma, N.; Xu, J.; Song, G.; Van der Eycken, E. V. *Chem.-Eur. J.* **2015**, *21*, 4908.
- [11] (a) Boominathan, S. S. K.; Hu, W.-P.; Senadi, G. C.; Vandavsi, J. K.; Wang, J.-J. *Chem. Commun.* **2014**, *50*, 6726.
- (b) Jiang, J.; Zou, H.; Dong, Q.; Wang, R.; Lu, L.; Zhu, Y.; He, W. *J. Org. Chem.* **2016**, *81*, 51.
- (c) Alijaar, N.; Malakar, C. C.; Conrad, J.; Beifuss, U. *J. Org. Chem.* **2015**, *80*, 10829.
- (d) Dhameliya, T. M.; Chourasiya, S. S.; Mishra, E.; Jadhavar, Bharatam, P. S. P. V.; Chakraborti, A. K. *J. Org. Chem.* **2017**, *82*, 10077.
- [12] (a) Wang, H.; Liu, J.; Qu, J.-P.; Kang, Y.-B. *J. Org. Chem.* **2020**, *85*, 3942.
- (b) Bao, K.; Li, F.; Liu, H.; Wang, Z.; Shen, Q.; Wang, J.; Zhang, W. *Sci. Rep.* **2015**, *5*, no. 10360.
- (c) Wu, Y.; Choy, P. Y.; Mao, F.; Kwong, F. Y. *Chem. Commun.* **2013**, *49*, 689.
- [13] (a) Cui, L.; He, Y.; Fan, X. *Chin. J. Chem.* **2012**, *30*, 992.
- (b) Fan, X.; He, Y.; Zhang, X.; Guo, S.; Wang, Y. *Tetrahedron* **2011**, *67*, 6369.
- [14] (a) Pridgen, L. N.; Shilcrat, S. C. *Synthesis* **1984**, 1048.
- (b) Jutzi, P.; Gilge, U. *J. Heterocycl. Chem.* **1982**, *20*, 1011.
- (c) Lester, R. P.; Bham, T.; Bousfield, T. W.; Lewis, W.; Camp, J. E. *J. Org. Chem.* **2016**, *81*, 12472.
- [15] (a) Liu, Y.; Zhang, Y.; Wan, J.-P. *J. Org. Chem.* **2017**, *82*, 8950.
- (b) Tu, Z.; Du, Y.; Cao, X.; Liu, Y. *Adv. Synth. Catal.* **2019**, *361*, 4989.
- (c) Chen, X.; Liu, Y. *Tetrahedron* **2018**, *74*, 3691.
- (d) Tian, S.; Wang, C.; Xia, J.; Wan, J.-P.; Liu, Y. *Adv. Synth. Catal.* **2021**, *363*, 4627.
- (e) Wang, L.; Liu, X.; Liu, J.-b.; Shen, J.; Chen, Q.; He, M.-Y. *Chem. Asian J.* **2018**, *13*, 765.
- [16] For selected recent reviews on C—H bond activation or functionalization, see:
- (a) Gandeepan, P.; Müller, T.; Zell, D.; Cera, G.; Warratz, S.; Ackermann, L. *Chem. Rev.* **2019**, *119*, 2192.
- (b) Xu, X.; Chen, D.; Wang, Z. *Chin. J. Org. Chem.* **2019**, *39*, 3338 (in Chinese).
- (徐鑫明, 陈德茂, 王祖利, 有机化学, **2019**, *39*, 3338.)
- (c) Tian, S.; Luo, T.; Zhu, Y.; Wan, J.-P. *Chin. Chem. Lett.* **2020**, *31*, 3073.
- (d) Zhao, B.; Liu, Y. *Synthesis* **2020**, *52*, 3211.
- (e) Yuan, X.; Yang, G.; Yu, B. *Chin. J. Org. Chem.* **2020**, *40*, 3620 (in Chinese).
- (袁晓亚, 杨国平, 於兵, 有机化学, **2020**, *40*, 3620.)
- (f) Luo, T.; Tian, S.; Wan, J.-P.; Liu, Y. *Curr. Org. Chem.* **2021**, *25*, 1180.
- [17] For some recent references on aerobic C—H transformation, see:
- (a) Boess, E.; Hoof, M. V.; Birdsall, S. L.; Klusmann, M. *J. Org. Chem.* **2020**, *85*, 1972.
- (b) Gui, Q.-W.; Teng, F.; Li, Z.-C.; Jin, X.-F.; Zhang, M.; Dai, J.-N.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Org. Chem. Front.* **2020**, *7*, 4026.
- (c) Bao, P.; Liu, F.; Lv, Y.; Yue, H.; Li, J.-S.; Wei, W. *Org. Chem. Front.* **2020**, *7*, 492.
- (d) Xie, L.-Y.; Liu, Y.-S.; Ding, H.-R.; Gong, S.-F.; Tan, J.-X.; He, J.-Y.; Cao, Z.; He, W.-M. *Chin. J. Catal.* **2020**, *41*, 1168.
- (e) Tanimoto, K.; Ohkado, R.; Lida, H. *J. Org. Chem.* **2019**, *84*, 14980.
- (f) Abdulkader, A.; Wang, R.; Mamat, M.; Liu, C. *Chin. J. Org. Chem.* **2020**, *40*, 1697 (in Chinese).
- (阿布力米提·阿布都卡德尔, 汪荣, 买买提·买买提, 刘晨江, 有机化学, **2020**, *40*, 1697.)
- (g) Sun, K.; Li, Y.; Feng, R.; Mu, S.; Wang, X.; Zhang, B. *J. Org. Chem.* **2020**, *85*, 1001.
- (h) Wang, Y.; Liu, Y. *Acta Chim. Sinica* **2019**, *77*, 418 (in Chinese).
- (王昱贇, 刘云云, 化学学报, **2019**, *77*, 418.)
- (i) Yao, B.; Wu, J.; Wang, Y.; Jiang, H. *Chin. J. Org. Chem.* **2020**, *40*, 3044 (in Chinese).
- (姚彪, 巫佳浩, 汪钰, 江焕峰, 有机化学, **2020**, *40*, 3044.)
- [18] Sun, A.; Prussia, A.; Zhan, W. Q.; Murray, E. E.; Doyle, J.; Cheng, L. T.; Yoon, J. J.; Radchenko, E. V.; Palyulin, V. A.; Compans, R. W.; Liotta, D. C.; Plemper, R. K.; Snyder, J. P. *J. Med. Chem.* **2006**, *49*, 5080.
- [19] (a) Pan W.-J.; Wang, Z.-X. *Asian J. Org. Chem.* **2018**, *7*, 1626.
- (b) Xie, P.; Huang, H. M.; Xie, Y. J.; Guo, S. M.; Xia, C. G. *Adv. Synth. Catal.* **2012**, *354*, 1692.
- [20] Ji, Y.-Y.; Zhang, Y.; Hu, Y.-Y.; Shao, L.-X. *Tetrahedron* **2015**, *71*, 6818.
- [21] Niu, Z.-J.; Li, L.-H.; Liu, X.-Y.; Liang, Y.-M. *Adv. Synth. Catal.* **2019**, *361*, 5217.
- [22] Nguyen, T. B.; Retailleau, P. *Org. Lett.* **2017**, *19*, 3887.

(Li, L.; Fan, Y.)