无过渡金属条件下可见光诱导 α -重氮酯的氧化合成 α -酮酯赵保丽^{a,b} 杨良凤^b 程凯^b 周丽云^{*,a} 万结平^{*,a}^(a) 江西师范大学化学化工学院 南昌 330022)^(b) 绍兴文理学院 浙江省精细化工与传统工艺绿色替代技术重点实验室 浙江绍兴 312000)

摘要 发展了一例空气氛围下氧气作为无污染氧化剂合成 α -酮酯的方法. 以易得的 α -重氮酯为起始原料, 该转化在采用曙红 Y 作为光敏剂的光诱导条件下进行. 反应在室温和空气氛围下操作, 底物兼容性好. 提供了一个绿色、简便合成结构多样的 α -酮酯的方法.

关键词 光催化; 无过渡金属; 空气氧化; α -重氮酯; α -酮酯

Visible Light Induced Oxidation of α -Diazo Esters for the Transition Metal-Free Synthesis of α -Keto EstersZhao, Baoli^{a,b} Yang, Liangfeng^b Cheng, Kai^b Zhou, Liyun^{*,a} Wan, Jie-Ping^{*,a}^(a) College of Chemistry and Chemical Engineering, Jiangxi Normal University, Nanchang 330022)^(b) Zhejiang Key Laboratory of Alternative Technologies for Fine Chemicals Process, Shaoxing University, Shaoxing, Zhejiang 312000)

Abstract Herein, an aerobic oxidation approach for the synthesis of α -keto esters using oxygen as a cheap and pollution-free oxidant has been developed. By using readily available α -diazo esters as starting materials, the transformation was achieved by the inducement of visible light with eosin Y as the light sensitizer. The reactions proceed at room temperature under air atmosphere, tolerating substrates with broad scope, thus providing a green and facile synthetic approach to structurally diverse α -keto esters.

Keywords photocatalysis; transition metal-free; aerobic oxidation; α -diazo esters; α -keto esters

1 Introduction

Visible light photoredox catalysis, which relies on the redox ability of excited-state of photosensitizer or photocatalyst to promote single-electron-transfer (SET) process upon visible-light irradiation, has drawn intensive research interests and received tremendous advances.^[1] Particularly, due to the incomparable advantages in atom economy, sustainability and facile operation, the photocatalytic oxidation reactions employing the oxygen as the oxidant has been demonstrated as powerful tools. A variety of different types of organic transformations, including the C—H bond functionalization,^[2] sulfur oxidation,^[3] carbon-carbon bond cleavage,^[4] intramolecular dehydrogenative elimination,^[5] difunctionalization of unsaturated carbon-carbon bonds,^[6] and a plethora of other reactions^[7] have all gained im-

portant progresses with photocatalysis. Inarguably, the known works have disclosed the promise in developing much more methods with high efficiency by combining visible light-based photocatalysis and oxygen-based oxidation.

On the other hand, α -keto esters are useful chemicals showing widespread application in synthetic chemistry and many related disciplines.^[8] The synthesis of α -keto esters thus occupies rather crucial position in the area of organic synthesis. In 2011, Jiao and co-workers^[9] have reported the synthesis of α -keto esters via oxidative esterification of benzyl halides by photocatalysis. Since then, the visible light-based synthesis of α -keto esters has won significant interests from synthetic chemists, which promoted the occurrence of many different visible light photocatalytic methods for the synthesis of these compounds. For in-

* Corresponding authors. E-mail: wanjiaping@jxnu.edu.cn; zly9510@126.com

Received November 12, 2021; revised December 1, 2021; published online December 8, 2021.

Project supported by the Natural Science Foundation of Jiangxi Province (No. 20202ACBL203006) and the Foundation of Zhejiang Educational Committee (No. Y202147050).

江西省自然科学基金(No. 20202ACBL203006)和浙江省教育厅科研计划(No. Y202147050)资助项目.

stance, our group^[10] has developed the synthetic approach toward α -keto esters by carbon-carbon double bond cleavage of enaminones. Wang and co-workers^[11] have established an alternative method via the reactions of bromoalkynes, alkynes and pure oxygen with visible light irradiation. In addition, the carbon-carbon bond cleavage of α -cyano ketones,^[12] the nitroarene mediated oxidation of α -diazo esters,^[13] the oxygenation of methylene C—H bonds on the methylene esters,^[14] as well as the oxidation of α -hydroxyl esters^[15] have been reported as practical methods as accesses to α -keto esters via visible light photocatalysis. Despite these notable advances, the utilization of pure oxygen as oxidant and/or the requirement of sensitive reagents or unstable starting materials raises up the requirement for more complementary method for the synthesis of such products. Because of their versatile reactivity, tolerance to moisture as well as the relative readily availability,^[16] we envisage that using α -diazo esters as starting materials might bring opportunity for devising photocatalytic synthesis of α -keto esters via milder and more facile operation. Here, based on the extensive concerns on the advantageous metal-free synthesis with visible light photocatalysis,^[17] we report our results in the visible light induced synthesis of α -keto esters via the oxidation of α -diazo esters by using Eosin Y as non-metal sensitizer with air as the source of oxygen, which would complement the presently known methods via simple and mild operation.^[18]

2 Results and discussion

Initially, α -phenyl diazoacetate **1a** (0.5 mmol) was treated with Rhodamine B (0.025 mmol) at room temperature in CH₃CN with the irradiation of 12 W blue LEDs, and α -keto ester **2m** was obtained in 68% yield (Table 1, Entry 1). No desired product was obtained in the absence of photocatalyst (PC) (Table 1, Entry 2). Later, performing the reaction using Rose Bengal and Eosin Y as PC was found to promote the product formation in 77%~83% yields (Table 1, Entries 3~4). The analogous reactions employing [Ru(bpy)₃]Cl₂ and Ir(ppy)₃ also led to the synthesis of product **2m** with lower yield (Table 1, Entries 5~6). For the solvent effect, common organic solvents such as CCl₄, CH₂Cl₂, and tetrahydrofuran (THF) gave poor results (Table 1, Entries 7~9). Evidently, aprotic polar medium was found to be effective for the α -keto ester formation wherein acetonitrile furnished better result than other solvents such as *N,N*-dimethylformamide (DMF), dimethyl sulfoxide (DMSO) and acetone (Table 1, Entries 10~12). In the section of comparing different light sources, the reactions under the irradiation of green LEDs, household light bulb, or red LEDs all gave inferior result (Table 1, Entries 13~15). Later, a control entry under pure oxygen gave the product with similar yield with the one under air (Table 1, Entry 16). Finally, the reaction conducted under dark atmosphere provided no product (Table 1, Entry 17), confirming that the light irradiation was indispensable for

Table 1 Optimization on reaction conditions^a

Entry	Photocatalyst	Solvent	Light source	Yield ^b /%
1	Rhodamine B	CH ₃ CN	Blue LED	68
2	—	CH ₃ CN	Blue LED	—
3	Rose Bengal	CH ₃ CN	Blue LED	33
4	Eosin Y	CH ₃ CN	Blue LED	83
5	[Ru(bpy) ₃]Cl ₂	CH ₃ CN	Blue LED	55
6	Ir(ppy) ₃	CH ₃ CN	Blue LED	60
7	Eosin Y	CCl ₄	Blue LED	39
8	Eosin Y	CH ₂ Cl ₂	Blue LED	36
9	Eosin Y	THF	Blue LED	28
10	Eosin Y	DMF	Blue LED	61
11	Eosin Y	DMSO	Blue LED	52
12	Eosin Y	acetone	Blue LED	67
13	Eosin Y	CH ₃ CN	Green LED	50
14	Eosin Y	CH ₃ CN	household bulb	63
15	Eosin Y	CH ₃ CN	Red LED	37
16 ^c	Eosin Y	CH ₃ CN	Blue LED	81
17	Eosin Y	CH ₃ CN	dark	0

^a Reaction condition: **1a** (0.5 mmol), photocatalyst (5 mol%), solvent (1 mL), and stirred for 6 h under air and light irradiation. ^b Isolated yield. ^c Reaction under pure oxygen (101 kPa).

the transformation.

With the optimized conditions in hand, the scope of the diazo esters was next examined. According to the obtained results (Table 2), diazo esters functionalized with both electron donating (OCF₃, SCH₃, *t*-Bu) and electron withdrawing (Br, Cl, CN) substituents on the *para*-position of the aryl groups were equally effective coupling partners, and the corresponding α -keto esters **2a**~**2f** were obtained in 53%~77% yields (Table 2). Diazo esters bearing functional groups on other positions of the phenyl ring such as *meta*-methyl (Table 2, **2g**), and *ortho*-substituted (Br, Cl, NO₂) (Table 2, **2h**~**2j**) diazo ester reacted also smoothly to give α -keto esters with comparable yields (61%~88%). Notably, CF₃-phenyl substrate was also well tolerated, and the corresponding α -keto esters **2k** was provided in 80% yield. Heteroaromatic structure such as pyridinyl (Table 2, **2l**) was also compatible to this synthetic protocol. Later, a series of α -phenyldiazoacetates derived from simple aliphatic alcohols were also employed to react with **1a**, which afforded products **2m**~**2t** in 52%~86% yields. Unfortunately, the reaction using methyl α -diazoacetate did not give the target product. When an α -diazoketone (α -diazoacetophenone) was employed to the standard catalytic conditions, the expected diazo oxygenation was not observed, either. Notably, the 1 mmol scale reaction for the synthesis of **2m** could give the product with 71% good yield.

Table 2 Scope of the photocatalytic α -keto ester synthesis

^a The yield in the parenthesis was given by 1 mmol-scale reaction.

To elucidate the reaction mechanism, control experiments were conducted. First, adding 2,2,6,6-tetramethylpiperidinoxy (TEMPO) or 2,6-di-*t*-butyl-*p*-cresol (BHT) to the model reaction under the standard conditions completely suppressed the formation of **2m** (Eqs. 1 and 2), supporting the reaction as a typical free radical prototype by photocatalysis. In addition, the reaction performed under nitrogen atmosphere gave only trace amount of the target product (Eq. 3), indicating that oxygen (air) was indispensable for this reaction. The results given by the control experiments suggested that the analogous reaction mechanism proposed by Wei *et al.*^[18a] was also involved in our work.

3 Conclusions

In conclusion, herein a metal-free photocatalytic oxidative reaction of α -diazo esters for the synthesis of structurally diverse α -keto esters is reported. By employing eosin Y as the sensitizer with blue LEDs irradiation, more than 20 α -keto esters have been synthesized with generally good to excellent yields. The simple operation, green reaction conditions as well as the broad substrate scope constitute the major advantages of present synthetic method.

4 Experimental section

4.1 General information

All materials were purchased from common commercial sources and used without further purification. The ¹H NMR spectra were recorded at 400 MHz using TMS as internal standard, and the frequency for NMR test was 100 MHz. All products are known compounds whose structures were assigned by comparing the NMR data with literature results. The products were isolated by column chromatography on a silica gel (300~400 mesh) using the mixture of petroleum and ethyl acetate as eluent.

4.2 General procedure for the synthesis of α -keto esters 2

The α -diazo ester **1** (0.5 mmol) and eosin Y (0.025 mmol) were placed in a Schlenk tube equipped with a stirring bar. The solvent CH₃CN (2 mL) was added under air atmosphere. The mixture was stirred and irradiated with 12 W green LEDs at room temperature for 6 h. Upon comple-

tion, the mixture was quenched with saturated sodium bicarbonate (10 mL) and extracted with ethyl acetate (5 mL $\times$ 3). The organic layers were combined and dried with anhydrous Na_2SO_4 . After filtration, the solution was concentrated *in vacuo*. The residue was subjected to silica gel column chromatography to provide pure product with the elution of mixed petroleum ether and ethyl acetate ($V:V=10:1$).

4.3 Procedure for the 1 mmol scale reaction

Diazo ester **1a** (1 mmol) and eosin Y (0.05 mmol) were placed in a Schlenk tube equipped with a stirring bar. The solvent CH_3CN (2 mL) was added under air atmosphere. The mixture was stirred and irradiated with 12 W green LEDs at room temperature for 6 h. Then, the mixture was quenched with saturated sodium bicarbonate (20 mL) and extracted with ethyl acetate (10 mL $\times$ 3). The organic layers were combined and concentrated *in vacuo*. The product was purified by flash column chromatography on silica gel with the elution of mixed petroleum ether and ethyl acetate ($V:V=10:1$) to afford target product **2m** with 71% yield.

Methyl 2-oxo-2-(4-(trifluoromethoxy)phenyl)acetate (**2a**):^[19a] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.13 (d, $J=8.2$ Hz, 2H), 7.34 (d, $J=8.2$ Hz, 2H), 4.00 (s, 3H).

Methyl 2-(4-(methylthio)phenyl)-2-oxoacetate (**2b**):^[19b] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.94 (d, $J=8.6$ Hz, 2H), 7.29 (dd, $J=10.6, 3.6$ Hz, 2H), 3.97 (s, 3H), 2.54 (s, 3H).

Methyl 2-(4-(*tert*-butyl)phenyl)-2-oxoacetate (**2c**):^[18] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.96 (d, $J=8.7$ Hz, 2H), 7.53 (d, $J=8.7$ Hz, 2H), 3.97 (s, 3H), 1.35 (s, 9H).

Methyl 2-(4-bromophenyl)-2-oxoacetate (**2d**):^[18] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.93 (d, $J=8.6$ Hz, 2H), 7.68 (d, $J=8.6$ Hz, 2H), 4.00 (s, 3H).

Methyl 2-(4-chlorophenyl)-2-oxoacetate (**2e**):^[19c] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.01~7.99 (m, 2H), 7.50 (d, $J=8.6$ Hz, 2H), 3.99 (s, 3H).

Methyl 2-(4-cyanophenyl)-2-oxoacetate (**2f**):^[19d] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.21 (d, $J=8.8$ Hz, 2H), 7.85 (d, $J=8.2$ Hz, 2H), 4.00 (s, 3H).

Methyl 2-oxo-2-(*m*-tolyl)acetate (**2g**):^[18] Colorless liquid. ^1H NMR (500 MHz, CDCl_3) δ : 7.16~7.10 (m, 2H), 6.84 (t, $J=7.4$ Hz, 1H), 6.79 (d, $J=8.2$ Hz, 1H), 3.79 (s, 3H), 2.22 (s, 3H).

Methyl 2-(2-bromophenyl)-2-oxoacetate (**2h**):^[19e] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.65 (dd, $J=8.0, 1.6$ Hz, 1H), 7.63~7.61 (m, 1H), 7.44~7.41 (m, 2H), 3.94 (s, 3H).

Methyl 2-(2-chlorophenyl)-2-oxoacetate (**2i**):^[18] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (dd, $J=7.7, 1.5$ Hz, 1H), 7.56~7.52 (m, 1H), 7.46~7.40 (m, 2H), 3.96 (s, 3H).

Methyl 2-(2-nitrophenyl)-2-oxoacetate (**2j**):^[19f] Pale yellow liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.20~8.17

(m, 1H), 7.94~7.86 (m, 2H), 7.73~7.71 (m, 1H), 3.83 (s, 3H).

Methyl 2-oxo-2-(3-(trifluoromethyl)phenyl)acetate (**2k**):^[19g] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.33 (s, 1H), 8.25 (d, $J=7.9$ Hz, 1H), 7.92 (d, $J=7.8$ Hz, 1H), 7.68 (t, $J=7.9$ Hz, 1H), 4.02 (s, 3H).

Ethyl 2-oxo-2-(pyridin-3-yl)acetate (**2l**):^[19h] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 9.26 (d, $J=1.6$ Hz, 1H), 8.86 (dd, $J=4.8, 1.7$ Hz, 1H), 8.37 (dt, $J=8.0, 2.0$ Hz, 1H), 7.49 (ddd, $J=8.0, 4.8, 0.6$ Hz, 1H), 4.48 (q, $J=7.2$ Hz, 2H), 1.45 (t, $J=7.2$ Hz, 3H).

Methyl 2-oxo-2-phenylacetate (**2m**):^[18] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.01 (d, $J=7.2$ Hz, 1H), 7.66 (d, $J=7.4$ Hz, 1H), 7.52 (t, $J=7.8$ Hz, 2H), 3.99 (s, 3H).

Propyl 2-oxo-2-phenylacetate (**2n**):^[10] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, $J=7.4$ Hz, 1H), 7.63 (t, $J=6.9$ Hz, 1H), 7.49 (t, $J=7.6$ Hz, 1H), 4.35 (t, $J=6.7$ Hz, 1H), 1.85~1.71 (m, 1H), 0.99 (t, $J=7.4$ Hz, 1H).

Cyclopentyl 2-oxo-2-phenylacetate (**2o**):^[10] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.91 (d, $J=7.2$ Hz, 2H), 7.55 (t, $J=7.4$ Hz, 1H), 7.42 (t, $J=7.7$ Hz, 2H), 5.40 (tt, $J=5.9, 2.9$ Hz, 1H), 1.94~1.85 (m, 2H), 1.82~1.75 (m, 2H), 1.73~1.63 (m, 2H), 1.60~1.49 (m, 2H).

Isopropyl 2-oxo-2-phenylacetate (**2p**):^[10] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.93 (d, $J=7.1$ Hz, 2H), 7.57 (t, $J=7.4$ Hz, 1H), 7.43 (t, $J=7.7$ Hz, 2H), 5.31~5.21 (m, 1H), 1.33 (d, $J=6.3$ Hz, 6H).

sec-Butyl 2-oxo-2-phenylacetate (**2q**):^[19i] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.88 (d, $J=7.1$ Hz, 2H), 7.55~7.48 (m, 1H), 7.38 (t, $J=7.7$ Hz, 2H), 1.70~1.47 (m, 1H), 1.26 (d, $J=6.3$ Hz, 3H), 0.85 (t, $J=7.5$ Hz, 3H).

Isopentyl 2-oxo-2-phenylacetate (**2r**):^[18] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, $J=7.2$ Hz, 2H), 7.46 (t, $J=7.4$ Hz, 1H), 7.32 (t, $J=7.7$ Hz, 2H), 4.26 (t, $J=6.8$ Hz, 2H), 1.59~1.53 (m, 1H), 1.48 (q, $J=6.8$ Hz, 2H), 0.77 (d, $J=6.6$ Hz, 6H).

tert-Pentyl 2-oxo-2-phenylacetate (**2s**):^[19j] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (dd, $J=8.0, 1.5$ Hz, 2H), 7.43~7.35 (m, 3H), 1.95~1.78 (m, 2H), 1.52 (s, 3H), 1.29 (t, $J=16.1$ Hz, 3H), 0.93 (t, $J=7.5$ Hz, 3H).

2-Methylallyl 2-oxo-2-phenylacetate (**2t**):^[19k] Colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ : 8.03 (d, $J=7.2$ Hz, 2H), 7.67 (t, $J=7.4$ Hz, 1H), 7.52 (t, $J=7.7$ Hz, 2H), 4.45 (s, 2H), 1.65 (s, 5H).

Supporting Information ^1H NMR spectra of products **2a**~**2t**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] For selected reviews, see: (a) Chen, Y.; Lu, L.-Q.; Yu, D.-G.; Zhu, C.-J.; Xiao, W.-J. *Sci. China Chem.* **2019**, *62*, 24.
(b) Yu, X.-Y.; Chen, J.-R.; Xiao, W.-J. *Chem. Rev.* **2021**, *121*, 506.
(c) Sun, K.; Xiao, F.; Yu, B.; He, W.-M. *Chin. J. Catal.* **2021**, *42*,

1921.
(d) Liu, Q.; Wu, L.-Z. *Nat. Chem. Rev.* **2017**, *4*, 359.
(e) Xie, J.; Jin, H.; Hashmi, A. S. K. *Chem. Soc. Rev.* **2017**, *46*, 5139.
(f) Courant, T.; Masson, G. *J. Org. Chem.* **2016**, *81*, 6945.
(g) Yi, R.-N.; He, W.-M. *Chin. J. Org. Chem.* **2021**, *41*, 1267 (in Chinese).
(易荣楠, 何卫民, 有机化学, **2021**, *41*, 1267.)
- [2] (a) Ge, L.; Wang, J.-H.; Lei, T.; Cheng, Y.-Y.; Zhou, C.; Chen, Y.-J.; Ye, C.; Chen, B.; Tung, C.-H.; Wu, L.-Z. *Org. Lett.* **2021**, *23*, 8082.
(b) Bao, P.; Liu, F.; Lü, Y.; Yue, H.; Li, J.-S.; Wei, W. *Org. Chem. Front.* **2020**, *7*, 492.
(c) Xie, L.-Y.; Peng, S.; Yang, L.-H.; Peng, C.; Lin, Y.-W.; Yu, X.; Cao, Z. *Green Chem.* **2021**, *23*, 374.
(d) Gao, Y.; Liu, Y.; Wan, J.-P. *J. Org. Chem.* **2019**, *84*, 2243.
(e) Yuan, W.; Huang, J.; Xu, X.; Wang, L.; Tang, X.-Y. *Org. Lett.* **2021**, *23*, 7139.
(f) Nagae, H.; Sakamoto, K.; Fujiwara, S.; Schindler, T.; Kon, Y.; Sato, K.; Okuda, J.; Mashima, K. *Green Chem.* **2021**, *57*, 11169.
(g) Zhou, K.; Yu, Y.; Lin, Y.-Y.; Li, Y.; Gong, L. *Green Chem.* **2020**, *22*, 4597.
(h) Shi, J.; Wei, W. *Chin. J. Org. Chem.* **2020**, *40*, 2170 (in Chinese).
(时建伟, 魏伟, 有机化学, **2020**, *40*, 2170.)
- [3] (a) Liu, Q.; Wang, L.; Yue, H.; Li, J.-S.; Luo, Z.; Wei, W. *Green Chem.* **2019**, *21*, 1609.
(b) Li, C.; Mizuno, N.; Murata, K.; Ishii, K.; Suenobu, T.; Yamaguchi, K.; Suzuki, K. *Green Chem.* **2020**, *22*, 3896.
(c) Liu, K.-J.; Deng, J.-H.; Yang, J.; Gong, S.-F.; Lin, Y.-W.; He, J.-Y.; Cao, Z.; He, W.-M. *Green Chem.* **2020**, *22*, 433.
(d) Jin, W.; Liu, C. *Chin. J. Org. Chem.* **2021**, *41*, 2148 (in Chinese).
(金伟伟, 刘晨江, 有机化学, **2021**, *41*, 2148.)
(e) Huang, Y.; Xin, Z.; Yao, W.; Hu, Q.; Li, Z.; Xiao, L.; Yang, B.; Zhang, J. *Chem. Commun.* **2018**, *54*, 13587.
- [4] (a) Ma, T.; Hua, J.; Bian, M.; Qin, H.; Lin, X.; Yang, X.; Liu, C.; Yang, Z.; Fang, Z.; Guo, K. *Org. Chem. Front.* **2022**, DOI: 10.1039/D1QO01087E.
(b) Cao, S.; Zhong, S.; Xin, L.; Wan, J.-P.; Wen, C. *ChemCatChem* **2015**, *7*, 1478.
(c) Xin, H.; Duan, X.-H.; Yang, M.; Zhang, Y.; Guo, L.-N. *J. Org. Chem.* **2021**, *86*, 8263.
(d) Dong, C.-L.; Huang, L.-Q.; Guan, Z.; Huang, C.-S.; He, Y.-H. *Adv. Synth. Catal.* **2021**, *363*, 3803.
- [5] (a) Ming, Z.-Y.; Li, K.-R.; Meng, F.-J.; Shi, L.; Jiang, W.-F. *Tetrahedron Lett.* **2020**, *61*, 152059.
(b) Karimi, M.; Sadeghi, S.; Mohebbi, H.; Azarkohosh, Z.; Safarifar, Y.; Mahjoub, A.; Heydari, A. *New Chem. J.* **2021**, *45*, 14024.
(c) Yu, K.; Zhang, H.; Su, C.; Zhu, Y. *Eur. J. Org. Chem.* **2020**, 1956.
(d) Zhang, Y.; Zhao, G.; Zhang, Y.; Huang, X. *Green Chem.* **2014**, *16*, 3860.
- [6] (a) Qian, L.; Zhang, K.; Wang, Z.; Li, H.; Lu, P.; Wang, Y. *J. Org. Chem.* **2021**, *86*, 7955.
(b) Gui, Q.-W.; Teng, F.; Li, Z.-C.; Xiong, Z.-Y.; Jin, X.-F.; Liu, H.-Y.; Lin, Y.-W.; Cao, Z.; He, W. *Chin. Chem. Lett.* **2021**, *32*, 1907.
(c) Sun, B.; Zhu, R.; Zhuang, X.; Shi, X.; Huang, P.; Yan, Z.; Yu, C.; Jin, C. *Org. Lett.* **2021**, *23*, 617.
(d) Jiang, Y.-Q.; Li, J.; Feng, Z.-W.; Xu, G.-Q.; Shi, X.; Ding, Q.-J.; Li, W.; Ma, C.-H.; Yu, B. *Adv. Synth. Catal.* **2020**, *362*, 2609.
(e) Zhang, Q.-B.; Yuan, P.-F.; Kai, L.-L.; Liu, K.; Ban, Y.-L.; Wang, X.-Y.; Wu, L.-Z. *Org. Lett.* **2019**, *21*, 885.
- [7] (a) Xu, H.; Zhang, Y.-F.; Lang, X. *Chin. Chem. Lett.* **2020**, *31*, 1520.
(b) Huang, Y.; Xie, W.; Luo, Y.; Fan, Q.; Zhu, X.; Liu, S.; Shi, X. *Chin. J. Org. Chem.* **2020**, *40*, 1281 (in Chinese).
(黄永康, 解文静, 罗永强, 范琪琦, 朱星亮, 刘世领, 施小新, 有机化学, **2020**, *40*, 1281.)
(c) He, S.; Chen, X.; Zeng, F.; Lu, P.; Peng, Y.; Qu, L.; Yu, B. *Chin. Chem. Lett.* **2020**, *31*, 1863.
(d) Ban, Y.-L.; Dai, J.-L.; Jin, X.-L.; Zhang, Q.-B.; Liu, Q. *Green Chem.* **2019**, *55*, 9701.
(e) Fu, M.; Ji, X.; Li, Y.; Deng, G.-J.; Huang, H. *Green Chem.* **2020**, *22*, 5594.
(f) Hosseini-Sarvari, M.; Bazyar, Z. *ChemistrySelect* **2020**, *5*, 8853.
(g) Han, A.; Sun, J.; Zhang, H.; Chuah, G.-K.; Jaenicke, S. *Chem-CatChem* **2019**, *11*, 6425.
(h) Herrera-Luna, J.; Díaz, D. D.; Abramov, A.; Encinas, S.; Jiménez, Pérez-Ruiz, R. *J. Org. Chem.* **2021**, *23*, 2320.
(i) Peng, X.-J.; He, H.-P.; Liu, Q.; She, K.; Zhang, B.-Q.; Wang, H.-S.; Tang, H.-T.; Pan, Y.-M. *Sci. China Chem.* **2021**, *64*, 753.
- [8] (a) Morgan, E. H.; Laurell, C. B. *Nature* **1963**, *197*, 921.
(b) Unger, F. M. *Adv. Carbohydr. Chem. Biochem.* **1981**, *38*, 323.
(c) Thorat, S. S.; Kataria, P.; Kontham, R. *Org. Lett.* **2018**, *20*, 872.
(d) Eftekhari B; Zirak, M. *Chem. Rev.* **2015**, *115*, 151.
- [9] Su, Y.; Zhang, L.; Jiao, N. *Org. Lett.* **2011**, *13*, 2168.
- [10] Yu, Q.; Zhang, Y.; Wan, J.-P. *Green Chem.* **2019**, *21*, 3436.
- [11] Ni, K.; Meng, L.-G.; Ruan, H.; Wang, L. *Chem. Commun.* **2019**, *55*, 8438.
- [12] Xu, C.; Zhang, N.-N.; Li, X.-J.; Ge, Y.-Q.; Diao, P.-H.; Guo, C. *Synlett* **2018**, *29*, 1065.
- [13] Roy, S.; Kumar, G.; Chatterjee, I. *Org. Lett.* **2021**, *23*, 6709.
- [14] Uygun, M.; Kuhlmann, J. H.; Pérez-Aguilar, M. C.; Piekarski, D. G.; Mancheño, O. G. *Green Chem.* **2021**, *23*, 3392.
- [15] Zhang, H.; Guo, T.; Wu, M.; Huo, X.; Tang, S.; Wang, X.; Liu, J. *Tetrahedron Lett.* **2021**, *67*, 152878.
- [16] For selected reviews on diazo compound-based synthesis, see: (a) Ciszewski, L. W.; Rybicka-Jasińska, K.; Gryko, D. *Org. Biomol. Chem.* **2019**, *17*, 432.
(b) Xia, Y.; Qiu, D.; Wang, J. *Chem. Rev.* **2017**, *117*, 13810.
(c) Chen, N.; He, M.; Zhou, T.; Zhu, Y.; Zhang, H.; Peng, S. *Synthesis* **2021**, *53*, 611.
- [17] (a) Tian, C.; Wang, Q.; Wang, X.; An, G.; Li, G. *J. Org. Chem.* **2019**, *84*, 14241.
(b) Yang, H.; Tian, C.; Qiu, D.; An, G.; Li, G. *Org. Chem. Front.* **2019**, *6*, 2365.
(c) Tian, H.; Yang, H.; Tian, C.; An, G.; Li, G. *Org. Lett.* **2020**, *22*, 7709.
(d) Shi, T.; Liu, Y.-T.; Wang, S.-S.; Lü, Q.-Y.; Yu, B. *Chin. J. Chem.* **2022**, *40*, 97.
(e) He, S.-Q.; Li, H.-C.; Chen, X.-L.; Krylov, I. B.; Terent'ev, A. O.; Qu, L.-B.; Yu, B. *Chin. J. Org. Chem.* **2021**, *41*, 4661 (in Chinese).
(贺树旗, 李昊聪, 陈晓岚, Krylov, I. B.; Terent'ev, A. O.; 屈凌波, 於兵, 有机化学, **2021**, *41*, 4661.)
(f) Wang, X.; Zhang, Y.; Sun, K.; Meng, J.; Zhang, B. *Chin. J. Org. Chem.* **2021**, *41*, 4588 (in Chinese).
(王薪, 张艳, 孙凯, 孟建萍, 张冰, 有机化学, **2021**, *41*, 4588)
(g) Cai, B.; Xuan, J. *Chin. J. Org. Chem.* **2021**, *41*, 4565 (in Chinese).
(蔡宝贵, 宣俊, 有机化学, **2021**, *41*, 4565.)
- [18] During the preparation of this manuscript, Wei *et al.* reported similar Eosin Y induced photocatalytic synthesis of α -keto esters:
(a) Liu, R.; Liu, Q.; Meng, H.; Ding, H.; Hao, J.; Ji, Z.; Yue, H.; Wei, W. *Org. Chem. Front.* **2021**, *8*, 1970.
For selected other recent references on air-based oxidation, see:
(b) Zhuang, W.; Zhang, X.; Huang, Q. *Chin. J. Org. Chem.* **2021**, *41*, 529 (in Chinese).
(庄伟辉, 张晓凤, 黄秋锋, 有机化学, **2021**, *41*, 529.)
(c) Wang, R.; Dong, X.; Zhang, Y.; Wang, B.; Xia, Y.; Abdulkader, A.; Xue, F.; Jin, W.; Liu, C. *Chem.-Eur. J.* **2021**, *27*, 14931.
(d) Wang, C.; Yao, Y.; Xie, J.; Wang, J.; Wang, F.; Zhang, J.; Tang, L. *Chin. J. Org. Chem.* **2021**, *41*, 370 (in Chinese).
(王聪, 姚瑶瑶, 谢珺, 王建塔, 王飞清, 张吉泉, 汤磊, 有机化学, **2021**, *41*, 370.)
(e) Tu, Z.; Zhao, B.; Wan, J.-P.; Wang, C.; Liu, Y. *Chin. J. Org.*

- Chem.* **2021**, *41*, 4780 (in Chinese).
(涂志, 赵保丽, 万结平, 王超莉, 刘云云, 有机化学, **2021**, *41*, 4780.)
- [19] (a) Jana, S.; Li, F.; Empel, C.; Verspeek, D.; Aseeva, P.; Koenigs, R. *M. Chem.-Eur. J.* **2020**, *26*, 2586.
(b) He, X.-L.; Gao, Y.-J.; Nie, J.; Sun, F. *Macromolecules* **2021**, *54*, 3854.
(c) Eckert, K. E.; Ashfeld, B. L. *Org. Lett.* **2018**, *20*, 2315.
(d) Zhang, Y.; Ye, D.; Shen, L.; Liang, K.-J.; Xia, C.-F. *Org. Lett.* **2021**, *23*, 7112.
(e) Roy, S.; Kumar, G.; Chatterjee, I. *Org. Lett.* **2021**, *23*, 6709.
(f) Son, J. H.; Zhu, J.-S.; Phuan, P. W.; Cil, O.; Teuthorn, A. P.; Ku, C. K.; Lee, S.-J.; Verkman, A. S.; Kurth, M. J. *J. Med. Chem.* **2017**, *60*, 2401.
(g) Tanaka, N.; Tsutsumi, R.; Uraguchi, D.; Ooi, T. *Chem. Commun.* **2017**, *53*, 6999.
(h) Murray, S. A.; Green, J. C.; Tailor, S. B.; Meek, P. S. J. *Angew. Chem., Int. Ed.* **2016**, *55*, 9065.
(i) Kumar, Y.; Jaiswal, Y.; Kumar, A. *J. Org. Chem.* **2016**, *81*, 12247.
(j) Liu, H.-K.; Lei, Y.-F.; Tian, P.-J.; Wang, H.; Zhao, X.; Li, Z.-T.; Zhang, D.-W. *J. Mater. Chem. A* **2021**, *9*, 6361.
(k) Mizota, I.; Tanaka, K.; Shimizu, M. *Tetrahedron Lett.* **2012**, *53*, 1847.

(Cheng, F.)