α -酮酸、异腈和水的电化学三组分合成 α -酮酰胺吴红谕^a 于贤勇^{*,b} 曹忠^{*,a}

(长沙理工大学化学化工学院 电力与交通材料保护湖南省重点实验室&细胞化学湖南省重点实验室 长沙 410114)

(湖南科技大学化学化工学院 理论有机化学与功能分子教育部重点实验室 湖南湘潭 411201)

摘要 电化学有机合成使用无痕的电子替代化学氧化还原试剂, 是一种环境友好的合成方法. 利用碘化钾作为电解质和催化剂, 乙腈和水的混合溶液作为溶剂, 在非分离式电解池中实现了 α -酮酸、异腈和水的电化学脱羧酰基化和接力水合反应, 高效合成了一系列的 α -酮酰胺化合物. 该电化学反应无需使用化学氧化剂, 无需过渡金属催化剂, 具有反应条件温和等优点.

关键词 电化学合成, 脱羧酰基化反应, 水合反应, 异腈, α -酮酰胺

Electrochemical Multicomponent Synthesis of α -Ketoamides from α -Oxocarboxylic Acids, Isocyanides and WaterWu, Hongyu^a Yu, Xianyong^{*,b} Cao, Zhong^{*,a}

(^a Hunan Provincial Key Laboratory of Materials Protection for Electric Power and Transportation, Hunan Provincial Key Laboratory of Cytochemistry, School of Chemistry and Chemical Engineering, Changsha University of Science and Technology, Changsha 410114)

(^b Key Laboratory of Theoretical Organic Chemistry and Functional Molecule of Ministry of Education, School of Chemistry and Chemical Engineering, Hunan University of Science and Technology, Xiangtan, Hunan 411201)

Abstract Electrochemical organic synthesis is considered as an environmentally benign method because it employs traceless electrons as redox agents, thereby avoiding the need for chemical oxidants. The electrochemical decarboxylative acylation of isocyanides and relay hydration reaction for the synthesis of α -ketoamides was developed. This reaction can be carried out under mild conditions in the absence of a chemical oxidant and a transition-metal catalyst.

Keywords electrochemical synthesis; decarboxylative acylation; hydration reaction; isocyanide; α -ketoamides

1 Introduction

Isocyanides is a highly valuable C(1) building blocks in current organic chemistry, which could be transform into heteroatomic molecules with electrophiles, nucleophiles and radicals.^[1] On the other hand, α -ketoamides are broadly used as versatile synthetic intermediates and precursors for complicated molecule production.^[2] They also serve as common scaffolds in a wide variety of natural products, pharmaceuticals and biologically active molecules.^[3] As a consequence, a myriad of research has been spent on the development of novel synthetic protocols to access this scaffold.^[4]

Among these methods, the most common one is the

amidation of α -oxocarboxylic acids^[5] and amines, which require equivalent amounts of carboxylic acid activators such as thionyl chloride,^[6] EDA·HCl^[7] or oxalyl chloride.^[8] In recent years, chemists have done a lot of remarkable work in this field. In 2019, Wei and Yue's group^[9] reported an efficient visible-light-induced method for synthesizing α -ketoamides from α -oxocarboxylic acids, isocyanides and water (Scheme 1a). Despite these remarkable advances, the development of novel approaches for the preparation of α -ketoamides is still necessary.

Over the past years, electrochemical organic synthesis^[10] has increasing attention because the electrochemical oxidation/reduction of reaction substrates can sustainably activate them for further manipulation in the absence of stoi-

* Corresponding authors. E-mail: caoz@csust.edu.cn; fine_chem@163.com

Received November 4, 2021; revised December 9, 2021; published online December 15, 2021.

Project supported by the National Natural Science Foundation of China (No. 21545010), the Natural Science Foundation of Hunan Province (No. 2020JJ4599) and the Science and Technology Innovation Program of Hunan Province (No. 2021RC5028).

国家自然科学基金(No. 21545010)、湖南省自然科学基金(No. 2020JJ4599)和湖南省科技创新计划(No. 2021RC5028)资助项目.

chiometric amounts of chemical redox reagents and in an even more energy-efficient fashion than visible-light photosynthesis.^[11] However, an electrochemical decarboxylative acylation of isocyanide with α -oxocarboxylic acid for constructing α -ketoamides has not been reported to date. Herein, we report the successful development of an electrochemical method for synthesizing α -ketoamides from α -oxocarboxylic acids, isocyanides and water. Advantageously, this process proceeds in a transition metal- and external oxidant-free fashion to deliver various functionalized α -ketoamides (Scheme 1b).

Scheme 1 Decarboxylative acylations of isocyanides with α -oxocarboxylic acids and H_2O

2 Results and discussion

We initiated our studies on the electrochemical decarboxylative acylation of tosylmethyl isocyanide (**2a**) with α -oxobenzeneacetic acid (**1a**) in an undivided cell equipped with a reticulated vitreous carbon (RVC) anode and a platinum (Pt) cathode. After extensive investigation of the reaction conditions, a combination of 20 mol% KI and MeCN/ H_2O (8 mL/2 mL) mixed solvent under 6 mA constant current electrolysis at ambient temperature provided α -ketoamide **3aa** in 86% GC-MS yield (Table 1, Entry 1). Inferior yields of **3aa** were obtained when other electrode system was used instead of RVC anode/Pt cathode system (Entries 2~6). A series of supporting electrolytes was then examined. Iodine salts, such as NH_4I and $n\text{Bu}_4\text{NI}$ gave lower yields of **3aa** (Entries 7, 8). No reaction occurred when non-iodine salts (LiBF_4 , $n\text{Bu}_4\text{NBF}_4$, Et_4NClO_4 and LiClO_4) replaced KI (Entries 9~12). Decreasing the loading of KI from 20 mol% to 10 mol% deleterious impact on this transformation (Entry 13). The experimental results suggested that iodide ion was crucial for this transformation (Entries 7~13). The electrochemical reaction was then carried out in different solvents like HFIP/ H_2O , DMF/ H_2O , THF/ H_2O , DMSO/ H_2O , EtOH/ H_2O and DMA/ H_2O (Entries 14~19). However, no improvement in yields was observed compared to the MeCN/ H_2O . The ratio of MeCN to water was also explored (Entries 1 and 20, 21) and the suitable ratio was observed at 8 mL to 2 mL. Both an increase and decrease of operating current led to a diminished reaction yield (Entries 22, 23). Control experiment showed that electric current was critical for this reaction to occur (Entry 24). No improvement in the transformation was observed when nitrogen gas was used instead of air (Entry 25).

Table 1 Screening the optimized reaction conditions^{a,b}

Entry	Variation from the standard reaction conditions	Yield ^b /%
1	None	86
2	RVC(+) Al(−) instead of RVC(+) Pt(−)	74
3	RVC(+) Ag(−) instead of RVC(+) Pt(−)	78
4	RVC(+) RVC(−) instead of RVC(+) Pt(−)	73
5	Pt(+) Pt(−) instead of RVC(+) Pt(−)	80
6	C(+) Pt(−) instead of Pt(+) Pt(−)	81
7	NH_4I instead of KI	64
8	$n\text{Bu}_4\text{NI}$ instead of KI	79
9	LiBF_4 instead of KI	N.R.
10	$n\text{Bu}_4\text{NBF}_4$ instead of KI	N.R.
11	Et_4NClO_4 instead of KI	N.R.
12	LiClO_4 instead of KI	N.R.
13	KI (10 mol%) instead of KI (20 mol%), 12 h	23
14	HFIP/ H_2O instead of MeCN/ H_2O	70
15	DMF/ H_2O instead of MeCN/ H_2O	26
16	THF/ H_2O instead of MeCN/ H_2O	21
17	DMSO/ H_2O instead of MeCN/ H_2O	Trace
18	EtOH/ H_2O instead of MeCN/ H_2O	Trace
19	DMA/ H_2O instead of MeCN/ H_2O	Trace
20	MeCN/ H_2O = 8 mL/1 mL	76
21	MeCN/ H_2O = 8 mL/3 mL	80
22	10 mA instead of 6 mA, 12 h	34
23	3 mA instead of 6 mA, 12 h	81
24	No electricity	N.R.
25	Under N_2 atmosphere	87

^a Conditions: Pt (50 mm × 8 mm × 0.1 mm) as the cathode and RVC as the anode, 20 mm of the electrode was immersed under the solvent, constant current = 6 mA, **1a** (0.3 mmol), **2a** (0.6 mmol), KI (20 mol%), MeCN/ H_2O (8/2 mL), r.t., air, 8 h. ^b Estimated by GC-MS using xylene as an internal reference.

Having the operationally simple optimal conditions, the generality of this electrochemical reaction was investigated (Table 2). Firstly, a series of α -oxobenzeneacetic acid substrates bearing electron-donating or electron-deficient functional groups at the *para*-position of the benzene ring underwent the decarboxylative acylation leading to the generation of the desired products (**3aa**~**3fa**) in good to excellent yields. The target α -ketoamides (**3ga** and **3ha**) derived from the *meta*- and *ortho*-methyl phenylglyoxylic acids were also obtained with good isolated yields. A bulkier tri-substituted phenylglyoxylic acid substrate is also effective to deliver the corresponding product (**3ia**) in 70% yield. Alkyl isocyanides with different carbon lengths and isomeric structures almost did not affect the reaction outcome (**3ab**~**3af**). Various functionalized benzyliisocyanides were tested, and 76%~88% yields of α -ketoamides (**3ag**~**3ak**) were obtained. Both 1-isocyano-4-methoxybenzene and bulky 2-isocyano-1,3-dimethylbenzene participated in the current reactions smoothly and gave the target products (**3al** and **3am**) in excellent yields.

Table 2 Reaction scope of the synthesis of α -ketoamides^{a,b}

$\text{R}^1\text{C}(=\text{O})\text{CO}_2\text{H} + \text{R}^2\text{NC} \xrightarrow[\text{MeCN/H}_2\text{O} (8:2, \text{V:V}), \text{r.t.}, \text{air}, 8 \text{ h}]{\text{RVC Pt, } I = 6 \text{ mA}, \text{KI (20 mol\%)}}$		$\text{R}^1\text{C}(=\text{O})\text{C}(=\text{O})\text{NHR}^2$
1	2	3
$\text{R} = \text{H}, 4\text{-Et}, 4\text{-OMe}, 4\text{-F}, 4\text{-Cl}, 4\text{-Br}$		
		3aa , 86% 3ba , 82% 3ca , 85% 3da , 78% 3ea , 75% 3fa , 77%
		3ga , 81% 3ha , 75% 3ia , 70% 3ab , 73%
		3ac , 75% 3ad , 78% 3ae , 85% 3af , 83% 3ag , 88% 3ah , 82%
		3ai , 80% 3aj , 78% 3ak , 76% 3al , 76% 3am , 70%

^a Conditions: Pt (50 mm × 8 mm × 0.1 mm) as the cathode, and RVC as anode, 20 mm of the electrode was immersed under the solvent, constant current = 6 mA, **1** (0.3 mmol), **2** (0.6 mmol), KI (20 mol%), MeCN/H₂O (8/2 mL), r.t., air, 8 h. ^b Isolated yields.

To gain insight into the reaction mechanism, some control experiments were carried out. Firstly, the electrochemical reaction was conducted in the presence of radical scavenger (TEMPO or BHT), only trace amount of **3aa** was detected (Scheme 2a). These experimental results suggested that a radical process might involve in the present reaction. No reaction occurred when molecular iodine was used instead of potassium iodide in the decarboxylative acylation reaction (Scheme 2b). Replacing the MeCN/H₂O mixed solvent with anhydrous MeCN as a single solvent resulted in no formation of **3aa** (Scheme 2c).

Scheme 2 Control experiments for mechanism study

Next, the cyclic voltammograms of all reactants were investigated. An oxidative peak of α -oxobenzeneacetic acid (**1a**) was observed at 2.6 V (vs. Ag/AgCl) in the presence of anhydrous acetonitrile as the sole solvent, whereas a lower potential oxidative peak was detected at 2.12 V with MeCN/H₂O as co-solvent. This result suggested that the oxidation potential of **1a** could be reduced by the addition of a small amount of water to acetonitrile. And the cyclic

voltammogram of KI appeared two oxidative peaks (1.08 and 1.46 V). The oxidative peaks corresponded to the oxidation of iodide to triiodide in an overall two-electron process (I^- to I_3^-) and triiodide to iodine in an overall one-electron process (I_3^- to I_2).

On the basis of the above-mentioned mechanistic investigations and the related published research studies in this field,^[9,12] a plausible reaction mechanism is proposed and shown in Scheme 3. Firstly, the anion (**A**) was easily generated from α -keto acid (**1**) along with the production of hydrogen gas at the surface of the platinum cathode. The anodic oxidation of iodide ion can generate the molecular iodine (I_2), which further reacted with the intermediate **A** to produce acyl hypoiodite (**B**) and the iodide anion. The intermediate **B** next underwent hemolytic dissociation to give

Scheme 3 Plausible reaction mechanism

aryloxy radical (C) and iodine radical. The decarboxylation of the radical C gave an acyl radical D, which added to the isocyanide (2) to form a radical intermediate E. Subsequently, the radical E coupled with iodine radical to generate intermediate F. The addition of water to the intermediate F gave the intermediate G, which underwent tautomerization to form the target α -ketoamide 3. However, we could not exclude the possibility of the anodic oxidation of iodide ion to generate an iodine radical, which might directly react with α -keto acid (1) to form aroyloxy radical (C).

3 Conclusions

In conclusion, we report a novel and efficient electrocatalytic protocol for synthesizing α -ketoamides through the multicomponent reaction α -oxocarboxylic acids, isocyanides and water. A broad range of substrates exhibit good functional group tolerance and provide the corresponding products in good yields via electrochemical decarboxylative acylation and hydration reaction. The mechanistic studies show the key role of water, which not only act as co-solvent and reactant but also decrease the oxidative potential. Potassium iodide as an invaluable reagent plays the role of oxidant and supporting electrolyte in the reaction.

4 Experimental section

4.1 General information

All reactions were carried using materials obtained from commercial suppliers and used without further purification unless otherwise noted. The instrument for electrolysis and cyclic voltammetry is IKA ElectraSyn 2.0. The anode electrode is reticulated vitreous carbon (RVC) and cathode electrode is platinum electrode (Pt, 50 mm $\times$ 8 mm $\times$ 0.1 mm). Thin layer chromatography (TLC) was performed on precoated aluminium-backed silica gel 60 F254 plates. TLC plates were visualized with ultraviolet light or I_2 . Flash column chromatography was performed using 200–300 mesh silica gel. 1H NMR, and ^{13}C NMR spectra were recorded on a Bruker Avance DRX-500 spectrometers; Mass spectra were performed on a spectrometer operating on ESI-TOF.

4.2 General experimental procedure for 3

α -Oxocarboxylic acids 1 (0.3 mmol), isocyanides 2 (0.6 mmol), KI (0.06 mmol), MeCN (8 mL) and H_2O (2 mL) were placed in an undivided reaction tube (10 mL). RVC (35 mm $\times$ 5 mm $\times$ 6 mm) as the anode and a platinum plate (Pt, 50 mm $\times$ 8 mm $\times$ 0.1 mm) as the cathode, and 20 mm of the electrodes were immersed in the reaction mixture. The electrolysis was carried out under air atmosphere at room temperature using a constant current of 6 mA for 8 h. After completion, the MeCN was removed, and the residue was extracted with dichloromethane (10 mL $\times$ 3). The organic layer was dried with anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was chromatographed through silica gel eluting with ethyl acetate/petroleum ether

to give the desired products 3.

2-oxo-2-Phenyl-*N*-(tosylmethyl)acetamide (3aa)^[9]: 1H NMR (500 MHz, $CDCl_3$) δ : 7.97 (d, J =7.5 Hz, 2H), 7.85 (t, J =6.5 Hz, 1H), 7.54 (d, J =8.0 Hz, 2H), 7.56 (t, J =7.0 Hz, 1H), 7.36 (t, J =7.5 Hz, 2H), 7.26 (d, J =8.0 Hz, 2H), 4.72 (d, J =7.0 Hz, 2H), 2.54 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 185.03, 159.81, 144.68, 133.90, 132.34, 131.45, 130.05, 129.07, 128.04, 127.55, 58.96, 20.71.

2-(4-Ethylphenyl)-2-oxo-*N*-(tosylmethyl)acetamide (3ba): 1H NMR (500 MHz, $CDCl_3$) δ : 7.98 (t, J =7.0 Hz, 1H), 7.89 (d, J =8.5 Hz, 2H), 7.73 (d, J =8.0 Hz, 2H), 7.24 (d, J =8.0 Hz, 2H), 7.16 (d, J =8.5 Hz, 2H), 4.72 (d, J =7.0 Hz, 2H), 2.64–2.59 (m, 2H), 2.34 (s, 3H), 1.16 (t, J =7.5 Hz, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 184.72, 160.29, 151.31, 144.56, 132.48, 130.31, 129.23, 129.03, 128.05, 127.12, 59.05, 28.12, 20.69, 13.98; HRMS (ESI) calcd for $C_{18}H_{20}NO_4S$ $[M+H]^+$ 346.1108, found 346.1104.

2-(4-Methoxyphenyl)-2-oxo-*N*-(tosylmethyl)acetamide (3ca): 1H NMR (500 MHz, $CDCl_3$) δ : 8.07 (d, J =8.5 Hz, 2H), 7.73 (t, J =8.0 Hz, 3H), 7.26 (t, J =8.0 Hz, 2H), 6.82 (d, J =9.0 Hz, 2H), 4.68 (d, J =7.0 Hz, 2H), 3.81 (s, 3H), 2.36 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 182.73, 164.09, 160.14, 144.62, 132.86, 132.30, 129.05, 128.01, 124.46, 112.91, 58.91, 54.61, 20.73; HRMS (ESI) calcd for $C_{17}H_{18}NO_5S$ $[M+H]^+$ 348.0090, found 348.0084.

2-(4-Fluorophenyl)-2-oxo-*N*-(tosylmethyl)acetamide (3da): 1H NMR (500 MHz, $CDCl_3$) δ : 8.13–8.10 (m, 2H), 7.74 (d, J =8.5 Hz, 3H), 7.27 (d, J =8.0 Hz, 2H), 7.04 (t, J =8.5 Hz, 2H), 4.69 (d, J =7.0 Hz, 2H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 183.02, 166.93, 159.52, 144.72, 133.24, 133.16, 132.44, 129.09, 128.02, 127.99, 115.01, 114.84, 58.96, 20.71; HRMS (ESI) calcd for $C_{16}H_{15}FNO_4S$ $[M+H]^+$ 336.0070, found 336.0078.

2-(4-Chlorophenyl)-2-oxo-*N*-(tosylmethyl)acetamide (3ea)^[9]: 1H NMR (500 MHz, $CDCl_3$) δ : 8.01 (d, J =8.0 Hz, 2H), 7.73 (t, J =8.0 Hz, 3H), 7.34 (d, J =7.5 Hz, 2H), 7.27 (d, J =7.5 Hz, 2H), 4.68 (d, J =6.5 Hz, 2H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 183.52, 159.34, 144.75, 140.84, 132.44, 131.55, 129.84, 129.10, 128.02, 127.99, 58.96, 20.71.

2-(4-Bromophenyl)-2-oxo-*N*-(tosylmethyl)acetamide (3fa): 1H NMR (500 MHz, $CDCl_3$) δ : 7.92 (t, J =8.0 Hz, 2H), 7.73 (d, J =8.0 Hz, 3H), 7.53–7.51 (m, 2H), 7.27 (d, J =8.0 Hz, 2H), 4.69 (d, J =7.0 Hz, 2H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 183.78, 159.32, 144.76, 132.42, 131.54, 130.99, 130.23, 129.80, 129.10, 128.01, 58.95, 20.72; HRMS (ESI) calcd for $C_{16}H_{14}BrNO_4S$ $[M+H]^+$ 395.9900, found 396.0277.

2-oxo-2-(*m*-Tolyl)-*N*-(tosylmethyl)acetamide (3ga): 1H NMR (500 MHz, $CDCl_3$) δ : 7.85–7.74 (m, 5H), 7.37 (d, J =7.5 Hz, 1H), 7.26 (t, J =7.5 Hz, 3H), 4.71 (d, J =7.0 Hz, 2H), 2.36 (s, 3H), 2.31 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 185.16, 159.82, 144.66, 137.44, 134.77, 132.37, 131.44, 130.31, 129.07, 128.03, 127.44, 127.38, 58.94, 20.72, 20.27; HRMS (ESI) calcd for $C_{17}H_{18}NO_4S$ $[M+H]^+$ 332.0951, found 332.0947.

2-oxo-2-(*o*-Tolyl)-*N*-(tosylmethyl)acetamide (3ha): 1H

NMR (500 MHz, CDCl₃) δ : 7.81 (t, J =6.5 Hz, 1H), 7.76 (d, J =10.0 Hz, 2H), 7.38~7.34 (m, 2H), 7.28 (d, J =8.0 Hz, 2H), 7.18 (d, J =8.0 Hz, 1H), 7.11 (t, J =8.0 Hz, 1H), 4.72 (d, J =7.5 Hz, 2H), 2.37 (d, J =8.5 Hz, 6H); ¹³C NMR (126 MHz, CDCl₃) δ : 188.11, 160.17, 144.69, 139.70, 132.36, 132.28, 131.12, 130.85, 130.61, 129.04, 128.09, 59.07, 20.72, 19.99; HRMS (ESI) calcd for C₁₇H₁₈NO₄S [M+H]⁺ 332.0951, found 332.0949.

2-Mesityl-2-oxo-*N*-(tosylmethyl)acetamide (**3ia**): ¹H NMR (500 MHz, CDCl₃) δ : 7.78 (s, 1H), 7.74 (d, J =8.0 Hz, 2H), 7.25 (d, J =8.0 Hz, 2H), 6.73 (s, 2H), 4.70 (d, J =7.0 Hz, 2H), 2.34 (s, 3H), 2.18 (s, 3H), 1.90 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ : 193.96, 158.67, 144.64, 139.65, 134.54, 132.46, 131.07, 129.03, 127.96, 127.65, 59.02, 20.66, 20.18, 18.31; HRMS (ESI) calcd for C₁₉H₂₂NO₄S [M+H]⁺ 360.1264, found 360.1268.

2-oxo-2-Phenyl-*N*-propylacetamide (**3ab**)^[13]: ¹H NMR (500 MHz, CDCl₃) δ : 8.27 (d, J =8.0 Hz, 2H), 7.55 (t, J =7.5 Hz, 1H), 7.41 (t, J =7.0 Hz, 2H), 7.19 (s, 1H), 3.32~3.28 (m, 2H), 1.60~1.55 (m, 2H), 0.92 (t, J =7.5 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.88, 160.69, 133.36, 132.32, 130.19, 127.46, 40.09, 21.58, 10.37.

N-Octyl-2-oxo-2-phenylacetamide (**3ac**)^[14]: ¹H NMR (500 MHz, CDCl₃) δ : 8.27 (d, J =8.0 Hz, 2H), 7.55 (d, J =7.5 Hz, 1H), 7.41 (t, J =7.5 Hz, 2H), 7.03 (s, 1H), 3.34~3.30 (m, 2H), 1.57~1.52 (m, 2H), 1.28~1.20 (m, 10H), 0.81 (t, J =6.5 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.91, 160.71, 133.32, 132.41, 130.21, 127.45, 38.47, 30.75, 28.28, 28.18, 28.15, 25.88, 21.61, 13.05.

N-Cyclohexyl-2-oxo-2-phenylacetamide (**3ad**)^[9]: ¹H NMR (500 MHz, CDCl₃) δ : 7.26 (d, J =7.5 Hz, 2H), 7.55 (t, J =7.5 Hz, 1H), 7.40 (d, J =8.0 Hz, 2H), 6.88 (s, 1H), 3.79 (d, J =9.5 Hz, 1H), 1.92 (d, J =11.5 Hz, 2H), 1.69 (d, J =13.5 Hz, 2H), 1.57 (t, J =13.5 Hz, 2H), 1.39~1.31 (m, 2H), 1.24~1.14 (m, 2H); ¹³C NMR (126 MHz, CDCl₃) δ : 187.10, 183.33, 133.28, 130.21, 127.43, 106.25, 47.46, 31.71, 24.41, 23.72.

Methyl (2-oxo-2-phenylacetyl)glycinate (**3ae**)^[9]: ¹H NMR (500 MHz, CDCl₃) δ : 8.26 (d, J =7.5 Hz, 2H), 7.57 (t, J =7.5 Hz, 1H), 7.49 (s, 1H), 7.42 (t, J =8.0 Hz, 2H), 4.12 (d, J =5.5 Hz, 2H), 3.74 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 185.77, 168.35, 160.80, 133.55, 132.09, 130.19, 127.54, 51.59, 40.07.

Ethyl (2-oxo-2-phenylacetyl)glycinate (**3af**)^[9]: ¹H NMR (500 MHz, CDCl₃) δ : 8.26 (d, J =8.0 Hz, 2H), 7.57 (t, J =7.5 Hz, 1H), 7.48 (s, 1H), 7.43~7.39 (m, 2H), 4.22~4.18 (m, 2H), 4.10 (d, J =5.5 Hz, 2H), 1.25 (t, J =7.5 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 185.83, 167.88, 160.80, 133.52, 132.12, 130.17, 127.53, 60.81, 40.23, 13.13.

N-Benzyl-2-oxo-2-phenylacetamide (**3ag**)^[9]: ¹H NMR (500 MHz, CDCl₃) δ : 8.29 (d, J =8.0 Hz, 2H), 7.56 (t, J =7.5 Hz, 1H), 7.42 (t, J =7.5 Hz, 2H), 7.31~7.24 (m, 5H), 5.67 (d, J =4.5 Hz, 1H), 4.51 (d, J =6.0 Hz, 2H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.49, 160.45, 136.03, 133.47, 132.25, 130.23, 127.85, 127.50, 126.90, 126.85, 42.47.

N-(4-Isopropylbenzyl)-2-oxo-2-phenylacetamide (**3ah**)^[14]: ¹H NMR (500 MHz, CDCl₃) δ : 8.29 (d, J =8.0 Hz,

2H), 7.56 (t, J =7.5 Hz, 1H), 7.41 (t, J =7.5 Hz, 2H), 7.28 (s, 1H), 7.19 (d, J =8.0 Hz, 2H), 7.15 (d, J =8.0 Hz, 2H), 4.47 (d, J =6.0 Hz, 2H), 2.86~2.81 (m, 1H), 1.17 (d, J =7.0 Hz, 6H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.55, 160.41, 147.64, 133.42, 133.36, 132.28, 130.21, 127.48, 127.00, 125.90, 42.24, 32.81, 22.96.

2-oxo-2-Phenyl-*N*-(4-(trifluoromethoxy)benzyl)acetamide (**3ai**): ¹H NMR (500 MHz, CDCl₃) δ : 8.28 (d, J =8.0 Hz, 2H), 7.57 (t, J =7.0 Hz, 1H), 7.42 (t, J =7.0 Hz, 3H), 7.29 (d, J =8.5 Hz, 2H), 7.13 (d, J =8.0 Hz, 2H), 4.50 (d, J =6.0 Hz, 2H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.33, 160.60, 147.79, 134.94, 133.57, 132.21, 130.24, 128.29, 127.55, 120.34, 41.68; HRMS (ESI) calcd for C₁₆H₁₃F₃NO₃ [M+H]⁺ 324.0842, found 324.0848.

N-(4-Chlorobenzyl)-2-oxo-2-phenylacetamide (**3aj**)^[15]: ¹H NMR (500 MHz, CDCl₃) δ : 8.29 (d, J =8.0 Hz, 2H), 7.57 (t, J =7.5 Hz, 1H), 7.42 (t, J =7.5 Hz, 2H), 7.36 (s, 1H), 7.26 (d, J =8.5 Hz, 2H), 7.20 (d, J =5.0 Hz, 2H), 4.47 (d, J =6.0 Hz, 2H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.31, 160.49, 134.60, 133.57, 132.66, 132.13, 130.23, 128.21, 127.96, 127.53, 41.74.

N-(4-Bromobenzyl)-2-oxo-2-phenylacetamide (**3ak**)^[14]: ¹H NMR (500 MHz, CDCl₃) δ : 8.29 (d, J =8.0 Hz, 2H), 7.57 (t, J =7.5 Hz, 1H), 7.42 (t, J =7.0 Hz, 4H), 7.34 (s, 1H), 7.15 (d, J =8.0 Hz, 2H), 4.46 (d, J =6.0 Hz, 2H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.40, 160.47, 135.12, 133.58, 132.14, 130.93, 130.24, 128.54, 127.54, 120.76, 41.80.

N-(4-Methoxyphenyl)-2-oxo-2-phenylacetamide (**3al**)^[15]: ¹H NMR (500 MHz, CDCl₃) δ : 8.80 (s, 1H), 8.34 (d, J =7.5 Hz, 2H), 7.56 (t, J =9.5 Hz, 3H), 7.44 (t, J =7.5 Hz, 2H), 6.86 (d, J =8.0 Hz, 2H), 3.75 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 186.58, 157.59, 155.99, 133.55, 132.13, 130.42, 128.74, 127.51, 120.45, 113.31, 54.47.

N-(2,6-Dimethylphenyl)-2-oxo-2-phenylacetamide (**3am**)^[9]: ¹H NMR (500 MHz, CDCl₃) δ : 8.34 (d, J =8.0 Hz, 3H), 7.59 (t, J =7.5 Hz, 1H), 7.45 (t, J =7.0 Hz, 2H), 7.12~7.05 (m, 3H), 2.23 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ : 187.72, 159.86, 135.15, 134.67, 133.25, 132.47, 131.40, 128.65, 128.40, 127.81, 18.50.

4.3 Cyclic voltammetry experiment

The conditions for CV experiment are as follow: Pt plate electrode, diameter glassy carbon disc working electrode, Ag wire reference electrode, scan rate=100 mV/s using 0.02 mol/L of ⁿBu₄NBF₄ as supporting electrolyte. (a) ⁿBu₄NBF₄ (0.02 mol/L) in dry MeCN (10 mL); (b) KI (0.02 mol/L) in dry MeCN (10 mL); (c) α -oxobenzeneacetic acid (0.02 mol/L) in dry MeCN (10 mL); (d) α -oxobenzeneacetic acid **1a** (0.02 mol/L) in MeCN (8 mL) and H₂O (2 mL); (e) tosylmethy isocyanide (**2a**) (0.02 mol/L) in dry MeCN (10 mL); (f) reaction mixture of KI, **1a**, **2a** (0.02 mol/L) in co-solvent of MeCN (8 mL) and H₂O (2 mL).

Supporting Information Copies of ¹H NMR and ¹³C spectra of the products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Zhang, B.; Studer, A. *Chem. Soc. Rev.* **2015**, *44*, 3505.
 (b) Song, B.; Xu, B. *Chem. Soc. Rev.* **2017**, *46*, 1103.
 (c) Wei, W.; Bao, P.; Yue, H.; Liu, S.; Wang, L.; Li, Y.; Yang, D. *Org. Lett.* **2018**, *20*, 5291.
 (d) Bao, W.-H.; Wang, Z.; Tang, X.; Zhang, Y.-F.; Tan, J.-X.; Zhu, Q.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, *30*, 2259.
 (e) Mao, Z.-Y.; Nie, X.-D.; Feng, Y.-M.; Si, C.-M.; Wei, B.-G.; Lin, G.-Q. *Chem. Commun.* **2021**, *57*, 9248.
 (f) Liu, Y.; Chen, X.-L.; Li, X.-Y.; Zhu, S.-S.; Li, S.-J.; Song, Y.; Qu, L.-B.; Yu, B. *J. Am. Chem. Soc.* **2021**, *143*, 964.
 (g) He, Y.; Wang, Y.-C.; Hu, K.; Xu, X.-L.; Wang, H.-S.; Pan, Y.-M. *J. Org. Chem.* **2016**, *81*, 11813.
 (h) He, Y.; Wang, Y.; Liang, X.; Huang, B.; Wang, H.; Pan, Y.-M. *Org. Lett.* **2018**, *20*, 7117.
 (i) Cao, M.; Liu, L.; Tang, S.; Peng, Z.; Wang, Y. *Adv. Synth. Catal.* **2019**, *361*, 1887.
 (j) Cao, M.; Fang, Y.-L.; Wang, Y.-C.; Xu, X.-J.; Xi, Z.-W.; Tang, S. *ACS Comb. Sci.* **2020**, *22*, 268.
 (k) Cao, M.; Teng, Q.-H.; Xi, Z.-W.; Liu, L.-Q.; Gu, R.-Y.; Wang, Y.-C. *Org. Biomol. Chem.* **2020**, *18*, 655.
- [2] (a) De Risi, C.; Pollini, G. P.; Zanirato, V. *Chem. Rev.* **2016**, *116*, 3241.
 (b) Rao, Y. S.; Latha, D. S.; Devunuri, N.; Almansour, A. I.; Arumugam, N.; Yaragorla, S. *Eur. J. Org. Chem.* **2020**, *2020*, 4134.
 (c) Mishra, A. A.; Chaurasia, S. R.; Bhanage, B. M. *New J. Chem.* **2020**, *44*, 10578.
- [3] (a) Chen, J.-C.; Uang, B.-J.; Lyu, P.-C.; Chang, J.-Y.; Liu, K.-J.; Kuo, C.-C.; Hsieh, H.-P.; Wang, H.-C.; Cheng, C.-S.; Chang, Y.-H.; Chang, M. D.-T.; Chang, W.-S. W.; Lin, C.-C. *J. Med. Chem.* **2010**, *53*, 4545.
 (b) Steuer, C.; Gege, C.; Fischl, W.; Heinonen, K. H.; Bartenschlager, R.; Klein, C. D. *Biorg. Med. Chem.* **2011**, *19*, 4067.
 (c) Sridhar, S. N. C.; Ginson, G.; Venkataramana Reddy, P. O.; Tantak, M. P.; Kumar, D.; Paul, A. T. *Biorg. Med. Chem.* **2017**, *25*, 609.
 (d) Du, F.-T.; Ji, J.-X. *Chem. Sci.* **2012**, *3*, 460.
 (e) Kumar, D.; Vemula, S. R.; Cook, G. R. *ACS Catal.* **2016**, *6*, 4920.
 (f) Majumdar, B.; Sarma, D.; Bhattacharya, T.; Sarma, T. K. *ACS Sustainable Chem. Eng.* **2017**, *5*, 9286.
 (g) Chen, J.-Y.; Wu, H.-Y.; Gui, Q.-W.; Han, X.-R.; Wu, Y.; Du, K.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Org. Lett.* **2020**, *22*, 2206.
- [5] (a) Pentead, F.; Lopes, E. F.; Alves, D.; Perin, G.; Jacob, R. G.; Lenardão, E. J. *Chem. Rev.* **2019**, *119*, 7113.
 (b) Zhang, J.; Wu, M.; Fan, J.; Xu, Q.; Xie, M. *Chem. Commun.* **2019**, *55*, 8102.
 (c) Shi, J.; Wei, W. *Chin. J. Org. Chem.* **2020**, *40*, 2170 (in Chinese).
 (d) 时建伟, 魏伟, 有机化学, **2020**, *40*, 2170.
 (e) Xie, L.-Y.; Bai, Y.-S.; Xu, X.-Q.; Peng, X.; Tang, H.-S.; Huang, Y.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Green Chem.* **2020**, *22*, 1720.
 (f) Xie, K.; Jiang, M.; Chen, X.; Lv, Q.; Yu, B. *Chin. J. Org. Chem.* **2021**, *41*, 4575 (in Chinese).
 (g) 谢坤宸, 江铭轩, 陈晓岚, 吕琪妍, 於兵, 有机化学, **2021**, *41*, 4575.
 (h) Zhu, H.-L.; Zeng, F.-L.; Chen, X.-L.; Sun, K.; Li, H.-C.; Yuan, X.-Y.; Qu, L.-B.; Yu, B. *Org. Lett.* **2021**, *23*, 2976.
- [6] (a) Mamillapalli, N. C.; Sekar, G. *Chem.-Eur. J.* **2015**, *21*, 18584.
 (b) Muthukumar, A.; Sekar, G. *J. Org. Chem.* **2018**, *83*, 8827.
- [7] Agarwal, R.; Liao, Y.; Lin, D.-J.; Yang, Z.-X.; Lai, C.-F.; Chen, C.-T. *Org. Chem. Front.* **2020**, *7*, 2505.
- [8] Zhu, Z.; Lv, X.; Anesini, J. E.; Seidel, D. *Org. Lett.* **2017**, *19*, 6424.
- [9] Lv, Y.; Bao, P.; Yue, H.; Li, J.-S.; Wei, W. *Green Chem.* **2019**, *21*, 6051.
- [10] (a) Jiang, Y.; Xu, K.; Zeng, C. *Chem. Rev.* **2018**, *118*, 4485.
 (b) Tang, H.-T.; Jia, J.-S.; Pan, Y.-M. *Org. Biomol. Chem.* **2020**, *18*, 5315.
 (c) Li, M.; Wang, R.; Hao, W.; Jiang, B. *Chin. J. Org. Chem.* **2020**, *40*, 1540 (in Chinese).
 (d) 李梦帆, 王榕, 郝文娟, 姜波, 有机化学, **2020**, *40*, 1540.
 (e) Cheng, X. *Chin. J. Org. Chem.* **2020**, *40*, 2600 (in Chinese).
 (f) 程旭, 有机化学, **2020**, *40*, 2600.
 (g) Yuan, Y.; Yang, J.; Lei, A. *Chem. Soc. Rev.* **2021**, *50*, 10058.
 (h) Chen, J.-Y.; Zhong, C.-T.; Gui, Q.-W.; Zhou, Y.-M.; Fang, Y.-Y.; Liu, K.-J.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Chin. Chem. Lett.* **2021**, *32*, 475.
 (i) Ye, Z.; Zhang, F. *Chin. J. Org. Chem.* **2021**, *41*, 241 (in Chinese).
 (j) 叶增辉, 张逢质, 有机化学, **2021**, *41*, 241.
 (k) Wu, Y.; Chen, J.-Y.; Ning, J.; Jiang, X.; Deng, J.; Deng, Y.; Xu, R.; He, W.-M. *Green Chem.* **2021**, *23*, 3950.
 (l) Chen, N.; Lai, X.; Xu, H. *Chin. J. Org. Chem.* **2021**, *40*, 2592 (in Chinese).
 (m) 陈娜, 赖小丽, 徐海超, 有机化学, **2021**, *40*, 2592.
 (n) Jiang, J.; Wang, Z.; He, W.-M. *Chin. Chem. Lett.* **2021**, *32*, 1591.
 (o) Tang, H.; Pan, Y. *Chin. J. Org. Chem.* **2021**, *41*, 435 (in Chinese).
 (p) 唐海涛, 潘英明, 有机化学, **2021**, *41*, 435.
 (q) Yi, R.-N.; Wu, Z.-L.; Ouyang, W.-T.; Wang, W.-F.; He, W.-M. *Tetrahedron Lett.* **2021**, *77*, 153257.
 (r) Zhang, M.; Zhao, P.; Yuan, H.; Zhang, A. a.; Zhang, W.; Zheng, L.; Wu, D.; Liu, L. *Chin. J. Org. Chem.* **2021**, *41*, 669 (in Chinese).
 (s) 张明亮, 赵聘, 袁浩, 张安, 张文宇, 郑琳琳, 吴冬青, 刘澜涛, 有机化学, **2021**, *41*, 669.
 (t) Sun, K.; Xiao, F.; Yu, B.; He, W.-M. *Chin. J. Catal.* **2021**, *42*, 1921.
 (u) Yang, Z.; Yu, Y.; Lai, L.; Zhou, L.; Ye, K.; Chen, F.-E. *Green Synth. Catal.* **2021**, *2*, 19.
- [11] (a) Chen, D.; Sun, Y.; Dong, D.; Han, Q.; Wang, Z. *Chin. J. Org. Chem.* **2020**, *40*, 4267 (in Chinese).
 (b) 陈德茂, 孙媛媛, 董道青, 韩晴晴, 王祖利, 有机化学, **2020**, *40*, 4267.
 (c) Gan, Z.; Li, G.; Yan, Q.; Deng, W.; Jiang, Y.-Y.; Yang, D. *Green Chem.* **2020**, *22*, 2956.
 (d) Wang, L.; Zhang, M.; Zhang, Y.; Liu, Q.; Zhao, X.; Li, J.-S.; Luo, Z.; Wei, W. *Chin. Chem. Lett.* **2020**, *31*, 67.
 (e) Li, W.; He, W. *Chin. J. Org. Chem.* **2021**, *41*, 4528 (in Chinese).
 (f) 李文艺, 何卫民, 有机化学, **2021**, *41*, 4528.
 (g) Zeng, F.-L.; Zhu, H.-L.; Chen, X.-L.; Qu, L.-B.; Yu, B. *Green Chem.* **2021**, *23*, 3677.
 (h) Gui, Q.-W.; Teng, F.; Li, Z.-C.; Xiong, Z.-Y.; Jin, X.-F.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Chin. Chem. Lett.* **2021**, *32*, 1907.
 (i) Wang, B.; Zou, L.; Wang, L.; Sun, M.; Li, P. *Chin. Chem. Lett.* **2021**, *32*, 1229.
 (j) Yi, R.; He, W. *Chin. J. Org. Chem.* **2021**, *41*, 1267 (in Chinese).
 (k) 易荣楠, 何卫民, 有机化学, **2021**, *41*, 1267.
 (l) Jin, W.; Liu, C. *Chin. J. Org. Chem.* **2021**, *41*, 2148 (in Chinese).
 (m) 金伟伟, 刘晨江, 有机化学, **2021**, *41*, 2148.
- [12] (a) Quinn, R. K.; Schmidt, V. A.; Alexanian, E. J. *Chem. Sci.* **2013**, *4*, 4030.
 (b) Wang, Q.-Q.; Xu, K.; Jiang, Y.-Y.; Liu, Y.-G.; Sun, B.-G.; Zeng, C.-C. *Org. Lett.* **2017**, *19*, 5517.
- [13] Huang, J.; Liang, B.; Chen, X.; Liu, Y.; Li, Y.; Liang, J.; Zhu, W.; Tang, X.; Li, Y.; Zhu, Z. *Org. Biomol. Chem.* **2021**, *19*, 4783.
- [14] Zhang, C.; Zong, X.; Zhang, L.; Jiao, N. *Org. Lett.* **2012**, *14*, 3280.
- [15] Zhang, Y.-K.; Wang, B. *Eur. J. Org. Chem.* **2019**, *2019*, 5732.

(Lu, Y.)